

Article

# Candidate Causal Variants at the 8p12 Breast Cancer Risk Locus Regulate *DUSP4*

Dylan M. Glubb <sup>1\*</sup>, Wei Shi <sup>1</sup>, Jonathan Beesley <sup>1</sup>, Laura Fachal <sup>2</sup>, Jayne-Louise Pritchard <sup>1</sup>, Karen McCue <sup>1</sup>, Daniel R. Barnes <sup>3</sup>, Antonis C. Antoniou <sup>3</sup>, Alison M. Dunning <sup>2</sup>, Douglas F. Easton <sup>2,3</sup>, Georgia Chenevix-Trench <sup>1</sup>

<sup>1</sup> Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, QLD 4006, Australia

<sup>2</sup> Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge CB1 8RN, UK

<sup>3</sup> Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Cambridge CB1 8RN, UK

\* Correspondence: dylan.glubb@qimrberghofer.edu.au; Tel.: +61-7-3845-3725

**Abstract:** Genome-wide association studies have revealed a locus at 8p12 that is associated with breast cancer risk. Fine-mapping of this locus identified 16 candidate causal variants (CCVs). However, as these variants are intergenic, their function is unclear. To map chromatin looping from this risk locus to a previously identified candidate target gene, *DUSP4*, we performed chromatin conformation capture analyses in normal and tumoral breast cell lines. We identified putative regulatory elements, containing CCVs, that loop to the *DUSP4* promoter region. Using reporter gene assays, we found that the risk allele of CCV rs7461885 reduced the activity of a *DUSP4* enhancer element, consistent with the function of *DUSP4* as a tumor suppressor gene. Furthermore, the risk allele of CCV rs12155535, located in another *DUSP4* enhancer element, was negatively correlated with looping of this element to the *DUSP4* promoter region, suggesting that this allele would be associated with reduced expression. These findings provide the first evidence that CCV risk alleles downregulate *DUSP4* expression, suggesting that this gene is a regulatory target of the 8p12 breast cancer risk locus.

**Keywords:** breast cancer risk; GWAS; candidate causal variant; chromatin conformation capture; reporter gene activity; enhancer; promoter

## 1. Introduction

Genome-wide association studies by the Breast Cancer Association Consortium (BCAC) and Consortium of Investigators of Modifiers of BRCA1/2 (CIMBA) have previously found genetic variants at 8p12 associated with breast cancer risk [1-4]. In the largest study, the rarer allele (A) of SNP rs9693444 was associated with a per-allele odds ratio (OR)=1.06 (95% CI 1.04-1.08;  $p=2 \times 10^{-21}$ ) [1]. Subsequent fine-mapping analysis of BCAC and CIMBA data at this locus revealed 16 candidate causal variants (CCVs; **Supplementary Table S1**), including rs9693444, which could not be statistically separated from each other using a likelihood ratio criterion of 100:1 [5]. The CCVs are located in an intergenic region  $\geq 300$  kb centromerically from *DUSP4* and  $\geq 285$  kb telomerically from the microRNA *MIR3148*.

A major obstacle in the follow-up of GWAS loci is the identification of target genes. Indeed, given the intergenic location of the CCVs at 8p12, it is unclear what function the risk variant(s) may have. To identify breast cancer GWAS target genes, we previously developed a pipeline called


INQUISIT (integrated expression quantitative trait and in silico prediction of GWAS targets) [1,5]. In our latest fine-mapping study, INQUISIT revealed seven target genes at the 8p12 breast cancer risk locus, of which only *DUSP4*, was a high confidence target [5]. This prediction was partly based on evidence of long-range regulatory chromatin interactions between CCVs and the *DUSP4* promoter which included: RNA polymerase II chromatin interaction analysis by paired-end tag sequencing (ChIA-PET) data in a breast cancer cell line [6]; Hi-C chromatin-interaction and bioinformatic analysis identifying an enhancer coincident with CCVs that interacts with the *DUSP4* promoter [7]; and capture Hi-C data from breast cancer cell lines [8].

The chromatin interaction data identified through INQUISIT suggest that CCVs could regulate *DUSP4* expression through enhancer-promoter interactions. This may be a common mechanism through which GWAS variants act (reviewed in [9]). Indeed, at a number of breast cancer GWAS risk loci, we have previously shown that CCVs in regulatory elements loop to promoters and regulate their activity [1,10,11]. In the current study, we map long-range chromatin interactions between the 8p12 risk locus and the promoter region of *DUSP4*. We provide evidence that a CCV, coincident with a putative regulatory element (PRE) that interacts with *DUSP4*, can regulate *DUSP4* promoter activity. Furthermore, we find a CCV in a different PRE that is associated with allele-specific chromatin looping to the *DUSP4* promoter region. These data suggest strongly that *DUSP4* is a target gene of breast cancer risk variation at the 8p12 locus.

## 2. Results

### 2.1. Chromatin conformation capture (3C) analyses confirm that the 8p12 risk locus physically interacts with the promoter region of *DUSP4*

To explore chromatin interactions between the 8p12 risk locus and the *DUSP4* promoter in breast cells, chromatin conformation capture (3C) analyses were conducted in breast cancer (MCF-7 and T47D) and normal mammary epithelial (Bre-80) cell lines. Some level of interaction with the *DUSP4* promoter was observed in all three cell lines across the 3C restriction fragments at the risk locus (**Figure 1**). Notably, this broad level of activity is not often observed in 3C analyses. For example, the 2q35 and 5q11.2 breast cancer risk loci are considerably closer ( $\leq 100$ kb) to corresponding promoter targets than 8p12 CCVs are to the *DUSP4* promoter region, yet many restriction fragments at these loci have no observable promoter interaction [10,11]. At the 8p12 locus, five 3C restriction fragments (#6, #8, #10, #11 and #12) with Hi-C and/or ChIA-PET evidence of chromatin looping to the *DUSP4* promoter demonstrated peaks of interaction in the breast cell lines (**Figure 1**), highlighting three fragments (#6, #11 and #12) which contain CCVs (three, two and three, respectively) for further analysis.




**Figure 1.** Candidate risk variants (CCVs) are located in putative regulatory elements (PREs) that interact with the *DUSP4* promoter. The figure shows the 8p12 breast cancer risk locus annotated with breast cell functional genomic data and 3C analyses of interactions between EcoRI fragments and the *DUSP4* promoter region (located  $\geq 300$  kb upstream) in Bre-80, MCF-7 and T-47D cells. Functional genomic data from normal mammary epithelial cells (HMEC and MCF10A) and breast cancer cell lines (MCF-7 and T-47D) in this figure include Hi-C analysis [7] and ENCODE data accessed from the UCSC Genome Browser: RNA polymerase II ChIA-PET, epigenetic marks characteristic of enhancers (DNaseI hypersensitivity sites and H3K4Me1, H3K4Me2 and H3K27Ac histone modifications), and transcription factor binding (Pol2, STAT3, MYC and FOS). Notably, all ChIA-PET anchors and the predicted Hi-C enhancer loop to the *DUSP4* promoter. For the 3C analyses, interaction frequencies were normalised to those of fragment 8, a common peak of interaction in all three cell lines. Interaction frequencies from three independent biological replicates are shown (error bars represent standard error of the mean). Fragments demonstrating interaction peaks and encompassing PREs with CCVs are bounded by boxes. The PRE regions cloned for reporter gene analysis are highlighted in pink.

## 2.2. CCV rs7461885 reduces the enhancer activity of PRE1 on the *DUSP4* promoter

From analysis of the 3C, ChIA-PET and epigenetic data at the 8p12 locus (**Figure 1**), CCVs in three putative regulatory elements (PRE1-3) were prioritized for reporter gene analyses to determine their effects on *DUSP4* promoter activity (**Figure 2**). PRE1, containing three CCVs, significantly enhanced *DUSP4* promoter activity by 2.3-2.5 fold in Bre-80 and MCF-7 cells, respectively (**Figure 2A** and **B**).


In these two cell lines, the risk-associated allele of rs7461885, whether as a single variant or in a haplotype with the risk alleles of two other CCVs, significantly reduced the PRE1 enhancer activity by 1.3-1.5 fold (**Figure 2A and B**). Analysis of rs7461885 using HaploReg [12] did not indicate any evidence for transcription factor motifs or binding; however, this variant is located in active histone marks (H3K4Me1, H3K4Me2 and H3K27Ac) in normal mammary epithelial cells (**Figure 1**). PRE2, containing two CCVs, significantly enhanced *DUSP4* promoter activity by 2.5-2.7 fold in MCF-7 and Bre-80 cells, respectively, but neither CCV affected this enhancer activity, nor did the haplotype containing both risk alleles (**Figure 2C and D**). PRE3, containing one CCV, significantly enhanced promoter activity by 1.3 fold in Bre-80 cells but had no significant effect in MCF-7 cells (**Figure 2E and F**). The CCV in PRE3 did not affect promoter activity in either cell line (**Figure 2E and F**). Reporter assays were also performed in T-47D cells but neither PREs nor CCVs significantly affected *DUSP4* promoter activity (**Supplementary Figure S1**).



**Figure 2.** PREs enhance *DUSP4* promoter activity in luciferase reporter assays and the risk allele of CCV rs7461885 reduces PRE1 enhancer activity. PRE1, PRE2 and PRE3 reference regions, containing protective allelic variants of CCVs, were cloned downstream of a *DUSP4* promoter luciferase construct for the creation of reference (ref) constructs. Risk allelic variants of CCVs were engineered into the constructs and are designated by the rs ID of the corresponding variant. Constructs containing risk allele haplotypes (hap) were also generated. Cells were transiently transfected with each of these constructs and assayed for luciferase activity after 24 h. Panels show back-transformed data for: (a) PRE1 activity in Bre-80 cells; (b) PRE1 activity in MCF-7 cells; (c) PRE2 activity in Bre-80 cells; (d) PRE2 activity in MCF-7 cells; (e) PRE3 activity in Bre-80 cells; and (f) PRE3 activity in MCF-7 cells. Error bars denote 95% confidence intervals of experiments performed in triplicate. P-values were determined two-way ANOVA followed by Dunnett's multiple comparisons test (\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  and \*\*\*\* $p < 0.0001$ ).

### 2.3 CCV rs1215535 (PRE2) is associated with allele-specific looping to the *DUSP4* promoter region

Of the three breast cell lines used in the 3C assay, only one (Bre-80) was heterozygous for CCVs and thus amenable for allele-specific 3C analysis. After assessing CCVs in the PREs and their proximity to the 3C restriction sites, we found that only PRE2 could be analysed for allele-specific looping using PCR-based Sanger sequencing. Using Bre-80 3C libraries, we found that there was a reduction of the risk (C) allele of CCV rs12155535 in PRE2 interactions (**Figure 3**), compared to the Bre-80 genomic DNA control. This finding indicates that the risk allele was negatively correlated with the PRE2 enhancer-*DUSP4* promoter region interaction, providing evidence that risk allele may be associated with diminished *DUSP4* expression through a reduction in enhancer looping to the *DUSP4* promoter region.



**Figure 3.** Allele-specific interaction between PRE2 and the *DUSP4* promoter region. Bre-80 cells are heterozygous for the rs12155535 CCV in PRE2. Allele-specific analysis using PCR-based Sanger sequencing of two independent 3C libraries indicated that sequence containing the protective (T) allele of rs12155535 preferentially interacted with the *DUSP4* promoter region.

### 3. Discussion

In this study, we used genetic approaches to identify functional CCVs and regulatory targets at the 8p12 breast cancer risk locus. Firstly, we performed 3C in normal and breast cancer cell lines to map chromatin interactions between PREs, containing CCVs, and the *DUSP4* promoter region. Although we cannot rule out interactions with other genes at this locus, these findings were consistent with available chromatin interaction analyses, and no other high-confidence targets were predicted by the INQUISIT pipeline in this region [5].

Secondly, we prioritised three PREs with evidence of interaction with the *DUSP4* promoter region (i.e. PRE1-3) for reporter gene analyses. These assays demonstrated that PRE1 and PRE2 acted as enhancers of *DUSP4* promoter activity in both normal and tumoral breast cell lines. PRE1 and PRE2 are located in areas of active chromatin as indicated by the presence of transcription factor binding and strong histone peaks characteristic of enhancers (**Figure 1**). CCVs in PRE1-3 were tested for effects on the *DUSP4* promoter and the risk allele of CCV rs7461885 was found to repress the *DUSP4* enhancer activity of PRE1. It is possible that other CCVs at the 8p12 locus, which were not assessed in the reporter gene assays, may also regulate *DUSP4*; however, the available epigenetic data suggested these variants were less likely to be functional.

Finally, we performed an allele-specific 3C analysis and found that the risk allele of CCV rs12155535 was present less often in PRE2-*DUSP4* promoter region interactions in comparison with the protective allele. As the reporter gene assays had demonstrated that PRE2 acted as a *DUSP4*

enhancer, the association of the rs12155535 risk allele with reduced PRE2-*DUSP4* interaction frequency was consistent with the negative effect of the risk allele of rs7461885 on *DUSP4* promoter activity.

The 3C and reporter gene findings indicate that *DUSP4* is a likely target gene and is downregulated by risk associated alleles at the 8p12 locus. However, expression quantitative trait loci data from mammary tissue (n=396) in the Genotype Tissue Expression database (version 8; <https://gtexportal.org/home/> accessed on 09/10/2019) and tumour tissue (n=799) in The Cancer Genome Atlas eQTL browser ([https://albertlab.shinyapps.io/tcga\\_eqtl/](https://albertlab.shinyapps.io/tcga_eqtl/) accessed on 9/10/2019) provided no evidence that CCVs associated with *DUSP4* expression in normal or tumour breast tissue. A lack of CCV eQTLs at cancer risk GWAS loci is a common finding [10,11,13,14] and could be due to several issues: i) the available eQTL studies may not have enough statistical power to detect modest effects on *DUSP4* expression; ii) the tissues used, especially tumour tissue, are heterogeneous, containing multiple cell types, and thus may mask CCV effects on *DUSP4* expression that occur in specific cell types; and iii) CCVs could have development or context specific effects on *DUSP4* expression that are not captured by the available studies.

*DUSP4* encodes a nuclear dual specificity protein phosphatase which inactivates ERK, JNK and p38 mitogen-activated protein kinases [15] and may also play a regulatory role in gene expression through chromatin binding [16]. *DUSP4* is frequently deleted in breast tumours [17,18] and cancer cell lines [19]. *DUSP4* knockdown enhances the formation of breast cancer stem cells [16,20] and increases the invasive ability of ER-positive breast cancer stem cells [20]. In ER-negative breast cancer cells, *DUSP4* knockdown increases the formation of mammospheres and the expression of cancer stem cell promoting cytokines [19]. Thus, the literature indicates that *DUSP4* acts as a tumor suppressor in breast cancer which is compatible with our finding of its negative regulation by a breast cancer risk allele.

## 4. Materials and Methods

### 4.1. Cell culture

Cell lines were stored in liquid nitrogen vapour phase with *Mycoplasma* testing and short tandem repeat profiling performed for cell authentication prior to storage. The immortalised mammary epithelial Bre-80 cell line was cultured as previously described [21]. The breast cancer cell lines MCF-7 and T-47D were cultured in RPMI 1640 medium supplemented with fetal calf serum, penicillin/streptomycin and 10 µg/ml insulin. All cell lines were cultured at 37°C in a humidified 5% CO<sub>2</sub> atmosphere.

### 4.2. 3C analysis

3C libraries were generated from breast cell lines using EcoRI as previously described [22]. 3C interactions were quantified by qPCR with primers designed with the EcoRI restriction fragments spanning the 8p12 risk locus (**Supplementary Table S2**). qPCR was performed using a RotorGene 6000 with a reaction mix containing MyTaq HS DNA polymerase and the addition of 5 mM Syto9. Thermal cycling was performed with an annealing temperature of 66 °C (20 s) and extension at 72 °C (30 s). Three independent 3C libraries were analysed by qPCR with each experiment quantified in duplicate. Two BAC clones (RP11-56L5 and RP11-833O14: BACPAC Resource Center) encompassing the *DUSP4* promoter region and the 8p12 breast cancer risk locus were used to create a library of ligation products to determine 3C primer efficiencies for normalization.

### 4.3. Reporter gene vector construction

A *DUSP4* promoter luciferase reporter construct was generated by inserting a 1650 bp of PCR-amplified DNA, containing the *DUSP4* transcription start (chr8:29,207,878-29,209,527; GRCh37), site into the KpnI and HindIII sites of a pGL3-Basic construct. The pGL3-Basic construct had been engineered to include AgeI and SbfI sites downstream of the luciferase gene [6]. A 2464 bp PRE1 region (chr88:29,527,323-29,530,033) was PCR generated using primers engineered with

BamHI and Sall sites for insertion downstream of the Firefly luciferase gene. A 980 bp PRE2 region (chr8:29522728-29523707) and a 981bp PRE3 region (chr8:29508759-29509739) were synthesised with terminal AgeI/Sall and BamHI/SbfI sites, respectively (Integrated DNA Technologies, Singapore) for cloning into the *DUSP4* promoter construct. For PRE1, the allelic variants of CCVs were introduced by overlap extension PCR. For PRE2 and 3, constructs containing allelic variants were also synthesised by Integrated DNA Technologies and cloned as above. All constructs were Sanger sequenced (QIMR Berghofer sequencing facility) to confirm variant incorporation. All PCR and sequencing primers are listed in **Supplementary Table S3**.

#### 4.4. Reporter gene vector construction

Bre-80, MCF-7 and T-47D cells were transfected with equimolar amounts of luciferase reporter constructs and 50 ng of the Renilla luciferase pRL-SV40 construct with Lipofectamine 2000. The total amount of transfected DNA was kept constant at 600 ng for each construct by adding pUC19 as a carrier plasmid. Luciferase activity was measured 24 h post-transfection by the Dual-Glo Luciferase Assay System. To correct for variation in transfection efficiency or cell lysate preparation, Firefly luciferase activity was normalised to that of the Renilla luciferase. All *DUSP4* promoter constructs had great activity than the negative control (empty pGL3-Basic construct). Data were log-transformed and statistical significance was assessed by two-way ANOVA followed by Dunnett's multiple comparisons test in GraphPad Prism (version 7.02).

#### 4.5. Allele-specific 3C analysis

DNA was amplified by PCR of two independent Bre-80 3C libraries using the *DUSP4* promoter bait primer and a reverse primer specific to the PRE2 fragment containing rs12155535 (**Supplementary Table S2**). The region containing rs12155535 was also amplified from Bre-80 genomic DNA as a control. Sanger sequencing (QIMR Berghofer sequencing facility) was performed to determine the alleles present in the interacting PRE2 fragment.

## 5. Conclusions

We have found the first evidence that breast cancer risk variation at the 8p12 locus downregulates *DUSP4* promoter activity and is negatively correlated with long-range chromatin looping interactions with the *DUSP4* promoter region. These findings are consistent with the role of *DUSP4* as a tumor suppressor in breast cancer and suggest that the effects of CCVs on breast cancer risk may be mediated by a reduction in *DUSP4* expression.

**Supplementary Materials:** The following are available online at [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1), Figure S1: Luciferase reporter gene analysis of PRE1-3 in T-47D cells, Table S1: 8p12 CCVs, Table S2: 3C primers, Table S3: Primers for generation of reporter gene constructs.

**Author Contributions:** Conceptualization, D.G. and G.C-T.; formal analysis, W.S., J.B. and D.G.; investigation, W.S., J-L.P., K.M., J.B. and D.G.; resources, D.E. and G.C-T.; data curation, D.G., L.F. and J.B.; writing—original draft preparation, D.G. and G.C-T.; writing—review and editing, D.G., W.S., J.B., L.F., J-L.P., K.M., D.B., A.A., A.D., D.E., and G.C-T.; visualization, D.G.; supervision, D.G., K.M., A.A., A.D., D.E., G.C-T.; project administration, G.C-T.; funding acquisition, G.C-T.

**Funding:** This research was funded by the National Health and Medical Research Council of Australis.

**Acknowledgments:** The authors acknowledge the BCAC and CIMBA consortia in providing pre-publication access to the fine-mapping data.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. Michailidou, K.; Lindstrom, S.; Dennis, J.; Beesley, J.; Hui, S.; Kar, S.; Lemacon, A.; Soucy, P.; Glubb, D.; Rostamianfar, A., et al. Association analysis identifies 65 new breast cancer risk loci. *Nature* **2017**, *551*, 92-94, doi:10.1038/nature24284.
2. Michailidou, K.; Beesley, J.; Lindstrom, S.; Canisius, S.; Dennis, J.; Lush, M.J.; Maranian, M.J.; Bolla, M.K.; Wang, Q.; Shah, M., et al. Genome-wide association analysis of more than 120,000 individuals identifies 15 new susceptibility loci for breast cancer. *Nat Genet* **2015**, *47*, 373-380, doi:10.1038/ng.3242.
3. Michailidou, K.; Hall, P.; Gonzalez-Neira, A.; Ghoussaini, M.; Dennis, J.; Milne, R.L.; Schmidt, M.K.; Chang-Claude, J.; Bojesen, S.E.; Bolla, M.K., et al. Large-scale genotyping identifies 41 new loci associated with breast cancer risk. *Nat Genet* **2013**, *45*, 353-361, 361e351-352, doi:10.1038/ng.2563
4. Mulligan, A.M.; Couch, F.J.; Barrowdale, D.; Domchek, S.M.; Eccles, D.; Nevanlinna, H.; Ramus, S.J.; Robson, M.; Sherman, M.; Spurdle, A.B., et al. Common breast cancer susceptibility alleles are associated with tumour subtypes in BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2. *Breast Cancer Res* **2011**, *13*, R110, doi:10.1186/bcr3052.
5. Fachal, L.; Aschard, H.; Beesley, J.; Barnes, D.R.; Allen, J.; Kar, S.; Pooley, K.A.; Dennis, J.; Michailidou, K.; Turman, C., et al. Fine-mapping of 150 breast cancer risk regions identifies 178 high confidence target genes. *bioRxiv* **2019**, 10.1101/521054, 521054, doi:10.1101/521054.
6. Li, G.; Ruan, X.; Auerbach, R.K.; Sandhu, K.S.; Zheng, M.; Wang, P.; Poh, H.M.; Goh, Y.; Lim, J.; Zhang, J., et al. Extensive promoter-centered chromatin interactions provide a topological basis for transcription regulation. *Cell* **2012**, *148*, 84-98, doi:10.1016/j.cell.2011.12.014.
7. Rao, S.S.; Huntley, M.H.; Durand, N.C.; Stamenova, E.K.; Bochkov, I.D.; Robinson, J.T.; Sanborn, A.L.; Machol, I.; Omer, A.D.; Lander, E.S., et al. A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **2014**, *159*, 1665-1680, doi:10.1016/j.cell.2014.11.021.
8. Beesley, J.; Sivakumaran, H.; Marjaneh, M.M.; Lima, L.G.; Hillman, K.M.; Kaufmann, S.; Tuano, N.; Hussein, N.; Ham, S.; Mukhopadhyay, P., et al. Chromatin interactome mapping at 139 independent breast cancer risk signals. *bioRxiv* **2019**, 10.1101/520916, 520916, doi:10.1101/520916.
9. Gallagher, M.D.; Chen-Plotkin, A.S. The Post-GWAS Era: From Association to Function. *Am J Hum Genet* **2018**, *102*, 717-730, doi:10.1016/j.ajhg.2018.04.002.
10. Glubb, D.M.; Maranian, M.J.; Michailidou, K.; Pooley, K.A.; Meyer, K.B.; Kar, S.; Carlebur, S.; O'Reilly, M.; Betts, J.A.; Hillman, K.M., et al. Fine-Scale Mapping of the 5q11.2 Breast Cancer Locus Reveals at Least Three Independent Risk Variants Regulating MAP3K1. *Am J Hum Genet* **2015**, *96*, 5-20.
11. Ghoussaini, M.; Edwards, S.L.; Michailidou, K.; Nord, S.; Cowper-Sal Lari, R.; Desai, K.; Kar, S.; Hillman, K.M.; Kaufmann, S.; Glubb, D.M., et al. Evidence that breast cancer risk at the 2q35 locus is mediated through IGFBP5 regulation. *Nat Commun* **2014**, *4*, 4999, doi:10.1038/ncomms5999.
12. Ward, L.D.; Kellis, M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* **2012**, *40*, D930-934, doi:10.1093/nar/gkr917.
13. O'Mara, T.A.; Glubb, D.M.; Kho, P.F.; Thompson, D.J.; Spurdle, A. Genome-wide association studies of endometrial cancer: Latest developments and future directions. *Cancer Epidemiol Biomarkers Prev* **2019**, *28*, 1095-1102.
14. Lawrenson, K.; Li, Q.; Kar, S.; Seo, J.H.; Tyrer, J.; Spindler, T.J.; Lee, J.; Chen, Y.; Karst, A.; Drapkin, R., et al. Cis-eQTL analysis and functional validation of candidate susceptibility genes for high-grade serous ovarian cancer. *Nat Commun* **2015**, *6*, 8234, doi:10.1038/ncomms9234.

15. Keyse, S.M. Dual-specificity MAP kinase phosphatases (MKPs) and cancer. *Cancer metastasis reviews* **2008**, *27*, 253-261, doi:10.1007/s10555-008-9123-1.
16. Boulding, T.; Wu, F.; McCuaig, R.; Dunn, J.; Sutton, C.R.; Hardy, K.; Tu, W.; Bullman, A.; Yip, D.; Dahlstrom, J.E., et al. Differential Roles for DUSP Family Members in Epithelial-to-Mesenchymal Transition and Cancer Stem Cell Regulation in Breast Cancer. *PLoS One* **2016**, *11*, e0148065, doi:10.1371/journal.pone.0148065.
17. Armes, J.E.; Hammet, F.; de Silva, M.; Ciciulla, J.; Ramus, S.J.; Soo, W.K.; Mahoney, A.; Yarovaya, N.; Henderson, M.A.; Gish, K., et al. Candidate tumor-suppressor genes on chromosome arm 8p in early-onset and high-grade breast cancers. *Oncogene* **2004**, *23*, 5697-5702, doi:10.1038/sj.onc.1207740.
18. Mazumdar, A.; Poage, G.M.; Shepherd, J.; Tsimelzon, A.; Hartman, Z.C.; Den Hollander, P.; Hill, J.; Zhang, Y.; Chang, J.; Hilsenbeck, S.G., et al. Analysis of phosphatases in ER-negative breast cancers identifies DUSP4 as a critical regulator of growth and invasion. *Breast Cancer Res Treat* **2016**, *158*, 441-454, doi:10.1007/s10549-016-3892-y.
19. Balko, J.M.; Schwarz, L.J.; Bhola, N.E.; Kurupi, R.; Owens, P.; Miller, T.W.; Gomez, H.; Cook, R.S.; Arteaga, C.L. Activation of MAPK pathways due to DUSP4 loss promotes cancer stem cell-like phenotypes in basal-like breast cancer. *Cancer Res* **2013**, *73*, 6346-6358, doi:10.1158/0008-5472.can-13-1385.
20. Gaggianesi, M.; Turdo, A.; Chinnici, A.; Lipari, E.; Apuzzo, T.; Benfante, A.; Sperduti, I.; Di Franco, S.; Meraviglia, S.; Lo Presti, E., et al. IL4 Primes the Dynamics of Breast Cancer Progression via DUSP4 Inhibition. *Cancer Res* **2017**, *77*, 3268-3279, doi:10.1158/0008-5472.can-16-3126.
21. Meyer, K.B.; O'Reilly, M.; Michailidou, K.; Carlebur, S.; Edwards, S.L.; French, J.D.; Prathalingham, R.; Dennis, J.; Bolla, M.K.; Wang, Q., et al. Fine-Scale Mapping of the FGFR2 Breast Cancer Risk Locus: Putative Functional Variants Differentially Bind FOXA1 and E2F1. *Am J Hum Genet* **2013**, *93*, 1046-1060, doi:10.1016/j.ajhg.2013.10.026.
22. Tan-Wong, S.M.; French, J.D.; Proudfoot, N.J.; Brown, M.A. Dynamic interactions between the promoter and terminator regions of the mammalian BRCA1 gene. *Proc Natl Acad Sci U S A* **2008**, *105*, 5160-5165, doi:10.1073/pnas.0801048105.