

Hearing loss and brain plasticity: The hyperactivity phenomenon

Björn Herrmann^{1,2,*} & Blake E. Butler^{3,4}

¹Rotman Research Institute
Baycrest, Toronto, ON, Canada

²Department of Psychology
University of Toronto, Toronto, ON, Canada

³Department of Psychology & The Brain and Mind Institute
University of Western Ontario, London, ON, Canada

⁴National Centre for Audiology
University of Western Ontario, London, ON, Canada

*Correspondence concerning this article should be addressed to Björn Herrmann, Rotman Research Institute, Baycrest, Toronto, ON, M6A 2E1, Canada.
E-mail: bherrmann@research.baycrest.org

Abstract

Many aging adults experience some form of hearing problems that may arise from auditory peripheral damage. However, it has been increasingly acknowledged that hearing loss is not only a dysfunction of the auditory periphery but results from changes within the entire auditory system, from periphery to cortex. Damage to the auditory periphery is associated with an increase in neural activity at various stages throughout the auditory pathway. Here, we review neurophysiological evidence of hyperactivity, auditory perceptual difficulties that may result from hyperactivity, and outline open conceptual and methodological questions related to the study of hyperactivity. We suggest that hyperactivity alters all aspects of hearing – including spectral, temporal, spatial hearing – and, in turn, impairs speech comprehension when background sound is present. By focusing on the perceptual consequences of hyperactivity and the potential challenges of investigating hyperactivity in humans, we hope to bring animal and human electrophysiologists closer together to better understand hearing problems in older adulthood.

Keywords: Hearing loss, aging, hyperactivity, excitability, loss of inhibition, neurophysiology, auditory perception, neural plasticity, speech processing

Introduction

Many aging adults experience some form of hearing problems (Cruickshanks et al., 1998; Feder et al., 2015; Goman and Lin, 2016; Helfer et al., 2017). The loss of sensitivity, particularly at high frequencies, comprises the traditional profile of ‘age-related hearing loss’ and is commonly associated with impairments of the auditory periphery, including damage of hair cells, spiral ganglion cells, and the stria vascularis (Gratton and Vázquez, 2003; Moore, 2007; Bao and Ohlemiller, 2010; Schmiedt, 2010; Dubno et al., 2013; Plack, 2014; Keithley, 2020). Other impairments include hearing sound in absence of an identifiable source (tinnitus; Anari et al., 1999; Eggermont and Roberts, 2004; McCormack et al., 2016), finding sounds at moderate intensities too loud and distracting (hyperacusis; Anari et al., 1999; Baguley, 2003; Parmentier and Andrés, 2010; Tyler et al., 2014), or experiencing difficulty comprehending speech when background sound is present (Pichora-Fuller, 2003; Pichora-Fuller et al., 2016; Presacco et al., 2019).

Although hearing loss in older adulthood typically arises from dysfunction of the auditory periphery (Gratton and Vázquez, 2003; Moore, 2007; Bao and Ohlemiller, 2010; Schmiedt, 2010; Dubno et al., 2013; Plack, 2014; Moser and Starr, 2016; Keithley, 2020), an increasing amount of evidence suggests that the behavioral consequences of hearing loss may reflect dysfunction of the entire auditory system. Peripheral damage induces neuroplastic changes in downstream brain regions of the auditory pathway, including the cochlear nucleus, superior olivary complex, inferior colliculus, medial geniculate nucleus (thalamus), and auditory cortex (Knipper et al., 2013; Auerbach et al., 2014; Zhao et al., 2016; Salvi et al., 2017). Perhaps the most prominent change that has been extensively studied in animal models (Knipper et al., 2013; Auerbach et al., 2014; Zhao et al., 2016; Salvi et al., 2017) and that is increasingly observed in older human adults with clinical or subclinical hearing loss (Lister et al., 2011; Alain et al., 2012; Bidelman et al., 2014; Herrmann et al., 2016; Presacco et al., 2016a; Herrmann et al., 2018; Presacco et al., 2019) is neuronal hyperactivity. Here, we use the term hyperactivity to include three related phenomena: 1) an increase in the rate at which neurons in the auditory pathway generate action potentials in the absence of stimulation (i.e., enhanced spontaneous firing rate); 2) an increase in synchronized activity between neurons; and 3) an increase in the number of action potentials and the magnitude of local field potentials generated by neurons in response to sound (hyperresponsivity; Figure 1).

Previous reviews have detailed the neurophysiological evidence for, and proposed mechanisms underlying hyperactivity in the auditory systems of animal models of hearing loss (Knipper et al., 2013; Auerbach et al., 2014; Eggermont, 2015; Zhao et al., 2016; Salvi et al., 2017). However, the full extent of the perceptual consequences of hyperactivity have not been described in as much depth. Moreover, studies conducted in humans – where the study of auditory system hyperactivity is typically limited to non-invasive measures of hyperresponsivity to sound – have not been commonly integrated within previous reviews. This lack of consolidation across

models is a critical oversight that limits our understanding of the causes of hearing impairments in older adults, and slows the development of appropriate treatments.

The current review delivers a detailed picture of the potential perceptual consequences of hyperactivity in the auditory system and provides a foundation from which researchers can develop targeted approaches for investigating hyperactivity in humans. The paper is divided into three sections that aim to convey the following messages: firstly, reduced acoustic input leads to hyperactivity in the central auditory system, regardless of the etiology or nature of the hearing loss (conductive/sensorineural); secondly, auditory system hyperactivity likely degrades spectral, temporal, and spatial processing and, in turn, speech perception; finally, the non-linear relationship between peripheral damage and hyperresponsivity in the auditory system presents unique challenges that should be considered in the design of human studies that target hyperactivity.

Hyperactivity in central auditory regions

In this section, we will review empirical observations of hyperactivity associated with different experimental manipulations (e.g., sound exposure and drug treatment) and age groups. We will focus on direct measures of hyperactivity in the central auditory system, such as increased spontaneous firing rates and/or correlated activity among neurons, measured with invasive neurophysiological approaches. We will further focus on indirect measures of hyperactivity, such as increased neural responses to supra-threshold sounds and enhanced behavioral sensitivity to electrical stimulation, measured using invasive or non-invasive methods. We will describe the animal literature and discuss related work from humans.

Hyperactivity associated with acoustic injury

Inducing acoustic injury to the auditory system of animals via sound exposure has a long tradition in investigations of hearing function and dysfunction (Saunders et al., 1985; Slepecky, 1986). Sound exposures of varied duration and intensity give rise to varying degrees of auditory peripheral pathology. High-intensity exposure (100 dB SPL or above) to pure tones or noise bursts causes mechanical and metabolic damage of the cochlea, including hair cell loss, damage to the stereocilia, tears in the basilar membrane, rupture of the organ of Corti, metabolic exhaustion, and more (Saunders et al., 1985; Slepecky, 1986). Less intense sound exposure may not cause as much mechanical damage, but may leave scars from degenerating hair cells, alter the morphology of cells, change micromechanical properties of hair cells, and cause metabolic damage (Saunders et al., 1985; Slepecky, 1986). Even sound exposure that does not appear to cause hair cell loss may still eliminate synapses that connect inner hair cells with auditory nerve fibers, potentially

impacting the efficiency with which sound waves are translated into electrical signals (Kujawa and Liberman, 2009; Liberman and Kujawa, 2017).




Figure 1: Simulated electrophysiological data of cortical activity. **A:** Simulated data show spontaneous activity (i.e., in absence of experimental acoustic stimulation) and stimulus-evoked activity for simulated neurons under normal and hyperactive conditions. Top rows show raster plots, where each dot reflects the occurrence of an action potential (spike). Spike-rate time courses are depicted on the bottom. **B:** Simulated spike times for several neurons under normal and hyperactive conditions. Synchronization among hyperactive neurons is enhanced (Note that schematic, simulated data are displayed).

High-intensity sound exposure not only leads to damage of auditory peripheral structures, but is also associated with plastic changes along the auditory neural pathway. Compared to a pre-exposure baseline or to control animals, sound-exposed animals show altered spontaneous firing rates in the cochlear nucleus, inferior colliculus, medial geniculate nucleus (thalamus), and auditory cortex (Zhang and Kaltenbach, 1998; Seki and Eggermont, 2003; Manzoor et al., 2012; Mulders and Robertson, 2013; Wang et al., 2013; Coomber et al., 2014; Kalappa et al., 2014; Eggermont, 2015; Figure 1A). Changes appear to be frequency specific, such that spontaneous firing rates of neurons that typically respond to the sound frequencies comprising the exposure stimulus are reduced, while neurons whose frequency sensitivities lie outside of those comprising the exposure stimulus become hyperactive. Increased spontaneous activity is often accompanied by enhanced synchronization among neurons (Seki and Eggermont, 2003; Norena and Eggermont, 2006; Eggermont, 2012; Eggermont and Tass, 2015; Figure 1B).

High-intensity exposure also changes sound-evoked responses. While acoustic injury can lead to smaller sound-evoked responses in neurons of the inferior colliculus of exposed animals compared to non-exposed controls (Popelár et al., 1987; Sun et al., 2012), hyperresponsivity is observed in neurons of the medial geniculate nucleus (thalamus) and auditory cortex (Popelár et al., 1987; Syka et al., 1994; Yang et al., 2011; Sun et al., 2012; Kalappa et al., 2014; Schormans et al., 2019; Wiczerzak et al., 2021; for a detailed review see Auerbach et al., 2014). Hyperactivity

has also been observed following low-intensity sound exposure, where mechanical damage to the cochlea is minimal. Weeks-long continuous sound exposure at intensities that are common in the everyday lives of humans (e.g., ~70 dB SPL) is associated with enhanced spontaneous firing rates in auditory cortex (Munguia et al., 2013) and temporary hyperresponsivity to sound in the inferior colliculus (Sheppard et al., 2018).

In sum, a large body of work has demonstrated that damage to the auditory periphery induced by sound exposure is associated with hyperactivity in the central auditory system and that the degree of damage and subsequent physiological changes depend on the nature of the exposure and the region of the auditory pathway under study.

Hyperactivity associated with pharmacological treatment

In contrast to the broad pattern of inner-ear pathology induced by many high-intensity sound-exposure protocols (Saunders et al., 1985; Slepecky, 1986), pharmacological manipulations that induce ototoxicity enable a fine-grained investigation of the changes in the central auditory system arising from damage to specific peripheral structures, while leaving others intact (but, see recent examples of sound-exposure protocols that specifically damage nerve synapses; Resnik and Polley, 2017; Asokan et al., 2018; Shaheen and Liberman, 2018).

Cisplatin and carboplatin are chemotherapeutic agents that have been shown to selectively impair outer and inner hair cells in rodent models, respectively (Qiu et al., 2000; Kaltenbach et al., 2002; Zhao et al., 2016). In both cases, ototoxicity is followed by an increase in auditory system hyperactivity. For example, cisplatin-induced damage of more than 50% of outer hair cells increases spontaneous firing in the dorsal cochlear nucleus of rodents (Kaltenbach et al., 2002; Rachel et al., 2002). Carboplatin-induced damage to 30-40% of inner hair cells is associated with hyperresponsivity to sound in auditory cortex (Qiu et al., 2000; Salvi et al., 2017). These enhanced responses vanish when 75% of inner hair cells are damaged (Qiu et al., 2000), suggesting that the relationship between the degree of cochlear damage and hyperresponsivity may be non-linear. We will discuss this point in more detail below.

Damage can be specifically restricted to cochlear nerve afferent synapses using ouabain without damaging hair cells in the cochlea (Schmiedt, 2010; Yuan et al., 2014; Chambers et al., 2016b; Chambers et al., 2016a; Resnik and Polley, 2017). Synapse degeneration is associated with a reduction of inputs to central brain regions and results in reduced sound-evoked responses as far along the pathway as the inferior colliculus. However, even with over 90% of synapses degenerated, auditory cortical responses to sound do not differ between ouabain-treated and control animals (Chambers et al., 2016b; consistent with carboplatin-induced damage Qiu et al., 2000), indicating that auditory cortex becomes hyperresponsive following afferent synapse loss.

Salicylates are a group of common anti-inflammatory drugs (e.g., aspirin) that, when administered in high doses, are associated with hearing loss and behavioral indices of hyperactivity, most prominently tinnitus and hyperacusis (Day et al., 1989; Yang et al., 2007; Radziwon et al., 2016; Radziwon et al., 2017; Salvi et al., 2020). A high dose of a salicylate is consistently associated with increased spontaneous firing rates and increased sound-evoked responses in auditory cortex compared to control animals (Yang et al., 2007; Sun et al., 2009; Stolzberg et al., 2012; Salvi et al., 2017). Stronger increases in neural activity are associated with systemic injections of salicylates than with local injections to the cochlea, suggesting that salicylate-induced hyperactivity may not result from changes in cochlear function, but from direct effects on the central auditory structures (Sun et al., 2009).

This small sample of studies demonstrates that damage resulting from ototoxic drugs leads to hyperactivity in the central auditory system. Over 100 drugs have known ototoxic effects that impair hearing in humans, including antibiotics, chemotherapeutics, antimalarials, antiepileptics, anti-inflammatories, and diuretics (Radziwon et al., 2016). However research on ototoxic effects in humans has thus far focused mainly on peripheral and midbrain auditory structures (see review Radziwon et al., 2016); investigations of auditory cortex – and potential hyperactivity – following ototoxic drugs have not yet been conducted.

Hyperactivity in the aged auditory system

Age-related hearing loss is not a uniquely human phenomenon; animals that grow up in relative quiet and are not exposed to noise nor treated with ototoxic drugs, nevertheless, develop inner-ear dysfunction as they age. Cochlear dysfunction in aged mammals results from a degeneration of hair cells, spiral ganglion cells, and the stria vascularis (Gratton and Vázquez, 2003; Moore, 2007; Bao and Ohlemiller, 2010; Schmiedt, 2010; Dubno et al., 2013; Plack, 2014; Keithley, 2020), although it appears that atrophy of the stria is the predominant factor related to aging (Schmiedt, 2010; but see also Wu et al., 2020). As with noise- and drug-induced damage, age-related peripheral impairments are accompanied by hyperactivity in the central auditory system. A given input to the auditory system activates a larger number of neurons along the auditory pathway (Walton et al., 2002; Herrmann et al., 2017; Parthasarathy et al., 2019), and spontaneous activity is increased in the inferior colliculus (Willott et al., 1988b; Parthasarathy et al., 2019) and auditory cortex of older compared to younger animals (Hughes et al., 2010; Juarez-Salinas et al., 2010; Overton and Recanzone, 2016; Ng and Recanzone, 2018).

In contrast to aged animal models, the degree of cochlear damage and subsequent hearing loss observed in older human adults reflects a combination of age-related degenerative changes, long-term sound exposure from daily activities (e.g., in train stations, restaurants), occasional episodes of high-intensity sound exposure (e.g., concerts, industrial noise), cardiovascular

problems (e.g., high blood pressure), and side effects from drugs with ototoxic properties (for reviews see Gates and Mills, 2005; Schmiedt, 2010; Zhao et al., 2016; Ibrahim and Llano, 2019). Hence, humans are exposed to various potential causes of auditory peripheral damage across their lifespans that may give rise to changes in neural activity. Accordingly, hyperresponsivity to sound has been observed in the auditory cortex of older compared to younger humans (e.g., Anderer et al., 1996; Amenedo and Díaz, 1999; Harkrider et al., 2005; Ross and Tremblay, 2009; Sörös et al., 2009; Lister et al., 2011; Alain et al., 2012; Herrmann et al., 2013a; Bidelman et al., 2014; Herrmann et al., 2018; Herrmann et al., 2019), although most of these studies did not aim to investigate age-related hyperresponsivity specifically. Moreover, older adults with clinical hearing loss (i.e., audiometric pure-tone threshold average of 25 dB HL or greater; Plack, 2014) show even greater increases in cortical responses to sound than those with normal hearing (Tremblay et al., 2003; Alain et al., 2014; Millman et al., 2017). These observations are consistent with those from noise-exposed, drug-treated, and aged animals, and suggest that auditory cortical hyperactivity in older humans arises subsequent to peripheral damage.

Hyperactivity associated with acoustic deprivation

The research reviewed thus far suggests that damage to auditory peripheral structures, whether due to sound exposure, drug treatment, or changes related to aging, is associated with hyperactivity in downstream brain regions. However, some researchers suggest that it may not be damage per se, but rather the resulting reduction of input to the central auditory system that leads to hyperactivity (Sanes and Kotak, 2011; Radziwon et al., 2016; Zhao et al., 2016; Parry et al., 2019). Investigating cortical responses to sound following conductive hearing loss may help disentangle these factors (Teichert et al., 2017; Parry et al., 2019). Conductive hearing loss arises from dysfunction of the middle-ear apparatus that normally transduces airborne sound pressure waves into mechanical vibrations before they reach the cochlea. Conductive hearing loss thus provides a model of reduced input to the auditory system in the absence of cochlear damage.

In adult mice, sound-evoked activity in auditory cortex is reduced immediately following the induction of conductive hearing loss through removal of the malleus of the middle ear. However sound-evoked responses appear to recover to normal levels within a few days despite the chronic reduction in peripheral input (Teichert et al., 2017). Conductive hearing loss induced during early development of the auditory system reduces inhibitory activity in auditory cortex, thereby unmasking excitation (Sanes et al., 2010; Sanes and Kotak, 2011; Takesian et al., 2012; but see also Rosen et al., 2012). In humans, electroencephalographic recordings from auditory cortex revealed that supra-threshold bone-conducted stimuli – the transmission of which is unaffected by middle-ear dysfunction – elicit larger responses in people with conductive hearing loss compared to people without (Parry et al., 2019). These data suggest that hyperactivity in

auditory cortex is the result of acoustic deprivation that occurs subsequent to peripheral dysfunction rather than arising from cochlear damage itself (cf. Bauer et al., 2008).

Mechanisms underlying hyperactivity

Neurons can be broadly categorized as being either excitatory or inhibitory, and maintaining a balance between excitation and inhibition is crucial for neural function (Wehr and Zador, 2003; Silver, 2010; Froemke and Martins, 2011; Isaacson and Scanziani, 2011; Shew and Plenz, 2013; Whitmire and Stanley, 2016). There are about four times more excitatory neurons than there are inhibitory neurons in auditory cortex (Ouellet and de Villers-Sidani, 2014), but inhibitory neurons are critical for regulating levels of excitation in order to avoid run-away excitation in neural circuits – a pattern of unconstrained excitation that is detrimental to signalling dynamics and which can lead to neuronal death via excitotoxicity (Shew and Plenz, 2013; Hattori et al., 2017; Imam and Hannan, 2017).

Different types of inhibitory neurons can be distinguished in auditory cortex. The most common express either somatostatin (SST), parvalbumin (PV), or vasoactive intestinal polypeptide (VIP). SST-expressing neurons tend to form synapses on dendrites of excitatory neurons, thereby suppressing neuronal inputs. PV-expressing neurons preferentially connect at the cell body of excitatory neurons and inhibit their output (Figure 2A; Ouda et al., 2015; Hattori et al., 2017). VIP-expressing neurons typically synapse with PV and SST neurons and, by inhibiting or disinhibiting them, regulate cortical excitation (Blackwell and Geffen, 2017; Wood et al., 2017). While we will review research on different types of inhibitory neurons (PV and SST specifically, as VIP neurons have been the focus of far less research) and their selective manipulation, we note that interneurons are integrated in complex interactive networks that make inferences about individual cell types difficult (Pfeffer et al., 2013).




Figure 2: Schematic representation of a loss of inhibition in microcircuits of auditory cortex. A: Normal microcircuit with intact somatostatin (SST) and parvalbumin (PV) inhibitory neurons, regulating the relation between synaptic input and spiking output of an excitatory neuron (EXC). **B:** Microcircuit marked by a loss of SST and PV inhibitory neurons, mirroring observations for the aged auditory cortex (Ouellet and de Villers-Sidani, 2014). A loss of inhibition is associated with increased spiking output.

Hyperactivity in the central auditory system is commonly associated with a loss of inhibition (Casparly et al., 2005; Casparly et al., 2008; Takesian et al., 2009; Sanes et al., 2010; Auerbach et al., 2014; Ouda et al., 2015; Salvi et al., 2017), but increased intrinsic excitation and increased sensitivity to excitatory neurotransmitters also contribute to auditory system hyperactivity (Kotak et al., 2005; Abolafia et al., 2011; Li et al., 2013; Li et al., 2015; Sturm et al., 2017; Teichert et al., 2017; Balaram et al., 2019). We focus here on the loss of inhibition, as its relationship with cortical hyperactivity has received the most empirical attention, and it has been observed in both animal models and humans (Casparly et al., 2008; Gao et al., 2015; Dobri and Ross, 2021).

Sound exposure, ototoxic drug treatment, aging, and acoustic deprivation are all associated with reduced inhibition in the central auditory system compared to control animals (Rabang et al., 2012; Takesian et al., 2012; Kamal et al., 2013; Ouellet and de Villers-Sidani, 2014; Resnik and Polley, 2017). For low-intensity sound (Zhou and Merzenich, 2012; Kamal et al., 2013) and ototoxic drug exposure (Resnik and Polley, 2017), this has been attributed to reductions in the number of PV inhibitory neurons. However, in aging animals, the number of both PV and SST interneurons in auditory cortex appears to decrease (Figure 2B; de Villers-Sidani et al., 2010; Martin del Campo et al., 2012; Kamal et al., 2013; Ouellet and de Villers-Sidani, 2014; Cisneros-Franco et al., 2018), whereas other types of inhibitory neurons remain relatively constant in number across the lifespan (Ouellet and de Villers-Sidani, 2014). While the loss inhibitory neurons in aging mammals may occur secondary to degradation of the auditory periphery, a loss of inhibition and hyperactivity could also be the consequence of metabolic vulnerabilities of inhibitory neurons associated with aging processes directly (Brown, 1984; Ibrahim and Llano, 2019; Rogalla and Hildebrandt, 2020). Inhibitory neurons are metabolically highly demanding, but the cellular energy metabolism supporting inhibitory neurons may decline with age. This may lead to a selective loss of inhibitory function in central auditory brain regions (Ibrahim and Llano, 2019), and thus presents a mechanism by which auditory hyperactivity may arise from the aging process in the absence of age-related cochlear degeneration (King et al., 2007; Rogalla and Hildebrandt, 2020).

That a loss of inhibition – whether through peripheral damage, aging, or a combination thereof – may be an important driver of hyperactivity in auditory circuits is further highlighted by direct manipulations of inhibition. Downregulation of PV- and SST-expressing interneurons function via optogenetic (Aizenberg et al., 2015; Natan et al., 2015; Natan et al., 2017) or pharmacological approaches (Wang et al., 2002; Llano et al., 2012) increases spontaneous firing rates and sound-evoked responses in auditory cortex. Conversely, drugs that increase inhibition reduce spontaneous activity and suppress sound-evoked responses (Manunta and Edeline, 1997; Kaur et al., 2004).

While quantifying inhibitory interneuron function non-invasively in humans is difficult, a loss of inhibition has been documented in older adults with hearing loss compared to those without using magnetic resonance spectroscopy (Gao et al., 2015; Dobri and Ross, 2021; but see also discussions in Ouda et al., 2015). The underlying changes in neural subtype distribution are unknown; one study observed no age-related changes in PV inhibitory neurons (Bu et al., 2003), but included only five adults with a median age of 50 and may thus have underestimated the magnitude of PV inhibitory neuron loss associated with aging. Due to inherent technical challenges, further advances in measuring inhibition and quantifying inhibitory neurons in humans will likely rely on indirect inference drawn from measures of hyperresponsivity. Given the large body of work in animals, we suggest that the role of reduced inhibition along the auditory pathway, particularly in auditory cortex, be considered in future human studies.

Summary

In this section, we provided evidence from animal and human studies converging on the idea that damage to the auditory periphery, whether as a result of noise exposure, ototoxicity, the aging process, or some combination thereof, leads to changes throughout the auditory pathway. The most pronounced of these changes is a shift in the balance of excitation and inhibition that results in hyperactivity of auditory brain structures that can be measured both at rest (as increased spontaneous firing rates and increased synchronization among neurons) or in the presence of sound (as hyperresponsivity). These functional changes are in accordance with changes that have been observed in the distribution of inhibitory neurons in the auditory system. Moreover, research suggests these structural and functional changes do not arise from cochlear damage per se, but rather, result from decreased input to the system occurring subsequent to that damage.

Hyperactivity and perception

In this section, we describe the changes in auditory perception that may be associated with a loss of inhibition and subsequent hyperactivity in the auditory pathway. Empirical research in animals has focused mostly on electrophysiological studies of neuronal function, as described above, with fewer studies designed to assess the relationship between hyperactivity and auditory perception using behavioral approaches. Moreover, very few targeted investigations of hyperactivity and perception have been undertaken in humans, with the exception of the role of hyperactivity in tinnitus and hyperacusis. As a consequence, the associations laid out here between hyperactivity and auditory perception reflect hypotheses that require additional empirical verification. We begin with a discussion of tinnitus and hyperacusis percepts, and then consider how hyperactivity might

affect complex sound perception by fundamentally altering the encoding of sound features to which auditory pathway neurons are normally highly sensitive.

Tinnitus and hypersensitivity to sound

The perceptual phenomena most closely associated with hyperactivity are: 1) tinnitus – the perception of sound in absence of an identifiable sound source, which has been attributed to enhanced spontaneous firing rates and increased synchrony among neurons in the central auditory system of animals following cochlear damage (Bauer et al., 2008; Eggermont, 2012; Kalappa et al., 2014; Eggermont, 2015); and 2) hyperacusis – the sensation that sounds at moderate intensities are too loud (Eggermont and Roberts, 2004; Auerbach et al., 2014; Thomas et al., 2019a), which has been linked to enhanced sound-evoked activity in regions along the auditory pathway, most prominently in auditory cortex (Auerbach et al., 2014). Indeed, tinnitus and hyperacusis comprise the focus of the majority of articles that review, model, or theorize on the perceptual consequences of auditory system hyperactivity and related phenomena (Eggermont and Roberts, 2004; Eggermont, 2012; Knipper et al., 2013; Zeng, 2013; Eggermont, 2015; Zhao et al., 2016; Sedley, 2019; Knipper et al., 2020; Sheppard et al., 2020).

Behavioral indices of tinnitus have been observed in animals after the type of high-intensity sound exposures known to induce auditory system hyperactivity (Hayes et al., 2014). For example, exposed animals are poorer at perceiving a gap in noise than animals that were not exposed, where the continuous phantom sound percept of tinnitus is assumed to mask the gap (Turner et al., 2006; but see Radziwon et al., 2015 for counter evidence). Additionally, sound-exposed or ototoxic drug-treated animals categorize periods of silence as containing noise more often than control animals (Stolzberg et al., 2013; Hayes et al., 2014). Whether the degree of tinnitus correlates with the degree of spontaneous activity in the central auditory system is unknown. Since age-related tinnitus is often accompanied by peripheral hearing loss (Baguley et al., 2013), disentangling the two factors can be challenging (Sedley, 2019), and some evidence suggests increased spontaneous activity may be insufficient to explain tinnitus (Hayes et al., 2021). Indeed, a recent human neuroimaging study suggests that cortical hyperactivity is less prevalent in individuals with hearing loss *and* tinnitus than in those with hearing loss alone (Koops et al., 2020). Other mechanisms underlying tinnitus have also been proposed, but are beyond the scope of the current review (see Weisz et al., 2005; Sedley et al., 2016; Sedley, 2019; Knipper et al., 2020; Hayes et al., 2021).

Behavioral work in animals is also consistent with hypersensitivity to sound following cochlear damage. Cats with cochlear lesions show lower behavioral detection thresholds for electrical stimulation of neurons in the cochlear nucleus, inferior colliculus, and medial geniculate nucleus compared to pre-lesion (Gerken, 1979). Moreover, perceptual thresholds for pure tones

are maintained after drug-induced loss of up to 95% of inner hair cells in rodents (Lobarinas et al., 2013; Chambers et al., 2016b), potentially due to enhanced sound-evoked responses in auditory cortex (Qiu et al., 2000). This suggests that hyperresponsivity of auditory neurons may provide the means to maintain sound awareness despite severe cochlear damage (Asokan et al., 2018).

In humans, tinnitus and hyperacusis are often comorbid (Hebert et al., 2013), suggesting a common underlying mechanism. The probability of experiencing both tinnitus (Schaette et al., 2012; Brotherton et al., 2019) and hyperacusis (Fromby et al., 2007; Fournier et al., 2014; Munro et al., 2014; Brotherton et al., 2016, 2017) are increased in adults following temporary sound deprivation via ear plugging. These effects typically vanish a few hours after the ear plug is removed (Schaette et al., 2012; Brotherton et al., 2016) and, while this pattern of results would be predicted by the onset of hyperactivity, the exact mechanisms underlying the perceptual changes are unknown. Functional imaging in humans further indicates hyperactivity in auditory cortex in people with tinnitus (Koops et al., 2020). Tinnitus, however, does not appear to affect perception of speech in noise, frequency discrimination, or temporal-modulation detection (Zeng et al., 2020, but see also Ivansic et al., 2017). There is also evidence of increased sound sensitivity in humans following age-related hearing loss; older adults with clinical hearing loss detect smaller deviations in sound amplitude than older adults without (Bacon and Gleitman, 1992; Füllgrabe et al., 2003; Ernst and Moore, 2012; Sek et al., 2015; Schlittenlacher and Moore, 2016; but see Rosen et al., 2012 for results in early developmental hearing loss).

Although the perceptual changes described here are consistent with hyperactivity, investigations directly linking hyperactivity of the auditory pathway to tinnitus and perceptual hypersensitivity to sound are needed.

Spectral hearing

The ability to accurately perceive sound frequency is critical for the discrimination between two sounds with different spectral profiles, the perception of emotional content of speech, and the segregation of speech from other concurrent sound. Accordingly, neurons along the auditory pathway are frequency-tuned, responding preferentially to some sound frequencies at the expense of others (Moore, 1987; Ramachandran et al., 1999; Miller et al., 2002; Bartlett et al., 2011; Schreiner et al., 2011; Noelle O'Connell et al., 2014). Brain regions along the auditory pathway are also organized tonotopically, at least macroscopically (Bandyopadhyay et al., 2010), with neurons tuned to similar sound frequencies being clustered spatially (Kandler et al., 2009; Da Costa et al., 2011; Hackett, 2011; Baumann et al., 2013; Moerel et al., 2014; Saenz and Langers, 2014; Brewer and Barton, 2016).

Accurate perception and discrimination of sound frequency depends critically on narrow frequency tuning – the strong preference of neurons for specific, characteristic frequencies. Neural

inhibition shapes this tuning of neurons by modulating the probability of sound frequencies above and below the characteristic frequency eliciting a response (Isaacson and Scanziani, 2011). In auditory cortex, frequency tuning appears to be mediated largely by PV- and SST- expressing inhibitory interneurons (Aizenberg et al., 2015; Kato et al., 2017). Pharmacological and optogenetic manipulations that specifically modulate the activity of these interneurons have revealed that an increase in inhibition narrows tuning (Kaur et al., 2004; Aizenberg et al., 2015), whereas a decrease in inhibition broadens tuning (Wang et al., 2002; Aizenberg et al., 2015). These changes in tuning have demonstrable perceptual consequences; increasing or decreasing PV inhibition improves or worsens frequency discrimination, respectively (Aizenberg et al., 2015; SST-expressing interneurons were not tested).




Figure 3: Schematic frequency-response areas. Normal, shifted, and broadened frequency tuning. (Note that schematic data are displayed).

While less is known about the perceptual consequences of a loss of inhibition and hyperactivity, aged animals and animals with peripheral injury due to sound exposure or drug treatment also exhibit broadened tuning or tuning with reduced specificity (Turner et al., 2005; Barsz et al., 2007; Izquierdo et al., 2008; Leong et al., 2011). Reorganization of tonotopic maps, where neurons shift their frequency preference to lower frequencies, has been observed as well (Figure 3; Willott, 1986; Willott et al., 1988a; Norena et al., 2003; Turner et al., 2005; de Villers-Sidani et al., 2010; Zhou and Merzenich, 2012; Chambers et al., 2016a; Resnik and Polley, 2017). Thus, cochlear pathology associated with aging and sound exposure gives rise to changes in neural tuning that are comparable to those occurring when inhibition is specifically suppressed (Wang et al., 2002; Aizenberg et al., 2015); this consistent with the loss of PV and SST interneurons described above (Martin del Campo et al., 2012; Kamal et al., 2013; Ouellet and de Villers-Sidani, 2014; Resnik and Polley, 2017). These changes are not limited to severe peripheral damage; frequency tuning of neurons in auditory cortex broadens even after a few weeks of sound exposure at levels that humans typically encounter (65-70 dB SPL; Zhou and Merzenich, 2012; Thomas et al., 2019b).

As described above, direct evidence of changes in inhibitory interneuron distribution and function is limited in aged humans and humans with hearing impairment. Accordingly, few studies have provided demonstrable links between hyperactivity and frequency perception. However, broadening of neural tuning is consistent with decreased frequency discrimination acuity and frequency-modulation detection observed in older adults with and without hearing loss (Turner and Nelson, 1982; Nelson and Freyman, 1986; Moore and Skrodzka, 2002; Clinard et al., 2010; Moore, 2014; Moore et al., 2019). Decreased frequency-modulation detection, in turn, is associated with reduced speech-in-noise perception (Holmes and Griffiths, 2019; Parthasarathy et al., 2020). Moreover, it has been reported that severe hearing loss leads to tonotopic map reorganization in human auditory cortex (Wolak et al., 2017; Koops et al., 2020), however, this finding is not consistent across studies (Saenz and Langers, 2014; Ouda et al., 2015). Tonotopic map reorganization (Mühlnickel et al., 1998) and broadened frequency tuning (Sekiya et al., 2017) have also been observed in individuals with tinnitus (albeit inconsistently; e.g. Koops et al., 2020). In contrast, frequency-specific adaptation in auditory cortex – a unique form of frequency tuning – appears to be unaltered in older compared to younger adults, despite concurrent hyperresponsivity that suggests a loss of inhibition (Herrmann et al., 2013a). Together, these studies suggest that, at least in some individuals, frequency tuning in auditory brain regions and the neuronal inhibition governing it are altered by peripheral damage in a way that would be expected to affect perception.

In sum, frequency-discrimination behavior in aged animals and animals with hearing loss is affected in a manner consistent with changes in the frequency-tuning of neurons in auditory cortex – changes that are likely the result of reduced inhibition and hyperactivity. Deficits in frequency discrimination behavior are well established in older humans, but the currently available data are insufficient to directly link these perceptual deficits to neural hyperactivity and altered frequency tuning.

Spatial hearing

Most auditory signals that reach a listener's ears consist of a mixture of sounds from spatially distinct sources. A listener must isolate a sound source of interest (e.g., speech) from the mixture and segregate it from sounds originating from other locations in order to track and attend it over time. Hearing loss and processes related to aging impair spatial hearing abilities (Brown, 1984; Abel et al., 2000; Koehnke and Besing, 2001), leading to a significant behavioral detriment that may be tied to auditory system hyperactivity.

The accurate perception of sound source location is thought to depend on narrow spatial tuning that relies on primary (King et al., 2007) and posterior-dorsal regions of auditory cortex (Rauschecker and Tian, 2000; Arnott et al., 2004; Rauschecker and Scott, 2009; Woods and Alain,

2009; Herrmann et al., 2011; Rauschecker, 2011). Neurons in posterior auditory cortex are spatially tuned such that they respond preferentially to sounds originating from one location at the expense of other locations (Figure 4; Woods et al., 2006; Juarez-Salinas et al., 2010).




Figure 4: Schematic of sound-evoked activity depicting spatial tuning for normal cortical activity and hyperactivity. Left: Schematic of a person positioned at the center of surrounding speakers. **Right:** Spatial tuning centered at -45 degrees. (Note that schematic data are displayed).

Electrophysiological recordings from monkeys demonstrate that age-related hyperactivity extends from primary auditory cortex into spatially-tuned posterior auditory cortex (Juarez-Salinas et al., 2010), suggesting a loss of inhibition in these regions as individuals age. Hyperactivity in posterior auditory cortex also coincides with broadened spatial tuning (Juarez-Salinas et al., 2010) that would be expected to decrement spatial acuity. Accordingly, reduced behavioral performance in spatial hearing task have been reported in aged rats (Brown, 1984) and humans (Abel et al., 2000; Koehnke and Besing, 2001) compared to younger listeners.

In sum, hyperactivity following peripheral damage appears throughout auditory cortex, including dorsal-stream regions that are crucial to spatial hearing. The loss of inhibition that underlies these changes also results in broadened spatial tuning of neurons in these dorsal areas, which may underlie behavioral decrements observed in aged humans. However, a direct link between hyperactivity and the perception of sound location is currently missing.

Temporal sensitivity

A sound originating from a given source is not only spectrally and spatially unique but also has a unique profile with which its amplitude and frequency fluctuate over time. In speech, for example, low- and high-frequency amplitude and frequency modulations reflect the envelope and fine-structure of a sound, respectively (Rosen, 1992). The relative contributions of envelope and fine structure to the perception of complex sounds like speech remains the topic of much research, and is beyond the scope of this review (but see Drullman, 1995; Lorenzi et al., 2006; Shamma and Lorenzi, 2013). Nevertheless, successful tracking of amplitude and frequency modulations across frequencies is considered crucial for the segregation of sounds from different sources and ultimately for speech comprehension in the presence of other sounds (Kerlin et al., 2010; Giraud and Poeppel, 2012; Edwards and Chang, 2013; Peelle and Davis, 2013).

Neurons along the auditory pathway are sensitive to acoustic periodicity, in that their activity synchronizes with a sound's amplitude and frequency fluctuations (Picton et al., 2003; Anderson et al., 2012; Herrmann et al., 2013b; Henry et al., 2014; Goossens et al., 2016; Parthasarathy et al., 2019). Electrophysiological work in animals and humans increasingly points to frequency-dependent changes in neural synchronization associated with aging and hearing loss. Auditory neurons synchronize more strongly with lower-frequency periodicities and less strongly with higher-frequency periodicities in aged compared to young rodents (Figure 5; Palombi et al., 2001; Schatteman et al., 2008; Overton and Recanzone, 2016; Herrmann et al., 2017; Parthasarathy et al., 2019). Similar changes have been observed following continuous exposure to sound at low-to-moderate intensities (~65 dB SPL; Zhou and Merzenich, 2012). Overall firing rate may also be increased in older compared to younger animals for sounds with periodic amplitude modulations, especially for slower modulation rates (Palombi et al., 2001; Walton et al., 2002). However, neural synchronization in inferior colliculus and thalamus appears to be reduced for all stimulation frequencies when over 95% of afferent fibers are eliminated through drug manipulations in animals (Chambers et al., 2016b; Chambers et al., 2016a). Both enhanced low-frequency (Purcell et al., 2004; Goossens et al., 2016; Presacco et al., 2016a; Herrmann et al., 2019) and decreased high-frequency synchronization (Purcell et al., 2004; Anderson et al., 2012; Clinard and Tremblay, 2013) have also been reported for older compared to younger humans, although low-frequency enhancements may not necessarily scale with the degree of peripheral hearing loss (Goossens et al., 2019; Herrmann et al., 2019; Presacco et al., 2019; but see Millman et al., 2017). Modeling and empirical measurements of neural activity suggests that a loss of inhibition could underlie this bidirectional change, although the effects of inhibition on synchronization appear complex and non-linear (Yang and Pollak, 1997; Backoff et al., 1999; Rabang et al., 2012).




Figure 5: Simulated neural activity depicting neural synchronization to periodicity in sound for normal activity and hyperactivity. Panel A and B show simulated activity for low- and high-frequency stimulus periodicities, respectively. **Top:** Stimulus periodicity and simulated activity time courses. **Bottom:** Frequency spectra (from a fast Fourier transform) of the simulated response time courses. Peaks

in the spectra indicate that neural activity synchronized with the periodicity in the acoustic signal. Peaks at harmonic frequencies emerge when time courses of synchronized neural activity are not fully sinusoidal. (Note that schematic, simulated data are displayed).

In line with these changes in neural synchronization, individuals with unilateral hearing impairment perceive low-frequency amplitude modulations (4–32 Hz) as fluctuating more strongly in their hearing-impaired compared to their non-impaired ear (Moore et al., 1996), while discrimination of higher-frequency pure tones (500-1000 Hz) declines with age and hearing loss (Turner and Nelson, 1982; Nelson and Freyman, 1986; Clinard et al., 2010). However, because the relationship between low-frequency and high-frequency periodicity cues is not fully understood, the behavioral consequences of a bidirectional shift in neural synchronization on real-world listening are hard to predict. Reduced higher-frequency synchronization has been associated with poorer speech-in-noise perception (Anderson and Kraus, 2010; but see Presacco et al., 2016a), but the mechanism underlying this behavioral deficit is unknown. Additionally, enhanced low-frequency synchronization has been linked to poor speech intelligibility, specifically when a modulated masker sound is present (Millman et al., 2017; Goossens et al., 2018), suggesting that hyperactivity makes ignoring irrelevant, fluctuating sounds more difficult.

Finally, neural and perceptual sensitivity to short gaps – that is, periods of silence – in a sound have long been considered a measure of temporal resolution of the auditory system (Frisina et al., 2001; Fitzgibbons and Gordon-Salant, 2010; Walton, 2010), and the ability to detect gaps appears to be a good predictor of speech comprehension abilities (Gordon-Salant and Fitzgibbons, 1993). Work in rodents shows reduced neural responses to gaps in the inferior colliculus of older compared to younger animals and in animals exposed to sound (Walton et al., 1998; Allen et al., 2003; Sturm et al., 2017). In humans, older adults exhibit enhanced responses to gaps in auditory cortex compared to younger listeners, indicating that the auditory system is hyperresponsive (Ross et al., 2010; Lister et al., 2011; but see Harris et al., 2012). Accordingly, behavioral gap-detection thresholds are typically larger for older compared to younger human adults (Glasberg et al., 1987; Schneider et al., 1994; Snell, 1997; Schneider et al., 1998; Snell and Frisina, 2000; Humes et al., 2009; Harris et al., 2012). However, sensorineural hearing loss may not additionally contribute to changes in gap-detection thresholds beyond effects of aging (Moore and Glasberg, 1988; Fitzgibbons and Gordon-Salant, 1996; Gordon-Salant and Fitzgibbons, 1999, but see Moore et al., 1992), suggesting that peripheral damage and associated hyperactivity may not contribute substantially to gap-detection performance. This is further corroborated by the observation that salicylate, a drug known to induce auditory system hyperactivity (see above), does not seem to affect behavioral gap-detection performance in animals (Radziwon et al., 2015).

Summary

The work reviewed in this section suggests that hyperactivity (or the loss of inhibition underlying it) may contribute to a variety of perceptual phenomena that are common among normally aging adults and individuals with hearing loss. Hyperactivity has long been hypothesized to underlie tinnitus and hyperacusis (Eggermont and Roberts, 2004; Knipper et al., 2013). But there is growing evidence that a loss of inhibition and hyperactivity also affect spectral, spatial, and periodicity processing, altering responses to complex auditory signals like speech, and increasing interference from background sounds (Millman et al., 2017; Goossens et al., 2018). Since listening in acoustically challenging environments relies heavily on these processes, it is not surprising that speech perception in noise declines with age and hearing loss (Pichora-Fuller, 2003; Presacco et al., 2016a, 2019). Hence, it is becoming increasingly clear that peripheral damage and associated plasticity along the auditory pathway fundamentally alter how sounds are represented neurally and how they are perceived. Yet, research linking hyperactivity to perception remains underdeveloped.

Open questions and ongoing challenges in the study of hyperactivity

Despite the increasing evidence of hyperactivity following auditory peripheral damage, and the clear hypotheses regarding how perceptual impairments could arise secondary to hyperactivity, open questions remain. In many cases, the types of investigations necessary to fill the knowledge gaps outlined in the previous two sections face considerable conceptual, experimental, and methodological challenges that we will attempt to highlight in the following sections. Indeed, a fulsome understanding of the role of inhibition and hyperactivity in reshaping neural representations of sound and subsequent behaviors following peripheral degeneration will require additional study in both animal models of hearing loss and in human listeners. Critically, the ability to integrate findings across models, and the development of innovative approaches to the measurement of hyperactivity are necessary to move this field forward.

Open question 1: What is the functional role of auditory system hyperactivity?

The loss of inhibition giving rise to hyperactivity along the auditory pathway is commonly referred to as a process that *compensates* for reduced inputs from damaged peripheral structures to central brain regions to maintain sensation (Caspary et al., 2008; Schaette and McAlpine, 2011; Knipper et al., 2013; Auerbach et al., 2014; Chambers et al., 2016a; Möhrle et al., 2016; Salvi et al., 2017). Homeostatic mechanisms are thought to underlie this compensatory adjustment in neuronal activity (Caspary et al., 2008; Sanes et al., 2010; Auerbach et al., 2014; Teichert et al., 2017), by stabilizing and maintaining excitability such that both low levels of excitability and run-away excitability are avoided. These homeostatic mechanisms operate across many levels, from synaptic

changes to changes at the whole-brain network level (Turrigiano, 1999; Turrigiano and Nelson, 2000; Turrigiano, 2012; Keck et al., 2017). Accordingly, the term ‘compensation’ has recently been adopted by human cognitive neuroscientists to describe, for example, enhanced sound-evoked activity observed in older people and people with hearing loss (Bidelman et al., 2014; Herrmann et al., 2016; Presacco et al., 2016b, a; Goossens et al., 2018, 2019).

The use of the term ‘compensation’ to describe homeostatic mechanisms can give rise to considerable confusion. The fields of psychology and cognitive neuroscience commonly use ‘compensation’ to refer to behavioral or neurological adaptations that arise to offset a perceptual deficit (Bäckman and Dixon, 1992; Salthouse, 1995). In the case of hearing loss, ‘compensation’ is thus commonly used to refer to a breadth of adaptive strategies that help maintain or restore the perception of sound. Homeostatic compensation, however, stabilizes and maintains neuronal excitability without consideration of perceptual consequences (Turrigiano, 1999; Turrigiano and Nelson, 2000; Turrigiano, 2012; Nahmani and Turrigiano, 2014). Indeed, homeostatic compensation may actually impair auditory perception (as described above). Thus, the term ‘compensation’ must be clearly defined when studying hyperactivity in sensory circuits.

Many observations of hyperactivity following peripheral damage could be considered to reflect homeostatic compensation, however, this idea is increasingly debated (Herrmann et al., 2016; Asokan et al., 2018; Cisneros-Franco et al., 2018; Ibrahim and Llano, 2019; Rogalla and Hildebrandt, 2020). Whereas enhanced spontaneous activity has been observed at essentially all levels of the auditory pathway following peripheral damage (Kaltenbach et al., 2002; Rachel et al., 2002; Eggermont, 2015; Parthasarathy et al., 2019), the effects on sound-evoked activity are not as straightforward. For example, sound-evoked neural activity can be suppressed in the inferior colliculus, and enhanced in auditory cortex following peripheral damage (Hughes et al., 2010; Auerbach et al., 2014; Chambers et al., 2016b; Herrmann et al., 2016; Zhao et al., 2016). If homeostatic mechanisms underlie these changes, this would suggest they are not equally successful across brain regions.

An alternative view arises from observed similarities between damage-induced hyperactivity in auditory brain regions and the activity levels in the developing auditory system (Cisneros-Franco et al., 2018). Reduced inhibition may return auditory brain regions to a state of increased neural plasticity (Herrmann et al., 2016; Hattori et al., 2017; Cisneros-Franco et al., 2018; Cisneros-Franco and de Villers-Sidani, 2019; Thomas et al., 2019b) that resembles early developmental periods during which inhibitory circuits have not yet matured and new connections between neurons can be formed efficiently (Cisneros-Franco et al., 2018). Indeed, a reduction of inhibition is thought to foster Hebbian learning (Hattori et al., 2017) and new neural connections are formed in auditory cortex following peripheral hearing loss that give rise to visual and somatosensory functions (Allman et al., 2009; Ptito et al., 2012). However, homeostatic and

Hebbian mechanisms may be intertwined at the molecular level (Turrigiano and Nelson, 2000) and disentangling them will present a significant future challenge.

Open question 2: How does lifelong hearing experience influence damage and hyperactivity?

Accumulated sound exposure over the lifespan is thought to significantly contribute to hearing loss in older people (Schmiedt, 2010; Wu et al., 2020). The exact nature of this relationship has been the focus of substantial interest recently based on the observations that even low-intensity sound exposure can damage inner hair cell synapses (Kujawa and Liberman, 2009; Liberman and Kujawa, 2017), and that this type of cochlear degeneration is not fully captured by standard hearing assessment tools in humans, such as pure-tone audiometry (Schaette and McAlpine, 2011; Plack et al., 2014; but see recent approaches Liberman et al., 2016; Mepani et al., 2020; Parthasarathy et al., 2020). As described above, peripheral damage – including that which arises from sound exposure – may lead to a loss of inhibition and hyperactivity in the auditory pathway (Auerbach et al., 2014; Zhao et al., 2016; Salvi et al., 2017). Concerts, urban streets, bars, restaurants, schools, train stations, hospitals, and industrial workplaces are only a few examples of acoustic environments that humans encounter regularly in which sound levels are comparable to those that induce hyperactivity in noise-exposed animals (Hopkins, 1994; Olsen, 1998; Tsai et al., 2009). We may thus expect many humans to exhibit hyperactivity along their auditory pathway reflecting frequent exposures.

However, it has also been demonstrated that, under certain circumstances, sound exposure may have a protective function. For example, routine exposure to sounds at low (~46 dB; Fukushima et al., 1990) or moderate-to-high intensities (e.g., 80 dB SPL; Canlon et al., 1988; Yoshida and Liberman, 2000; Niu et al., 2004, 2007; Oliver et al., 2011) – often referred to as enriched acoustic environments – can reduce the damage induced by subsequent traumatic, high-intensity sounds and improve auditory processing. In fact, the protective effects of auditory enrichment can even mitigate sound-induced peripheral damage when presented *after* high-intensity exposure (Niu et al., 2004; Norena and Eggermont, 2006; Noreña et al., 2006; Niu et al., 2007). In humans, enriched acoustic environments have also been shown to reduce perceptual hypersensitivity to sound (Norena and Chery-Croze, 2007) and have been used to treat tinnitus (Henry et al., 2006; Jastreboff, 2007).

This presents a paradox: how can enriched environmental exposure exert a protective function while noise exposure at the same amplitude and duration results in considerable peripheral damage? The answer may lie in the sound structure. Exposure to unmodulated, continuous noise leads to hyperactivity and increased plasticity in auditory cortex, whereas exposure to sound with more naturalistic, structured properties, such as amplitude modulation, does not (Zhou and Merzenich, 2012; Thomas et al., 2019b). Enriched environmental exposure to a temporally

complex sound, compared to a less complex sound, has also been shown to improve temporal processing acuity in the auditory midbrain (Dziorny et al., 2020). As such, it is possible that humans' everyday environments may protect them (to some extent) from occasional high-intensity sound exposure, and that this could explain why investigations of the impact of recreational sound exposure on hearing function have revealed mixed results (Prendergast et al., 2017a; Prendergast et al., 2017b; but see Liberman et al., 2016; Imam and Hannan, 2017). It may ultimately be critical to distinguish between sounds that may be more harmful for the auditory system, such as occupational or industrial sounds with broad-band, unmodulated, continuous properties, and sounds like music and speech, which elicit structured, correlated neural activity which may be less harmful (Thomas et al., 2019b; Dziorny et al., 2020).

Open question 3: How does hyperactivity change as function of hearing loss severity?

Hyperactivity in human auditory cortex is most commonly measured indirectly as hyperresponsivity to sound (e.g., Alain et al., 2014; Bidelman et al., 2014; Herrmann et al., 2016). However, a decrease in inhibition and hyperresponsivity changes as a function of the degree of auditory deprivation (Qiu et al., 2000; Resnik and Polley, 2017; Figure 6). It may be tempting to interpret decreased responsivity to sound following severe cochlear damage as evidence against hyperactivity in auditory brain regions. However, one must be mindful that peripheral damage also affects signal generation at the cochlea. Animals with a mild to moderate noise-induced or age-related hearing loss show hyperresponsivity in auditory cortex (Hughes et al., 2010; Auerbach et al., 2014). However, as the degree of peripheral damage increases – for example, when 80-90% of hair cells are lost – hyperresponsivity is offset by the reduction in signal transduction, and cortical responses are no larger than those observed in unexposed animals (Qiu et al., 2000; Chambers et al., 2016b; Salvi et al., 2017). However, in the extreme case of complete peripheral damage, signal transduction is lost and auditory cortex becomes non-responsive to sound (Figure 6).




Figure 6: Hypothetical depiction of auditory cortical activity as a function of cochlear damage. The solid line reflects cortical activity measured indirectly using sound stimulation. The dashed line reflects cortical activity measured directly via spontaneous neuronal activity or indirectly via electric stimulation of auditory cortex.

Thus, hyperresponsivity to sound may only be a useful proxy for hyperactivity when sufficient peripheral integrity exists to facilitate sound transduction. Indeed, behavioral thresholds for electrical stimulation of auditory thalamic neurons (bypassing peripheral damage) are improved after complete, or almost complete peripheral damage compared to a pre-damage control (Gerken, 1979), suggesting hyperresponsivity in auditory circuits that would not be evident using acoustic stimulation. Unfortunately, techniques that measure spontaneous neural activity or behavioral responses to electrical stimulation are not typically applicable in humans.

How long hyperactivity in auditory cortex persists following complete auditory deprivation is also unclear. Based on research on other biological tissue (Cannon and Rosenblueth, 1949), hyperactivity of neurons in auditory cortex is expected to persist for at least several weeks after complete denervation. However, this is likely an underestimate, as neurons in auditory cortex are integrated into larger networks, including non-auditory brain regions, that continue to deliver synaptic inputs to auditory cortex following hearing loss. Indeed, the persistence of non-auditory inputs to auditory brain regions is hypothesized to underlie the preservation of normal patterns of connectivity in auditory cortex, even following profound early-onset hearing loss (Chabot et al., 2015; Butler et al., 2016; Butler et al., 2018). Moreover, the way in which excitability changes over time is also likely dependent on the degree of peripheral damage and subsequent hearing loss.

In sum, current measures of hyperresponsivity to sound are likely non-linear, such that the absence of *measurable* hyperresponsivity associated with aging or hearing loss may not imply the absence of hyperactivity nor of a loss of inhibition. This is particularly relevant for studies of older human adults and people with hearing loss that often rely on non-invasive recordings of event-related potentials like the P1 and N1 (e.g., Laffont et al., 1989; Anderer et al., 1996; Sörös et al., 2009; Bidelman et al., 2014; Stothart and Kazanina, 2016; Henry et al., 2017) that originate from auditory cortex (Näätänen and Picton, 1987; Huotilainen et al., 1998; Maess et al., 2007; Herrmann et al., 2018). The development of measures designed to capture hyperactivity more directly from human listeners is an important first step towards understanding how hyperactivity is shaped by the degree of hearing loss.

Open question 4: How do experimental design choices impact measures of hyperactivity?

In addition to its dependence on the degree of peripheral damage, hyperresponsivity to sound also varies as a function of stimulus characteristics. For example, as described above, hyperactive auditory neurons in aged and noise-exposed animals show reduced neural synchronization specifically for higher-frequency periodicities (Zhou and Merzenich, 2012; Herrmann et al., 2017). Studies using high-frequency periodic stimuli may thus underestimate hyperactivity, and observe different effects compared to studies that present stimuli with low-frequency periodicities.

Another example involves the relationship between neural responsivity and the temporal presentation dynamics of auditory stimuli. Neurons in the auditory pathway undergo adaptation – a reduction in neural response to sound due to sustained stimulation (Whitmire and Stanley, 2016) – and require time to recover full responsivity after responding to sound. This recovery time appears to be shortened in older compared to younger adults (de Villers-Sidani et al., 2010; Mishra et al., 2014; Herrmann et al., 2016; Herrmann et al., 2019). Thus, sounds presented in short succession may lead to inputs to a larger proportion of adapted neurons in individuals with normal hearing compared to aged individuals or individuals with hearing loss. As a result, studies with different stimulation rates may confound hyperactivity and neural recovery times, and may come to different conclusions.

This short list of examples is not exhaustive – it simply serves as a reminder that a loss of inhibition and increased activity in the central auditory system may not always manifest as response enhancements to sound and that careful consideration of experimental design is required for the meaningful study of neural inhibition, hyperactivity, and perception.

Summary

In this section, we reviewed open questions in the study of hyperactivity, and outlined some of the challenges inherent to their study. First, we highlighted the ongoing debate surrounding the relative contributions of homeostatic mechanisms that stabilize excitation and Hebbian mechanisms that support the formation of new connections between neurons to hyperactivity in the auditory system. Second, we discussed the paradox that sound exposure can have damaging *and* protective effects on peripheral auditory function, suggesting that spectral and temporal properties of exposure sounds may play an underappreciated role in whether and how sound exposure impairs hearing. Finally, we highlighted the non-linear relationship between peripheral damage, hyperactivity, and hyperresponsivity to sound and between experimental factors and hyperresponsivity to sound. These non-linearities pose a challenge for investigations of how hyperactivity interacts with the degree of hearing loss in humans, where hyperactivity is measured non-invasively as hyperresponsivity to sound.

Conclusions

Hearing loss in older adulthood is associated with a broad range of perceptual impairments that include a loss of sensitivity, tinnitus, hyperacusis, problems locating sounds, difficulties with speech comprehension in noise, and more. Traditionally, many of these perceptual deficits have been associated with damage to the cochlea in the auditory periphery. However, an increasing amount of evidence, mostly from studies in animals, suggests that hearing dysfunction results from

changes in the entire auditory system, from the periphery to cortex. The most prominent change following peripheral degeneration is an increase in activity levels of downstream brain regions in the auditory pathway. In this paper, we have a) reviewed the causes and neurophysiological mechanisms underlying hyperactivity, showing that hyperactivity does not arise from cochlear damage per se, but rather results from decreased input to the system occurring subsequent to that damage; b) suggested that hyperactivity alters all aspects of auditory perception – including spectral, spatial, and temporal hearing – and, in turn, speech comprehension abilities when background sound is present; and c) discussed the need for further understanding of the functional role of hyperactivity, and outlined some of the open questions in the study of hyperactivity in humans. Our hope is that this review provides a foundation from which researchers can develop targeted approaches for investigating hyperactivity in humans.

Declarations

Funding

BH was supported by the Canada Research Chair program.

Conflicts of interest/Competing interests

The authors have no conflict of interest/competing interests.

Availability of data and materials

Not applicable.

Code availability

Not applicable.

Authors' contributions

BH had the idea for the article. BH and BEB performed the literature search. BH drafted the manuscript. BH and BEB critically revised the manuscript.

Ethics approval

Not applicable.

Consent to participate

Not applicable.

Consent for publication

BH and BEB consent to the publication.

References

- Abel SM, Giguere C, Consoli A, Papsin BC (2000) The effect of aging on horizontal plane sound localization. *Journal of the Acoustical Society of America* 108:743-752.
- Abolafia JM, Vergara R, Arnold MM, Reig R, Sanchez-Vives MV (2011) Cortical auditory adaptation in the awake rat and the role of potassium currents. *Cerebral Cortex* 21:977-990.
- Aizenberg M, Mwilambwe-Tshilobo L, Briguglio JJ, Natan RG, Geffen MN (2015) Bidirectional Regulation of Innate and Learned Behaviors That Rely on Frequency Discrimination by Cortical Inhibitory Neurons. *PLoS Biology* 13:e1002308.
- Alain C, McDonald K, Van Roon P (2012) Effects of age and background noise on processing a mistuned harmonic in an otherwise periodic complex sound. *Hearing Research* 283:126-135.
- Alain C, Roye A, Salloum C (2014) Effects of age-related hearing loss and background noise on neuromagnetic activity from auditory cortex. *Frontiers in Systems Neuroscience* 8:Article 8.
- Allen PD, Burkard RF, Ison JR, Walton JP (2003) Impaired gap encoding in aged mouse inferior colliculus at moderate but not high stimulus levels. *Hearing Research* 186:17-29.
- Allman B, Keniston LP, Meredith MA (2009) Adult deafness induces somatosensory conversion of ferret auditory cortex. *Proceedings of the National Academy of Sciences* 106:5925–5930.
- Amenedo E, Díaz F (1999) Ageing-related changes in the processing of attended and unattended standard stimuli. *NeuroReport* 10:2383-2388.
- Anari M, Axelsson A, Eliasson A, Magnusson L (1999) Hypersensitivity to sound: Questionnaire data, audiometry and classification. *Scandinavian Audiology* 28:219-230.
- Anderer P, Semlitsch HV, Saletu B (1996) Multichannel auditory event-related brain potentials: effects of normal aging on the scalp distribution of N1, P2, N2 and P300 latencies and amplitudes. *Electroencephalography and clinical Neurophysiology* 99:458-472.
- Anderson S, Kraus N (2010) Objective Neural Indices of Speech-in-Noise Perception. *Trends in Amplification* 14:73-83.
- Anderson S, Parbery-Clark A, White-Schwoch T, Kraus N (2012) Aging Affects Neural Precision of Speech Encoding. *The Journal of Neuroscience* 32:14156-14164.
- Arnott SR, Binns MA, Grady CL, Alain C (2004) Assessing the auditory dual-pathway model in humans. *NeuroImage* 22:401-408.
- Asokan MM, Williamson RS, Hancock KE, Polley DB (2018) Sensory overamplification in layer 5 auditory corticofugal projection neurons following cochlear nerve synaptic damage. *Nature Communications* 9:2468.

- Auerbach BD, Rodrigues PV, Salvi RJ (2014) Central gain control in tinnitus and hyperacusis. *Frontiers in Neurology* 5:Article 206.
- Bäckman L, Dixon RA (1992) Psychological Compensation: A Theoretical Framework. *Psychological Bulletin* 112:259-283.
- Backoff PM, Palombi PS, Caspary DM (1999) γ -Aminobutyric acidergic and glycinergic inputs shape coding of amplitude modulation in the chinchilla cochlear nucleus. *Hearing Research* 134:77-88.
- Bacon SP, Gleitman RM (1992) Modulation detection in subjects with relatively flat hearing losses. *J Speech Hear Res* 35:642-653.
- Baguley DM (2003) Hyperacusis. *Journal of the Royal Society of Medicine* 96:582-585.
- Baguley DM, McFerran D, Hall AD (2013) Tinnitus. *The Lancet* 382:1600-1607.
- Balaram P, Hackett TA, Polley DB (2019) Synergistic Transcriptional Changes in AMPA and GABAA Receptor Genes Support Compensatory Plasticity Following Unilateral Hearing Loss. *Neuroscience* 407:108-119.
- Bandyopadhyay S, Shamma SA, Kanold PO (2010) Dichotomy of functional organization in the mouse auditory cortex. *Nature Neuroscience* 13:361-368.
- Bao J, Ohlemiller KK (2010) Age-related loss of spiral ganglion neurons. *Hearing Research* 264:93-97.
- Barsz K, Wilson WW, Walton JP (2007) Reorganization of receptive fields following hearing loss in inferior colliculus neurons. *Neuroscience* 147:532-545.
- Bartlett EL, Sadagopan S, Wang X (2011) Fine frequency tuning in monkey auditory cortex and thalamus. *Journal of Neurophysiology* 106:849-859.
- Bauer CA, Turner JG, Caspary DM, Myers KS, Brozoski TJ (2008) Tinnitus and inferior colliculus activity in chinchillas related to three distinct patterns of cochlear trauma. *J Neurosci Res* 86:2564-2578.
- Baumann S, Petkov CI, Griffiths TD (2013) A unified framework for the organization of the primate auditory cortex. *Frontiers in Systems Neuroscience* 7:Article 11.
- Bidelman GM, Villafuerte JW, Moreno S, Alain C (2014) Age-related changes in the subcortical encoding and categorical perception of speech. *Neurobiology of Aging* 35:2526-2540.
- Blackwell JM, Geffen MN (2017) Progress and challenges for understanding the function of cortical microcircuits in auditory processing. *Nature Communications* 8:2165.
- Brewer AA, Barton B (2016) Maps of the Auditory Cortex. *Annual Review of Neuroscience* 39:385-407.

- Brotherton H, Plack CJ, Schaette R, Munro KJ (2016) Time course and frequency specificity of sub-cortical plasticity in adults following acute unilateral deprivation. *Hearing Research* 341:210-219.
- Brotherton H, Plack CJ, Schaette R, Munro KJ (2017) Using acoustic reflex threshold, auditory brainstem response and loudness judgments to investigate changes in neural gain following acute unilateral deprivation in normal hearing adults. *Hearing Research* 345:88-95.
- Brotherton H, Turtle C, Plack CJ, Munro KJ, Schaette R (2019) Earplug-induced changes in acoustic reflex thresholds suggest that increased subcortical neural gain may be necessary but not sufficient for the occurrence of tinnitus. *Neuroscience* 407:192-199.
- Brown CH (1984) Directional hearing in aging rats. *Experimental Aging Research* 10:35-38.
- Bu J, Sathyendra V, Nagykerly N, Geula C (2003) Age-related changes in calbindin-D28k, calretinin, and parvalbumin-immunoreactive neurons in the human cerebral cortex. *Experimental Neurology* 182:220-231.
- Butler BE, Chabot NA, Lomber SG (2016) Quantifying and comparing the pattern of thalamic and cortical projections to the posterior auditory field in hearing and deaf cats. *The Journal of Comparative Neurology* 524:3042-3063.
- Butler BE, de la Rua A, Ward-Able T, Lomber SG (2018) Cortical and thalamic connectivity to the second auditory cortex of the cat is resilient to the onset of deafness. *Brain Structure and Function* 223:819-835.
- Canlon B, Borg E, Flock A (1988) Protection against noise trauma by pre-exposure to a low-level acoustic stimulus. *Hearing Research* 34:197-200.
- Cannon WB, Rosenblueth A (1949) *The supersensitivity of denervated structures: A law of denervation*. New York: The Macmillan Company.
- Caspary DM, Schatteman TA, Hughes LF (2005) Age-related changes in the inhibitory response properties of dorsal cochlear nucleus output neurons: role of inhibitory inputs. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 25:10952-10959.
- Caspary DM, Ling L, Turner JG, Hughes LF (2008) Inhibitory neurotransmission, plasticity and aging in the mammalian central auditory system. *The Journal of Experimental Biology* 211:1781-1791.
- Chabot NA, Butler BE, Lomber SG (2015) Differential modification of cortical and thalamic projections to cat primary auditory cortex following early-and late-onset deafness. *The Journal of Comparative Neurology* 529:2297-2232.
- Chambers AR, Salazar JJ, Polley DB (2016a) Persistent Thalamic Sound Processing Despite Profound Cochlear Denervation. *Frontiers in Neural Circuits* 10:Article 72.

- Chambers AR, Resnik J, Yuan Y, Whitton JP, Edge AS, Liberman MC, Polley DB (2016b) Central Gain Restores Auditory Processing following Near-Complete Cochlear Denervation. *Neuron* 89:867-879.
- Cisneros-Franco JM, de Villers-Sidani É (2019) Reactivation of critical period plasticity in adult auditory cortex through chemogenetic silencing of parvalbumin-positive interneurons. *Proceedings of the National Academy of Sciences* 116:26329-26331.
- Cisneros-Franco JM, Ouellet L, Kamal B, de Villers-Sidani E (2018) A Brain without Brakes: Reduced Inhibition Is Associated with Enhanced but Dysregulated Plasticity in the Aged Rat Auditory Cortex. *eNeuro* 5:e0051-0018.2018.
- Clinard CG, Tremblay KL (2013) Aging Degrades the Neural Encoding of Simple and Complex Sounds in the Human Brainstem. *Journal of the American Academy of Audiology* 24:590-599.
- Clinard CG, Tremblay KL, Krishnan AR (2010) Aging alters the perception and physiological representation of frequency: Evidence from human frequency-following response recordings. *Hearing Research* 264:48-55.
- Coomber B, Berger JI, Kowalkowski VL, Shackleton TM, Palmer AR, Wallace MN (2014) Neural changes accompanying tinnitus following unilateral acoustic trauma in the guinea pig. *European Journal of Neuroscience* 40:2427-2441.
- Cruikshanks KJ, Wiley TL, Tweed TS, Klein BEK, Klein R, Mares-Perlman JA, Nondahl DM (1998) Prevalence of Hearing Loss in Older Adults in Beaver Dam, Wisconsin. *American Journal of Epidemiology* 148:879-886.
- Da Costa S, van der Zwaag W, Marques JP, Frackowiak RSJ, Clarke S, Saenz M (2011) Human Primary Auditory Cortex Follows the Shape of Heschl's Gyrus. *The Journal of Neuroscience* 31:14067-14075.
- Day RO, Graham GG, Bieri D, Brown M, Cairns D, Harris G, Hounsell J, Platt-Hepworth S, Reeve R, Sambrook PN, Smith J (1989) Concentration-response relationships for salicylate-induced ototoxicity in normal volunteers. *British Journal of Clinical Pharmacology* 28:695-702.
- de Villers-Sidani E, Alzghoul L, Zhou X, Simpson KL, Lin RCS, Merzenich MM (2010) Recovery of functional and structural age-related changes in the rat primary auditory cortex with operant training. *Proceedings of the National Academy of Sciences* 107:13900-13905.
- Dobri SGJ, Ross B (2021) Total GABA level in human auditory cortex is associated with speech-in-noise understanding in older age. *NeuroImage* 225:117474.
- Drullman R (1995) Temporal envelope and fine structure cues for speech intelligibility. *The Journal of the Acoustical Society of America* 97:585-592.

- Dubno JR, Eckert MA, Lee F-S, Matthews LJ, Schmiedt RA (2013) Classifying Human Audiometric Phenotypes of Age-Related Hearing Loss from Animal Models. *Journal of the Association for Research in Otolaryngology* 14:687-701.
- Dziorny AC, Scott LL, Luebke AE, Walton JP (2020) Rescuing auditory temporal processing with a novel augmented acoustic environment in a mouse model of congenital SNHL. *BioRxiv*.
- Edwards E, Chang EF (2013) Syllabic (~2-5 Hz) and fluctuation (~1-10 Hz) ranges in speech and auditory processing. *Hearing Research* 305:113-134.
- Eggermont JJ (2012) The neural synchrony model of tinnitus. In: *The Neuroscience of Tinnitus*.
- Eggermont JJ (2015) Animal models of spontaneous activity in the healthy and impaired auditory system. *Frontiers in Neural Circuits* 9:19.
- Eggermont JJ, Roberts LE (2004) The neuroscience of tinnitus. *Trends in Neurosciences* 27:676-682.
- Eggermont JJ, Tass PA (2015) Maladaptive neural synchrony in tinnitus: origin and restoration. *Frontiers in neurology* 6:29-29.
- Ernst SM, Moore BCJ (2012) The role of time and place cues in the detection of frequency modulation by hearing impaired listeners. *The Journal of the Acoustical Society of America* 131:4722-4731.
- Feder K, Michaud D, Ramage-Morin P, McNamee J, Beaugard Y (2015) Prevalence of hearing loss among Canadians aged 20 to 79: Audiometric results from the 2012/2013 Canadian Health Measures Survey. *Health Reports* 26:18-25.
- Fitzgibbons PJ, Gordon-Salant S (1996) Auditory temporal processing in elderly listeners. *J Am Acad Audiol* 7:183-189.
- Fitzgibbons PJ, Gordon-Salant S (2010) Behavioral Studies With Aging Humans: Hearing Sensitivity and Psychoacoustics. In: *The Aging Auditory System* (Gordon-Salant S, Frisina RD, Popper AN, Fay RR, eds), pp 111-134. New York, USA: Springer-Verlag.
- Fournier P, Schönwiesner M, Hébert S (2014) Loudness modulation after transient and permanent hearing loss: implications for tinnitus and hyperacusis. *Neuroscience* 283:64-77.
- Frisina DR, Frisina RD, Snell KB, Burkard R, Walton JP, Ison JR (2001) Auditory Temporal Processing during Aging. In: *Functional Neurobiology of Aging* (Hof PR, Mobbs CV, eds), pp 565-579. San Diego: Academic Press.
- Fromme RC, Martins ARO (2011) Spectrotemporal dynamics of auditory cortical synaptic receptive field plasticity. *Hearing Research* 279:149-161.
- Fromby C, Sherlock LP, Gold SL, Hawley ML (2007) Adaptive Recalibration of Chronic Auditory Gain. *Seminars in Hearing* 28:295-302.
- Fukushima N, White P, Harrison RV (1990) Influence of acoustic deprivation on recovery of hair cells after acoustic trauma. *Hearing Research* 50:107-118.

- Füllgrabe C, Meyer B, Lorenzi C (2003) Effect of cochlear damage on the detection of complex temporal envelopes. *Hearing Research* 178:35-43.
- Gao F, Wang G, Ma W, Ren F, Li M, Dong Y, Liu C, Liu B, Bai X, Zhao B, Edden RAE (2015) Decreased auditory GABA+ concentrations in presbycusis demonstrated by edited magnetic resonance spectroscopy. *NeuroImage* 106:311-316.
- Gates GA, Mills JH (2005) Presbycusis. *Lancet* 366:1111–1120.
- Gerken GM (1979) Central denervation hypersensitivity in the auditory system of the cat. *The Journal of the Acoustical Society of America* 66:721-727.
- Giraud A-L, Poeppel D (2012) Cortical oscillations and speech processing: emerging computational principles and operations. *Nature Neuroscience* 15.
- Glasberg BR, Moore BC, Bacon SP (1987) Gap detection and masking in hearing-impaired and normal-hearing subjects. *J Acoust Soc Am* 81:1546-1556.
- Goman AM, Lin FR (2016) Prevalence of Hearing Loss by Severity in the United States. *American Journal of Public Health* 106:1820-1822.
- Goossens T, Vercammen C, Wouters J, van Wieringen A (2016) Aging Affects Neural Synchronization to Speech-Related Acoustic Modulations. *Frontiers in Aging Neuroscience* 8:Article 133.
- Goossens T, Vercammen C, Wouters J, Van Wieringen A (2018) Neural envelope encoding predicts speech perception performance for normal-hearing and hearing-impaired adults. *Hearing Research* 370:189-200.
- Goossens T, Vercammen C, Wouters J, Van Wieringen A (2019) The association between hearing impairment and neural envelope encoding at different ages. *Neurobiology of Aging* 74:202-212.
- Gordon-Salant S, Fitzgibbons PJ (1993) Temporal Factors and Speech Recognition Performance in Young and Elderly Listeners. *Journal of Speech, Language, and Hearing Research* 36:1276-1285.
- Gordon-Salant S, Fitzgibbons PJ (1999) Profile of auditory temporal processing in older listeners. *Journal of speech, language, and hearing research : JSLHR* 42:300-311.
- Gratton MA, Vázquez AE (2003) Age-related hearing loss: current research. *Current Opinion in Otolaryngology & Head and Neck Surgery* 11:367–371.
- Hackett TA (2011) Information flow in the auditory cortical network. *Hearing Research* 271:133-146.
- Harkrider AW, Plyler PN, Hedrick MS (2005) Effects of age and spectral shaping on perception and neural representation of stop consonant stimuli. *Clinical Neurophysiology* 116:2153-2164.

- Harris KC, Wilson S, Eckert MA, Dubno JR (2012) Human evoked cortical activity to silent gaps in noise: effects of age, attention, and cortical processing speed. *Ear Hear* 33:330-339.
- Hattori R, Kuchibhotla KV, Froemke RC, Komiyama T (2017) Functions and dysfunctions of neocortical inhibitory neuron subtypes. *Nature Neuroscience* 20:1199-1208.
- Hayes SH, Radziwon KE, Stolzberg DJ, Salvi RJ (2014) Behavioral models of tinnitus and hyperacusis in animals. *Frontiers in Neurology* 5:179.
- Hayes SH, Schormans AL, Sigela G, Beha K, Herrmann B, Allman BL (2021) Uncovering the contribution of enhanced central gain and altered cortical oscillations to tinnitus generation. *Progress in Neurobiology* 196:101893.
- Hebert S, Fournier P, Noreña A (2013) The Auditory Sensitivity is Increased in Tinnitus Ears. *The Journal of Neuroscience* 33:2356-2364.
- Helfer KS, Merchant GR, Wasiuk PA (2017) Age-Related Changes in Objective and Subjective Speech Perception in Complex Listening Environments. *Journal of Speech, Language, and Hearing Research* 60:3009-3018.
- Henry JA, Schechter MA, Zaugg TL, Griest S, Jastreboff PJ, Vernon JA, Kaelin C, Meikle MB, Lyons KS, Stewart BJ (2006) Clinical Trial to Compare Tinnitus Masking and Tinnitus Retraining Therapy. *Acta Oto-Laryngologica Supplementum* 556:64-69.
- Henry MJ, Herrmann B, Obleser J (2014) Entrained neural oscillations in multiple frequency bands co-modulate behavior. *Proceedings of the National Academy of Sciences* 111:14935-14940.
- Henry MJ, Herrmann B, Kunke D, Obleser J (2017) Aging affects the balance of neural entrainment and top-down neural modulation in the listening brain. *Nature Communications* 8:15801.
- Herrmann B, Parthasarathy A, Bartlett EL (2017) Aging affects dual encoding of periodicity and envelope shape in rat inferior colliculus neurons. *European Journal of Neuroscience* 45:299-311.
- Herrmann B, Maess B, Johnsrude IS (2018) Aging Affects Adaptation to Sound-Level Statistics in Human Auditory Cortex. *The Journal of Neuroscience* 38:1989-1999.
- Herrmann B, Buckland C, Johnsrude IS (2019) Neural signatures of temporal regularity processing in sounds differ between younger and older adults. *Neurobiology of Aging* 83:73-85.
- Herrmann B, Henry MJ, Scharinger M, Obleser J (2013a) Auditory filter width affects response magnitude but not frequency specificity in auditory cortex. *Hearing Research* 304:128-136.
- Herrmann B, Henry MJ, Grigutsch M, Obleser J (2013b) Oscillatory Phase Dynamics in Neural Entrainment Underpin Illusory Percepts of Time. *The Journal of Neuroscience* 33:15799-15809.
- Herrmann B, Henry MJ, Johnsrude IS, Obleser J (2016) Altered temporal dynamics of neural adaptation in the aging human auditory cortex. *Neurobiology of Aging* 45:10-22.

- Herrmann B, Maess B, Hahne A, Schröger E, Friederici AD (2011) Syntactic and auditory spatial processing in the human temporal cortex: An MEG study. *NeuroImage* 57:624-633.
- Holmes E, Griffiths TD (2019) 'Normal' hearing thresholds and fundamental auditory grouping processes predict difficulties with speech-in-noise perception. *Scientific Reports* 9:16771.
- Hopkins J (1994) Orchestrating an Indoor City: Ambient Noise Inside a Mega-Mall. *Environment and Behavior* 26:785-812.
- Hughes LF, Turner JG, Parrish JL, Caspary DM (2010) Processing of broadband stimuli across A1 layers in young and aged rats. *Hearing Research* 264:79-85.
- Humes LE, Busey TA, Craig JC, Kewley-Port D (2009) The effects of age on sensory thresholds and temporal gap detection in hearing, vision, and touch. *Attention, Perception, & Psychophysics* 71:860-871.
- Huotilainen M, Winkler I, Alho K, Escera C, Virtanen J, Ilmoniemi RJ, Jääskeläinen IP, Pekkonen E, Näätänen R (1998) Combined mapping of human auditory EEG and MEG responses. *Electroencephalography and clinical Neurophysiology* 108:370-379.
- Ibrahim BA, Llano DA (2019) Aging and Central Auditory Disinhibition: Is It a Reflection of Homeostatic Downregulation or Metabolic Vulnerability? *Brain Sciences* 9:351.
- Imam L, Hannan SA (2017) Noise-induced hearing loss: a modern epidemic? *British Journal of Hospital Medicine* 78:286-290.
- Isaacson JS, Scanziani M (2011) How Inhibition Shapes Cortical Activity. *Neuron* 72:231-243.
- Ivansic D, Guntinas-Lichius O, Müller B, Volk GF, Schneider G, Dobel C (2017) Impairments of Speech Comprehension in Patients with Tinnitus-A Review. *Front Aging Neurosci* 9:224.
- Izquierdo MA, Gutiérrez-Conde PM, Merchán MA, Malmierca MS (2008) Non-plastic reorganization of frequency coding in the inferior colliculus of the rat following noise-induced hearing loss. *Neuroscience* 154:355-369.
- Jastreboff MM (2007) Sound therapies for tinnitus management. *Progress in Brain Research* 166:435-440.
- Juarez-Salinas DL, Engle JR, Navarro XO, Recanzone GH (2010) Hierarchical and Serial Processing in the Spatial Auditory Cortical Pathway Is Degraded by Natural Aging. *The Journal of Neuroscience* 30:14795-14804.
- Kalappa BI, Brozoski TJ, Turner JG, Caspary DM (2014) Single unit hyperactivity and bursting in the auditory thalamus of awake rats directly correlates with behavioural evidence of tinnitus. *The Journal of Physiology* 592:5065-5078.
- Kaltenbach JA, Rachel JD, Mathog TA, Zhang J, Falzarano PR, Lewandowski M (2002) Cisplatin-induced Hyperactivity in the Dorsal Cochlear Nucleus and Its Relation to Outer Hair Cell Loss: Relevance to Tinnitus. *Journal of Neurophysiology* 88:699-714.

- Kamal B, Holman C, de Villers-Sidani E (2013) Shaping the aging brain: role of auditory input patterns in the emergence of auditory cortical impairments. *Frontiers in Systems Neuroscience* 7:Article 52.
- Kandler K, Clause A, Noh J (2009) Tonotopic reorganization of developing auditory brainstem circuits. *Nature Neuroscience* 12:711-717.
- Kato HK, Asinof SK, Isaacson JS (2017) Network-Level Control of Frequency Tuning in Auditory Cortex. *Neuron* 95:412-423.
- Kaur S, Lazar R, Metherate R (2004) Intracortical Pathways Determine Breadth of Subthreshold Frequency Receptive Fields in Primary Auditory Cortex. *Journal of Neurophysiology* 91:2551-2567.
- Keck T, Toyozumi T, Chen L, Doiron B, Feldman DE, Fox K, Gerstner W, Haydon PG, Hübener M, Lee H-K, Lisman JE, Rose T, Sengpiel F, Stellwagen D, Stryker MP, Turrigiano GG, van Rossum MC (2017) Integrating Hebbian and homeostatic plasticity: the current state of the field and future research directions. *Philosophical Transactions of the Royal Society B* 372:20160158.
- Keithley EM (2020) Pathology and mechanisms of cochlear aging. *Journal of Neuroscience Research* 98:1674-1684.
- Kerlin JR, Shahin AJ, Miller LM (2010) Attentional Gain Control of Ongoing Cortical Speech Representations in a “Cocktail Party”. *The Journal of Neuroscience* 30:620-628.
- King AJ, Bajo VM, Bizley JK, Campbell RAA, Nodal FR, Schulz AL, Schnupp JWH (2007) Physiological and behavioral studies of spatial coding in the auditory cortex. *Hearing Research* 229:106-115.
- Knipper M, Van Dijk P, Nunes I, Rüttiger L, Zimmermann U (2013) Advances in the neurobiology of hearing disorders: Recent developments regarding the basis of tinnitus and hyperacusis. *Progress in Neurobiology* 111:17-33.
- Knipper M, van Dijk P, Schulze H, Mazurek B, Krauss P, Scheper V, Warnecke A, Schlee W, Schwabe K, Singer W, Braun C, Delano PH, Fallgatter AJ, Ehlis A-C, Searchfield GD, Munk MHJ, Baguley DM, Rüttiger L (2020) The Neural Bases of Tinnitus: Lessons from Deafness and Cochlear Implants. *The Journal of Neuroscience* 40:7190-7202.
- Koehnke J, Besing JM (2001) The Effects of Aging on Binaural and Spatial Hearing. *Seminars in Hearing* 22:241-254.
- Koops EA, Renken RJ, Lanting CP, van Dijk P (2020) Cortical Tonotopic Map Changes in Humans Are Larger in Hearing Loss Than in Additional Tinnitus. *The Journal of Neuroscience* 40:3178-3185.
- Kotak VC, Fujisawa S, Lee FA, Karthikeyan O, Aoki C, Sanes DH (2005) Hearing Loss Raises Excitability in the Auditory Cortex. *The Journal of Neuroscience* 25:3908–3918.

- Kujawa SG, Liberman MC (2009) Adding Insult to Injury: Cochlear Nerve Degeneration after “Temporary” Noise-Induced Hearing Loss. *The Journal of Neuroscience* 29:14077-14085.
- Laffont F, Bruneau N, Roux S, Agar N, Minz M, Cathala HP (1989) Effects of age on auditory evoked responses (AER) and augmenting-reducing. *Clinical Neurophysiology* 19:15-23.
- Leong UC, Barsz K, Allen PD, Walton JP (2011) Neural correlates of age-related declines in frequency selectivity in the auditory midbrain. *Neurobiology of Aging* 32:168-178.
- Li S, Choi V, Tzounopoulos T (2013) Pathogenic plasticity of Kv7.2/3 channel activity is essential for the induction of tinnitus. *Proceedings of the National Academy of Sciences of the United States of America* 110:9980-9985.
- Li S, Kalappa BI, Tzounopoulos T (2015) Noise-induced plasticity of KCNQ2/3 and HCN channels underlies vulnerability and resilience to tinnitus. *Elife* 4.
- Liberman MC, Kujawa SG (2017) Cochlear synaptopathy in acquired sensorineural hearing loss: Manifestations and mechanisms. *Hearing Research* 349:138-147.
- Liberman MC, Epstein MJ, Cleveland SS, Wang H, Maison SF (2016) Toward a Differential Diagnosis of Hidden Hearing Loss in Humans. *PLoS ONE* 11:e0162726.
- Lister JJ, Maxfield ND, Pitt GJ, Gonzalez VB (2011) Auditory evoked response to gaps in noise: older adults. *International Journal of Audiology* 50:211-225.
- Llano DA, Turner J, Caspary DM (2012) Diminished Cortical Inhibition in an Aging Mouse Model of Chronic Tinnitus. *The Journal of Neuroscience* 32:16141-16148.
- Lobarinas E, Salvi R, Ding D (2013) Insensitivity of the audiogram to carboplatin induced inner hair cell loss in chinchillas. *Hearing Research* 302:113-120.
- Lorenzi C, Gilbert G, Héloïse C, Garnier S, Moore BCJ (2006) Speech perception problems of the hearing impaired reflect inability to use temporal fine structure. *Proceedings of the National Academy of Sciences* 103:18866-18869.
- Maess B, Jacobsen T, Schröger E, Friederici AD (2007) Localizing pre-attentive auditory memory-based comparison: Magnetic mismatch negativity to pitch change. *NeuroImage* 37:561-571.
- Manunta Y, Edeline J-M (1997) Effects of Noradrenaline on Frequency Tuning of Rat Auditory Cortex Neurons. *European Journal of Neuroscience* 9:833-847.
- Manzoor NF, Licari FG, Klapchar M, Elkin RL, Gao Y, Chen G, Kaltenbach JA (2012) Noise-induced hyperactivity in the inferior colliculus: its relationship with hyperactivity in the dorsal cochlear nucleus. *Journal of Neurophysiology* 108:976-988.
- Martin del Campo HN, Measor KR, Razak KA (2012) Parvalbumin immunoreactivity in the auditory cortex of a mouse model of presbycusis. *Hearing Research* 294:31-39.
- McCormack A, Edmondson-Jones M, Somerset S, Hall DA (2016) A Systematic Review of the Reporting of Tinnitus Prevalence and Severity. *Hearing Research* 337:70-79.

- Mepani AM, Kirk SA, Hancock KE, Bennett K, de Gruttola V, Liberman MC, Maison SF (2020) Middle Ear Muscle Reflex and Word Recognition in "Normal-Hearing" Adults: Evidence for Cochlear Synaptopathy? . *Ear & Hearing* 41:25-38.
- Miller LM, Escabí MA, Read HL, Schreiner CE (2002) Spectrotemporal receptive fields in the lemniscal auditory thalamus and cortex. *Journal of Neurophysiology* 87:516-527.
- Millman RE, Mattys SL, Gouws AD, Prendergast G (2017) Magnified Neural Envelope Coding Predicts Deficits in Speech Perception in Noise. *The Journal of Neuroscience* 37:7727-7736.
- Mishra J, de Villers-Sidani E, Merzenich MM, Gazzaley A (2014) Adaptive Training Diminishes Distractibility in Aging across Species. *Neuron* 84:1091-1103.
- Moerel M, De Martino F, Formisano E (2014) An anatomical and functional topography of human auditory cortical areas. *Frontiers in Neuroscience* 8:Article 225.
- Möhrle D, Ni K, Varakina K, Bing D, Lee SC, Zimmermann U, Knipper M, Rüttiger L (2016) Loss of auditory sensitivity from inner hair cell synaptopathy can be centrally compensated in the young but not old brain. *Neurobiology of Aging* 44:173-184.
- Moore BC, Glasberg BR (1988) Gap detection with sinusoids and noise in normal, impaired, and electrically stimulated ears. *J Acoust Soc Am* 83:1093-1101.
- Moore BCJ (2007) *Cochlear Hearing Loss: Physiological, Psychological and Technical Issues*. West Sussex, England: John Wiley & Sons, Ltd.
- Moore BCJ (2014) *Auditory Processing of Temporal Fine Structure: Effects of Age and Hearing Loss*. Singapore: World Scientific.
- Moore BCJ, Skrodzka E (2002) Detection of frequency modulation by hearing-impaired listeners: Effects of carrier frequency, modulation rate, and added amplitude modulation. *The Journal of the Acoustical Society of America* 111:327-335.
- Moore BCJ, Peters RW, Glasberg BR (1992) Detection of temporal gaps in sinusoids by elderly subjects with and without hearing loss. *The Journal of the Acoustical Society of America* 92:1923-1932.
- Moore BCJ, Wojtczak M, Vickers D (1996) Effect of loudness recruitment on the perception of amplitude modulation. *Journal of the Acoustical Society of America* 100:481-489.
- Moore BCJ, Mariathasan S, Şek AP (2019) Effects of Age and Hearing Loss on the Discrimination of Amplitude and Frequency Modulation for 2- and 10-Hz Rates. *Trends Hear* 23:2331216519853963.
- Moore DR (1987) Physiology of higher auditory system. *British Medical Bulletin* 43:856–870.
- Moser T, Starr A (2016) Auditory neuropathy — neural and synaptic mechanisms. *Nature Reviews Neurology* 12:135-149.
- Mühlnickel W, Elbert T, Taub E, Flor H (1998) Reorganization of auditory cortex in tinnitus. *Proceedings of the National Academy of Sciences* 95:10340-10343.

- Mulders WHAM, Robertson D (2013) Development of hyperactivity after acoustic trauma in the guinea pig inferior colliculus. *Hearing Research* 298:104-108.
- Munguia R, Pienkowski M, Eggermont JJ (2013) Spontaneous firing rate changes in cat primary auditory cortex following long-term exposure to non-traumatic noise: Tinnitus without hearing loss? *Neuroscience Letters* 546:46-50.
- Munro KJ, Turtle C, Schaette R (2014) Plasticity and modified loudness following short-term unilateral deprivation: Evidence of multiple gain mechanisms within the auditory system. *The Journal of the Acoustical Society of America* 135:315-322.
- Näätänen R, Picton TW (1987) The N1 wave of the human electric and magnetic response to sound: a review and an analysis of the component structure. *Psychophysiology* 24:375-425.
- Nahmani M, Turrigiano GG (2014) Adult cortical plasticity following injury: Recapitulation of critical period mechanisms? *Neuroscience* 283:4-16.
- Natan RG, Rao W, Geffen MN (2017) Cortical Interneurons Differentially Shape Frequency Tuning following Adaptation. *Cell Reports* 21:878-890.
- Natan RG, Briguglio JJ, Mwilambwe-Tshilobo L, Jones SI, Aizenberg M, Goldberg EM, Geffen MN (2015) Complementary control of sensory adaptation by two types of cortical interneurons. *eLife* 4:e09868.
- Nelson DA, Freyman RL (1986) Psychometric functions for frequency discrimination from listeners with sensorineural hearing loss. *The Journal of the Acoustical Society of America* 79:799-805.
- Ng C-W, Recanzone GH (2018) Age-Related Changes in Temporal Processing of Rapidly-Presented Sound Sequences in the Macaque Auditory Cortex. *Cerebral Cortex* 28:3775-3796.
- Niu X, Tahera Y, Canlon B (2004) Protection against Acoustic Trauma by Forward and Backward Sound Conditioning. *Audiology & Neuro-otology* 9:265-273.
- Niu X, Tahera Y, Canlon B (2007) Environmental enrichment to sound activates dopaminergic pathways in the auditory system. *Physiology & Behavior* 92:34-39.
- Noelle O'Connell M, Barczak A, Schroeder CE, Lakatos P (2014) Layer Specific Sharpening of Frequency Tuning by Selective Attention in Primary Auditory Cortex. *The Journal of Neuroscience* 34:16496-16508.
- Norena AJ, Eggermont JJ (2006) Enriched acoustic environment after noise trauma abolishes neural signs of tinnitus. *NeuroReport* 17:559-563.
- Norena AJ, Chery-Croze S (2007) Enriched acoustic environment rescales auditory sensitivity. *NeuroReport* 18:1251-1255.
- Norena AJ, Tomita M, Eggermont JJ (2003) Neural Changes in Cat Auditory Cortex After a Transient Pure-Tone Trauma. *Journal of Neurophysiology* 90:2387-2401.

- Noreña AJ, Gourévitch B, Aizawa N, Eggermont JJ (2006) Spectrally enhanced acoustic environment disrupts frequency representation in cat auditory cortex. *Nature Neuroscience* 7:932-939.
- Oliver DL, Izquierdo MA, Malmierca MS (2011) Persistent effects of early augmented acoustic environment on the auditory brainstem. *Neuroscience* 184:75-87.
- Olsen WO (1998) Average Speech Levels and Spectra in Various Speaking/Listening Conditions: A Summary of the Pearson, Bennett, & Fidell (1977) Report. *American Journal of Audiology* 7:21-25.
- Ouda L, Profant O, Syka J (2015) Age-related changes in the central auditory system. *Cell and Tissue Research* 361:337-358.
- Ouellet L, de Villers-Sidani E (2014) Trajectory of the main GABAergic interneuron populations from early development to old age in the rat primary auditory cortex. *Frontiers in Neuroanatomy* 8:Article 40.
- Overton JA, Recanzone GH (2016) Effects of aging on the response of single neurons to amplitude-modulated noise in primary auditory cortex of rhesus macaque. *Journal of Neurophysiology* 115:2911-2923.
- Palombi PS, Backoff PM, Caspary DM (2001) Responses of young and aged rat inferior colliculus neurons to sinusoidally amplitude modulated stimuli. *Hearing Research* 153:174-180.
- Parmentier FBR, Andrés P (2010) The Involuntary Capture of Attention by Sound Novelty and Postnovelty Distraction in Young and Older Adults. *Experimental Psychology* 57:68-76.
- Parry LV, Maslin MRD, Schaette R, Moore DR, Munro KJ (2019) Increased auditory cortex neural response amplitude in adults with chronic unilateral conductive hearing impairment. *Hearing Research* 372:10-16.
- Parthasarathy A, Herrmann B, Bartlett EL (2019) Aging alters envelope representations of speech-like sounds in the inferior colliculus. *Neurobiology of Aging* 73:30-40.
- Parthasarathy A, Hancock KE, Bennett K, DeGruttola V, Polley DB (2020) Bottom-up and top-down neural signatures of disordered multi-talker speech perception in adults with normal hearing. *eLife* 9:e51419.
- Peelle JE, Davis MH (2013) Neural oscillations carry speech rhythm through to comprehension. *Frontiers in Psychology* 3:Article 320.
- Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M (2013) Inhibition of Inhibition in Visual Cortex: The Logic of Connections Between Molecularly Distinct Interneurons *Nature Neuroscience* 16:1068-1076.
- Pichora-Fuller MK (2003) Processing speed and timing in aging adults: psychoacoustics, speech perception, and comprehension. *International Journal of Audiology* 42:S59-S67.

- Pichora-Fuller MK, Kramer SE, Eckert MA, Edwards B, Hornsby BWY, Humes LE, Lemke U, Lunner T, Matthen M, Mackersie CL, Naylor G, Phillips NA, Richter M, Rudner M, Sommers MS, Tremblay KL, Wingfield A (2016) Hearing Impairment and Cognitive Energy: The Framework for Understanding Effortful Listening (FUEL). *Ear & Hearing* 37 Suppl 1:5S-27S.
- Picton TW, John SM, Dimitrijevic A, Purcell DW (2003) Human auditory steady-state responses. *International Journal of Audiology* 42:177-219.
- Plack CJ (2014) *The sense of hearing*. New York, USA: Psychology Press.
- Plack CJ, Barker D, Prendergast G (2014) Perceptual Consequences of “Hidden” Hearing Loss. *Trends in Hearing* 18:1-11.
- Popelár J, Syka J, Berndt H (1987) Effect of noise on auditory evoked responses in awake guinea pigs. *Hearing Research* 26:239-247.
- Prendergast G, Guest H, Munro KJ, Kluk K, Leger A, Hall AD, Heinz MG, Plack CJ (2017a) Effects of noise exposure on young adults with normal audiograms I: Electrophysiology. *Hearing Research* 344:68-81.
- Prendergast G, Millman RE, Guest H, Munro KJ, Kluk K, Dewey RS, Hall DA, Heinz MG, Plack CJ (2017b) Effects of noise exposure on young adults with normal audiograms II: Behavioral measures. *Hearing Research* 356:74-86.
- Presacco A, Simon JZ, Anderson S (2016a) Evidence of degraded representation of speech in noise, in the aging midbrain and cortex. *Journal of Neurophysiology* 116:2346-2355.
- Presacco A, Simon JZ, Anderson S (2016b) Effect of informational content of noise on speech representation in the aging midbrain and cortex. *Journal of Neurophysiology* 116:2356-2367.
- Presacco A, Simon JZ, Anderson S (2019) Speech-in-noise representation in the aging midbrain and cortex: Effects of hearing loss. *PLoS ONE* 14:e0213899.
- Ptito M, Kupers R, Lomber S, Pietrini P (2012) Sensory Deprivation and Brain Plasticity. *Neural Plasticity* 2012:Article ID 810370.
- Purcell DW, John SM, Schneider BA, Picton TW (2004) Human temporal auditory acuity as assessed by envelope following responses. *The Journal of the Acoustical Society of America* 116:3581-3593.
- Qiu C, Salvi R, Ding D, Burkard R (2000) Inner hair cell loss leads to enhanced response amplitudes in auditory cortex of unanesthetized chinchillas: evidence for increased system gain. *Hearing research* 139:153-171.
- Rabang CF, Parthasarathy A, Venkataraman Y, Fisher ZL, Gardner SM, Bartlett EL (2012) A computational model of inferior colliculus responses to amplitude modulated sounds in young and aged rats. *Frontiers in Neural Circuits* 6:77.

- Rachel JD, Kaltenbach JA, Janisse J (2002) Increases in Spontaneous Neural Activity in the Hamster Dorsal Cochlear Nucleus Following Cisplatin Treatment: A Possible Basis for Cisplatin-Induced Tinnitus Hearing Research 164:206-214.
- Radziwon K, Hayes SH, Sheppard AM, Ding D, Salvi R (2016) Drug-Induced Tinnitus. In: Tinnitus: Clinical and research perspectives (Baguley DM, Fagelson M, eds), pp 89-109: Plural Publishing.
- Radziwon KE, Stolzberg DJ, Urban ME, Bowler RA, Salvi RJ (2015) Salicylate-Induced Hearing Loss and Gap Detection Deficits in Rats. *Frontiers in Neurology* 6.
- Radziwon KE, Holfoth D, Lindner J, Kaier-Green Z, Bowler R, Urban M, Salvi R (2017) Salicylate-induced hyperacusis in rats: Dose- and frequency-dependent effects. *Hearing Research* 350:133-138.
- Ramachandran R, Davis KA, May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats. I. Classification based on frequency response maps. *Journal of Neurophysiology* 82:152-163.
- Rauschecker JP (2011) An expanded role for the dorsal auditory pathway in sensorimotor control and integration. *Hearing Research* 271:16-25.
- Rauschecker JP, Tian B (2000) Mechanisms and streams for processing of “what” and “where” in auditory cortex. *Proceedings of the National Academy of Sciences* 97:11800-11806.
- Rauschecker JP, Scott SK (2009) Maps and streams in the auditory cortex: nonhuman primates illuminate human speech processing. *Nature Neuroscience* 12:718-724.
- Resnik J, Polley DB (2017) Fast-spiking GABA circuit dynamics in the auditory cortex predict recovery of sensory processing following peripheral nerve damage. *eLife* 6:e21452.
- Rogalla MM, Hildebrandt KJ (2020) Aging But Not Age-Related Hearing Loss Dominates the Decrease of Parvalbumin Immunoreactivity in the Primary Auditory Cortex of Mice. *eNeuro* 7:ENEURO.0511-0519.2020.
- Rosen MJ, Sarro EC, Kelly JB, Sanes DH (2012) Diminished behavioral and neural sensitivity to sound modulation is associated with moderate developmental hearing loss. *PLoS One* 7:e41514.
- Rosen S (1992) Temporal Information in Speech: Acoustic, Auditory and Linguistic Aspects. *Philosophical Transactions: Biological Sciences* 336:367-373.
- Ross B, Tremblay KL (2009) Stimulus experience modifies auditory neuromagnetic responses in young and older listeners. *Hearing Research* 248:48-59.
- Ross B, Schneider B, Snyder JS, Alain C (2010) Biological Markers of Auditory Gap Detection in Young, Middle-Aged, and Older Adults. *PLOS ONE* 5:e10101.
- Saenz M, Langers DRM (2014) Tonotopic mapping of human auditory cortex. *Hearing Research* 307:42-52.

- Salthouse TA (1995) Refining the concept of psychological compensation. In: *Compensating for psychological deficits and declines: Managing losses and promoting gains* (Dixon RAaB, L., ed), pp 21-34: Lawrence Erlbaum Associates, Inc. .
- Salvi R, Sun W, Ding D, Chen G-D, Lobarinas E, Wang J, Radziwon K, Auerbach BD (2017) Inner Hair Cell Loss Disrupts Hearing and Cochlear Function Leading to Sensory Deprivation and Enhanced Central Auditory Gain. *Frontiers in Neuroscience* 10:Article 621.
- Salvi RJ, Auerbach BD, Lau C, Chen Y-C, Manohar S, Liu X, Ding D, Chen G-D (2020) Functional Neuroanatomy of Salicylate- and Noise-Induced Tinnitus and Hyperacusis. In: *Current Topics in Behavioral Neurosciences* (Geyer MA, Ellenbroek BA, Marsden CA, Barnes TRE, Andersen SL, Paulus MP, eds). Berlin, Heidelberg: Springer.
- Sanes DH, Kotak VC (2011) Developmental plasticity of auditory cortical inhibitory synapses. *Hearing Research* 279:140-148.
- Sanes DH, Sarro EC, Takesian AE, Aoki C, Kotak VC (2010) Regulation of Inhibitory Synapse Function in the Developing Auditory CNS. In: *Developmental Plasticity of Inhibitory Circuitry* (Pallas SL, ed), pp 43-69. Boston, MA: Springer US.
- Saunders JC, Dear SP, Schneider ME (1985) The anatomical consequences of acoustic injury: A review and tutorial. *The Journal of the Acoustical Society of America* 78:833-860.
- Schaette R, McAlpine D (2011) Tinnitus with a Normal Audiogram: Physiological Evidence for Hidden Hearing Loss and Computational Model. *The Journal of Neuroscience* 31:13452-13457.
- Schaette R, Turtle C, Munro KJ (2012) Reversible induction of phantom auditory sensations through simulated unilateral hearing loss. *PLoS ONE* 7:e35238.
- Schatteman TA, Hughes LF, Caspary DM (2008) Age-related loss of temporal processing: Altered responses to amplitude modulated tones in rat dorsal cochlear nucleus. *Neuroscience* 154:329-337.
- Schlittenlacher J, Moore BCJ (2016) Discrimination of amplitude-modulation depth by subjects with normal and impaired hearing. *Journal of the Acoustical Society of America* 140:3487-3495.
- Schmiedt RA (2010) The Physiology of Cochlear Presbycusis. In: *The Aging Auditory System*. Springer Handbook of Auditory Research (Gordon-Salant S, Frisina R, Popper A, Fay R, eds), pp 9-38. New York, USA: Springer.
- Schneider B, Speranza F, Pichora-Fuller MK (1998) Age-related changes in temporal resolution: envelope and intensity effects. *Canadian journal of experimental psychology = Revue canadienne de psychologie experimentale* 52:184-191.
- Schneider BA, Pichora-Fuller MK, Kowalchuk D, Lamb M (1994) Gap detection and the precedence effect in young and old adults. *J Acoust Soc Am* 95:980-991.

- Schormans AL, Typlt M, Allman B (2019) Adult-Onset Hearing Impairment Induces Layer-Specific Cortical Reorganization: Evidence of Crossmodal Plasticity and Central Gain Enhancement. *Cerebral Cortex* 29:1875-1888.
- Schreiner CE, Froemke RC, Atencio CA (2011) Spectral Processing in Auditory Cortex. In: *The Auditory Cortex* (Winer JA, Schreiner CE, eds), pp 275-308. USA: Springer.
- Sedley W (2019) Tinnitus: Does Gain Explain? . *Neuroscience* 407:213-228.
- Sedley W, Friston KJ, Gander PE, Kumar S, Griffiths TD (2016) An Integrative Tinnitus Model Based on Sensory Precision. *Trends in Neurosciences* 39.
- Sek A, Baer T, Crinnion W, Springgay A, Moore BCJ (2015) Modulation masking within and across carriers for subjects with normal and impaired hearing. *The Journal of the Acoustical Society of America* 138:1143-1153.
- Seki S, Eggermont JJ (2003) Changes in spontaneous firing rate and neural synchrony in cat primary auditory cortex after localized tone-induced hearing loss. *Hearing Research* 180:28-38.
- Sekiya K, Takahashi M, Murakami S, Kakigi R, Okamoto H (2017) Broadened population-level frequency tuning in the auditory cortex of tinnitus patients. *Journal of Neurophysiology* 117:1379-1384.
- Shaheen LA, Liberman MC (2018) Cochlear Synaptopathy Changes Sound-Evoked Activity Without Changing Spontaneous Discharge in the Mouse Inferior Colliculus *Frontiers in Systems Neuroscience* 12:Article 59.
- Shamma S, Lorenzi C (2013) On the balance of envelope and temporal fine structure in the encoding of speech in the early auditory system. *The Journal of the Acoustical Society of America* 133:2818-2833.
- Sheppard A, Stocking C, Ralli M, Salvi RJ (2020) A review of auditory gain, low-level noise and sound therapy for tinnitus and hyperacusis. *International Journal of Audiology* 59:5-15.
- Sheppard AM, Liu X, Ding D, Salvi RJ (2018) Auditory central gain compensates for changes in cochlear output after prolonged low-level noise exposure. *Neuroscience Letters* 687:183-188.
- Shew WL, Plenz D (2013) The Functional Benefits of Criticality in the Cortex. *The Neuroscientist* 19:88-100.
- Silver RA (2010) Neuronal arithmetic. *Nature Reviews Neuroscience* 11:474-489.
- Slepecky N (1986) Overview of mechanical damage to the inner ear: noise as a tool to probe cochlear function. *Hearing Research* 22:307-321.
- Snell KB (1997) Age-related changes in temporal gap detection. *J Acoust Soc Am* 101:2214-2220.
- Snell KB, Frisina DR (2000) Relationships among age-related differences in gap detection and word recognition. *J Acoust Soc Am* 107:1615-1626.

- Sörös P, Treisman IK, Manemann E, Lütkenhöner B (2009) Auditory temporal processing in healthy aging: a magnetoencephalographic study. *BMC Neuroscience* 10:34.
- Stolzberg D, Chrostowski M, Salvi RJ, Allman B (2012) Intracortical circuits amplify sound-evoked activity in primary auditory cortex following systemic injection of salicylate in the rat. *Journal of Neurophysiology* 108:200-214.
- Stolzberg D, Hayes SH, Kashanian N, Radziwon K, Salvi RJ, Allman BL (2013) A Novel Behavioral Assay for the Assessment of Acute Tinnitus in Rats Optimized for Simultaneous Recording of Oscillatory Neural Activity. *Journal of Neuroscience Methods* 219:224-232.
- Stothart G, Kazanina N (2016) Auditory perception in the aging brain: the role of inhibition and facilitation in early processing. *Neurobiology of Aging* 47:23-24.
- Sturm JJ, Zhang-Hooks YX, Roos H, Nguyen T, Kandler K (2017) Noise Trauma-Induced Behavioral Gap Detection Deficits Correlate with Reorganization of Excitatory and Inhibitory Local Circuits in the Inferior Colliculus and Are Prevented by Acoustic Enrichment. *The Journal of Neuroscience* 37:6314-6330.
- Sun W, Deng A, Jayaram A, Gibson B (2012) Noise exposure enhances auditory cortex responses related to hyperacusis behavior. *Brain Research* 1485:108-116.
- Sun W, Lu J, Stolzberg D, Gray L, Deng A, Lobarinas E, Salvi RJ (2009) Salicylate Increases the Gain of the Central Auditory System. *Neuroscience* 159:325-334.
- Syka J, Rybalko N, Popelár J (1994) Enhancement of the auditory cortex evoked responses in awake guinea pigs after noise exposure. *Hearing Research* 78:158-168.
- Takesian AE, Kotak VC, Sanes DH (2009) Developmental hearing loss disrupts synaptic inhibition: implications for auditory processing. *Future Neurology* 4:331-349.
- Takesian AE, Kotak VC, Sanes DH (2012) Age-dependent effect of hearing loss on cortical inhibitory synapse function. *Journal of Neurophysiology* 107:937-947.
- Teichert M, Liebmann L, Hübner CA, Bolz J (2017) Homeostatic plasticity and synaptic scaling in the adult mouse auditory cortex. *Scientific Reports* 7:17423.
- Thomas ME, Guercio GD, Drudik KM, de Villiers-Sidani É (2019a) Evidence of Hyperacusis in Adult Rats Following Non-traumatic Sound Exposure. *Frontiers in Systems Neuroscience* 13:Article 55.
- Thomas ME, Friedman NHM, Cisneros-Franco JM, Ouellet L, de Villiers-Sidani E (2019b) The Prolonged Masking of Temporal Acoustic Inputs with Noise Drives Plasticity in the Adult Rat Auditory Cortex. *Cerebral Cortex* 29:1032-1046.
- Tremblay KL, Piskosz M, Souza P (2003) Effects of age and age-related hearing loss on the neural representation of speech cues. *Clinical Neurophysiology* 114:1332-1343.
- Tsai K-T, Lin M-D, Chen Y-H (2009) Noise mapping in urban environments: A Taiwan study. *Applied Acoustics* 70:964-972.

- Turner CW, Nelson DA (1982) Frequency discrimination in regions of normal and impaired sensitivity. *Journal of Speech and Hearing Research* 25:34-41.
- Turner JG, Hughes LF, Caspary DM (2005) Affects of Aging on Receptive Fields in Rat Primary Auditory Cortex Layer V Neurons. *Journal of Neurophysiology* 94:2738-2747.
- Turner JG, Brozoski TJ, Bauer CA, Parrish JL, Myers K, Hughes LF, Caspary DM (2006) Gap Detection Deficits in Rats With Tinnitus: A Potential Novel Screening Tool. *Behavioral Neuroscience* 120:188-195.
- Turrigiano GG (1999) Homeostatic plasticity in neuronal networks: the more things change, the more they stay the same. *Trends in Neurosciences* 22:221-227.
- Turrigiano GG (2012) Homeostatic Synaptic Plasticity: Local and Global Mechanisms for Stabilizing Neuronal Function. *Cold Spring Harbor Perspectives in Biology* 4:a005736.
- Turrigiano GG, Nelson SB (2000) Hebb and homeostasis in neuronal plasticity. *Current Opinion in Neurobiology* 10:358-364.
- Tyler RS, Pienkowski M, Roncancio ER, Jun HJ, Brozoski T, Dauman N, Coelho CB, Andersson G, Keiner AJ, Cacace AT, Martin N, Moore BCJ (2014) A Review of Hyperacusis and Future Directions: Part I. Definitions and Manifestations. *American Journal of Audiology* 23:402-419.
- Walton JP (2010) Timing is everything: Temporal processing deficits in the aged auditory brainstem. *Hearing Research* 264:63-69.
- Walton JP, Frisina RD, O'Neill WE (1998) Age-Related Alteration in Processing of Temporal Sound Features in the Auditory Midbrain of the CBA Mouse. *The Journal of Neuroscience* 18:2764-2776.
- Walton JP, Simon H, Frisina RD (2002) Age-Related Alterations in the Neural Coding of Envelope Periodicities. *Journal of Neurophysiology* 88:565-578.
- Wang F, Zuo L, Hong B, Han D, Range EM, Zhao L, Sui Y, Guo W, Liu L (2013) Tonotopic reorganization and spontaneous firing in inferior colliculus during both short and long recovery periods after noise overexposure. *Journal of Biomedical Science* 20:91.
- Wang J, McFadden SL, Caspary DM, Salvi RJ (2002) Gamma-aminobutyric acid circuits shape response properties of auditory cortex neurons. *Brain Research Interactive* 944:219-231.
- Wehr M, Zador AM (2003) Balanced inhibition underlies tuning and sharpens spike timing in auditory cortex. *Nature* 426:442-446.
- Weisz N, Moratti S, Meinzer M, Dohrmann K, Elbert T (2005) Tinnitus Perception and Distress Is Related to Abnormal Spontaneous Brain Activity as Measured by Magnetoencephalography. *PloS Medicine* 2:e153.
- Whitmire CJ, Stanley GB (2016) Rapid Sensory Adaptation Redux: A Circuit Perspective. *Neuron* 92:298-315.

- Wieczerzak KB, Patel SV, MacNeil H, Scott KE, Schormans AL, Hayes SH, Herrmann B, Allman BL (2021) Differential Plasticity in Auditory and Prefrontal Cortices, and Cognitive-Behavioral Deficits Following Noise-Induced Hearing Loss. *Neuroscience* 455:1-18.
- Willott JF (1986) Effects of aging, hearing loss, and anatomical location on thresholds of inferior colliculus neurons in C57BL/6 and CBA mice. *Journal of Neurophysiology* 56:391-408.
- Willott JF, Parham K, Hunter KP (1988a) Response properties of inferior colliculus neurons in young and very old CBA/J mice. *Hearing Research* 37:1-14.
- Willott JF, Parham K, Hunter KP (1988b) Response properties of inferior colliculus neurons in middle-aged C57BL/6J mice with presbycusis. *Hearing Research* 37:15-27.
- Wolak T, Ciesla K, Lorens A, Kochanek K, Lewandowska M, Rusiniak M, Pluta A, Wojcik J, Skarzynski H (2017) Tonotopic organisation of the auditory cortex in sloping sensorineural hearing loss. *Hearing Research* 355:81-96.
- Wood KC, Blackwell JM, Geffen MN (2017) Cortical inhibitory interneurons control sensory processing. *Current Opinion in Neurobiology* 46:200-207.
- Woods DL, Alain C (2009) Functional imaging of human auditory cortex. *Current Opinion in Otolaryngology & Head and Neck Surgery* 17:407-411.
- Woods TM, Lopez SE, Long JH, Rahman JE, Recanzone GH (2006) Effects of stimulus azimuth and intensity on the single-neuron activity in the auditory cortex of the alert macaque monkey. *Journal of Neurophysiology* 96:3323-3337.
- Wu P-z, O'Malley JT, de Gruttola V, Liberman MC (2020) Age-Related Hearing Loss Is Dominated by Damage to Inner Ear Sensory Cells, Not the Cellular Battery That Powers Them. *The Journal of Neuroscience* 40:6357-6366.
- Yang G, Lobarinas E, Zhang L, Turnerc J, Stolzberg D, Salvi R, Sun W (2007) Salicylate induced tinnitus: Behavioral measures and neuralactivity in auditory cortex of awake rats. *Hearing Research* 226:244-253.
- Yang L, Pollak GD (1997) Differential Response Properties to Amplitude Modulated Signals in the Dorsal Nucleus of the Lateral Lemniscus of the Mustache Bat and the Roles of GABAergic Inhibition. *Journal of Neurophysiology* 77:324-340.
- Yang S, Weiner BD, Zhang LS, Cho S-J, Bao S (2011) Homeostatic plasticity drives tinnitus perception in an animal model. *Proceedings of the National Academy of Sciences* 108:14974-14979.
- Yoshida N, Liberman MC (2000) Sound conditioning reduces noise-induced permanent threshold shift in mice. *Hearing Research* 148:213-219.
- Yuan Y, Shi F, Yin Y, Tong M, Lang H, Polley DB, Liberman MC, Edge AS (2014) Ouabain-induced Cochlear Nerve Degeneration: Synaptic Loss and Plasticity in a Mouse Model of Auditory Neuropathy *Journal of the Association for Research in Otolaryngology* 15:31-43.

- Zeng F-G (2013) An active loudness model suggesting tinnitus as increased central noise and hyperacusis as increased nonlinear gain. *Hearing Research* 295:172-179.
- Zeng FG, Richardson M, Turner K (2020) Tinnitus Does Not Interfere with Auditory and Speech Perception. *The Journal of Neuroscience* 40:6007-6017.
- Zhang JS, Kaltenbach JA (1998) Increases in spontaneous activity in the dorsal cochlear nucleus of the rat following exposure to high-intensity sound. *Neuroscience Letters* 250:197-200.
- Zhao Y, Song Q, Li X, Li C (2016) Neural Hyperactivity of the Central Auditory System in Response to Peripheral Damage. *Neural Plasticity* 2016:2162105.
- Zhou X, Merzenich MM (2012) Environmental noise exposure degrades normal listening processes. *Nature Communications* 3:843.