

Article

# Comparison of three skin sampling methods and two media for culturing *Malassezia* yeast

Abdourahim Abdillah <sup>1,2</sup>, Saber Khelaifia <sup>2</sup>, Didier Raoult <sup>2,3</sup>, Fadi Bittar <sup>2,3</sup>, Stéphane Ranque <sup>1,2,\*</sup>

<sup>1</sup> Aix-Marseille Université, APHM, IRD, VITROME, IHU Méditerranée Infection, Marseille, France; .

[abdourahim15@live.fr](mailto:abdourahim15@live.fr) (A.A.); [stephane.ranque@univ-amu.fr](mailto:stephane.ranque@univ-amu.fr) (S.R.)

<sup>2</sup> IHU-Méditerranée Infection, Marseille, France; [khelaifia\\_saber@yahoo.fr](mailto:khelaifia_saber@yahoo.fr) (S.K.);

[didier.raoult@mediterranee-infection.com](mailto:didier.raoult@mediterranee-infection.com) (D.R.)

<sup>3</sup> Aix-Marseille Université, APHM, MEPHI, IHU Méditerranée Infection, Marseille, France;

[fadi.BITTAR@univ-amu.fr](mailto:fadi.BITTAR@univ-amu.fr) (F.B.)

\* Correspondence: [stephane.ranque@univ-amu.fr](mailto:stephane.ranque@univ-amu.fr)

**Abstract:** *Malassezia* is lipid-dependent commensal yeast of the human skin. The different culture media and skin sampling methods used to grow these fastidious yeasts are a source of heterogeneity in culture-based epidemiological study results. This study aimed to compare the performances of three methods of skin sampling, and two culture media for the detection of *Malassezia* yeasts by culture from the human skin. Three skin sampling methods, namely sterile gauze, dry swab and Transwab™ with transport medium, were applied on 10 healthy volunteers. Each sample was further inoculated onto either the novel FastFung medium or the reference Dixon agar for the detection of *Malassezia* spp. by culture. At least one colony of *Malassezia* spp. grew on 93/300 (31%) of the cultures, corresponding to 150 samplings. The positive culture rate was 67%, 18%, and 15% ( $P < 10^{-3}$ ), for samples collected with sterile gauze, Transwab™, and dry swab, respectively. The positive culture rate was 62% and 38% ( $P < 0.003$ ) by using the FastFung and the Dixon media, respectively. Our results showed that sterile gauze rubbing skin sampling followed by inoculation on FastFung medium should be implemented in the routine clinical laboratory procedure for *Malassezia* spp. cultivation.

**Keywords:** *Malassezia*; isolation; culture media; skin sampling

## 1. Introduction

The *Malassezia* yeasts are basidiomycetes of the Ustilaginomycotina subphylum and Malasseziomycetes class [1]. These yeasts are lipid-dependent because comparative genomic analyses have revealed that all *Malassezia* lack the gene encoding the synthesis of fatty acids [2]. They are part of the normal eukaryotic microbiota of the skin in humans and other warm-blooded animals, and it is estimated that about 75-98% of healthy subjects carry these yeasts [3]. However, under certain circumstances, *Malassezia* species can be involved in relatively common skin diseases, including pityriasis versicolor, seborrheic dermatitis, and folliculitis. They also cause sporadic severe systemic infections in neonates and immunocompromised individuals with parenteral nutrition [4]–[6].

*Malassezia* spp. are not able to grow on the mycological media routinely used in the clinical laboratory because of their lipid-dependency; they necessitate using specific culture media, such as Dixon agar or modified Dixon agar, Leeming-Notmann agar or modified Leeming-Notmann agar, etc. Yet specific media are relatively seldom used and partially explain why *Malassezia* spp. has been detected via metagenomics but not via culture, especially in the human gut [7], [8] or respiratory tract [9], [10]. Moreover, the lack of standardization of skin sampling procedure and culture medium used to grow these fastidious yeasts is a source of heterogeneity in culture-based study results [5], that

should be avoided. Many authors believe that the lack of agreement between findings of culture-based epidemiological studies is mainly due to the varieties of methods used [11].

To overcome these limitations, this study aimed to compare the performances of three methods of skin sampling, namely sterile gauze, dry swab and Transwab™ with transport medium, and two culture media, the FastFung medium and the reference Dixon agar, for the detection of *Malassezia* yeasts by culture from the human skin. Our results showed that sterile gauze is the best technique to recover *Malassezia* yeasts from the skin for culture and that the FastFung medium is more efficient than Dixon agar for isolation and cultivation of *Malassezia* spp. from the skin.

## 2. Methods

### 2.1 Study participants

A total of 10 healthy volunteers (9 males and 1 female, aged from 25-56 years; mean age: 31.7 years; median age: 27.5 years) at the IHU-MI (Institut Hospitalo-Universitaire Méditerranée Infection, Marseille, France) were included. The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to and the appropriate ethical review committee approval has been received. All participants gave their informed consent and completed a form with age, gender whether or not systemic or local antifungal treatment has been taken in the last 15 days after sampling. Participants with a history of antifungal treatment were not included. This study protocol was approved by the CPP Ile de France II (N°19.05.29.69947 RIPH3) the 21/10/2019. Each volunteer included gave an informed written consent to participate in the study.

### 2.2 Culture media

The Dixon agar (Mediaproducs BV, Groningen, Netherlands) containing 0.2 g/L chloramphenicol was used, as the reference medium for *Malassezia* spp. detection by culture. The FastFung medium (4.3% Schædler agar, 2% peptone, 1% glucose, 1% malt extract, 0.5% ox-bile, 0.5% Tween 60, 0.2% oleic acid and 0.25% glycerol [pH6], each from Sigma-Aldrich, Saint-Quentin Fallavier, France), developed at the Institut Hospitalo-Universitaire Méditerranée Infection (IHU-MI), has been derived from the Schædler agar as previously described [12] (Bittar et al. submitted).

### 2.3 Sample collection and cultures

Skin samples were collected at five different body sites (left and right nasolabial folds, left and right retroauricular folds, and presternal skin) using three sampling methods: dry cotton swabs (DCS) (Greiner Bio-One, Courtaboeuf, France), swabs with Amies liquid transport medium (Transwab®, ELITech, Puteaux, France); and sterile gauze (non-woven sterile swabs 10 cm x 10 cm, Laboratoire SYLAMÉD, Paris, France). The three sampling methods were applied successively in a random order, which was noted, at each sampling site for 5 to 10 seconds. A total of 15 samples were collected for each participant. Each sample was then plated in parallel, in a random order that was noted, on Dixon agar (Mediaproducs BV) and FastFung medium supplemented with 0.1 g/L vancomycin (Sandoz, Levallois-Perret, France) and 0.03 g/L colistin (Sigma-Aldrich). The gauze was directly applied onto the plate. Unused dry cotton swabs, Transwab®, and sterile gauze were also plated on both media as negative controls. All plates were incubated aerobically at 30°C for one week and examined daily for *Malassezia* spp. growth.

### 2.4 Colony identification


Colonies of *Malassezia* spp. were identified via MALDI-TOF Mass Spectrometry (MALDI Biotyper, Bruker Daltonics) with a reference spectra library supplemented with in house (including *Malassezia* spp.) reference spectra, following a previously published procedure [13].

### 2.5 Statistical analysis

Statistical analyses were done with the SAS 9.4 software (SAS Institute Inc., Cary, NC, USA). The effect of different sampling and culture conditions on the number of CFUs (Colony Forming Units) of *Malassezia* spp. was tested with a negative binomial regression generalized linear model (Proc. Genmod), using generalized estimating equations to account for the non-independence of the measurements carried out on the same participant. The influence of the variables on the positivity (presence/absence) of the *Malassezia* spp. culture was tested by logistic regression (Proc. Logistics) allowing on the participant effect. Two-sided tests were used; and a *P*-value < 0.05 was considered statistically significant.

### 3. Results

A total of 93/300 (31%) of cultures, corresponding to 150 samplings, were positive for *Malassezia*, i.e. allowed the growth of at least one colony of *Malassezia* spp. We found a statistically significant independent effect ( $P < 10^{-3}$ ) of the sampling methods (Figure 1A); with the highest performance for sterile gauze totaling 67% (62/93) positive cultures compared to 18% (17/93) with Transwab™ and 15% (14/93) with the dry swab. We also found a statistically significant ( $P < 0.003$ ) independent effect of the culture medium, with the highest performance of the FastFung medium with 62% (58/93) of positive cultures compared to 38% (35/93) on Dixon agar medium (Figure 1B). A total of 1 082 colonies of *Malassezia* spp. were isolated and identified by MALDI-TOF Mass Spectrometry, including 455 (42%) *M. globosa*, 424 (40%) *M. sympodialis*, and 203 (18%) *M. restricta*. Noteworthy, the majority of the colonies of *M. globosa* and *M. restricta*, which are species of clinical interest and fastidious culture, were isolated on the FastFung medium (Figure 1C). The isolated species were heterogeneously distributed according to sampling body sites, as illustrated in Figure 1D.



**Figure 1. *Malassezia* species isolation.** Relative contribution to *Malassezia* spp. positive culture of each sampling technique (A) and isolation media (B). (C) Relative contribution (CFU) of the Dixon agar and FastFung medium to the isolation of each detected *Malassezia* species. (D) Relative abundance (CFU) of each detected *Malassezia* species in the positive culture according to the body site sampled.

#### 4. Discussion

Our findings demonstrate that rubbing the skin with a sterile gauze that is further inoculated onto a FastFung plate is the most efficient method for cultivating *Malassezia* yeast from the skin. Whereas a growing number of fungal metagenomics studies highlight a strong representation of *Malassezia* spp. among the fungal communities of various microbiomes, comparative studies of skin sampling techniques and culture-based *Malassezia* spp. isolation media are lacking. Though, the standardization of an effective and simple *Malassezia* spp. isolation procedure is crucial to enhance the patients' diagnosis in the clinical laboratory and to homogenize and allow the comparison of distinct epidemiological studies results.

In the present study, we compared three skin sampling techniques (2 by swabbing and 1 by rubbing with a sterile gauze) on healthy volunteers and we found that rubbing with a sterile gauze was the most efficient method for recovering *Malassezia* yeast from the skin for culture, with a four times greater (67%) positive culture rate by using sterile gauzes than that by using either dry swabs (15%) or Transwab™ (18%) (Figure. 1A). Swabbing has been used in several studies [3], [14-15] and it is considered as an alternative option to other methods such as scraping. Our findings question the reliability and efficacy of the swabbing technique not only in epidemiological studies, but also for the diagnosis of patients in whom a *Malassezia* yeast infection is suspected. Noteworthy, our results regarding the effectiveness of sampling via sterile gauze are in line with those of Ilahi *et al.* [16], who used a similar sampling technique and found an overall positive culture rate of 75.71% on 70 patients, including pityriasis versicolor patients (28/28); breeders, most of them presenting skin lesions (15/17); and healthy controls (10/25). The sterile gauze technique presents many advantages, including simplicity, practicability, efficacy, cost-effectiveness, and non-invasiveness.

Furthermore, the FastFung medium proved more efficient than the reference Dixon medium for isolation and growth of *Malassezia* yeast, with 62% *vs.* 38% of positive culture rates, respectively (Figure 1B). Our 38% positive culture rate with the Dixon medium was similar to the one observed by Ilahi *et al.* [16], who found a 40% of positive culture in healthy controls with the modified Dixon agar. This indicates that the difference observed in the present study is chiefly explained by the relatively enhanced efficacy of the FastFung medium. Noteworthy, the FastFung medium efficiently grew *M. globosa* and *M. restricta* (Figure 1C), both species that are considered to be among the most fastidious yeast to culture within the *Malassezia* genus [17–19]. Moreover, several studies showed that *M. globosa* and *M. restricta* are the most frequently involved species in pityriasis versicolor and seborrheic dermatitis, respectively [15], [20-21]. The enhanced performances of the FastFung medium compared to the Dixon medium may at least in part be due to some of the Schaedler agar components. In line with this hypothesis was that our previous finding that this medium showed an enhanced efficiency to grow and isolate a variety of clinical fungi compared to the Sabouraud reference medium (Bittar *et al.*, submitted).

#### 5. Conclusions

Notwithstanding our study limitations, which include the relatively small number of participants studied and the lack of patients with *Malassezia*-associated diseases, our findings show that sterile gauze is an effective and reliable sampling technique for recovering *Malassezia* yeasts from the human skin for culture and that the efficacy for *Malassezia* culture of the novel FastFung medium is enhanced when compared to the reference Dixon medium. We thus propose implementing sterile gauze rubbing skin sampling followed by inoculation on FastFung medium in the routine clinical laboratory procedure for *Malassezia* spp. cultivation in both clinical and epidemiological studies.

**Author Contributions:**

Abdourahim Abdillah: Conceptualization; Data curation; Investigation; Formal analysis; Methodology; Visualization; Writing the original draft.

Saber Khelaifia: Conceptualization; Methodology; Data curation; Investigation; Writing the original draft.

Didier Raoult: Conceptualization; Funding acquisition; Methodology; Resources; Supervision; Writing - review & editing.

Fadi Bittar: Conceptualization; Methodology; Supervision; Writing - review & editing.

Stéphane Ranque: Conceptualization; Methodology; Investigation; Formal analysis; Resources; Supervision; Writing - review & editing.

**Funding:** This research was funded by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the French National Research Agency under the program “Investissements d’avenir”, reference ANR-10-IAHU-03, the Région Provence Alpes Côte d’Azur and European funding FEDER PRIM1.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results”.

**References**

- [1] Q. Wang, B. Theelen, M. Groenewald, F. Bai, and T. Boekhout, “Moniliellomycetes and Malasseziomycetes, two new classes in Ustilaginomycotina,” *Persoonia*, pp. 41–47, 2014.
- [2] G. Wu *et al.*, “Genus-Wide Comparative Genomics of *Malassezia* Delineates Its Phylogeny, Physiology, and Niche Adaptation on Human Skin,” *PLOS*, pp. 1–26, 2015.
- [3] B. H. Oh, Y. C. Song, Y. W. Lee, Y. B. Choe, and K. J. Ann, “Comparison of Nested PCR and RFLP for Identification and Classification of *Malassezia* Yeasts from Healthy Human Skin,” *Ann. Dermatol.*, vol. 21, no. 4, pp. 352–357, 2009.
- [4] E. Chryssanthou, U. Broberger, and B. Petrini, “*Malassezia pachydermatis* fungaemia in a neonatal intensive care unit,” *Acta Paediatr. Int. J. Paediatr.*, vol. 90, no. 3, pp. 323–327, 2001.
- [5] G. Gaitanis, P. Magiatis, M. Hantschke, I. D. Bassukas, and A. Velegraki, “The *Malassezia* Genus in Skin and Systemic Diseases,” *Clin. Microbiol. Rev.*, pp. 106–141, 2012.
- [6] R. Iatta *et al.*, “Bloodstream infections by *Malassezia* and *Candida* species in critical care patients,” *Med. Mycol.*, no. December 2013, pp. 264–269, 2014.
- [7] T. A. Auchtung *et al.*, “Investigating Colonization of the Healthy Adult Gastrointestinal Tract by Fungi,” *mSphere*, vol. 3, no. 2, pp. 1–16, 2018.
- [8] S. Raimondi *et al.*, “Longitudinal Survey of Fungi in the Human Gut: ITS Profiling, Phenotyping, and Colonization,” *Front. Microbiol.*, vol. 10, no. July, pp. 1–12, 2019.
- [9] L. Delhaes *et al.*, “The airway microbiota in cystic fibrosis: A complex fungal and bacterial community-implications for therapeutic management,” *PLoS One*, vol. 7, no. 4, 2012.
- [10] S. D. Willger *et al.*, “Characterization and quantification of the fungal microbiome in serial samples from individuals with cystic fibrosis,” *Microbiome*, pp. 1–15, 2014.

- [11] T. Jagielski, E. Rup, A. Ziólkowska, K. Roeske, A. B. Macura, and J. Bielecki, "Distribution of *Malassezia* species on the skin of patients with atopic dermatitis, psoriasis, and healthy volunteers assessed by conventional and molecular identification methods," *BMC Dermatol.*, vol. 14, 2014.
- [12] I. Hamad, S. Ranque, E. I. Azhar, and M. Yasir, "Culturomics and Amplicon-based Metagenomic Approaches for the Study of Fungal Population in Human Gut Microbiota," *Sci. Rep.*, no. November, pp. 1–8, 2017.
- [13] C. Cassagne *et al.*, "Routine identification and mixed species detection in 6,192 clinical yeast isolates," *Med. Mycol.*, vol. 54, no. 3, pp. 256–265, 2016.
- [14] J. A. Omodo-Eluk, K. P. Baker, and H. Fuller, "Comparison of two sampling techniques for the detection of *Malassezia pachydermatis* on the skin of dogs with chronic dermatitis," *Vet. J.*, vol. 165, no. 2, pp. 119–124, 2003.
- [15] Y. Won Lee *et al.*, "Distribution of *malassezia* species on the scalp in Korean seborrheic dermatitis patients," *Ann. Dermatol.*, vol. 23, no. 2, pp. 156–161, 2011.
- [16] A. Ilahi, I. Hadrich, S. Neji, H. Trabelsi, F. Makni, and A. A. Ayadi, "Real-Time PCR Identification of Six *Malassezia* Species," *Curr. Microbiol.*, vol. 0, no. 0, p. 0, 2017.
- [17] P. Honnavar, G. S. Prasad, A. Ghosh, S. Dogra, and S. Handa, "*Malassezia arunalokei* sp. nov., a Novel Yeast Species Isolated from Seborrheic Dermatitis Patients and Healthy Individuals from India," *J. Clin. Microbiol.*, vol. 54, no. 7, pp. 1826–1834, 2016.
- [18] C. M. Gemmer, Y. M. DeAngelis, B. Theelen, T. Boekhout, and T. L. Dawson, "Fast, noninvasive method for molecular detection and differentiation of *Malassezia* yeast species on human skin and application of the method to dandruff microbiology," *J. Clin. Microbiol.*, vol. 40, no. 9, pp. 3350–3357, 2002.
- [19] S. Triana *et al.*, "Lipid metabolic versatility in *Malassezia* spp. yeasts studied through metabolic modeling," *Front. Microbiol.*, vol. 8, no. SEP, pp. 1–18, 2017.
- [20] N. Morishita, Y. Sei, and T. Sugita, "Molecular analysis of *Malassezia* microflora from patients with pityriasis versicolor," *Mycopathologia*, pp. 61–65, 2006.
- [21] M. Kaur, T. Narang, M. Bala, S. Gupte, P. Aggarwal, and A. Manhas, "Study of the distribution of *Malassezia* species in patients with pityriasis versicolor and healthy individuals in Tertiary Care Hospital, Punjab," *Indian J. Med. Microbiol.*, vol. 31, pp. 270–274, 2013.