

**Table 2.** Outcomes of the MVI IOL in the studied group by type of risk factor.

<b>Parameter</b>	<b>No risk obese MVI IOL</b>		<b>High risk obese MVI IOL</b>	
	On-term pregnancies (n=17) X± SD*	Late-term pregnancies (n=25) X± SD	Preeclampsia (n=12) X± SD	Diabetes Mellitus (n=20) X± SD
Gestational age (weeks)	39.9 ± 0.9	41.3 ± 0.2	38.9 ± 0.6	38.9 ± 0.6
Misoprostol action (h)	12.34 ± 5.7	11.0 ± 5.9	12.3 ± 6.6	10.8 ± 3.9
Time from induction to delivery (h)	18.75 ± 8.47	16.78 ± 5.57	17.37 ± 7.42	16.62 ± 4.69
Initial Bishop score	2.2 ± 0.8	1.28 ± 1.5	2.7 ± 1.1	1.7 ± 1.2
One-min* Apgar score	8.5 ± 0.8	8.56 ± 0.7	8.41 ± 0.8	8.4 ± 1.2
Five-min* Apgar score	9.2 ± 0.7	9.08 ± 0.6	9 ± 0.6	9 ± 0.6
Fetal weight (grams)	3205.2 ± 434.4	3450 ± 301.4	3417.5 ± 478.8	3417.5 ± 478.8

\*X± SD - Data are expressed as mean ± standard deviation or n (%) unless otherwise specified, \* min –minute