

Review

Physical-Activity-Based Interventions Targeting Overweight and Obesity among University Students – A Systematic Review

Julia Pfisterer^{1,†}, Constantin Rausch^{1,†}, Doreen Wohlfarth^{1,†}, Philip Bachert¹, Darko Jekauc^{1,*} and Kathrin Wunsch^{1,*}¹ Institute of Sports and Sports Science, Karlsruhe Institute of Technology, Karlsruhe, Germany

* Correspondence: darko.jekauc@kit.edu; phone: + 49 721 608 – 45725; kathrin.wunsch@kit.edu; phone: + 49 721 608 – 45431

† These authors contributed equally to this work

Abstract: Overweight and obesity, including its prevalence and consequences reflect a leading public health problem. Studies have already shown that physical activity leads to a reduction in body weight in children and adults. However, the university setting has rarely been investigated. The aim of this review is therefore to examine and summarize the effectiveness of physical-activity-based interventions to reduce obesity and overweight in university students. Three databases (PubMed, Scopus & Web of Science) were searched for relevant studies published in English between January 2010 and February 2022. Quantitative studies, conducting a physical-activity-based intervention with overweight or obese university students and reporting changes in BMI, were included. Data were described in a narrative synthesis. Eleven of 16 included studies reported a significant reduction in BMI. However, all studies except one were able to demonstrate some BMI improvements, whereas all studies reported significant changes in at least one health-related indicator. Aerobic exercises were able to demonstrate the greatest reductions in BMI. This review is the first systematic presentation on the effectiveness of physical-activity-based interventions on overweight and obese university students. Future work should reconsider BMI as the primary outcome because it is prone to bias. More interventions are needed to improve strategies.

Keywords: physical-activity-based interventions; overweight; obesity; BMI; change; university students; tertiary education; review

1. Introduction

It is already known that physical activity is an essential component of combating overweight and obesity. The positive effects of physical activity on physical and mental health have been demonstrated in numerous studies and reviews (1–3). Nonetheless, in 2016 the WHO identified that 28 % of the adults (>18 years) worldwide are not physically active enough. This means that these adults are physically active for less than 150 min per week at moderate intensity or less than 75 min at a vigorous intensity (4). Additionally, the rate of overweight and obese individuals increased sharply in recent decades and has developed into a leading public health problem. Being overweight causes cardiovascular diseases, diabetes, musculoskeletal disorders, and different types of cancer (5). According to the WHO, the prevalence of obesity has almost tripled since 1975. In 2016, 1.9 billion people aged 18 years and older were overweight, of which 650 million individuals were considered obese. Accordingly, in 2016 39% of the adult population worldwide was overweight and 13% was classified as obese (6). Depending on the geographical region, the classification of overweight and obese differs; for more details see WHO (6, 7).

Even though existing evidence shows that higher education levels seem to be associated with a lower probability of being overweight or obese (8), the prevalence among university students is substantially high (9). In the ACHA-National College Health Assessment (ACHA-NCHA) III by the American College Health Association (ACHA), 38,1% of

undergraduate students were classified as overweight or obese (10). Peltzer et al. investigated the prevalence of overweight and obesity among 15,746 university students from 22 different countries and concluded that 22% were overweight and obese (9). Additionally, when compared to the general population, weight gain is five times higher among university students (11). Especially in the first year of college, university students tend to gain weight, which is referred to as the "Freshman 15" phenomenon. Freshman 15 is the myth that college students would gain 15 pounds (6.8kg) during their first year of university (12). Vadeboncoeur et al. conducted a meta-analysis of 22 studies on this topic and concluded that 60.9% of students gained on average 7.5 pounds (3.38kg) weight during their first year of college (13). This meta-analysis failed to confirm the myth of the Freshman 15, but weight gain in the first year of university was significant. These numbers are alarming given that university students tend to engage in unhealthy habits like insufficient physical activity. The ACHA-NCHA survey, for example, demonstrated that less than half of the US college students (42.1%) met the Guidelines for Active Adults in Spring 2021, meaning being physically active on at least two days a week at a moderate or vigorous intensity and demanding all major muscle groups (10). Similar figures have already been observed by Irwin in a review of 35,747 students, stating that half of the students were insufficiently physically active and consequently at higher risk for health problems (14). Furthermore, Keating et al. found in their meta-analysis that around 30 to 50% of college students were not physically active enough to reach any health benefits (15). Due to the abovementioned reasons, university as a setting should be allocated a significant role in public health promotion.

Considering the high prevalence of overweight and obesity among university students, the associated health risks, as well as insufficient physical activity, tailored interventions are needed to address this vulnerable population. The effectiveness of interventions including physical activity on the overweight and obese population has already been investigated in different age groups such as adults and children (16–18). Even though Plotnikoff et al. examined the effectiveness of interventions targeting weight-related behaviors among university students in their meta-analysis (19), these results cannot be generalized to overweight or obese students. Thus, there exists no systematic review, which focuses on physical-activity-based interventions for overweight and obese students. In light of findings that increasing physical activity is an important approach to reducing weight and that current reviews and meta-analyses neglect the population of overweight and obese university students, this systematic review aims to investigate the effectiveness of physical-activity-based interventions targeting overweight and obesity among university students.

2. Materials and Methods

This study was carried out in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement guidelines (20).

2.1. Eligibility criteria

Eligible characteristics were based on different items of the PI(C)OC scheme (Population, Intervention, Comparison (optional), Outcomes, and Context) (21). Overall, selection criteria were kept general at the outset to provide an overview of the topic and to ensure finding all relevant sources. The further specific narrowing was done in a later phase.

2.2. Types of participants

Studies that recruited overweight ($> 25 \text{ kg/m}^2$) or obese ($> 30 \text{ kg/m}^2$) university students in terms of body mass index (BMI) as well as studies where the authors declared that the population was overweight/obese were included. Other characteristics such as

age, gender, socioeconomic, and ethnic characteristics of the students were not further considered or narrowed down.

2.3. Types of intervention

Interventions that were eligible for inclusion had to be conducted in a tertiary educational setting, where university students were recruited to perform a physical-activity-based program. Anthropometric characteristics (height and weight) of the enrolled participants were measured, which could be used for the calculation of the BMI. All interventions encompassing any physical movement were considered for this review. Thus, interventions implementing a fitness, exercise, or sports program were equally taken into account as interventions fostering light physical activity, such as walking. Type, duration, guidance, and the number of sessions were not further defined to provide a comprehensive overview of the current study situation. Intervention studies implementing online programs, using digital communication or technical devices such as activity trackers to foster physical activity among university students were considered and not further specified in their study design for the abovementioned reason. Interventions without any physical activity component, for instance, focusing solely on diet to target overweight and obesity in university students were not taken into consideration.

2.4. Types of studies

All quantitative study designs conducting an intervention (including randomized controlled trials [RCT], controlled clinical trials [CCT], cohort analytics, case-control, or cohort studies) were included in this review.

2.5. Types of outcome measures

To measure the change in students' BMI due to physical activity, an analysis of BMI was the primary outcome measure of the intervention. Changes in other anthropometric characteristics such as body fatness, blood counts, etc. were accepted as additional outcome documentation.

In order to estimate and compare the effectiveness of the interventions, effect sizes were calculated using Psychometrica (22). The effect size d_{ppc2} according to Morris (23) of studies using an RCT or CCT design in which the control group [CG] did not change their behavior was calculated with formula 3. Formula 4 was used for cohort designs or RCTs and CCTs where the CG changed their behavior. Since none of the studies reported inter-correlations, the effect size d_{av} of Cumming (24) was used, a pragmatic approach often used in meta-analysis.

2.6. Information sources and search

To examine the effectiveness of physical-activity-based interventions among university students, a structured electronic search of three databases was conducted based on the PRISMA guidelines (20). PubMed, Scopus, and Web of Science were specified as suitable databases as they comprehensively cover the fields of interest regarding the research question. Only articles published in peer-reviewed journals between January 2010 and February 2022 and written in English were considered for the review. The following keywords and operators were used following the PI(C)OC scheme in all databases. Since the *Population* and the *Context* are congruent, these two terms are merged and searched as follows: students OR university OR college OR tertiary education. Using the NOT operator, some terms relating to the population or context were excluded to restrict the search: adults OR elementary OR children. *Interventions* were reflected with the keywords as follows: intervention OR treatment OR health promotion OR physical activity* OR exercise OR training OR sport OR fitness. The *Outcome* was searched with the help of: overweight OR obesity OR BMI. The search terms had to be adapted regarding title, abstract and full-text search for the specific databases. The appendix contains the complete and final search

terms of all three databases. In addition, authors were contacted via *ResearchGate* to obtain eligible studies that were not publicly available. Furthermore, while processing the data extraction, the bibliography of each study was searched for additional potentially relevant studies. Searches began on the first of December 2020 and ended with the last search query in March 2022.

2.7. Study selection

A pre-selection of studies was conducted by each author individually through title screening, saving those which included relevant keywords of inclusion criteria in *Citavi 6.11* (Swiss Academic Software GmbH, Wädenswil, Switzerland) to avoid duplications. In the next step, all authors first performed abstract screenings of an equitable number of the pre-selected studies individually to check for inclusion and exclusion characteristics. Secondly, each abstract was again discussed by all authors together to avoid omitting relevant studies. In consequence, studies without an intervention design, without targeting a university students' population, and without reporting, BMI results were removed. In the last step, studies were checked for inclusion and exclusion criteria more thoroughly by full-text analysis in triple approval to reduce bias.

2.8. Data collection and data details

Data extraction was performed in a standardized manner, with each author examining an equal number of studies individually and extracting all relevant information from the respective studies into spreadsheet format. Based on the results of data extraction, the authors discussed ambiguities and jointly decided which studies should ultimately be included for analysis in the review.

2.9. Quality assessment

Quality assessment for the selected studies was conducted by using the EPHPP Quality Assessment Tool for quantitative studies (Effective Public Healthcare Panacea Project, 25). This assessment tool is an elaborated instrument for evaluating the quality of a study and can be used in a wide range of health-related topics to develop recommendations for study findings. It includes several items helping to classify a study according to its quality from one (strong) to three (weak). Each rating is described for: (1) selection bias (2) design (3) confounders (4) blinding (5) data collection methods (6) withdrawal and dropouts (25).

Every author rated each full text individually at first. Results of the individual ratings were then discussed by all authors together and in case of any ambiguities, a joint decision for a final rating was made (26). Full-text analysis, data extraction, and quality assessment were completed by April 2022.

2.10. Synthesis of results

The synthesis of results was based on the Guidance on the Conduct of Narrative Synthesis in Systematic Reviews (26). This approach by Popay et al. uses a general framework to describe the four elements of the narrative synthesis process which are presented as follows: Developing a theoretical model and preliminary syntheses, exploring relationships within and between studies, and assessing the robustness of synthesis.

Following Plotnikoff et al. (27), it was analyzed whether the selected studies based their intervention on a theoretical model, focusing especially on social cognitive theories as they can be used to explain health-related behavior, including physical activity (28). To develop a preliminary synthesis, the authors identified similarities and differences among study findings, partially using charts and tables. In the process of analyzing the results of each study, the authors yielded with the help of their methods further new insights. In the next step, these preliminary results were then discussed in detail and compiled in group sessions. Based on the direction and magnitude of the effects, associations and patterns between the interventions were identified, which needed to be explained in the third step.

To this end, the authors analyzed heterogeneity across studies, such as variability in outcomes, study designs, study populations, interventions, and settings. This step of the analysis is characterized by the relationship of outcomes within and between studies. Finally, each outcome of the included studies was evaluated to be able to assess the strength of the synthesis product. As already described before, the results of the quality assessment were considered and could contribute to the overall assessment of the review results. With the help of the quality assessment, it could be identified if results can be generalized and transferred to other populations or contexts.

3. Results

A total of 3231 hits were obtained in the PubMed, Scopus, and Web of Science databases using the previously defined search terms. Individual screening per database for matching keywords in title and abstract led to a selection of 93 studies. The exclusion of studies based on the selection criteria that did not include an intervention design, student population, or physical activity ultimately led to a large reduction in eligible studies.

By using the literature management program *Citavi 6.11* (Swiss Academic Software GmbH, Wädenswil, Switzerland), duplicates were immediately removed, leaving 93 studies that were subjected to a more detailed abstract screening. In particular, studies were excluded that did not correspond to an intervention or did not examine a student sample. Web-based interventions, i.e. interventions that promote physical activity via online communication, were equally shortlisted, resulting in 36 studies after the abstract screening. In some cases, no information was available on the BMI of the group after the intervention, as the focus of the studies was on other dependent variables, therefore 15 studies were excluded. Thus, 21 studies were still available for the full-text analysis. Of these, two additional studies were identified through the full-text analysis that used online communication to motivate physical activity but did not conduct an actual training program. Full-text access was not available for three other studies. As no feedback was received after contacting the authors of these three studies via *ResearchGate*, none of them could be included. The abovementioned five studies were therefore excluded, resulting in 16 studies that met the specified inclusion criteria. The forward and backward search of the bibliography via the website *Connected Papers* did not yield any additional sources. Figure 1 shows the literature search process.




Figure 1: Flow chart of study selection process

3.1. Study characteristics

3.1.1. Methods

The 16 studies that were ultimately included in the review were all published in English between 2012 and 2021. The interventions differed in their study design in seven CCTs (29–35), four RCTs (36–39) and five cohort studies with a single group and pre-post design (40–44). The interventions were conducted in different countries around the world, including four in the USA (30, 40, 41, 39), four in China (36, 33, 38, 34), two in Korea (32, 42) and one each in the United Emirates (29), Indonesia (31), South Africa (43), Iran (37), Jamaica (44) and in Saudi Arabia (35).

3.1.2. Subjects

A total of 969 students were included in the final analyses. Sample sizes varied from ten to 300 participants, with only three of the 16 studies having a sample size of at least 100 (30, 33, 38). Only DiFrancisco-Donoghue et al. (30) ensured an equal gender distribution in their sample. Chen et al., (36), Pacholek et al., (35) and Sun et al. (38) studied only male subjects, while the other interventions involved either exclusively female or predominantly female subjects. In eight studies, the main inclusion criterion was a BMI of at least

over 25 (36, 30, 31, 40, 41, 43, 37, 38). Ha & So (32) and Wei-wei & Ling (34) used a body fat percentage of more than 30%, while Siqiang (33) used a bodyweight of more than 58 kg to declare their sample as overweight. Winters-Stone et al. (39) recruited students who were overweight or were at risk of being obese ($BMI > 22 \text{ kg/m}^2$) and in the remaining studies, being overweight was not an inclusion criterion. Nonetheless, Roopchand-Martin et al. (44) used a BMI above 25 to divide the sample into an overweight and a normal weight group for analysis in addition to the total group. Pacholek et al. (35) divided the sample into two overweight groups to analyze the change in overweight students based on the two different exercise programs. Dalibalta et al. (29), in turn, used the same group allocation to use the overweight group as the intervention group (IG) and the normal weight group as the CG. Lee et al. (42) has not described its main inclusion criteria in detail. Nonetheless, the sample was declared inactive and had in total a BMI over 25 kg/m^2 (25.14 kg/m^2).

3.1.3. Interventions

Subjects in all interventions received a physical-activity-based program that varied in type, intensity, and frequency. The training sessions for the IGs included aerobic training (40, 41, 33), a combination of aerobic and resistance training (35, 34), the combination of strength and aerobic training (32), high-intensity interval training (31, 37, 38), high-intensity circuit training (42), a specific exercise or sport, namely dancing, Tae-Bo, Whole Body Vibration (WBV), volleyball (43, 44, 39, 35) or an unspecified training program (29, 30). It should be noted that the studies by Gifari et al. (31) and Wei-wei & Ling (34) included an additional nutrition program in their IGs, however, these were analyzed separately. Chen et al. (36) carried out a cardio-based cycling program. The CGs mostly received either a different exercise program (36, 29, 38) or instructions to maintain their usual activity level and eating behavior (32, 37, 34, 39). In half of all included interventions, exercise was provided three times a week (29, 31, 32, 42, 43, 37, 34, 39), while in the other interventions sessions ranged from two to six times a week. The majority of the interventions were limited to moderate intensity or a heart rate between 50% and 85% of maximum heart rate (29, 31, 32, 40–43, 37, 44, 33, 38, 34). Only one intervention worked at a low intensity of 40 % $VO_{2\text{max}}$ (36) and one with a subjective feeling with maximal effort (35). Four of the 16 interventions successively increased the load by either increasing the training duration (44) or the intensity of the training sessions after a certain number of weeks (42, 43, 39). The intervention duration varied between studies from four weeks (42, 35) to 39 weeks (30). The most common intervention duration was twelve weeks (36, 32, 40, 41, 33, 38), followed by eight weeks (29, 31, 37).

3.1.4. Outcomes

All included studies provided information on the change in BMI after the intervention and for this purpose collected anthropometric parameters of the subjects. These included body weight and height for the calculation of BMI. In addition, all studies also examined other dependent variables such as body composition, blood pressure, blood lipid levels, or behavioral factors. Thus, BMI was not the sole factor of interest in any study. However, for this review, the necessary information on the change in BMI could be extracted from all studies. Study characteristics are summarized in detail in Table A1.

Table A1: Study characteristics of the included studies

3.3. Risk of bias within the studies

Five studies were rated as strong in the Global Rating using the EPHPP tool (30, 32, 37–39), seven received a moderate (36, 31, 40, 41, 43, 33, 34) and four weak ratings (29, 42, 35, 44). For the Selection Bias category, only Gifari et al. (31) received a strong rating as they described the selection of their sample as randomized. Even though the studies by

Pacholek et al. (35) and Wei-wei & Ling (34) declared a randomization process in their study population, they had a homogenous sample in terms of sex. In addition, the remaining 13 studies did not explicitly state a randomization process. Furthermore, Roopchand-Martin et al. (44) did not report how individuals were selected and what percentage of selected individuals agreed to participate and were thus assigned a weak rating. Thus, all studies, except the study by Roopchand-Martin et al. (44) and Gifari et al. (31), received a moderate rating. In sum, eleven out of 16 studies did not describe what percentage of the selected individuals ultimately gave their consent to participate in the intervention (29, 30, 32, 40–42, 37, 35, 44, 33, 34). Strong ratings were predominantly assigned to study design, as eleven of the included studies were either RCTs (36–39) or CCTs (29–35), which should receive the best rating according to the specifications of the assessment tool. The five cohort studies without CGs (40–44) were thus rated as moderate. Information on confounders differed greatly between the studies so that half of the 16 included studies were given the highest rating, whereas the other half received the lowest. The highest ratings were allocated when there were no crucial differences between groups before the intervention (36, 30, 32, 37, 33, 38, 34, 39). The weak ratings were assigned when 1) group differences existed before the intervention and it was not described if relevant confounders were controlled (29), 2) there was only one group due to the study design (40–44) or 3) control of confounders was not described (31, 35). For blinding, Chen et al. (36) and Lee et al. (42) received a weak rating, since either the awareness of the outcome assessors was not described or the participants and/or assessors were aware of the exposure. The remaining 14 studies received a moderate overall rating on blinding, as none but one study described the blinding process. Only Winters-Stone et al. (39) provided information on the blinding of test administrators and subjects but stated that the research question was communicated to the subjects, thus only a moderate ranking was assigned here as well. BMI was collected in all studies using the standardized methods of weight recording and measuring height. Consequently, the methods of data collection were all rated as strong. Withdrawals and drop-outs were described in detail in eight out of 16 studies, with reasons and/or numbers given (36, 30, 32, 41, 43, 37, 44, 39). Moreover, eleven of the 16 studies had a follow-up rate of 80 % or greater (36, 30–32, 40, 41, 43, 37, 44, 38, 39), so that strong ratings were assigned here. Five of the 16 studies (29, 42, 35, 33, 34) did not provide any information in this regard and thus received a weak ranking. Table 2 shows the Global Rating, as well as its composition for each study

Table 2: Assessment of study quality using the Quality Assessment Tool for quantitative Studies (EPHPP)

Study	Selection Bias	Study Design	Con-founders	Blinding	Data collection methods	Withdrawals and Dropouts	Global Rating
Chen et al. (2021)	2	1	1	3	1	1	2
Dalibalta et al. (2017)	2	1	3	2	1	3	3
DiFrancisco-Donoghue et al. (2018)	2	1	1	2	1	1	1
Gifari et al. (2021)	1	1	3	2	1	1	2
Ha & So (2012)	2	1	1	2	1	1	1
Joseph et al. (2014)	2	2	3	2	1	1	2
Joseph et al. (2016)	2	2	3	2	1	1	2
Lee et al. (2021)	2	2	3	3	1	3	3

Mathunjwa et al. (2013)	2	2	3	2	1	1	2
Moravveji et al. (2019)	2	1	1	2	1	1	1
Study	Selection Bias	Study Design	Con- founders	Blinding	Data collection methods	Withdrawals and Dropouts	Global Rating
Pacholek et al. (2021)	2	1	3	2	1	3	3
Roopchand-Martin et al. (2015)	3	2	3	2	1	1	3
Siqiang (2018)	2	1	1	2	1	3	2
Sun et al. (2020)	2	1	1	2	1	1	1
Wei-Wie & Ling (2021)	2	1	1	2	1	3	2
Winters-Stone et al. (2020)	2	1	1	2	1	1	1

Ratings: 1= strong. 2 = moderate. 3 = weak.

3.3. Results of the individual studies

Eleven out of the 16 included studies reported a significant reduction in BMI within IGs after the intervention at a predetermined significance level of five percent (36, 29, 31, 40, 42, 43, 38, 34, 37, 33, 39). Even though the studies by DiFrancisco-Donoghue et al. (30); Ha & So (32); Joseph et al. (41); Pacholek et al. (35); Roopchand-Martin et al. (44) did not produce significant BMI reductions, they showed some decrease in BMI ranging from -0.2 to -0.9 as well. The IG of the study by Roopchand-Martin (44) had the smallest decrease with -0.2, whereas Joseph et al. (41) had the highest non-significant reduction in BMI of -0.9. It should be noted that DiFrancisco-Donoghue et al. (30) showed a BMI reduction only for female participants of the Fitbit-Plus group (-0.5) and the male Fitbit-Only group (-0.2), but did not report any p-values and therefore no conclusion can be drawn on the significance. Pacholek et al. (35) reported a reduction of -0.3 in BMI for the volleyball IG. Finally, Ha & So (32) showed a slight decrease in BMI in the IG of -0.21.

Of the eleven included studies which used a RCT or CCT design, eight reported significant results in terms of changes in BMI (36, 29, 31, 37, 33, 38, 34, 39). However, since the CGs in five studies using a RCT or CCT design, changed their behavior as well, only four studies could be taken into account for a comparison between the IG and CG (37, 34, 39, 32). Here, three of four studies found significant differences between the CG and IG (37, 34, 39). In the five studies in which the CGs changed their behavior (36, 29, 30, 33, 38), three studies reported significant BMI reductions in the IG and CG (29, 33, 38). Chen et al. (36) only reported significant reductions in the IG but not in the CG. DiFrancisco-Donoghue et al. (30) gave insufficient information on significance and just reported the total differences in body composition for the overweight population. The studies by Gifari et al. (31) and Pacholek et al. (35) were declared as CCTs but did not include a CG in their design. Like the abovementioned studies, where the CGs changed their behavior, these studies used three (31) or two (35) IGs to compare the effectiveness of the intervention. Pacholek et al. (35) found a non-significant reduction in BMI in both groups, whereas in the study by Gifari et al (31) one group significantly decreased their BMI.

In four of the seven included CCTs significant improvements in BMI were reported (29, 33, 34, 31). Dalibalta et al. (29) conducted the same exercise program with both the IGs

($n = 14$, MBMI = 28.7, SD = 3.27) and the CG ($n = 32$, MBMI = 21.6, SD = 2.05). Groups differed regarding the average BMI, so there was an overweight IG with a BMI of at least 25 and a normal weight CG with a BMI of less than 25. Both groups received an exercise training program that was not described in more detail. After eight weeks, BMI had decreased significantly in both the CG (MBMI = 21.1, SD = 2.00, $d_{av} = .247$) and the IG (MBMI = 28.1, SD = 2.79, $d_{av} = .198$). However, BMI was reduced by a greater extent in the normal weight group (2.31 %) than in the overweight group (2.09 %). Siqiang (33) chose aerobic exercise for the IG ($n = 50$, MBMI = 25.5, SD = 2.70), while the CG ($n = 50$, MBMI = 26.8, SD = 3.40) only had to take calcium pyruvate twice a day. After twelve weeks of intervention, there was a great significant reduction in BMI in the overweight CG (MBMI = 24.1, SD = 2.90, $d_{av} = .857$) and an even greater reduction in the IG (MBMI = 21.6, SD = 3.10, $d_{av} = 1.345$). Wei-wei & Ling (34) combined aerobic exercises and resistance training for their IG ($n = 20$, MBMI = 22.06, SD = 0.98) and asked the CG ($n = 20$, MBMI = 22.54, SD = 0.71) to maintain their behavior. After 16 weeks, BMI within the IG was significantly reduced by 1.9 % (MBMI = 21.64, SD = 0.85, $d_{ppc2} = .641$) whereas the CG showed an increase in BMI of 0.62 % (MBMI = 22.68, SD = 1.11). Gifari et al. (31) combined HIIT exercises with pre-meal water intake (PWI) for one of their IGs ($n = 9$, MBMI = 26.6, SD = 3.6) and exclusively HIIT exercises for the other IG (MBMI = 25.9, SD = 2.4). The HIIT + PWI group significantly reduced its BMI by 1.95 % (MBMI = 25.2, SD = 3.2, $d_{av} = .152$), whereas the exclusively HIIT group failed to significantly reduce the BMI with a decrease of 0.38% (MBMI = 25.8, SD = 2.4, $d_{av} = .042$). Small non-significant decreases can be seen for the IGs of DiFrancisco-Donoghue et al. (30), Pacholek et al. (35), and Ha & So (32).

All four included RCTs reported significant results (36–39). Chen et al. (36) used a combination of low-intensity cycling and blood flow restriction training (BFRT) by wearing air pressure belts for their IG ($n = 18$, MBMI = 30.10, SD = 0.95). The IG was compared with a CG ($n = 19$, MBMI = 30.30, SD = 1.08) also conducting low-intensity cycling but without performing BFRT. After the intervention period of twelve weeks, BMI was reduced within both groups. The IG was able to significantly reduce their BMI by 3.65% (MBMI = 29.0, SD = 1.79, $d_{av} = .803$), whereas the BMI in the CG was non-significantly reduced by 1.32% (MBMI = 29.90, SD = 1.44, $d_{av} = .317$). Moravveji et al. (37) used the principle of successive weekly increases in training units. A 1200 m run was completed three times per week and increased by 400 m each week. Subjects were divided into 1) a continuous group ($n = 10$, MBMI = 26.77, SD = 1.53), completing the 1200 m without a break, 2) an interval group ($n = 12$, MBMI = 27.57, SD = 1.02), which ran 400 m three times with a two-minute break of walking between runs, and 3) a CG ($n = 9$, MBMI = 27.57, SD = 1.62). The CG was asked to maintain its normal behavior and was instructed not to participate in any other training program during the intervention period. After eight weeks, BMI significantly decreased in both the continuous group (MBMI = 26.08, SD = 1.21, $d_{ppc2} = .704$) and the interval group (MBMI = 26.82, SD = 0.98, $d_{ppc2} = .896$). The CG, on the other hand, experienced a slight increase in BMI of 1.70% (MBMI = 28.04, SD = 1.46). Sun et al. (38) performed high-intensity interval training with the IG ($n = 150$, MBMI = 30.85, SD = 3.79) and aerobic training with the CG ($n = 150$, MBMI = 30.54, SD = 3.86). After twelve weeks of the intervention, there was a significant reduction in BMI within the IG (MBMI = 28.17, SD = 2.89, $d_{av} = .802$) and the CG (MBMI = 28.45, SD = 2.76, $d_{av} = .631$). The RCT by Winters-Stone et al. (39) differed somewhat from the other studies in its training program. Training sessions in the IG were performed using a whole-body vibration method ($n = 37$, MBMI = 28.24, 95% CI [23.87; 32.60]). The degree of vibration gradually increased in the first six weeks and was adjusted to 50 Hz in weeks 7-24. The CG ($n = 40$, MBMI = 28.27, 95% CI [26.48; 30.07]) was expected to maintain its usual eating behavior and physical activity level throughout the intervention period. Even though a slight increase in BMI of 0.22 (95% CI [-0.59; 1.03]) was found at a non-significant level in the IG, when analyzing only participants with high adherence (MBMI = 27.02, 95% CI [22.34; 31.71]) a small significant reduction of .04% could be found. As Winters-Stones et al. (39) provided insufficient information, no effect size could be calculated.

Three of the five included cohort studies declared a significant reduction in BMI (40, 42, 43). The subjects in the study of Joseph et al. (40) ($n = 15$, MBMI = 33.27, SD = 6.53) participated in aerobic training in the form of walking and were also given the choice between exercising by themselves and cardiovascular group training. After the intervention of twelve weeks, the average BMI was reduced by 2.10% (MBMI = 32.57, SD = 6.62, $d_{av} = .106$). Lee et al. (42) conducted high-intensity circuit training ($n = 10$, MBMI = 25.14, SD = 2.51) for four weeks, three times per week, and could reduce the BMI of their intervention population by 3.58% (MBMI = 24.24, SD = 2.60, $d_{av} = .352$). The training sessions of Mathunjwa et al. (43) consisted of Tae-Bo lessons in which subjects participated three times a week for ten weeks. After five weeks of moderate-intensity, training was increased to high intensity in weeks six to ten. The BMI of the group ($n = 60$, MBMI = 32.26, SD = 5.65) had decreased by 6.66% (MBMI = 30.11, SD = 5.46, $d = .2$). Joseph et al. (41) used an internet-enhanced approach combined with real-life physical activity interventions for twelve weeks on a moderate level ($n = 25$, MBMI = 33.80, SD = 5.70). Even though results were not significant, a decrease in BMI was shown after the intervention (MBMI = 32.90, SD = 6.00, $d_{av} = .154$). Roopchand-Martin et al. (44) conducted a dance training on the XBOX Kinect using the program Just Dance. During the six-week intervention, dance duration was increased by 15 minutes every two weeks, whereas the frequency was reduced by one session per week, to maintain participation rates. After six weeks of moderate training the BMI of the subjects ($n = 24$, MBMI = 30.51, SD = 5.18) decreased on a non-significant level by 0.66% (MBMI = 30.31, SD = 5.39, $d_{av} = .038$). Results of the included studies can be seen in more detail in Table A3.

Table A3: Individual results of included studies (change of BMI)

3.3. Synthesis of results

Behavioral interventions are characterized by their complexity and variability. Therefore, the included studies are distinguished by considerable heterogeneity in terms of methods, participants, intervention approaches, and other characteristics. As mentioned by Popay et al. (26) a primary reason for choosing a narrative approach is to investigate the differences in the included studies. Physical-activity-based interventions vary substantially across their intervention approaches. Thus, the focus of this systematic review is the presentation and description of the study characteristics, the quality, and outcomes, as well as a qualitative analysis of the included studies.

Theory-based interventions have been shown to have a positive impact on physical activity for adolescents (27). Of the studies included in this review, only the study by Joseph et al. (41) was theory-driven and based its intervention on the Social Cognitive Theory. Since only one study used a theoretical framework, no conclusion can be drawn on whether the results of the individual study can be attributed to the theoretical background. Likewise, as only two of the included studies used a web-based approach and did not yield any significant results (30, 41), no conclusion can be drawn about the effectiveness of technological devices or web-based approaches.

In all studies that included a CG in their study design, excluding DiFrancisco-Donoghue et al. (30) and Dalibalta et al. (29), the percentage of reduction in BMI was always higher in the IG. Although not all studies showed significant effects on BMI, all included interventions reported at least one significantly improved health-related indicator. Those health-related indicators included improvements in body fat percentage, step counts, sedentary behavior, maximal oxygen consumption, resting heart rate, or serum biomarkers.

However, it is noticeable that all studies except Winter-Stone et al. (39), showed a decrease in BMI in at least one of their IGs, suggesting a tendency towards weight loss. Of note, although Winter-Stone et al. (39) found no BMI reductions when all participants in the IG were included in the analysis, when controlling for high adherence of at least 80% of intervention sessions, there was a significant improvement. This may indicate that high participant adherence may be an important factor in the success of an intervention.

Furthermore, the study by Gifari et al. (31) produced significant BMI improvements of 1.95% in their combination group, which included pre-meal water intake and high-intensity training. Considering that the effect was lower when conducting only HIIT training (-0.38%), and there was no reduction in the group consuming only water before meals (0.38%), the combination of both approaches could positively influence the intervention effects.

The largest percentage reductions in BMI in the IGs were achieved in the studies by Siqiang (33), Sun et al. (38), and Mathunjwa et al. (43), with a decrease of 15.29%, 8.69%, and 6.66% respectively. In comparison to the other studies, except Winters-Stone et al. (39) and Wei-Wei & Ling (34), these interventions had a relatively large sample size, ranging from 60 to 300, with all participants being overweight. Conversely, the four studies, which reported p-values and did not produce any significant reductions in BMI, had a sample size smaller than 29 (41, 35, 44, 32). This small sample size may have resulted in the studies being underpowered and therefore did not show significant improvements.

The three studies that reached the largest BMI reductions (43, 33, 38) conducted only aerobic exercises. Since aerobic exercises tend to have a higher energy expenditure than resistance training, it is considered to be more effective in reducing body weight and fat mass (45) and could therefore explain the positive impacts on BMI changes. Conversely, the study by Ha & So (32) for example, implemented resistance training in their intervention, resulting in a decreased body fat percentage and an increased muscle mass, which leads to small changes in body weight and thus no significant improvements in BMI could be seen.

It is noticeable that only one of the included studies reported effect sizes (43). It was possible to calculate effect sizes for another 13 studies, which should be seen as approximate values. Half of the 14 studies did not reach the recommended threshold by Cohen (46) for a small effect of .3 (29, 31, 32, 40, 41, 35, 44). Furthermore, although Mathunjwa (43) did reach one of the largest percentage BMI reductions, only a small effect can be reported here. One explanation could be the rather large variance of BMI in their sample before as well as after the intervention. For Siqiang (33) and Sun et al. (38), on the other hand, the calculated effect sizes show a strong effect and thus appear to be consistent with the reported high percentage of BMI reduction. Here, the standard deviation was substantially lower in contrast to the study by Mathunjwa (43). Interestingly, although the highest BMI reductions do not necessarily coincide with the strongest effects, the highest effect sizes are also evident in those studies that implemented interventions involving aerobic exercise (36, 37, 33, 38).

Taking into account the study quality only three of eleven studies with significant results received a strong (37–39), six a moderate (36, 31, 40, 43, 33, 34), and two a weak global rating (29, 42). This leads to the assumption that the results shown above should be interpreted cautiously and should be considered when generalizing the results. Since the included studies provided highly homogenous samples, only one strong rating for selection bias could be assigned (31), so generalization of the results to different populations should be done with caution. Significant BMI reductions were achieved in five of six studies with a strong rating in study design and that used solely aerobic exercise in their intervention (36, 31, 37, 33, 38). Nevertheless, no clear patterns regarding study design, intervention duration, frequency, and intensity can be identified in the totality of included studies

4. Discussion

The current review identified 16 studies that examined the effects of physical-activity-based intervention programs to reduce BMI in students within the tertiary sector. Although only eleven studies declared significant reductions in BMI, all studies except one showed at least some reduction in BMI, indicating a trend toward the effectiveness of the intervention programs. In the majority of the studies, BMI was not the primary outcome. Therefore, the effectiveness of the intervention is additionally reinforced by the fact that

at least one health-related indicator was significantly improved in each study (e.g. body fat percentage, weight loss, lipid levels, resting heart rate, maximal oxygen uptake, less sedentary behavior or a higher step count). The greatest improvements in BMI were achieved in studies that performed exclusively aerobic exercises in their intervention and had a relatively large sample size ($N > 60$). This can also be supported by considering the effect sizes, with the strongest effects observed in studies involving aerobic exercise. No clear pattern regarding significant BMI reductions could be derived from study design. Additionally, no distinct trends in duration, frequency, and intensity of the intervention programs could be observed because of substantial heterogeneity among studies showing significant improvements in BMI. Since only three studies with significant results received a strong global rating and most of the studies had a highly homogenous sample, which consisted mostly of female participants, the results should be interpreted cautiously and global generalizability should be questioned. To account for innovative developments in intervention opportunities, studies in which no traditional training sessions were conducted were explicitly included. In these studies, attempts were made to promote the participants' physical activity with the use of websites and/or messages. However, the two included web-based interventions failed to report significant results in terms of BMI.

Between 1980 and 2008, the prevalence of obesity nearly doubled (6). Overweight and obesity is a current topic in all different age groups. To avoid an even more overweight and obese population, it is important to act now. There are already many approaches to increase physical activity and reduce BMI. Positive effects of such interventions have been confirmed on adults and children (47, 2, 48–53). Long-term findings on physical activity were highlighted by Reiner et al. (2) in their review, which included overweight and obesity in adolescents and adults: Over five years, weight change was examined concerning physical activity. It was found that those who reduced their daily physical activity gained considerable weight. Individuals who maintained their activity did not gain weight and, in turn, individuals who increased activity lost weight. Hankinson et al. (48) studied the change in BMI of adults older than 20 years with high habitual activity over 20 years. They found a smaller increase in BMI, waist circumference, and weight per year compared to subjects with low habitual activity. Furthermore, maintaining a higher level of activity in adulthood was able to reduce the subjects' weight gain over their lifetime. Morano et al. (52) also found improved physical activity with a simultaneous decrease in BMI in children. The review by Brown et al. (49) showed that children aged six to 12 years and adolescents up to 18 years have a lower risk of obesity from physical activity alone. By adding physical activity, they could show a low to moderate reduction in BMI in the age from six to 18 years (49). In summary, the results of these studies and reviews suggest that higher levels of physical activity lead to lower weight gain or reduce the BMI of the subjects. This systematic review could confirm that physical activity can help to reduce BMI in obese and overweight students. Although not all studies of this systematic review show significant results, it can be concluded that physical activity can improve BMI in overweight and obese students in tertiary institutions. Accordingly, the systemic review can be integrated into the current state of research, as eleven out of 16 studies were able to demonstrate a significant reduction in BMI among overweight and obese students and only one study was unable to reduce the BMI of the participants through physical activity. Nonetheless, the prevalence of physical inactivity has risen sharply in recent years (54), even in early childhood and adolescence (53). As already mentioned, there are numerous interventions designed to counteract the risk factors of physical inactivity like excessive body fat percentage, overweight and obesity, unhealthy lipid profile or high blood pressure etc., through physical activity. All these risk factors could result in cardiovascular diseases, strokes, etc. or even in an increased morbidity (54, 55). Since physical inactivity is especially prevalent among university students, interventions designed to mitigate these risk factors are necessary. Interventions and reviews have already demonstrated that the above mentioned risk factors could be significantly reduced through physical activity both in children as in adults (49, 50, 53, 52). In addition to

BMI, the physical-activity-based interventions in this systematic review have shown further improvements in risk factors of physical inactivity. In line with the current literature, this systematic review could therefore demonstrate that physical activity is useful in reducing or even eliminating the risk factors associated with physical inactivity. Relaxation techniques should be used as an additional tool to implement the approach of comprehensive health promotion. Moore & Cunningham (56) stated in their systematic review that exposure to higher stress was associated with poorer nutrition and additionally higher body weight. Conversely, a higher level of stress could be an obstacle to weight loss. In addition, Stults-Kolehmainen & Sinha (57) were able to demonstrate in their systematic review, that objective stress (e.g. life events), as well as subjective stress (e.g. distress), impair the efforts to be physically active. On the other hand, physical activity is known to be stress-buffering, as regular physical activity can buffer the negative consequences of stress on health, which is postulated by the stress-buffering hypothesis (58, 59). Given that the target population of this systematic review was university students who are particularly exposed to high levels of stress (60, 61), relaxation strategies should be considered for a comprehensive treatment of the increasing prevalence of obesity, overweight and physical inactivity.

At the level of the individual studies used for this review, some limitations can be identified. Particularly, the insufficient and homogeneous sample size is frequently cited, which may impair the informative value of the results and makes generalization to the general population difficult. Another limitation arises from the inadequate control over the intensity in some studies. Therefore, all studies which did not control the intensity with objective indicators (e.g., heart rate) impair generalizability in terms of highest effectiveness according to the intensity. This issue could be additionally exacerbated by the varying adherence to the training program. Even though eleven of the included studies showed significant improvements after the intervention, it should be considered that the intervention periods of these studies lasted only up to four months. Therefore, it is not possible to determine any long-term effects of the conducted interventions. Concerning statistical analysis, only one of the included studies reported effect sizes in addition to the significance level (43). Since p-values always refer to the sample size in contrast to effect sizes (46), effect sizes should be reported more frequently to facilitate detecting patterns across studies and to infer the effectiveness of the included studies. Even though effect sizes could be computed with *Psychometrica* for the remaining studies, they can only be seen as approximate values. At the review level, it is important to note that the heterogeneity of the included studies in terms of sample size, gender, type and duration of physical activity, duration of the intervention, and BMI of subjects aggravated to identify patterns across the studies. Additionally only 31% of the included studies were of strong quality, so generalization should be taken with caution. Since there was no possibility to translate studies that were in a language other than English or German, the search was limited to the English language from the beginning. In addition, limited access to full text of nine potentially suitable studies made it difficult to select appropriate studies. Due to these organizational limitations, there is a possibility that relevant literature could not be included in this review. The choice of BMI as the dependent variable of the intervention proved to be the biggest limitation and bias. Frequently, studies had to be excluded because they reported on other body characteristics than BMI, such as body fat levels. As the relationship between BMI and body fat mass is nonlinear, individuals having the same BMI may have different body compositions (e.g. varying body fat percentage or muscle mass; 62). Thus, with regard to the effectiveness of physical activity, the main concern is whether a reduction in BMI is associated with a reduced fat mass at all. Or conversely, especially for novice exercisers, whether an increase in muscle mass does not initially translate into an increased BMI.

The present review was deliberately limited to the aspect of physical activity. But since a promising feature of the intervention studies by Brown et al. (49) and Martin et al. (53) examining children and adolescents is the implementation of a program combining

physical activity and dietary changes, future reviews should additionally include the aspect of nutrition. Additionally, as the BMI is prone to bias, it should be reconsidered as an appropriate variable to capture the change in body characteristics of a physical activity intervention for future reviews. Future research should ensure larger sample sizes as well as representativeness of the sample to facilitate generalization. To be able to assess long-term effects, longer intervention periods need to be conducted. Combining different training programs for future research might also be a promising approach for reaching better effects. Similarly, as university students are especially exposed to high levels of stress which in turn is known to impede weight loss, relaxation strategies should be considered in future interventions. Web-based approaches, using for example messaging or informative websites, and technical devices, such as activity trackers, are promising approaches to foster physical activity. They can not only facilitate sampling and flexibility as an organizational aspect but also help to supervise the intensity of physical activity during the implementation. To be able to set optimal stimuli for participants, more research and interventions with a study design based on a theoretical approach are needed to identify key concepts and to be able to better understand the mechanisms of change behind the intervention effects. In general, more intervention studies of physical activity in overweight and obese university students are needed to gain insight into optimal types of interventions, their frequency, and duration and to explore more contemporary and sustainable opportunities.

5. Conclusions

Since all studies except one showed some reduction in BMI with eleven of 16 studies reporting significant improvements, a tendency toward the effectiveness of the interventions was demonstrated. However, no clear patterns could be discerned in terms of study design, intensity, and frequency regarding the intervention methods showing significant results or the strongest effects. There was only an observable trend, that the type of physical activity may be critical, as aerobic exercises achieved the highest reduction in BMI. However, the university as a setting proves to be ideal for several reasons. Due to the rising number of students, a large population from different ethnic and socioeconomic backgrounds can be reached. Furthermore, within universities students are surrounded by experts, facilities, and resources that can facilitate building a healthy lifestyle. Given that overweight, obesity and physical inactivity is a highly relevant issue and their prevalence continues to increase, further high-quality, large-sample studies on this topic are needed to gain a comprehensive understanding of the effectiveness of weight loss interventions in students.

Author Contributions: Conceptualization, J.P., C.R., D.W., D.J. and P.B.; methodology, J.P., C.R. and D.W.; validation, J.P., C.R. and D.W.; formal analysis, P.B., D.J., K.W.; investigation, P.B., D.J., K.W.; resources, J.P., C.R. and D.W.; data curation, T.E. and C.N.; writing—original draft preparation, C.R., D.W.; writing—review and editing, J.P., C.R., D.W.; visualization, J.P., C.R., D.W.; supervision, P.B., D.J., K.W.; project administration, J.P., C.R., D.W.; All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: None.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Rebar, A. L.; Stanton, R.; Geard, D.; Short, C.; Duncan, M. J.; Vandelanotte, C. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychology Review* **2015**, *9* (3), 366–378. DOI: 10.1080/17437199.2015.1022901.

2. Reiner, M.; Niermann, C.; Jekauc, D.; Woll, A. Long-term health benefits of physical activity--a systematic review of longitudinal studies. *BMC public health* **2013**, *13*, 813. DOI: 10.1186/1471-2458-13-813.
3. Warburton, D. E. R.; Nicol, C. W.; Bredin, S. S. D. Health benefits of physical activity: the evidence. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* **2006**, *174* (6), 801–809. DOI: 10.1503/cmaj.051351.
4. WHO. *Physical activity*. <https://www.who.int/news-room/fact-sheets/detail/physical-activity>.
5. Chu, D.-T.; Minh Nguyet, N. T.; Dinh, T. C.; Thai Lien, N. V.; Nguyen, K.-H.; Nhu Ngoc, V. T.; Tao, Y.; Le Son, H.; Le, D.-H.; Nga, V. B.; Jurgoński, A.; Tran, Q.-H.; van Tu, P.; Pham, V.-H. An update on physical health and economic consequences of overweight and obesity. *Diabetes & metabolic syndrome* **2018**, *12* (6), 1095–1100. DOI: 10.1016/j.dsx.2018.05.004.
6. WHO. *Obesity and overweight*. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> (accessed 2021-03-03).
7. WHO. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet* **2004**, *363* (9403), 157–163. DOI: 10.1016/S0140-6736(03)15268-3.
8. Devaux, M.; Sassi, F.; Church, J.; Cecchini, M.; Borgonovi, F. Exploring the Relationship Between Education and Obesity. *OECD Journal: Economic Studies* **2011**, *2011* (1), 1–40. DOI: 10.1787/eco_studies-2011-5kg5825v1k23.
9. Peltzer, K.; Pengpid, S.; Samuels, T. A.; Özcan, N. K.; Mantilla, C.; Rahamefy, O. H.; Wong, M. L.; Gasparishvili, A. Prevalence of overweight/obesity and its associated factors among university students from 22 countries. *International journal of environmental research and public health* **2014**, *11* (7), 7425–7441. DOI: 10.3390/ijerph110707425.
10. ACHA. *American College Health Association - National College Health Assessment III: Undergraduate Student Reference Group*. Executive Summary Spring 2021 (accessed 2021-08-02).
11. Levitsky, D. A.; Halbmaier, C. A.; Mrdjenovic, G. The freshman weight gain: a model for the study of the epidemic of obesity. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* **2004**, *28* (11), 1435–1442. DOI: 10.1038/sj.ijo.0802776.
12. Brown, C. The information trail of the 'Freshman 15'--a systematic review of a health myth within the research and popular literature. *Health information and libraries journal* **2008**, *25* (1), 1–12. DOI: 10.1111/j.1471-1842.2007.00762.x.
13. Vadeboncoeur, C.; Townsend, N.; Foster, C. A meta-analysis of weight gain in first year university students: is freshman 15 a myth? *BMC obesity* **2015**, *2*, 22. DOI: 10.1186/s40608-015-0051-7.
14. Irwin, J. D. Prevalence of University Students' Sufficient Physical Activity: A Systematic Review. *Perceptual and Motor Skills* **2004**, *98*, 927–943. DOI: 10.2466/pms.98.3.927-943.
15. Keating, X. D.; Guan, J.; Piñero, J. C.; Bridges, D. M. A meta-analysis of college students' physical activity behaviors. *Journal of American college health : J of ACH* **2005**, *54* (2), 116–125. DOI: 10.3200/JACH.54.2.116-126.

16. Jakicic, J. M.; Rogers, R. J.; Collins, A. M.; Jackson, R. Strategies for Physical Activity Interventions in the Treatment of Obesity. *Endocrinology and metabolism clinics of North America* **2020**, *49* (2), 289–301. DOI: 10.1016/j.ecl.2020.02.004.
17. Samdal, G. B.; Eide, G. E.; Barth, T.; Williams, G.; Meland, E. Effective behaviour change techniques for physical activity and healthy eating in overweight and obese adults; systematic review and meta-regression analyses. *The international journal of behavioral nutrition and physical activity* **2017**, *14* (1), 1–14. DOI: 10.1186/s12966-017-0494-y.
18. Miguel-Etayo, P. de; Bueno, G.; Garagorri, J. M.; Moreno, L. A. Interventions for Treating Obesity in Children. *World review of nutrition and dietetics* **2013** (108), 98–106. DOI: 10.1159/000351493.
19. Plotnikoff, R. C.; Costigan, S. A.; Williams, R. L.; Hutchesson, M. J.; Kennedy, S. G.; Robards, S. L.; Allen, J.; Collins, C. E.; Callister, R.; Germov, J. Effectiveness of interventions targeting physical activity, nutrition and healthy weight for university and college students: a systematic review and meta-analysis. *The international journal of behavioral nutrition and physical activity* **2015**, *12*, 45. DOI: 10.1186/s12966-015-0203-7.
20. Liberati, A.; Altman, D. G.; Tetzlaff, J.; Mulrow, C.; Gøtzsche, P. C.; Ioannidis, J. P. A.; Clarke, M.; Devereaux, P. J.; Kleijnen, J.; Moher, D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS medicine* **2009**, *6* (7), e1000100. DOI: 10.1371/journal.pmed.1000100.
21. Schardt, C.; Adams, M. B.; Owens, T.; Keitz, S.; Fontelo, P. Utilization of the PICO framework to improve searching PubMed for clinical questions. *BMC medical informatics and decision making* **2007**, *7*, 16. DOI: 10.1186/1472-6947-7-16.
22. Lenhard, W.; Lenhard, A. *Psychometrica: Computation of effect sizes*; Lenhard, W.; Lenhard, A. https://www.psychometrica.de/effect_size.html.
23. Morris, S. B. Estimating Effect Sizes From Pretest-Posttest-Control Group Designs. *Organizational Research Methods* **2008**, *11* (2), 364–386. DOI: 10.1177/1094428106291059.
24. Cumming, G. *Understanding The New Statistics: Effect Sizes, Confidence Intervals, and Meta-Analysis*, 1st ed.; Routledge, 2011. DOI: 10.4324/9780203807002.
25. Effective Public Healthcare Panacea Project. *Quality Assessment Tool for Quantitative Studies Dictionary*. https://www.ehphp.ca/PDF/QADictionary_dec2009.pdf.
26. Popay, J.; Roberts, H.; Sowden, A.; Petticrew, M.; Arai, L.; Rodgers, M.; Britten, N.; Roen, K.; Duffy, S. *Guidance on the Conduct of Narrative Synthesis in Systematic Reviews: A Product from the ESRC Methods Programme*. https://www.researchgate.net/publication/308362339_Guidance_on_the_conduct_of_narrative_synthesis_in_systematic_reviews#fullTextFileContent (accessed 2021-08-16).
27. Plotnikoff, R. C.; Costigan, S. A.; Karunamuni, N.; Lubans, D. R. Social cognitive theories used to explain physical activity behavior in adolescents: a systematic review and meta-analysis. *Preventive medicine* **2013**, *56* (5), 245–253. DOI: 10.1016/j.ypmed.2013.01.013.
28. Conner, M., Norman, P., Eds. *Predicting health behaviour: Research and practice with social cognition models*, 2. ed.; Open Univ. Press, 2005.

29. Dalibalta, S.; Mirshafiei, F.; Davison, G. Exercise intervention on cardiovascular disease risk factors in a university population in the United Arab Emirates. *International journal of adolescent medicine and health* **2017**, *30* (6). DOI: 10.1515/ijamh-2016-0132.
30. DiFrancisco-Donoghue, J.; Jung, M.-K.; Stangle, A.; Werner, W. G.; Zwibel, H.; Happel, P.; Balentine, J. Utilizing wearable technology to increase physical activity in future physicians: A randomized trial. *Preventive medicine reports* **2018**, *12*, 122–127. DOI: 10.1016/j.pmedr.2018.09.004.
31. Gifari, N.; Hardinsyah, H.; Martianto, D.; Kuswari, M. Effect of high-intensity interval training and pre-meal water consumption on lipid profile in overweight and obese students. *Mal J Nutr* **2021**, *27* (1), 159–168. DOI: 10.31246/mjn-2020-0034.
32. Ha, C. H.; So, W. Y. Effects of Combined Exercise Training on Body Composition and Metabolic Syndrome Factors. *Iranian Journal of Public Health* **2012**, *41* (8), 20–26. DOI: 10.22334/jbhost.v6i2.217.s47.
33. Siqiang, G. Experimental study of aerobic exercise on the weight loss effect of obese female college students. *biomedicalresearch* **2018**. DOI: 10.4066/biomedicalresearch.29-17-828.
34. Wei-wei, Z.; Ling, Y. Study on the Exercise and Nutrition Intervention on Body Composition and Fitness of Female College Students with Invisible Obesity. *E3S Web Conf.* **2021**, *233*, 2038. DOI: 10.1051/e3sconf/202123302038.
35. Pacholek, M.; Zemková, E.; Arnolds, K.; Šagát, P. The Effects of a 4-Week Combined Aerobic and Resistance Training and Volleyball Training on Fitness Variables and Body Composition on STEAM Students. *Applied Sciences* **2021**, *11* (18), 8397. DOI: 10.3390/app11188397.
36. Chen, Y.; Ma, C.; Wang, J.; Gu, Y.; Gao, Y. Effects of 40% of Maximum Oxygen Uptake Intensity Cycling Combined with Blood Flow Restriction Training on Body Composition and Serum Biomarkers of Chinese College Students with Obesity. *International journal of environmental research and public health* **2021**, *19* (1). DOI: 10.3390/ijerph19010168.
37. Moravveji, A.; Sayyah, M.; Shamsnia, E.; Vakili, Z. Comparing the prolonged effect of interval versus continuous aerobic exercise on blood inflammatory marker of Visfatin level and body mass index of sedentary overweight/fat female college students. *AIMS public health* **2019**, *6* (4), 568–576. DOI: 10.3934/publichealth.2019.4.568.
38. Sun, J.; Cheng, W.; Fan, Z.; Zhang, X. Influence of high-intensity intermittent training on glycolipid metabolism in obese male college students. *Annals of palliative medicine* **2020**, *9* (4), 2013–2019. DOI: 10.21037/apm-20-1105.
39. Winters-Stone, K. M.; Medysky, M. E.; Stoyles, S.; Bumgarner, L.; Witzke, K. A brief whole-body vibration intervention to avoid weight gain in college students: A randomized controlled pilot trial. *Journal of American college health : J of ACH* **2020**, *1*–9. DOI: 10.1080/07448481.2020.1784179.
40. Joseph, R. P.; Casazza, K.; Durant, N. H. The effect of a 3-month moderate-intensity physical activity program on body composition in overweight and obese African American college females. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* **2014**, *25* (10), 2485–2491. DOI: 10.1007/s00198-014-2825-z.
41. Joseph, R. P.; Pekmezi, D.; Dutton, G. R.; Cherrington, A. L.; Kim, Y.-I.; Allison, J. J.; Durant, N. H. Results of a Culturally Adapted Internet-Enhanced Physical Activity Pilot Intervention for Overweight and Obese Young

- Adult African American Women. *Journal of transcultural nursing : official journal of the Transcultural Nursing Society* **2016**, *27* (2), 136–146. DOI: 10.1177/1043659614539176.
42. Lee, J.-S.; Yoon, E.-S.; Jung, S.-Y.; Yim, K.-T.; Kim, D.-Y. Effect of high-intensity circuit training on obesity indices, physical fitness, and browning factors in inactive female college students. *Journal of exercise rehabilitation* **2021**, *17* (3), 207–213. DOI: 10.12965/jer.2142260.130.
43. Mathunjwa, M. L.; Semple, S.; Du Preez, C. A 10-week aerobic exercise program reduces cardiometabolic disease risk in overweight/obese female African university students. *Ethnicity & disease* **2013**, *23* (2), 142–148.
44. Roopchand-Martin, S.; Nelson, G.; Gordon, C.; Sing, S. Y. A pilot study using the XBOX Kinect for exercise conditioning in sedentary female university students. *Technology and health care : official journal of the European Society for Engineering and Medicine* **2015**, *23* (3), 275–283. DOI: 10.3233/THC-150899.
45. Church, T. Exercise in obesity, metabolic syndrome, and diabetes. *Progress in cardiovascular diseases* **2011**, *53* (6), 412–418. DOI: 10.1016/j.pcad.2011.03.013.
46. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*, 2nd ed.; Lawrence Erlbaum Associates, 1998.
47. Goodpaster, B. H.; Delany, J. P.; Otto, A. D.; Kuller, L.; Vockley, J.; South-Paul, J. E.; Thomas, S. B.; Brown, J.; McTigue, K.; Hames, K. C.; Lang, W.; Jakicic, J. M. Effects of diet and physical activity interventions on weight loss and cardiometabolic risk factors in severely obese adults: a randomized trial. *JAMA* **2010**, *304* (16), 1795–1802. DOI: 10.1001/jama.2010.1505.
48. Hankinson, A. L.; Daviglius, M. L.; Bouchard, C.; Carnethon, M.; Lewis, C. E.; Schreiner, P. J.; Liu, K.; Sidney, S. Maintaining a high physical activity level over 20 years and weight gain. *JAMA* **2010**, *304* (23), 2603–2610. DOI: 10.1001/jama.2010.1843.
49. Brown, T.; Moore, T. H.; Hooper, L.; Gao, Y.; Zayegh, A.; Ijaz, S.; Elwenspoek, M.; Foxen, S. C.; Magee, L.; O'Malley, C.; Waters, E.; Summerbell, C. D. Interventions for preventing obesity in children. *The Cochrane database of systematic reviews* **2019**, *7*, CD001871. DOI: 10.1002/14651858.CD001871.pub4.
50. Corbett, D. B.; Fennell, C.; Peroutky, K.; Kingsley, J. D.; Glickman, E. L. The effects of a 12-week worksite physical activity intervention on anthropometric indices, blood pressure indices, and plasma biomarkers of cardiovascular disease risk among university employees. *BMC research notes* **2018**, *11* (1), 80. DOI: 10.1186/s13104-018-3151-x.
51. Romero-Blanco, C.; Rodríguez-Almagro, J.; Onieva-Zafra, M. D.; Parra-Fernández, M. L.; Del Prado-Laguna, M. C.; Hernández-Martínez, A. Physical Activity and Sedentary Lifestyle in University Students: Changes during Confinement Due to the COVID-19 Pandemic. *International journal of environmental research and public health* **2020**, *17* (18). DOI: 10.3390/ijerph17186567.
52. Morano, M.; Robazza, C.; Bortoli, L.; Rutigliano, I.; Ruiz, M. C.; Campanozzi, A. Physical Activity and Physical Competence in Overweight and Obese Children: An Intervention Study. *International journal of environmental research and public health* **2020**, *17* (17). DOI: 10.3390/ijerph17176370.
53. Martin, A.; Booth, J. N.; Laird, Y.; Sproule, J.; Reilly, J. J.; Saunders, D. H. Physical activity, diet and other behavioural interventions for improving cognition and school achievement in children and adolescents with obesity or overweight. *The Cochrane database of systematic reviews* **2018**, *1*, CD009728. DOI: 10.1002/14651858.CD009728.pub3.

-
54. WHO. *Physical activity*. https://www.who.int/health-topics/physical-activity#tab=tab_1 (accessed 2021-11-25).
 55. Cunningham, C.; O' Sullivan, R.; Caserotti, P.; Tully, M. A. Consequences of physical inactivity in older adults: A systematic review of reviews and meta-analyses. *Scandinavian journal of medicine & science in sports* **2020**, *30* (5), 816–827. DOI: 10.1111/sms.13616.
 56. Moore, C. J.; Cunningham, S. A. Social position, psychological stress, and obesity: a systematic review. *Journal of the Academy of Nutrition and Dietetics* **2012**, *112* (4), 518–526. DOI: 10.1016/j.jand.2011.12.001.
 57. Stults-Kolehmainen, M. A.; Sinha, R. The effects of stress on physical activity and exercise. *Sports medicine (Auckland, N.Z.)* **2014**, *44* (1), 81–121. DOI: 10.1007/s40279-013-0090-5.
 58. Cohen, S.; Wills, T. A. Stress, social support, and the buffering hypothesis. *Psychological bulletin* **1985**, *98* (2), 310–357. DOI: Review.
 59. Gerber, M.; Pühse, U. Review article: do exercise and fitness protect against stress-induced health complaints? A review of the literature. *Scandinavian journal of public health* **2009**, *37* (8), 801–819. DOI: 10.1177/1403494809350522.
 60. Regehr, C.; Glancy, D.; Pitts, A. Interventions to reduce stress in university students: a review and meta-analysis. *Journal of affective disorders* **2013**, *148* (1), 1–11. DOI: 10.1016/j.jad.2012.11.026.
 61. Wunsch, K.; Fiedler, J.; Bachert, P.; Woll, A. The Tridirectional Relationship among Physical Activity, Stress, and Academic Performance in University Students: A Systematic Review and Meta-Analysis. *International journal of environmental research and public health* **2021**, *18* (2). DOI: 10.3390/ijerph18020739.
 62. Rothman, K. J. BMI-related errors in the measurement of obesity. *International journal of obesity (2005)* **2008**, *32* Suppl 3, S56-9. DOI: 10.1038/ijo.2008.87.

Appendix A

Table A1: Study characteristics of the included studies

Source	Design & Theoretical Framework	Population*	Intervention	Duration & Frequency	Intensity
Chen et al. (2021), China	Randomized Controlled Trial	N = 37 [0/37] IG (n=18) CG (n=19)	IG: cycling combined with blood flow restriction training (BFRT) by wearing air pressure belts CG: cycling without BFRT	12 weeks (2x per week, 3 sessions/time with 15 min/session)	IG: low-intensity cycling (40% VO _{2max}) with BFRT exercises; 1-min rest between sessions CG: low-intensity cycling (40% VO _{2max}) without BFRT exercises; 1-min rest between sessions
Dalibalta et al. (2017), United Arab Emirates	Controlled Clinical Trial	N = 46 [40/14] IG (n=14) CG (n=32)	IG (BMI ≥ 25): Exercise program (not further defined) CG (BMI ≤ 25): Exercise program (not further defined)	8 weeks (3x per week with 60 min/session)	moderate-high intensity (not further defined)
DiFrancisco-Donoghue et al. (2018), USA	Controlled Clinical Trial	N = 113 [60/67] Fitbit Plus group (n=35) Fitbit Only group (n=38) CG (n=40)	Fitbit Plus group: wrist activity trackers, participation in weekly mentored walks/runs, weekly emails (Sunday) offering fitness challenges and feedback on step count Fitbit Only group: wrist activity trackers and instructions how to use them CG: No wrist activity trackers	39 weeks Fitbit Plus group: 10,000 steps daily (30-45 min walking)	not mentioned
Source	Design & Theoretical Framework	Population*	Intervention	Duration & Frequency	Intensity

Source	Design & Theoretical Framework	Population*	Intervention	Duration & Frequency	Intensity
Gifari et al. (2021), Indonesia	Controlled Clinical Trial	N = 27 [22/27] PWI (n=9) HIIT (n=9) PWI + HIIT (n=9)	PWI: pre-meal water consumption (PWI); 600 mL before each mealtime with a total water intake of 1.8 L/day HIIT: high-intensity interval training (HIIT) HIIT + PWI: combination of PWI and HIIT	8 weeks (3x per week with 18 min/session)	70-85% of HR-max
Ha & So (2012), Korea	Controlled Clinical Trial	N = 16 [16/16] IG (n=7) CG (n=9)	IG: Combination of aerobic exercise (treadmill running) and resistance exercise (whole body) CG: Maintain normal sedentary activities	12 weeks (3x per week with 80 min/session)	Treadmill running: 60-80% of HRR Resistance training: 3 sets of 10-15 RM
Joseph et al. (2014), USA	Cohort study (1 group & Pre-Post-Design)	N = 15 [15/15]	aerobic exercise (1) Walking an indoor track (2) Option between exercise on their own or participate in a cardiovascular-based group exercise (Zumba, kickboxing, aerobics)	12 weeks (4x per week with 30-60min/session)	moderate (50-70% of HR)
Joseph et al. (2016), USA	Cohort study (1 group & Pre-Post-Design) Social Cognitive Theory (SCT)	N = 25 [25/25]	An Internet-enhanced physical activity program consisted of 1) aerobic exercise and 2) an Internet-based application as a promotion tool (1) Walking an indoor track (2) Option between exercise on their own or participate in a cardiovascular-based group exercise (Zumba, kickboxing, aerobics)	12 weeks (4x per week with 30-60min/session)	moderate (50-70% of HR)

Source	Design & Theoretical Framework	Population*	Intervention	Duration & Frequency	Intensity
Lee et al. (2021), Korea	Cohort study (1 group & Pre-Post-Design)	N = 10 [10/not mentioned]	high intensity Circuit Training 40 min circuit training: 5 min warm-up, 30 min exercise (full body workout), 5 min cool-down	4 weeks (3x per week with 40 min/session)	Week 1-2: 60-70 % HRR (RPE Scale = 13-14) Week 2-4: 65-80 % HRR (RPE Scale = 14-18)
Mathunjwa et al. (2013), South Africa	Cohort study (1 group & Pre-Post-Design)	N = 60 [60/60]	Tae-bo (body fitness exercises combining the moves of taekwondo, karate, boxing and hip-hop dancing)	10 weeks (3x per week with 60min/session)	Week 1-5: moderate (Borg RPE Scale = 11-13, 135 bpm) Week 6-10: high (Borg RPE Scale = 14-16, 150 bpm)
Moravveji et al. (2019), Iran	Randomized Controlled Trial	N = 31 [31/31]; continuous group (n=10) interval group (n=12) CG (n=9)	1200m-run (each week increase by 400m) continuous group: 1200m without rest intervals interval group: 3 x 400m, interspaced with a walking period of 2 minutes CG: maintain their behavior, should not participate in training programs	8 weeks (3x per week)	60-75% of maximum HR
Pacholek et al. (2021), Saudi Arabia	Controlled Clinical Trial	N = 28 [0/not mentioned] COM (n=14) VOL (n=14)	COM: combination program (aerobic & resistance) VOL: volleyball program (isolated skill exercises, small-sided games & volleyball games)	4 weeks (4x per week with 50 min/session)	subjective feelings with maximal effort (not further defined)

Source	Design & Theoretical Framework	Population*	Intervention	Duration & Frequency	Intensity
Roopchand-Martin et al. (2015), Jamaica	Cohort study (1 group & Pre-Post-Design)	N = 24 [24/13]	Dance intervention utilizing the XBOX Kinect 360 and Just Dance 4	6 weeks (week 1-2: 5x per week with 30min/session; week 3-4: 4x per week with 45min/session; week 4-6: 3x per week with 60min/session)	Moderate (Borg RPE Scale = 12-14)
Siqiang (2018), China	Controlled Clinical Trial	N = 100 [100/100] IG (n=50) CG (n=50)	IG: aerobic exercise divided into two (i.e. rope skipping, swimming) CG: Intake of Calcium pyruvate (2x per day)	12 weeks (4x per week with 60min/session. Divided into two: half an hour in the morning, half an hour in the afternoon)	Moderate (120-150 bpm)
Sun et al. (2020), China	Randomized Controlled Trial	N = 300 [0/300] IG (n=150) CG (n=150)	IG: High-intensity interval training (not further defined) CG: aerobic exercise (not further defined)	12 weeks (5x per week)	IG: 4 minutes with 85% VO _{2max} /HR=174 bpm then 2 minutes with 50% VO _{2max} , followed by 5 minutes relaxation (repeated 5x) CG: 40 minutes with 60% of VO _{2max} /HF 140=bpm
Wei-Wei & Ling (2021), China	Controlled Clinical Trial	N = 60*** [60/60] CG (n=20) IG (n=20) IG + nutrition** (n=20)	IG: combination of aerobic exercises and resistance training IG + nutrition: resistance training and peer nutrition health education method CG: maintain their behavior	16 weeks (3x per week with 60min/session)	aerobic exercises: 30-80% of max HR, according to stage of the training resistance training: 65-70% of 1-RM, 3 rounds, repetitions not further defined

Winters-Stone et al. (2020), USA	Randomized Controlled Trial	N = 77	IG: Whole Body Vibration (WBV) training CG: Shouldn't change their physical activity or dietary habits	24 weeks (3x per week with 20min/session)	week 1-3: 30 Hz
		[58/not mentioned]			week 4: 35 Hz
		IG (n=37)			week 5: 40 Hz
		CG (n=40)			week 6: 45 Hz
					week 7-24: 50 Hz

Sample size contains the number of subjects included in the final analysis. IG = Intervention group. CG = Control group. HR = Heart Rate. HRR = Heart Rate Reserve. RM = Repetition Maximum. RPE = Rating of Perceived Exertion. VO_{2max} = maximum oxygen uptake. bpm = beats per minute.* Population characteristics are divided into the number of female and number of overweight people [number of female/number of overweight]. **

The results of this group is not considered in Table A3, because of the inclusion of a nutrition intervention. *** Since no drop-outs were reported, the exact number of participants can only be assumed

Table A3: Individual results of the included studies (change of BMI)

Study	Controlgroup		Difference pre-post (SD)	Difference pre-post (%)	ES / Significance	Interventiongroup(s)		Difference pre-post (SD or CI)	Difference pre-post (%)	ES / Significance
	Baseline	Postintervention				Baseline	Postintervention			
	Mean BMI (SD)	Mean BMI (SD)				Mean BMI (SD)	Mean BMI (SD)			
Chen et al. (2021)	30.30 (1.08)	29.90 (1.44)	-0.40	-1.32	$d_{av} = .317$	30.10 (0.95)	29.0 (1.79)	-1.10	-3.65	$d_{av} = .803^*$
Dalibalta et al. (2017)	21.60 (2.05)	21.10 (2.00)	-0.50	-2.31	$d_{av} = .247^*$	28.70 (3.27)	28.10 (2.79)	-0.60	-2.09	$d_{av} = .198^*$
DiFrancisco- Donoghue et al (2018)			Men: -0.30 (0.50) Women: -0.20 (1.20)					Fitbit-Plus men: 0.20 (1.70) Fitbit-Plus women: -0.50 (0.50) Fitbit-Only men: -0.10 (2.10) Fitbit-Only women: 0.40 (2.30)		
Gifari et al. (2021)						PWI: 26.6 (3.6) HIIT: 25.9 (2.4) HIIT + PWI: 25.7 (3.4)	PWI: 26.7 (3.6) HIIT: 25.8 (2.4) HIIT + PWI: 25.2 (3.2)	PWI: 0.1 HIIT: -0.1 HIIT + PWI: -0.05	PWI: 0.38 HIIT: -0.38 HIIT + PWI: -1.95	PWI: $d_{av} = .028$ HIIT: $d_{av} = .042$ HIIT + PWI: d_{av} = .152*
Ha & So (2012)	24.18 (1.63)	24.38 (1.66)	0.20	0.83		24.97 (2.73)	24.76 (3.01)	-0.21	-0.84	$d_{ppc2} = .179$
Joseph et al. (2014)						33.27 (6.53)	32.57 (6.62)	-0.70	-2.10	$d_{av} = .106^*$
Study	Controlgroup					Interventiongroup(s)				

	Baseline		Difference pre-post (SD)	Difference pre-post (%)	ES / Significance	Baseline		Difference pre-post (SD or CI)	Difference pre-post (%)	ES / Significance
	Mean BMI (SD)	Postintervention Mean BMI (SD)				Mean BMI (SD)	Postintervention Mean BMI (SD)			
Joseph et al. (2016)						33.80 (5.70)	32.90 (6.00)	-0.90	-2.66	$d_{av} = .154$
Lee et al. (2021)						25.14 (2.51)	24.24 (2.60)	-0.9 (0.35)	-3.58	$d_{av} = .352^*$
Mathunjwa et al. (2013)						32.26 (5.65)	30.11 (5.46)	-2.15	-6.66	$d = .2^* (1)$
Moravveji et al. (2019) (2)	27.57 (1.62)	28.04 (1.46)	0.47	1.70		Continuos Group: 26.77 (1.53)	Continuous Group: 26.08 (1.21)	Continuos Group: -0.69	Continuous Group: -2.58	Continuos Group: $d_{ppc2} = .704^* (2)$
						Interval Group: 27.57 (1.02)	Interval Group: 26.82 (0.98)	Interval Group: -0.75	Interval Group: -2.72	Interval Group: $d_{ppc2} = .896^* (2)$
Pacholek et al. (2021)						COM: 31.30 (8.61) VOL: 26.50 (7.71)	COM: 31.30 (8.36) VOL: 26.20 (7.45)	COM: 0 VOL: -0.3	COM: 0 VOL: -1.13	COM: $d_{av} = 0$ VOL: $d_{av} = .04$
Roopchand- Martin et al. (2015)						30.51 (5.18)	30.31 (5.39)	-0.2	-0.66	$d_{av} = .038$
Siqiang (2018)	26.80 (3.40)	24.10 (2.90)	-2.70	-10.07	$d_{av} = .857^*$	25.50 (2.70)	21.60 (3.10)	-3.90	-15.29	$d_{av} = 1.345^*$
Study	Controlgroup			Interventiongroup(s)						

	Baseline	Postintervention	Difference	Difference	ES / Significance	Baseline	Postintervention	Difference	Difference	ES / Significance
	Mean BMI (SD)	Mean BMI (SD)	pre-post (SD)	pre-post (%)		Mean BMI (SD)	Mean BMI (SD)	pre-post (SD or CI)	pre-post (%)	
Sun et al. (2020)	30.54 (3.86)	28.45 (2.76)	-2.09	-6.84	$d_{av} = .631^*$	30.85 (3.79)	28.17 (2.89)	-2.68	-8.69	$d_{av} = .802^*$
Wei-Wei & Ling (2021) (3)	22.54 (0.71)	22.68 (1.11)	0.14	0.62		22.06 (0.98)	21.64 (0.85)	-0.42	-1.90	$d_{ppc2} = .641^*$
Winters- Stone et al. (2020)	28.27 (26.48;30.07)	not available (4)	0.64 (0.31;0.96)	2.26		Total: 28.24 (23.87;32.60) High Adherence: 27.02 (22.34;31.71) (5)	not available (4)	Total: 0.22 (-0.59; 1.03) High Adherence: -0.01 (-1.63; 0.86)	Total: 0.78 High Adherence: -0.04	Total: $p = .094$ High Adherence: $p = .026^*$

Determined level of significance $p = .05$ (* $p < .05$). SD = Standard Deviation. ES = Effect size (d_{av} sensu Cumming, 2012; d_{ppc2} sensu Morris, 2008). (1) Mathunjwa et al. (2013) reported effect sizes by their own (2) Inconsistent reported significance in table 2 and text. (3) nutrition group is excluded. (4) Winters-Stone et al. (2020) reported only a 6-month change. (5) Participants with high adherence ($\geq 80\%$ of prescribed sessions).

