
Review

Historic and Prehistoric Epidemics: An Overview of Sources Available for the Study of Ancient Pathogens

Antoinette C. van der Kuyl^{1,2}

¹ Amsterdam UMC location University of Amsterdam, Department of Medical Microbiology and Infection Prevention, Laboratory of Experimental Virology, Meibergdreef 9, Amsterdam, The Netherlands; a.c.vanderkuyl@amsterdamumc.nl; Tel.: +31-205-666-778

² Amsterdam Institute for Infection and Immunity, Infectious diseases, Amsterdam, The Netherlands

Abstract: Since life on earth developed, parasitic microbes have thrived. Increases in host numbers, or the conquest of a new species, provides an opportunity for such a pathogen to enjoy, before host defense systems kick in, a similar upsurge in reproduction. Outbreaks – caused by ‘endemic’ pathogens – and epidemics – caused by ‘novel’ pathogens – have thus been creating chaos and destruction since prehistorical times. To study such (pre)historic epidemics, recent advances in the ancient DNA field, applied to both archeological and historical remains, have helped tremendously to elucidate the evolutionary trajectory of pathogens. These studies offered new and unexpected insights in the evolution of, for instance, smallpox virus, hepatitis B virus and the plague-causing bacterium *Yersinia pestis*. Furthermore, burial patterns and historical publications can help in tracking down ancient pathogens. Another source of information is our genome, where selective sweeps in immune-related genes relate to past pathogen attacks, while multiple viruses have left their genomes behind for us to study. This review will discuss the sources available to investigate (pre)historic diseases, as molecular knowledge of historic and prehistoric pathogens may help us understand the past, the present, and prepare us for future epidemics.

Keywords: prehistory; history; archeobiology; paleovirology; EVEs; ancient DNA; paleogenomics; pathology collections; historic publications

1. Introduction

The most obvious evidence of parasites preying on earthly life forms can be found in our chromosomes, where various viruses have left their genomes as a testimony of past infections (see [1]). Proof of the infectivity of these so-called endogenous viruses can be found in recent and/or ongoing invasions of the host germ line, for instance in mice, cats, and koalas, as well as in ancient between-hosts viral transmission (reviewed in [2-4]), and the resurrection of endogenous human retroviruses to their infectious counterparts [5,6]. In addition, selection in immunity- and infection-related host genes illustrates the ongoing battle with pathogens, although of course here the causative pathogen is likely to remain anonymous.

Ancient burial patterns also can display evidence of past infectious disease outbreaks, especially when there is no evidence of external violence to the remains. When many die around the same time, people are either carefully buried together in a same grave, or bodies are hastily thrown into quickly dug graves. Likewise, since historic times, our ancestors have put pen to paper to describe epidemics affecting society, for instance, the ancient Greek described the multiple plagues of their times, while an inscription of the local word for pestilence on 1338-1339 CE century tomb stones, in what is now Kyrgyzstan, pointed to the likelihood of plague victims being buried there [7]. Another example can be found in the relatively accurate descriptions of a smallpox-like disease can be found in Chinese texts dating to the Sui and Tang dynasties, 581-907 CE [8]. Such information can be important to understand past epidemics, or to identify places to hunt for ancient pathogens, but the sources themselves are less helpful in identifying the infectious causes, as they rely

heavily on clinical symptoms, which are rarely diagnostic, and surely do not provide evidence for the involvement of a particular variant.

Although all of the above information is important in understanding our past, it was not until the development of ancient DNA techniques and understanding of DNA degradation mechanisms, that the actual detection of ancient pathogens in (pre)historical samples became feasible. Some studies focus on PCR detection of a specific pathogen in a historical sample, guided by clinical symptoms such as skin eruptions in a mummy suggestive of smallpox [9], mass burials coinciding with described plague epidemics [10], the plague of Athens [11] or the likelihood of infection in victims interred during the 1918 influenza pandemic [12]. Nowadays, modern sequencing techniques enable the retrieval of partial or complete ancient pathogen genomes (reviewed in [13]). In addition, genomic shotgun data often contain pathogen DNA, enabling identification and assembly of viral and bacterial genomes, as exemplified for instance by hepatitis B virus (HBV) DNA in Neolithic, Bronze Age and medieval sequence samples [14-16].

This review will emphasize the importance of studying past pathogens by giving an overview of sources available to researchers for the study of historic and prehistoric diseases, and associated epidemics, rather than providing an exhaustive overview of ancient pathogen studies. Coping with the struggles of retrieving and authenticating ancient DNA, as well as with sequence analysis of ancient microbial genomes, are the subjects of some excellent, recent reviews [13,17-19].

2. Materials and Methods

The PubMed database (www.ncbi.nlm.nih.gov/pubmed), Google Scholar (scholar.google.nl), and Google Books (books.google.nl) were manually searched with appropriate terms such as "ancient DNA", "archeovirology", "endogenous retrovirus", "EVE", "paleogenomics", and eventually a second term appropriate to the topic, such as "pathogen", "historic", "prehistoric", "evolution", "dating", etc. Abstracts were inspected for relevance; from the selected abstracts, only those of which a full-length publication could be retrieved were read and included. In addition, references in the retrieved papers were inspected and used when appropriate. The purpose of the current paper is to provide a comprehensive overview of the field, not an exhaustive review, so not all papers retrieved will be discussed.

3. Results

As a strict definition of the terms "pandemic", "epidemic", and "outbreak" does not exist, but a more general description of all three would be "a sudden increase in disease cases above the normal, endemic, disease burden in a certain area, community or population", either caused by a novel pathogen, an increased transmissibility or virulence of an established pathogen or any other important variation in the behavior of the pathogen, or an increase in host susceptibility or exposure (see also [20]). To reduce confusion, the term 'epidemic' will be used here as a synonym for the historical concepts of "plague" and "pestilence", and as a synonym of the term "outbreak". It must be noted that not all ancient pathogens are thus epidemic pathogens in a strict sense, and that epidemic pathogens can persist and thus become endemic, while endemic pathogens often are able to cause — small — outbreaks. Examples of the latter would be influenza virus, human parvovirus B19, measles virus, and *Yersinia pestis* in medieval and early modern times. Twenty-first century endemic pathogens, which are widespread, have been introduced in the human population a long time ago, and cause few or no outbreaks at present, would for instance be *Plasmodium* species — causing malaria —, intestinal parasites, Mycobacteria — causing tuberculosis and leprosy —, *Helicobacter pylori*, HBV, and herpesviruses. For an overview of the sources of ancient pathogens discussed below, see Table 1.

Table 1. Sources available for the study of ancient pathogens and historical epidemics.

Source	What can be retrieved? ¹	What is difficult to determine?	Level of study
Contemporary genomic DNA	Germ line infiltrating (retro)viruses	Disease symptoms	Pathogen, transmission dynamics
	Prevalence of protective alleles	Pathogen	Indirect evidence of epidemic
Archeological & historical sites	Bone and teeth samples for DNA analysis; burial patterns may assist in selection	Pathogen, disease symptoms	Epidemic
Archeological & historical samples	Ancient DNA	Disease symptoms ² Prevalence	Pathogen
Pathology collections	Ancient DNA & RNA	Prevalence	Pathogen
Historical publications	Outbreaks & disease symptoms	Pathogen identification	Epidemic

¹ Approximate age of the infection can be estimated from all sources

² Bone or skin lesions due to chronic *Mycobacterium* or *Treponema* infection, however, can reveal pathology. Chronic infections with *Plasmodium* spp. may induce hyperostotic skull lesions ('pitting'), which are, however, not diagnostic.

3.1 Contemporary genomic DNA as a source of ancient pathogens and epidemics

Genomes from living organisms contain the remnants of the ongoing battle between pathogens and hosts, both as integrated viruses and a selective sweeps in immune-related genes. To look for an easy-accessible source of the most ancient pathogens, all we have to do to study those carefully propagated leftovers of the past, is to sequence and analyze the corresponding DNA stretches in modern chromosomes. For the older viruses, those invading host genomes over 50-100 million years ago (mya), most of their genomes are likely to be only useful in retracing evolution by descent, rather than informing us on putative transmissions or epidemics caused by those parasites. However, when sufficient endogenous viral elements (EVEs) are present in a species, or in related species to reconstruct a detailed evolutionary history, their past transmission patterns may become visible, especially since more and more complete genomes are being sequenced and are accessible from public databases. Comparison of Simian Endogenous Retrovirus (SERV) genomes in Old World monkey species, for instance, showed that this virus was spreading among African and Asian monkeys less than 8 million years ago (mya) [3], while tracking down single-stranded RNA virus-related integrations suggested that around 40 mya, ancestral mammals suffered from a heavy load of borna- and filovirus infections [21,22]. Of course, virulence and pathology cannot be estimated from endogenous viral genomes alone, although similarity to modern pathogens could suggest it. In theory, transmission of ancient viruses could have resulted in little or no morbidity and mortality, which would then disqualify them from being the cause of bona fide epidemics. However, witnessing recent, successful novel introductions of microbes into human populations or domestic animal populations provide ample evidence of those microorganisms being real pathogens, with at least part of a population experiencing disease. It is not likely that this would have been different in the past.

Modern genomes can likewise show evidence of ancient battles by rapid local evolution, or by selection, positive or negative, of existing alleles from genes involved in either im-

munity or in the replication cycle of ancient parasites. Examples are, for instance, the acquisition and fast evolution of the APOBEC3G gene in primates (reviewed in [23]), and the dominance of a dysfunctional Toll-like receptor 1 variant associated with protection against leprosy in people of European descent [24]. A 32-base pair deletion in the CCR5 gene, already found in prehistoric Europeans, has likewise been attributed to such a selective sweep, although the causative pathogen remains a mystery after bubonic plague was ruled out [25-27]. However, selection of certain immunity-related alleles, such as those encoding Ficolin-2, NLRP14, and HLA-DRB1*13, involved in the response to pathogenic bacteria, was obvious from a genetic comparison of thirty-six 16th century plague victims buried in Ellwangen, Germany, with modern inhabitants of that city [28]. It is therefore likely that the plague pandemics did influence the genetic make-up of surviving populations. High carrier frequencies for pyrin – an inflammasome protein – mutations seen in Mediterranean populations have indeed been associated with resistance to *Y. pestis* [29]. Other allelic selections associated with epidemic pathogens are for instance the highly prevalent FUT2 – an $\alpha(1,2)$ fucosyltransferase involved in H type 1 blood group antigen synthesis – non-secretor alleles with resistance to noro- and rotavirus infections [30-32], IFITM3 alleles – encoding an interferon-inducible transmembrane protein – with influenza morbidity and mortality [33], DARC gene – encoding a chemokine receptor – variation with malaria resistance [34], a tyrosine kinase 2 (TYK2) polymorphism with clinical TB [35], and IFNL4 – encoding an antiviral interferon – pseudogenization with hepatitis C virus clearance [36]. In a Bangladeshi population, multiple genomic regions were found to be selected in association with cholera susceptibility [37]. These stretches for instance encode potassium channel genes, nuclear factor κ B (NF- κ B) signaling components and proteins of the NF- κ B/inflammasome-dependent pathway [37]. Glucose-6-phosphate dehydrogenase (G6PD) deficiency, beta thalassemia- and sickle-cell disease inducing mutations, prevalent in Africa and the Mediterranean, are all linked to malaria resistance (reviewed in [38]). For a more complete overview of pathogen-induced selection on human genetics, see: [39,40].

Due to improved technology and decreased costs, the number of complete genomes is steeply increasing nowadays, which will greatly enhance our knowledge on ancient host-microbe battles.

3.2 Archeological and historical sites as sources of ancient pathogens and epidemics

Archeological and historical burial sites can be good sources to look for patterns of epidemics and their causative pathogens in the distant past. For instance, a sudden increase in the number of individuals interred, or a high frequency of adolescents and young adults among the deceased, abnormal burials as demonstrated by multiple or mass graves, bodies stocked in layers, the use of lime, un-coffined burials, and/or other sloppy burial practices can all point to a major disaster, possibly in the form of an epidemic, especially when there is no evidence of physical trauma [41,42]. Such locations have been termed “mortality crisis” graves [43]. Grave artifacts, such as coins and clothing, ¹⁴C dating, and written evidence in the form of tombstones, or parish records, can be used to date the burial grounds. For instance, three mass burials discovered in the center of Ellwangen, Germany, concerning a total of 101 skeletal remains buried closely together, with other evidence, such as the majority being children, a characteristic of later plague epidemics, suggested that the site harbored 16th-17th century plague casualties [44]. Indeed, the presence of *Yersinia pestis* DNA was confirmed in dental pulp extracts [44]. The remains were also used to investigate genetic variation of the victims [28]. In a plethora of other studies, teeth and bone samples from putative 14th-18th plague “pits” in many locations in Eurasia have been confirmed to contain *Y. pestis* DNA [7,43-47]. From the 17th century onwards, direct written evidence for plague epidemics and dedicated graveyards is common [7,43,47]. Retrieval of *Y. pestis* DNA from victims of earlier epidemics, such as the 6th century ‘Justinian plague’, also relied on evidence for local mortality crises, such as when a “striking number” of double and multiple burials was seen in a grave-yard in Bavaria, Germany, collective burials were

found in Valencia, Spain, and an existing ditch was used as an emergency cemetery near Bourges in France [48,49]. In contrast, burial patterns in Neolithic and Bronze Age graves have not been found to be useful in this regard, likely due to a lower population density, so that samples from those periods are usually tested at random for pathogens [13,50-52]. Similarly, in Africa, where mass burial sites, which could represent plague pits, have not been found yet, it has been speculated that two- to fourfold simultaneous interments may already be indicative of epidemic disaster, as could be a population reduction resulting in abandonment of settlements, infrastructure decay, or fluctuations in industrial activities [53,54]. Of course, such developments could easily be attributed to other causes, but epidemics should certainly be considered when no war or other type of major violence is imminent.

In contrast to plague emergency burials, for other common medieval diseases, such as leprosy, dedicated leprosaria had been erected from medieval times onwards to house and isolate the large numbers of patients. Leprosy, or Hansen's disease, is caused by *Mycobacterium leprae*. It runs a chronic, slow course, leaving sufficient time to bury the dead peacefully in cemeteries within the institution. Ancient DNA analysis of remains from medieval European leprosaria, resulted in retrieval of partial and complete genomes of *M. leprae*, with surprisingly high genetic diversity [55-57]. Complete *M. leprae* genomes had likewise been retrieved from bones collected from a 10th-12th century British leprosarium [58], confirming that dedicated hospital sites are fertile hunting grounds for research on pathogen evolution. Evidence from ancient remains for another chronic disease, namely tuberculosis (TB), which is caused by another Mycobacterial species complex, *Mycobacterium tuberculosis*, also shows substantial worldwide genetic diversity of this ancient pathogen in the past compared with modern isolates [59-62]. Detection of *M. tuberculosis* DNA and mycolic acids from a double burial of a ~25-year old female and a ~12-month old infant dating to around 9000 years BCE in what is now Israel, demonstrate that infection with the bacterium could be rapidly fatal in the past [63]. The child's skeleton showed extensive bone pathology indicative of disseminated neonatal TB, to which young children are highly susceptible when infected by someone with contagious pulmonary TB [63].

Other locations where emergency burials might be expected are battlefields and sites of temporary army settlements. There, death from infectious diseases is equally likely as dying from violence, as military camps are, and have been, sites of mass crowding, stress, poor hygiene, poor nutrition, as well as the gathering of soldiers from different backgrounds. In the absence of battle injuries, mass graves near army quarters may be an indication of an epidemic of some sort. Local archives, for instance, suggested that mass graves excavated near Vilnius, Lithuania, contained the remains of soldiers from Napoleon's Great Army, retreating from Moscow in December 1812 [64]. The deceased had been quickly interred, usually before rigor mortis had set in, and were buried close to each other. DNA from the louse-borne pathogens *Bartonella quintana*, the causative agent of trench fever, and of typhus-inducing *Rickettsia prowazekii*, was found in up to one-third of the soldiers, with the first also being found in three lice samples, indicating that these diseases had had a great impact on those surviving the combats of Napoleon's Russian campaign [64]. DNA of the above mentioned louse-borne pathogens, was also found in soldiers buried in a mass grave near the besieged city of Douai in northern France, dating to 1710-1712 CE [65], while *B. quintana* and also *Y. pestis* DNA were retrieved from two French multiple burial sites in Bondy, dating to the 11th-15th centuries CE, and putatively containing plague victims buried closely together without coffins [66]. Finding ancient DNA from louse-borne diseases in mass graves together with *Y. pestis* DNA, has led to the hypothesis that in medieval times, the spread of human louse- and not rat flea-borne *Y. pestis* was the source of the plague epidemics [67]. DNA from another louse-borne pathogen, *Borrelia recurrentis*, the bacterial cause of relapsing fever, was detected in dental pulp from an individual buried in a 15th century CE double grave in Oslo, Norway [68], while paleo-serology

was able to identify seven *Borrelia* infections, most likely *B. recurrentis*, as *Borrelia* spp. inducing Lyme disease do not cause epidemics, and one *B. quintana* seropositive tooth from a 16th-17th century military garrison cemetery in Auxi-le-Château, France [69], again indicating that louse-borne pathogens were common in historical Europe.

Lack of war injuries, a contemporary chronicle of a 'pestilence' suffered by both besiegers and besieged during an attack on Barcelona, Spain, in 1652, and the presence of both shallow graves as well as large pits with dozens hastily buried with their boots still on, led to the discovery of *Salmonella enterica* Paratyphi C DNA in one of the victims, suggesting enteric fever could have been the cause of the documented pestilence, and not *Yersinia pestis*, as assumed earlier [70]. A related *S. enterica* Paratyphi C strain had been identified before as the likely cause of an epidemic in colonial Mexico during 1545-1550 CE, where a local cemetery witnessed "catastrophic mortality", demonstrated by stacks of corpses, and an overrepresentation of adolescents and young adults [71,72]. Historical accounts from medieval Lübeck, now in Germany, recorded two epidemics in 1350 CE and 1359 CE, which were likely Black Death outbreaks, together with at least four other unspecified pestilences occurring there during the 14th century (see: [73]). Subsequent sequence analysis of the remains found in two multiple burial pits next to the, still in use, Heiligen Geist Hospital, showed *S. enterica* Paratyphi C in eight of them, implicating enteric fever as a cause of death [73]. Interestingly, no *S. enterica* Paratyphi C DNA was found in teeth and bones from two adjoining mass burial graves, which are supposed to be slightly older, neither were reads corresponding to *Y. pestis* DNA, leaving the question open as to what caused the death of those victims [73]. Virulent precursor strains to the modern enteric fever-causing *Salmonellae* may already have been involved in ancient epidemics, as the detection of DNA of the so-called *S. enterica* Para C lineage was found in 3000 BCE Bronze Age remains from multiple simultaneous burials, mainly of infants, children and young adults, in a cemetery near Xinjiang, China [74]. The first documented case of *S. enterica* Paratyphi C in Europe dates to 800 years ago, when a young female, who most likely died of enteric fever, was buried in Norway [75]. Her single burial in cemetery near a church, however, did not suggest an ongoing epidemic.

So, investigation of historical sites, guided by abnormal burial patterns is a fruitful way of biological sample selection, with the associated benefit that there is already support for the epidemic potential of any pathogen retrieved. A disadvantage is that a mortality crisis to guide ancient DNA research is mainly apparent in densely populated regions, implying a focus on more recent times. However, in the permafrost in northeastern Siberia, an unusual burial of five bodies in a single grave drew the attention of archeologists as individual interment was the standard practice in late 17th - early 18th century Yakutia [76]. Variola virus (VARV) DNA fragments could be retrieved from one of the corpses, suggesting that these people had died from smallpox, and that the simultaneous deaths indeed did cause some sort of a crisis in their community.

3.3 Archeological and historical samples as a source of ancient pathogens

The dawn of ancient DNA research several decades ago, saw molecular biologists begging for pieces of valuable or, at least in the beginning, not so valuable, archeological and historical specimens to serve as a possible source of ancient DNA. Pathogen DNA retrieved from such unique objects cannot on its own cover epidemics, but it can significantly contribute to our knowledge of the earliest features and evolution of microbes known to be able to cause epidemics. (Electron) microscopy, targeted PCR amplification, employing microarrays, and microbial reads being present as a byproduct of deep sequencing have all been used to detect ancient microbes or their DNA in (pre)historic samples. Identification of a pathogen can already give valuable information on its appearance in history, but to learn something about virulence and genetic variation, more information is needed. For-

tunately, with the advance of ancient DNA techniques, generating complete or near-complete ancient pathogen genomes is increasingly feasible (for a review, see [13]). The results from such studies are already changing our view on ancient pathogens and epidemics. For instance, using multiple specimens collected from variable periods and geography, it was shown for *Y. pestis*, *M. leprae*, *M. tuberculosis*, VARV and HBV that the genetic variation of these pathogens was much larger in the past.

Variant strains of *Treponema pallidum*, the bacterial cause of syphilis, yaws and bejel (endemic syphilis), depending on the subspecies, were shown to be widespread in early modern times in both Europe and Colonial Mexico [77,78]. As a result, the mystery, as to what was the cause of a late 15th century epidemic in Europe, described in 16th century medical literature as an acute, severe venereal disease, has not yet been solved, yet. Was it due to the introduction of *T. pallidum pallidum* from the New World or elsewhere, or due to the introduction of a novel strain of that bacterium, or was the disease not syphilis at all, as, until the 19th century, the various venereal diseases were often confused with each other and with leprosy, while the symptoms described in those early manuscripts are very different from the chronic, relatively mild disease that syphilis soon became [78-81]. As *T. pallidum* sequences have been successfully obtained from both bone and teeth, it is likely that in the near future more samples will be located so that further bacterial genomes can be added to help solve the puzzle. Especially analyzing a well-dated sample from the early years of the epidemic, 1495~1545 CE, would help significantly.

Interesting results were also obtained after sequencing ancient VARV, the causative agent of smallpox, which was shown to have arisen as late as the 16th or 17th century, as VARV isolated from earlier samples, dating to the ~600-1050 CE, belonged to a different viral clade, which was not a direct ancestor to the later VARV [82]. It is not clear whether or not this Viking-age VARV did cause similar lethality as smallpox, as its genome did not have the gene reductions associated with virulence in orthopoxviruses. However, as VARV does not cause chronic infections, it was suggested that the victims positive for VARV DNA had died during an acute infection, which would be indicative of a fatal disease associated with the virus [82]. VARV sequences detected in lung tissue from one in five family members buried together in a single emergency grave in the Siberian permafrost in the late 17th or early 18th century, suggest that the virus by then was able to cause severe disease, although skin pustules were not observed, and phylogenetic analysis suggested that the virus belonged to a progenitor strain to modern VARV [76]. For HBV, many more viral genotypes were found in Neolithic and medieval samples than are currently known [14-16]. Also, the modern geographical distribution of HBV genotypes was found to be significantly different from that in the past [15,16]. So, this currently widespread virus, like the extinct VARV, also had a much broader distribution and genetic variation than estimated from modern samples.

Another virus with a worldwide distribution and a long-term association with humans is parvovirus B19 (B19V), a highly-infectious, blood-borne DNA virus capable of inducing multiple severe pathological conditions besides being the agent of the fifth disease in children. Regional prevalence of B19V varies, and outbreaks of the virus are common. After having first been identified in 70-year old bones from Finland [83], B19V DNA was also found in ancient dental and skeletal remains from Eurasia and Greenland dating to ~500-6900 years ago, confirming a lengthy presence in humans [84]. Three B19V genomes, reconstructed from dental remains present in mass graves near a colonial hospital in Mexico City, which had been used to bury epidemic victims during the 16th-18th centuries, illustrate an association of the virus with deadly epidemics in the past [85].

Human samples used to isolate ancient pathogen DNA from range from bone, teeth – both dental pulp and dental plaque –, dried/mummified skin and other – wet, fixed or frozen – tissue samples, feces, and even chewed birch pitch. Gut contents and paleofeces,

also known as coprolites, have been used to detect both DNA and microscopic evidence of intestinal parasite infections, for instance, *Ascaris* eggs were found in stool from a latrine, and in mummies dating to 14th century Europe, and 16th-17th century South Korea [86,87]. Helminth eggs and DNA were even detected in sediment samples taken from under the pelvic area of skeletons [88-93]. In addition, paleofeces can be used to reconstruct the intestinal microbiome and the intestinal virome, while dental plaque can offer insights in the oral microbiome [94-101]. A not much studied, but interesting source of ancient pathogens could possibly be found in collections of ancient Egyptian canopy jar holders, in which internal organs, and later even heads, from the deceased were preserved during the mummification process [102]. Canopy jars, containing liver, lung, stomach and intestine, have been part of the ancient Egyptian funerary practice since approximately 2600 BCE until the Roman period [102]. Indeed, calcified *Schistosoma haematobium* eggs have been identified in mummy liver, pointing to a long-term presence of bilharzia in Egypt [102,103].

With all samples, showing the presence of pathogens is the first step, but complete or near-complete microbial genomic information, ideally from multiple samples, is needed to understand the pathogen's evolution and role in ancient epidemics. Especially for bacteria, virulence genes should be identified, as bacteria without those are generally benign, so that the presence of a particular species does not necessarily indicate pathogenicity. For instance, the acquisition of virulence factors plays an important role in the evolution of *Y. pestis*, developments which have been retraced helped by ancient pathogen DNA studies. Investigation of bacterial genomes from various archeological sites showed the common ancestor of historic and current Black Death-causing *Y. pestis* strain originates from 14th century Central Asia, in what is now Kyrgyzstan [7]. Around 1338-1339 CE, there, a low-complexity variant equipped with a few extra virulence factors, as compared to its Neolithic predecessors, entered human society, never to leave again [7,43,50,51,104].

3.4. Pathology collections as a source of historical pathogens

Pathology collections, containing actual body parts, tissue or blood samples of those who lived less than a few centuries ago, the oldest of which can be found in museums, and the more recent in biobanks, medical archives or hospital freezers, are a valuable source when looking for historical pathogens in "wet" samples. As such collections are likely to contain only the most interesting specimens, it is improbable that epidemics can be reconstructed by analyzing their samples, yet information regarding the genetic make-up of an example microbe from a documented outbreak can be very valuable. For instance, analysis of an intestinal sample collected from a 1849 victim of the second cholera pandemic showed that *Vibrio cholerae* in the specimen indeed belonged to, as assumed, the classical O1 biotype, which was replaced in the 20th century by the El Tor O1 biotype [105]. Both a plasma sample and a lymph node biopsy from 1959 and 1960, respectively, stored in what is now the Democratic Republic of the Congo, have been used to amplify human immunodeficiency virus type 1 (HIV-1) genomic fragments [106,107]. The results indicated that the virus was not only definitely present by then, but had also diversified into the genetic subtypes seen at present, implying that its origin in humans dates back to at least the early 20th century. Formalin-fixed, paraffin-embedded lung tissue from victims of the 1918 influenza virus pandemic, as well as lung tissue biopsies taken from those buried in permafrost, have likewise suggested the probable origin — novel reassortment of avian and human flu viruses —, characteristics — hemagglutinin (HA) and polymerase genes contributed to its virulence — and genetic variation — evidence for early adaptation to humans in HA — from the causative agent of the deadliest pandemic in human history [12,108-111]. Influenza virus sequences obtained from a formalin-fixed, ethanol-preserved North American wild goose collected in 1917, however, suggested that the 1918 HA segment was not directly acquired from birds, but had likely already circulated for some time in humans or other mammals [112]. Poliovirus (PV) RNA fragments were retrieved from fifty-year old formalin-fixed, paraffin-embedded spinal cord samples of Norwegian poliomyelitis fatalities dating to 1951-1952 [113]. Subsequent sequencing suggested that the outbreak could most likely be

attributed to wild type PV1. A next RNA virus sequenced from fixed autopsy material is measles morbillivirus (MeV), one of the most infectious viruses known. As MeV infection induces live-long sterilizing immunity, the virus needs large groups of naïve hosts so as to sustain itself. Although an effective vaccine is available, measles continues to break out, as it did in the past. After retrieval of an almost complete MeV genome from a 1912 formalin-fixed lung specimen, Düx et al. calculated the entry of MeV, likely from a bovine precursor, into the human population as early as the 6th century BCE, a time period that saw the rise of big cities, supplying the virus with its needed number of targets [114].

In recent years, RNA and DNA extraction from formalin- or alcohol-fixed samples, which has been much more challenging than nucleic acid isolation from frozen or fluid samples as formaldehyde crosslinks nucleic acids and proteins, has been improved, potentially unlocking archival pathology collections for routine infectious diseases testing [115-121].

Table 2. Ancient pathogen nucleic acid retrieved from human sources.

Biological source	Pathogen characteristics	Ancient pathogen detected	Reference
Bone	- Blood-borne - Targeting bone*	HBV	[9,15]
		<i>Mycobacterium leprae</i>	[55,56,58]
		<i>Mycobacterium tuberculosis</i>	[59-61,63]
		Parvovirus B19	[84]
		<i>Salmonella enterica</i> Paratyphi C	[75]
		<i>Treponema pallidum</i>	[77,78,122]
		<i>Yersinia pestis</i>	[123]
Dried tissue, such as mummified skin or organs	- Tissue tropism	VARV	[82]
		<i>Helicobacter pylori</i>	[124,125]
		HBV	
		Human papilloma virus, HPV	[126]
Frozen/fixed – wet – tissue, incl. blood	-Tissue tropism	<i>Mycobacterium tuberculosis</i>	[62,127,128]
		VARV	[129,130]
		HCV	[131]
		<i>Helicobacter pylori</i>	[132,133]
		HIV	[106,107]
		Influenza virus	[108,109,111]
		Measles virus	[114]
Tooth, dental pulp	- Blood-borne	Poliovirus	[113]
		VARV	[76]
		<i>Vibrio cholerae</i>	[105]
		<i>Bartonella quintana</i>	[64-66,134,135]
		<i>Bordetella pertussis</i>	[136]
		<i>Borrelia recurrentis</i>	[68]
		<i>Haemophilus influenzae</i>	[137]
		HBV	[14,15]
		<i>Mycobacterium leprae</i>	[55,56]
		<i>Mycobacterium tuberculosis</i>	[61]
		Parvovirus B19	[84,85]
<i>Rickettsia prowazekii</i>	[64,65]		
<i>Salmonella enterica</i> Paratyphi C	[70,72,73,75]		
<i>Treponema pallidum</i> spp.	[78]		
<i>Yersinia pestis</i>	[10,44,47,48,50,138]		

		VARV	[82]
Tooth, dental calculus	- Presence in oral cavity	<i>Mycobacterium leprae</i> Oral microbiome, including pathogens	[139] [95-97,100,140-143]
Paleofeces	-Tropism for gastro-intestinal tract	Intestinal microbiome, including pathogens	[94,96,98,99,101,140,144-147]
		Helminths	[86,87]
Other: Birch pitch	- Presence in the oral cavity/expression in saliva	Epstein Barr virus, EBV (human gammaherpesvirus 4)	[148]
Other: Calcified nodules	- Presence in genitourinary tissue	<i>Gardnerella vaginalis</i> <i>Staphylococcus saprophyticus</i>	[149]
	-Presence in abdominal/pelvic tissue	<i>Brucella melitensis</i>	[150]

* Blood-borne pathogens can also end up in bone, where nucleic acids adsorb to, and are protected by, apatite crystals composing the mineral part of bones (and teeth) [151].

3.5. Historical publications as a source of past epidemics

From the texts and inscriptions of the early human civilizations till contemporary scientific publications, ever since the invention of writing, people have documented evidence of disease and disaster. For instance, the ancient Greek tragedian Sophocles and historian and general Thucydides have described the Plagues of Thebes (430-420 BCE) and Athens (430-426 BCE), respectively, while accounts of the Justinian Plague (541-543 CE) destroying the Roman empire, appeared in Arabic, Greek, Latin and Syriac works [152]. Several ancient Egyptian “medical” papyri, such as the Ebers, Smith, and Hearst papyri, describe disease symptoms and epidemics which have been interpreted to refer to current pathogens [153]. For —critical— reviews of ancient epidemics in literature, see: [153-155].

Outnumbering the ancient writings mentioning epidemic diseases are present papers speculating on the causative microorganisms from these accounts. However, not only are disease symptoms hardly diagnostic, translating or interpreting historical descriptions may be equally challenging. For instance, the Septuagint (LXX) and the Masoretic Text (MT), used in Old Testament translations, differ significantly as to the account of the plague of Ashdod (1190, or 1141 BCE), described in Samuel 5:6-12. The coincident arrival of mice — or rats? — is not described in MT, although later on in both texts, five golden rats — or mice — have to be supplied as a guild offering to free, both young and old, town’s inhabitants from the plague [156]. Disease symptoms of that plague are nowadays translated as tumors — or boils — of the groin, which together with the rats would make an excellent case for bubonic plague as a putative cause (discussed in [157]). However, would those rats be mice, the case would be much weaker. Regarding pathology, the original texts mention “boils” or “sores” on the anus [156], which is quite unlike *Y. pestis* infection as we know it.

Descriptions of the Plague of Athens and the Justinian Plague inspired researchers to explore their actual cause. After analysis of three teeth samples from a mass burial pit at Kerameikos, Greece, the likely cause of the first was determined to be typhoid fever due to *Salmonella enterica* infection [11]. That claim was refuted by others based on subsequent phylogenetic analysis of the initial results [158]. Teeth samples from individuals buried in Bavaria, Germany at the time of the Justinian epidemic, which affected large parts of Europe, were confirmed by two ancient DNA laboratories to contain DNA from an early,

independent, lineage of *Yersinia pestis*, making it the first plague outbreak in history [48,138]. Ethiopian sources from the 13th-15th centuries also do contain references to epidemics, possibly of plague, one of which wiped out so many people, that “no one was left to bury the death” [53,159]. Arabic, Egyptian, Syriac and Sudanese texts likewise refer to epidemic diseases, which may or may not have been plague [53]. The 16th century Mexican “catastrophe” burials, which tested positive for *S. enterica* serovar Paratyphi C DNA, were backed up by evidence from Aztec painted manuscripts, the Codex Telleriano-Remensis and the Codex en Cruz, in which drawings of stacks of corpses and of figurines displaying symptoms of nose bleeds (epistaxis) and a body rash, respectively, are believed to represent an epidemic starting around 1544 in New Spain [71]. Indeed, epistaxis, and a so-called ‘rose spot’ rash are clinical symptoms of enteric fever, although they are not the most prominent ones. Age, geography, and the specific causative organism are known to widely influence the signs and complications of enteric fever [160,161].

From around the 16th century onwards, parish registers in Europe kept track of baptisms, marriages, and deaths, and can be used as sources of unexpected or increased mortality in a neighborhood. Often, the churchyards related to the registers are known, and could be a source of ancient DNA, although many of them, and the skeletal remains, have been destroyed over time. Sometimes, records provide the cause of death, for instance, death registers from Bergen op Zoom, the Netherlands, provide that information from 1779 onwards. From that source, for instance, we learn that there was a particular severe outbreak of smallpox in the summer of 1790, with more than half of fatalities in July and August of that year being due to young children dying from it [162]. In the winter of 1797-1798, smallpox raged again, with even higher mortality this time. In the autumn of 1807, there was a deadly epidemic of something called “nervous fever” (*Febris nervosa epidemica*), most likely enteric fever, to which about 45% of adult deaths were attributed. Interestingly, no casualties were seen in children. Likewise, from the 16th century onwards, medical doctors in Europe and the USA started to prepare lists and publications on health and disease from their own observations. Many 19th century publications were, for instance, devoted to cholera, then a new disease that was spreading around the world and causing deadly pandemics (see [163]). From medical publications listing the annual diseases prevalent in town and villages, it is obvious that in these times of poverty, crowding, malnutrition and lack of vaccines, infectious diseases were common. A problem with the historical descriptions, however, is that the names of the maladies, or the symptoms listed, do not always ring a bell in our present times. *Rubeola*, *scarlatina* and *febres intermittentes* likely indicate measles, scarlet fever and malaria, respectively, but what conditions would nowadays best describe *morbi inflammatorii pectoris* and *erysipelas intestinorum*, as given in 19th century publications [164]. In addition, biological samples connected to the reports are likely to be rare. An exception is the analysis of a wet intestinal sample derived from a 1849 victim of the second cholera pandemic in the USA, from which a complete genome of *Vibrio cholerae* has been reconstructed [105].

In general, historical publications and other written evidence can be helpful in tracing pathology, sites and samples of past epidemics, but as those texts are not easy to translate into current medical language, and there is usually no direct connection with human biological samples, there is only limited use for them in ancient pathogen research.

3.5.1 A historical enigma: Sweating sickness of 15th and 16th century Europe

An epidemic that came and went between 1485 and 1551 is one given the name of ‘English sweating sickness’, ‘sweating disease’ or ‘Sudor Anglicus’ [165]. Its first appearance was indeed in London, England after the battle of Bosworth. Four more episodes followed in 1508, 1517, 1529 and 1551, of which the 1529 outbreak reached far into mainland Europe. The disease was also described from the Netherlands and Belgium in 1489 [165]. Between

the major epidemics, cases were reported from several European countries [165]. Symptoms included high fever, headache, hyperhidrosis, palpitations, stomach ache, dyspnea, madness, and many more. The estimated mortality rate was 30-50% [165,166]. John Caius of Cambridge, then president of the Royal College of Physicians, published in 1552 an account of the sweating disease, the first monograph entirely devoted to an infectious disease (see [167]). Over the years, the causative agent of sweating disease has been proposed, and mostly dismissed, to be *Bacillus anthracis* (the cause of anthrax), *Vibrio cholerae*, influenza virus, or a hantavirus, amongst others [165,166,168-171]. Cheshire et al. have argued that the etiological agent of sweating disease would likely be a virus affecting the autonomic nervous system [166]. Interestingly, autonomic dysfunction, sometimes with hyperhidrosis, is a prominent feature of acute SARS-CoV-2 infection [172,173]. Rhabdoviruses, a virus family to which for instance rabies virus belongs, may likewise be a possibility to be considered. Bovine ephemeral fever virus (BEFV), an arthropod-borne rhabdovirus, for instance induces rapid pulse and tachycardia accompanied by high fever and respiratory symptoms in cattle, commonly affecting adults over young animals, and fat bulls over lean steers [174,175]. Listlessness, paralysis, coma and death may occur [174]. Little known rhabdoviruses, like members of the genus tibrovirus, are increasingly being detected in humans [176,177]. Serological surveys in African populations have shown high exposure rates to rhabdoviruses [178], illustrating that such viruses have probably been spreading for some time.

As for samples, John Caius describes in his monograph the death and burial in 1551 of the Duke of Suffolk, Henry Brandon, aged 15, and his brother Charles, aged 13-14, due to sweating disease. As their graves in Buckden, Cambridgeshire, England, are still there, the remains could possibly be subjected to ancient DNA analysis. However, current techniques are not optimized yet for ancient RNA analysis from bone or teeth [179], should sweating disease have been caused by an RNA virus. In contrast to most rhabdoviruses, tibroviruses are able to cause a viremia, allowing them in theory to be isolated from dental pulp. Another opportunity to define a cause for sweating disease could perhaps be found in victims of the "Picardy sweat" or "suette miliare", a disease with some resemblance to the English sweating disease, which caused 196 outbreaks in Europe between 1718 and 1874, when the illness similarly disappeared [171]. However, no samples belonging to Picardy sweat patients have been traced yet, although a few names of victims are known.

4. Discussion

Investigating epidemic pathogens of the past has already resulted in changing views on the history of infectious diseases and on the evolution of microbes. For instance, it has been confirmed that most of the "catastrophe" burials in Eurasia in the past 700 years or so were indeed due to fatal *Yersinia pestis* infections, making *Y. pestis* without doubt the most prominent epidemic pathogen in history, at least on that continent. A second important epidemic pathogen turned out to be *Salmonella enterica* serovar Paratyphi C and precursor lineages. Enteric fever epidemics due to *S. Paratyphi C* were seen in Bronze Age China, in medieval northern Europe and in 16th century Mexico and Spain [70,72-74]. Whereas plague is now limited to around ~100-3000 cases per year, infections caused by *S. Typhi*, *S. Paratyphi A*, *B*, and *C*, the only four of the ~2600 *S. enterica* serovars able to infect systemically, are still a major problem in low income countries, with *S. Typhi* and *S. Paratyphi A* causing over 14 million cases annually [161,180]. The Para C lineage likely arose in Neolithic Europe from pig *Salmonellae* [75,181], and has thus been able to maintain its virulence in humans over millennia. It is, however, less prevalent at present, which could be partly due to host genetic selection, for instance at the HLA-DRB1 locus [182,183]. Next major historical pathogens, at least in Europe, proved to be louse-borne microbes, namely the agents of epidemic typhus, trench fever, relapsing fever, and possibly plague, which has been speculated to have been transmitted by lice, too [67]. Louse-borne diseases essentially affect the poor, and thrive during war, crowding, and famine. In the ancient DNA

studies, trench fever-inducing *Bartonella quintana*, and typhus-inducing *Rickettsia prowazekii* were mainly found in association with the military, illustrating that the living conditions in garrisons and during army campaigns in history offered plenty of opportunities for lice to breed and spread. Both pathogens continue to cause problems to this day, as for instance exemplified by a 1997 large outbreak of epidemic typhus and trench fever in Burundi, a 1998 epidemic typhus outbreak in a Russian mental nursing home, and a high prevalence of *B. quintana* infections seen among the homeless in the USA [184-186].

Regarding the sources for the study of these historic epidemics, burial patterns have been proven to be highly informative to locate pathogens. Multiple burials, with signs of a crisis and without evidence of physical damage to the remains, were often found to be associated with epidemic pathogens in Europe and Asia. The number of people per grave, from two to dozens, likely correlates with the local population density, so that a double burial can already indicate epidemic disaster in a lowly populated area. The biological sources that are found in graves are usually limited to bones and teeth, implying that the pathogens involved should be targeting bone, or be present in blood. The pathogens listed with bone and dental pulp sources (Table 2), for instance, *Y. pestis* and *S. enterica* Paratyphi C, are indeed able to cause a bacteremia. But the absence of sequences of these two bacterial species in one of the largest mass burial graves from 14th century Lübeck [73] suggests that another pathogen could be involved in these epidemics. One that may not be present in blood at all. Looking at current pandemic pathogens, such as human immunodeficiency virus (HIV), Ebola virus (EBOV), and SARS-CoV-2, the first two viruses do target immune system cells, but the latter can hardly be found in blood, and would not likely be picked up should our descendants ever investigate the surplus of burials dating to 2020-2022 CE. Another problem with putative SARS-CoV-2, and EBOV, retrieval is that they are RNA viruses. Ancient RNA techniques, particularly when isolating RNA from sources other than “wet” tissue, are still under development. However, remarkable results, including the sequences of RNA virus genomes from 750-1000 years old grains and maize cobs, have already been published [179,187-189]. Human genomes from all over the world show strong selective sweeps resulting from extensive adaptations to RNA virus infections, suggesting that this class of pathogens was as important in past epidemics as it is now [190]. Therefore, it is important for the study of (pre)historic epidemics that ancient RNA techniques are further developed so that they can be reliably deployed. DNA viruses left a much less significant imprint on the human genome [190]. In contrast to bone and teeth, from so-called “wet” samples, liquid, fixed or frozen, either stored in freezers or in permafrost, should be able to deliver nucleic acids from any pathogen targeting that tissue or organ. Unfortunately, such samples are in general not much older than ~300 years, except for permafrost remains, and so cannot routinely be used to track epidemics from a distant past.

Although burial patterns are a good predictor of where to find evidence for (pre)historic epidemics, and dental pulp is an excellent source of ancient DNA, the recovery of pathogen genomes is limited to those present in blood, or targeting bone. Samples of specific tissue or organs, which can be used to find other pathogens, are in general much younger, so that their use is limited. Collections of Egyptian canopy jars, containing mummified organs, or the dried organs present in some —natural— mummies, could be a putative source of DNA from pathogens with a tissue tropism other than blood or bone. However, success in extracting DNA from such sources is generally lower, and RNA virus recovery is currently a challenge from most samples. Similarly, tracking pathogens in our genome will result in finding remnants of viruses that have the capacity to enter the germ line or the early embryo (see [191]). Those will mainly have DNA genomes and express integrases, in other words, they will chiefly be retroviruses. Any evidence of true RNA or non-integrating DNA viruses is fragmentary at best, and may hardly suffice to define prehistoric epidemics. For instance, fragments of borna- and filoviruses, but not complete genomes, are part of vertebrate genomes [21,192]. Such EVEs can only tell us that at the time of integration,

there must have been plenty of borna- and filoviruses around, but details are hard to come by after so much time.

At present, burial patterns in relation to epidemics have been mostly investigated in Eurasia, but it is not unlikely that the increased detection of large settlements, for instance, the pre-Hispanic sites discovered in South and Middle America [193,194], will provide fertile new hunting grounds for pathogens. At one Amazonian site, 103 burials were recently discovered, though none of them was reported to show signs of emergency [194]. Still, large parts of the world are still uncharted territory with regard to (pre)historic epidemics. For instance, very little research has been done in Africa, Australia, and North America. More limitations apply when combining burials with pathogens, as for instance cultures that do not bury their dead cannot be investigated this way.

Historical documents can be used to trace places of interest, or be used to add details, but clinical descriptions of diseases are in general too unspecific or too indecipherable to be of any real use. Furthermore, there are far more descriptions of epidemics in the last two millennia than there will ever be archeological sites. On the other hand, some sites with mass graves have already been discovered, but are still lacking a pathogen, such as the victims from an unknown epidemic in 4-5th century Florence, and 14th century Lübeck [73,89].

Paleopathology has been used to guide ancient pathogen research as well, with sometimes surprising outcomes. For instance, a mid-16th century Italian child mummy with skin pustules suggestive of smallpox, was found to contain not VARV, but HBV sequences. This finding implied that the child had died from another condition, possibly Gianotti-Crosti syndrome, although that is, as far as we know, a self-limiting malady [9]. In contrast, a 300-year old Siberian mummy did harbor VARV reads, but there was no evidence of skin lesions, while for the mummy of Egyptian pharaoh Ramses V, which displays extensive pustules on especially its face and lower abdomen, no VARV has been found to date [76,153]. So, relying on skin pathology, which in the recent past always was a clear sign of smallpox, proved to be problematic when trying to connect pathology and pathogen in historic specimens.

Next to providing time and location, pathology descriptions are one of the main features of old epidemical accounts. However, because clinical symptoms are mostly too general to be diagnostic of a certain disease, and interpretation of historic writings is often debatable, those descriptions are only useful as additional information at best. Another drawback is that medical symptoms as a result of infection may change over time as host and pathogen adapt, so that what we see now may not be identical to historic disease expression. Similarly, patterns of mortality may change when immunity is retained in the survivors of a first wave. Infections may then turn into childhood diseases, a pattern seen for instance in later plague epidemics [44]. In contrast, older age is an important risk factor for mortality in coronavirus epidemics, such as COVID-19, where the middle aged and the elderly have died in much larger numbers than the younger generations [195,196]. Such a skew in demographics may be important to keep in mind when investigating burial sites. Interestingly, the English sweating disease was reported to have killed preferentially middle aged men, with few deaths observed in children, women and the elderly [166,169].

A prerequisite for the advancement of our knowledge of ancient diseases, is that collaborations between archeologists and molecular biologists should be increasingly sought and perpetuated, since, despite some struggles, both fields have much to gain from each other [197,198]. Digitizing medical and museum collections may also open up these sources of old pathogens to global researchers [111]. The current development of non-destructive techniques can play an elementary role in gaining access to these resources [199].

Funding: This research received no external funding.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Katzourakis, A. Editorial overview: Paleovirology: the genomic fossil record, and consequences of ancient viral infections. *Current opinion in virology* **2017**, *25*, ix-xi, doi:10.1016/j.coviro.2017.08.002.
2. Katzourakis, A.; Gifford, R.J. Endogenous viral elements in animal genomes. *PLoS genetics* **2010**, *6*, e1001191, doi:10.1371/journal.pgen.1001191.
3. van der Kuyl, A.C. Contemporary distribution, estimated age, and prehistoric migrations of Old World monkey retroviruses. *Epidemiologia* **2021**, *2*, 46-67, doi:10.3390/epidemiologia2010005.
4. Chiu, E.S.; VandeWoude, S. Endogenous retroviruses drive resistance and promotion of exogenous retroviral homologs. *Annu Rev Anim Biosci* **2021**, *9*, 225-248, doi:10.1146/annurev-animal-050620-101416.
5. Dewannieux, M.; Harper, F.; Richaud, A.; Letzelter, C.; Ribet, D.; Pierron, G.; Heidmann, T. Identification of an infectious progenitor for the multiple-copy HERV-K human endogenous retroelements. *Genome research* **2006**, *16*, 1548-1556, doi:10.1101/gr.5565706.
6. Lee, Y.N.; Bieniasz, P.D. Reconstitution of an infectious human endogenous retrovirus. *PLoS pathogens* **2007**, *3*, e10, doi:10.1371/journal.ppat.0030010.
7. Spyrou, M.A.; Musralina, L.; Gneccchi Ruscone, G.A.; Kocher, A.; Borbone, P.G.; Khartanovich, V.I.; Buzhilova, A.; Djansugurova, L.; Bos, K.I.; Kühnert, D., et al. The source of the Black Death in fourteenth-century central Eurasia. *Nature* **2022**, 10.1038/s41586-022-04800-3, doi:10.1038/s41586-022-04800-3.
8. Zhang, Y.M.; Niu, Y.H. [Understanding of smallpox in ancient China]. *Zhonghua Yi Shi Za Zhi* **2021**, *51*, 294-301, doi:10.3760/cma.j.cn112155-20200607-0008.
9. Patterson Ross, Z.; Klunk, J.; Fornaciari, G.; Giuffra, V.; Duchêne, S.; Duggan, A.T.; Poinar, D.; Douglas, M.W.; Eden, J.S.; Holmes, E.C., et al. The paradox of HBV evolution as revealed from a 16th century mummy. *PLoS pathogens* **2018**, *14*, e1006750, doi:10.1371/journal.ppat.1006750.
10. Drancourt, M.; Aboudharam, G.; Signoli, M.; Dutour, O.; Raoult, D. Detection of 400-year-old *Yersinia pestis* DNA in human dental pulp: an approach to the diagnosis of ancient septicemia. *Proceedings of the National Academy of Sciences of the United States of America* **1998**, *95*, 12637-12640, doi:10.1073/pnas.95.21.12637.
11. Papagrigorakis, M.J.; Yapijakis, C.; Synodinos, P.N.; Baziotopoulou-Valavani, E. DNA examination of ancient dental pulp incriminates typhoid fever as a probable cause of the Plague of Athens. *Int J Infect Dis* **2006**, *10*, 206-214, doi:10.1016/j.ijid.2005.09.001.
12. Basler, C.F.; Reid, A.H.; Dybing, J.K.; Janczewski, T.A.; Fanning, T.G.; Zheng, H.; Salvatore, M.; Perdue, M.L.; Swayne, D.E.; García-Sastre, A., et al. Sequence of the 1918 pandemic influenza virus nonstructural gene (NS) segment and characterization of recombinant viruses bearing the 1918 NS genes. *Proceedings of the National Academy of Sciences of the United States of America* **2001**, *98*, 2746-2751, doi:10.1073/pnas.031575198.
13. Spyrou, M.A.; Bos, K.I.; Herbig, A.; Krause, J. Ancient pathogen genomics as an emerging tool for infectious disease research. *Nat Rev Genet* **2019**, *20*, 323-340, doi:10.1038/s41576-019-0119-1.
14. Krause-Kyora, B.; Susat, J.; Key, F.M.; Kühnert, D.; Bosse, E.; Immel, A.; Rinne, C.; Kornell, S.C.; Yepes, D.; Franzenburg, S., et al. Neolithic and medieval virus genomes reveal complex evolution of hepatitis B. *eLife* **2018**, *7*, doi:10.7554/eLife.36666.
15. Mühlemann, B.; Jones, T.C.; Damgaard, P.B.; Allentoft, M.E.; Shevnina, I.; Logvin, A.; Usmanova, E.; Panyushkina, I.P.; Boldgiv, B.; Bazartseren, T., et al. Ancient hepatitis B viruses from the Bronze Age to the Medieval period. *Nature* **2018**, *557*, 418-423, doi:10.1038/s41586-018-0097-z.
16. Kocher, A.; Papac, L.; Barquera, R.; Key, F.M.; Spyrou, M.A.; Hübner, R.; Rohrlach, A.B.; Aron, F.; Stahl, R.; Wissgott, A., et al. Ten millennia of hepatitis B virus evolution. *Science (New York, N.Y.)* **2021**, *374*, 182-188, doi:10.1126/science.abi5658.

17. Bos, K.I.; Kühnert, D.; Herbig, A.; Esquivel-Gomez, L.R.; Andrades Valtueña, A.; Barquera, R.; Giffin, K.; Kumar Lankapalli, A.; Nelson, E.A.; Sabin, S., et al. Paleomicrobiology: diagnosis and evolution of ancient pathogens. *Annu Rev Microbiol* **2019**, *73*, 639-666, doi:10.1146/annurev-micro-090817-062436.
18. Arning, N.; Wilson, D.J. The past, present and future of ancient bacterial DNA. *Microb Genom* **2020**, *6*, doi:10.1099/mgen.0.000384.
19. Duchêne, S.; Ho, S.Y.W.; Carmichael, A.G.; Holmes, E.C.; Poinar, H. The recovery, interpretation and use of ancient pathogen genomes. *Curr Biol* **2020**, *30*, R1215-r1231, doi:10.1016/j.cub.2020.08.081.
20. Rosenberg, C.E. What is an epidemic? AIDS in historical perspective. *Bull Hist Med* **2020**, *94*, 563-577, doi:10.1353/bhm.2020.0082.
21. Belyi, V.A.; Levine, A.J.; Skalka, A.M. Unexpected inheritance: multiple integrations of ancient bornavirus and ebolavirus/marburgvirus sequences in vertebrate genomes. *PLoS pathogens* **2010**, *6*, e1001030, doi:10.1371/journal.ppat.1001030.
22. Kawasaki, J.; Kojima, S.; Mukai, Y.; Tomonaga, K.; Horie, M. 100-My history of bornavirus infections hidden in vertebrate genomes. *Proceedings of the National Academy of Sciences of the United States of America* **2021**, *118*, doi:10.1073/pnas.2026235118.
23. Uriu, K.; Kosugi, Y.; Ito, J.; Sato, K. The battle between retroviruses and APOBEC3 Genes: Its past and present. *Viruses* **2021**, *13*, doi:10.3390/v13010124.
24. Wong, S.H.; Gochhait, S.; Malhotra, D.; Pettersson, F.H.; Teo, Y.Y.; Khor, C.C.; Rautanen, A.; Chapman, S.J.; Mills, T.C.; Srivastava, A., et al. Leprosy and the adaptation of human toll-like receptor 1. *PLoS pathogens* **2010**, *6*, e1000979, doi:10.1371/journal.ppat.1000979.
25. Hummel, S.; Schmidt, D.; Kremeyer, B.; Herrmann, B.; Oppermann, M. Detection of the CCR5-Delta32 HIV resistance gene in Bronze Age skeletons. *Genes Immun* **2005**, *6*, 371-374, doi:10.1038/sj.gene.6364172.
26. Kremeyer, B.; Hummel, S.; Herrmann, B. Frequency analysis of the delta32ccr5 HIV resistance allele in a medieval plague mass grave. *Anthropol Anz* **2005**, *63*, 13-22.
27. Vargas, A.E.; Cechim, G.; Correa, J.F.; Gomes, P.A.; Macedo, G.S.; de Medeiros, R.M.; Perotoni, G.; Rauber, R.; Villodre, E.S.; Chies, J.A. Pros and cons of a missing chemokine receptor--comments on "Is the European spatial distribution of the HIV-1-resistant CCR5-D32 allele formed by a breakdown of the pathocenosis due to the historical Roman expansion?" by Eric Faure and Manuela Royer-Carenzi (2008). *Infect Genet Evol* **2009**, *9*, 387-389, doi:10.1016/j.meegid.2009.01.001.
28. Immel, A.; Key, F.M.; Szolek, A.; Barquera, R.; Robinson, M.K.; Harrison, G.F.; Palmer, W.H.; Spyrou, M.A.; Susat, J.; Krause-Kyora, B., et al. Analysis of genomic DNA from medieval plague victims suggests long-term effect of *Yersinia pestis* on human immunity genes. *Molecular biology and evolution* **2021**, *38*, 4059-4076, doi:10.1093/molbev/msab147.
29. Park, Y.H.; Remmers, E.F.; Lee, W.; Ombrello, A.K.; Chung, L.K.; Shilei, Z.; Stone, D.L.; Ivanov, M.I.; Loeven, N.A.; Barron, K.S., et al. Ancient familial Mediterranean fever mutations in human pyrin and resistance to *Yersinia pestis*. *Nat Immunol* **2020**, *21*, 857-867, doi:10.1038/s41590-020-0705-6.
30. Lindesmith, L.; Moe, C.; Marionneau, S.; Ruvoen, N.; Jiang, X.; Lindblad, L.; Stewart, P.; LePendou, J.; Baric, R. Human susceptibility and resistance to Norwalk virus infection. *Nat Med* **2003**, *9*, 548-553, doi:10.1038/nm860.
31. Thorven, M.; Grahn, A.; Hedlund, K.O.; Johansson, H.; Wahlfrid, C.; Larson, G.; Svensson, L. A homozygous nonsense mutation (428G-->A) in the human secretor (FUT2) gene provides resistance to symptomatic norovirus (GGII) infections. *Journal of virology* **2005**, *79*, 15351-15355, doi:10.1128/jvi.79.24.15351-15355.2005.

32. Imbert-Marcille, B.M.; Barbé, L.; Dupé, M.; Le Moullac-Vaidye, B.; Besse, B.; Peltier, C.; Ruvoën-Clouet, N.; Le Pendu, J. A FUT2 gene common polymorphism determines resistance to rotavirus A of the P[8] genotype. *The Journal of infectious diseases* **2014**, *209*, 1227-1230, doi:10.1093/infdis/jit655.
33. Everitt, A.R.; Clare, S.; Pertel, T.; John, S.P.; Wash, R.S.; Smith, S.E.; Chin, C.R.; Feeley, E.M.; Sims, J.S.; Adams, D.J., et al. IFITM3 restricts the morbidity and mortality associated with influenza. *Nature* **2012**, *484*, 519-523, doi:10.1038/nature10921.
34. McManus, K.F.; Taravella, A.M.; Henn, B.M.; Bustamante, C.D.; Sikora, M.; Cornejo, O.E. Population genetic analysis of the DARC locus (Duffy) reveals adaptation from standing variation associated with malaria resistance in humans. *PLoS genetics* **2017**, *13*, e1006560, doi:10.1371/journal.pgen.1006560.
35. Kerner, G.; Laval, G.; Patin, E.; Boisson-Dupuis, S.; Abel, L.; Casanova, J.L.; Quintana-Murci, L. Human ancient DNA analyses reveal the high burden of tuberculosis in Europeans over the last 2,000 years. *Am J Hum Genet* **2021**, *108*, 517-524, doi:10.1016/j.ajhg.2021.02.009.
36. Key, F.M.; Peter, B.; Dennis, M.Y.; Huerta-Sánchez, E.; Tang, W.; Prokunina-Olsson, L.; Nielsen, R.; Andrés, A.M. Selection on a variant associated with improved viral clearance drives local, adaptive pseudogenization of interferon lambda 4 (IFNL4). *PLoS genetics* **2014**, *10*, e1004681, doi:10.1371/journal.pgen.1004681.
37. Karlsson, E.K.; Harris, J.B.; Tabrizi, S.; Rahman, A.; Shlyakhter, I.; Patterson, N.; O'Dushlaine, C.; Schaffner, S.F.; Gupta, S.; Chowdhury, F., et al. Natural selection in a bangladeshi population from the cholera-endemic ganges river delta. *Sci Transl Med* **2013**, *5*, 192ra186, doi:10.1126/scitranslmed.3006338.
38. Sallares, R.; Bouwman, A.; Anderung, C. The spread of malaria to Southern Europe in antiquity: New approaches to old problems. *Medical History* **2004**, *48*, 311-328, doi:10.1017/S0025727300007651.
39. Karlsson, E.K.; Kwiatkowski, D.P.; Sabeti, P.C. Natural selection and infectious disease in human populations. *Nat Rev Genet* **2014**, *15*, 379-393, doi:10.1038/nrg3734.
40. Quintana-Murci, L. Dangerous liaisons: human genetic adaptation to infectious agents. *C R Biol* **2021**, *343*, 297-309, doi:10.5802/crbio.30.
41. Soren, D. Can archaeologists excavate evidence of malaria? *World Archaeology* **2003**, *35*, 193-209.
42. Fornaciari, A. Environmental microbial forensics and archaeology of past pandemics. *Microbiol Spectr* **2017**, *5*, doi:10.1128/microbiolspec.EMF-0011-2016.
43. Spyrou, M.A.; Keller, M.; Tukhbatova, R.I.; Scheib, C.L.; Nelson, E.A.; Andrades Valtueña, A.; Neumann, G.U.; Walker, D.; Alterauge, A.; Carty, N., et al. Phylogeography of the second plague pandemic revealed through analysis of historical *Yersinia pestis* genomes. *Nature communications* **2019**, *10*, 4470, doi:10.1038/s41467-019-12154-0.
44. Spyrou, M.A.; Tukhbatova, R.I.; Feldman, M.; Drath, J.; Kacki, S.; Beltrán de Heredia, J.; Arnold, S.; Sitdikov, A.G.; Castex, D.; Wahl, J., et al. Historical *Y. pestis* genomes reveal the European Black Death as the source of ancient and modern plague pandemics. *Cell host & microbe* **2016**, *19*, 874-881, doi:10.1016/j.chom.2016.05.012.
45. Haensch, S.; Bianucci, R.; Signoli, M.; Rajerison, M.; Schultz, M.; Kacki, S.; Vermunt, M.; Weston, D.A.; Hurst, D.; Achtman, M., et al. Distinct clones of *Yersinia pestis* caused the black death. *PLoS pathogens* **2010**, *6*, e1001134, doi:10.1371/journal.ppat.1001134.
46. Bos, K.I.; Schuenemann, V.J.; Golding, G.B.; Burbano, H.A.; Waglechner, N.; Coombes, B.K.; McPhee, J.B.; DeWitte, S.N.; Meyer, M.; Schmedes, S., et al. A draft genome of *Yersinia pestis* from victims of the Black Death. *Nature* **2011**, *478*, 506-510, doi:10.1038/nature10549.
47. Guellil, M.; Rinaldo, N.; Zedda, N.; Kersten, O.; Gonzalez Muro, X.; Stenseth, N.C.; Gualdi-Russo, E.; Bramanti, B. Bioarchaeological insights into the last plague of Imola (1630-1632). *Sci Rep* **2021**, *11*, 22253, doi:10.1038/s41598-021-98214-2.

48. Harbeck, M.; Seifert, L.; Hänsch, S.; Wagner, D.M.; Birdsell, D.; Parise, K.L.; Wiechmann, I.; Grupe, G.; Thomas, A.; Keim, P., et al. Yersinia pestis DNA from skeletal remains from the 6(th) century AD reveals insights into Justinianic Plague. *PLoS pathogens* **2013**, *9*, e1003349, doi:10.1371/journal.ppat.1003349.
49. Keller, M.; Spyrou, M.A.; Scheib, C.L.; Neumann, G.U.; Kröpelin, A.; Haas-Gebhard, B.; Pfüffgen, B.; Haberstroh, J.; Ribera, I.L.A.; Raynaud, C., et al. Ancient Yersinia pestis genomes from across Western Europe reveal early diversification during the First Pandemic (541-750). *Proceedings of the National Academy of Sciences of the United States of America* **2019**, *116*, 12363-12372, doi:10.1073/pnas.1820447116.
50. Rasmussen, S.; Allentoft, M.E.; Nielsen, K.; Orlando, L.; Sikora, M.; Sjögren, K.G.; Pedersen, A.G.; Schubert, M.; Van Dam, A.; Kapel, C.M., et al. Early divergent strains of Yersinia pestis in Eurasia 5,000 years ago. *Cell* **2015**, *163*, 571-582, doi:10.1016/j.cell.2015.10.009.
51. Rascovan, N.; Sjögren, K.G.; Kristiansen, K.; Nielsen, R.; Willerslev, E.; Desnues, C.; Rasmussen, S. Emergence and spread of basal lineages of Yersinia pestis during the Neolithic Decline. *Cell* **2019**, *176*, 295-305.e210, doi:10.1016/j.cell.2018.11.005.
52. Andrades Valtueña, A.; Neumann, G.U.; Spyrou, M.A.; Musralina, L.; Aron, F.; Beisenov, A.; Belinskiy, A.B.; Bos, K.I.; Buzhilova, A.; Conrad, M., et al. Stone Age Yersinia pestis genomes shed light on the early evolution, diversity, and ecology of plague. *Proceedings of the National Academy of Sciences of the United States of America* **2022**, *119*, e2116722119, doi:10.1073/pnas.2116722119.
53. Chouin, G. Reflections on plague in African history (14th–19th c.). In *Afriques*, 2018; Vol. 09.
54. Gallagher, D.E.; Dueppen, S.A. Recognizing plague epidemics in the archaeological record of West Africa. In *Afriques*, 2018; Vol. 09.
55. Schuenemann, V.J.; Singh, P.; Mendum, T.A.; Krause-Kyora, B.; Jäger, G.; Bos, K.I.; Herbig, A.; Economou, C.; Benjak, A.; Busso, P., et al. Genome-wide comparison of medieval and modern Mycobacterium leprae. *Science (New York, N.Y.)* **2013**, *341*, 179-183, doi:10.1126/science.1238286.
56. Schuenemann, V.J.; Avanzi, C.; Krause-Kyora, B.; Seitz, A.; Herbig, A.; Inskip, S.; Bonazzi, M.; Reiter, E.; Urban, C.; Dangvard Pedersen, D., et al. Ancient genomes reveal a high diversity of Mycobacterium leprae in medieval Europe. *PLoS pathogens* **2018**, *14*, e1006997, doi:10.1371/journal.ppat.1006997.
57. Pfrengle, S.; Neukamm, J.; Guellil, M.; Keller, M.; Molak, M.; Avanzi, C.; Kushniarevich, A.; Montes, N.; Neumann, G.U.; Reiter, E., et al. Mycobacterium leprae diversity and population dynamics in medieval Europe from novel ancient genomes. *BMC Biol* **2021**, *19*, 220, doi:10.1186/s12915-021-01120-2.
58. Mendum, T.A.; Schuenemann, V.J.; Roffey, S.; Taylor, G.M.; Wu, H.; Singh, P.; Tucker, K.; Hinds, J.; Cole, S.T.; Kierzek, A.M., et al. Mycobacterium leprae genomes from a British medieval leprosy hospital: towards understanding an ancient epidemic. *BMC Genomics* **2014**, *15*, 270, doi:10.1186/1471-2164-15-270.
59. Jaeger, L.H.; de Souza, S.M.; Dias, O.F.; Iñiguez, A.M. Mycobacterium tuberculosis complex in remains of 18th-19th century slaves, Brazil. *Emerging infectious diseases* **2013**, *19*, 837-839, doi:10.3201/eid1905.120193.
60. Bos, K.I.; Harkins, K.M.; Herbig, A.; Coscolla, M.; Weber, N.; Comas, I.; Forrest, S.A.; Bryant, J.M.; Harris, S.R.; Schuenemann, V.J., et al. Pre-Columbian mycobacterial genomes reveal seals as a source of New World human tuberculosis. *Nature* **2014**, *514*, 494-497, doi:10.1038/nature13591.
61. Müller, R.; Roberts, C.A.; Brown, T.A. Biomolecular identification of ancient Mycobacterium tuberculosis complex DNA in human remains from Britain and continental Europe. *Am J Phys Anthropol* **2014**, *153*, 178-189, doi:10.1002/ajpa.22417.
62. Kay, G.L.; Sergeant, M.J.; Zhou, Z.; Chan, J.Z.; Millard, A.; Quick, J.; Szikossy, I.; Pap, I.; Spigelman, M.; Loman, N.J., et al. Eighteenth-century genomes show that mixed infections were common at time of peak tuberculosis in Europe. *Nature communications* **2015**, *6*, 6717, doi:10.1038/ncomms7717.

63. Hershkovitz, I.; Donoghue, H.D.; Minnikin, D.E.; Besra, G.S.; Lee, O.Y.; Gernaey, A.M.; Galili, E.; Eshed, V.; Greenblatt, C.L.; Lemma, E., et al. Detection and molecular characterization of 9,000-year-old *Mycobacterium tuberculosis* from a Neolithic settlement in the Eastern Mediterranean. *PloS one* **2008**, *3*, e3426, doi:10.1371/journal.pone.0003426.
64. Raoult, D.; Dutour, O.; Houhamdi, L.; Jankauskas, R.; Fournier, P.E.; Ardagna, Y.; Drancourt, M.; Signoli, M.; La, V.D.; Macia, Y., et al. Evidence for louse-transmitted diseases in soldiers of Napoleon's Grand Army in Vilnius. *The Journal of infectious diseases* **2006**, *193*, 112-120, doi:10.1086/498534.
65. Nguyen-Hieu, T.; Aboudharam, G.; Signoli, M.; Rigeade, C.; Drancourt, M.; Raoult, D. Evidence of a louse-borne outbreak involving typhus in Douai, 1710-1712 during the war of Spanish succession. *PloS one* **2010**, *5*, e15405, doi:10.1371/journal.pone.0015405.
66. Tran, T.N.; Forestier, C.L.; Drancourt, M.; Raoult, D.; Aboudharam, G. Brief communication: co-detection of *Bartonella quintana* and *Yersinia pestis* in an 11th-15th burial site in Bondy, France. *Am J Phys Anthropol* **2011**, *145*, 489-494, doi:10.1002/ajpa.21510.
67. Barbieri, R.; Drancourt, M.; Raoult, D. The role of louse-transmitted diseases in historical plague pandemics. *The Lancet. Infectious diseases* **2021**, *21*, e17-e25, doi:10.1016/s1473-3099(20)30487-4.
68. Guellil, M.; Kersten, O.; Namouchi, A.; Bauer, E.L.; Derrick, M.; Jensen, A.; Stenseth, N.C.; Bramanti, B. Genomic blueprint of a relapsing fever pathogen in 15th century Scandinavia. *Proceedings of the National Academy of Sciences of the United States of America* **2018**, *115*, 10422-10427, doi:10.1073/pnas.1807266115.
69. Oumarou Hama, H.; Barbieri, R.; Guirou, J.; Chenal, T.; Mayer, A.; Ardagna, Y.; Signoli, M.; Aboudharam, G.; Raoult, D.; Drancourt, M. An outbreak of relapsing fever unmasked by microbial paleoserology, 16th century, France. *Am J Phys Anthropol* **2020**, *173*, 784-789, doi:10.1002/ajpa.24138.
70. de-Dios, T.; Carrión, P.; Olalde, I.; Llovera Nadal, L.; Lizano, E.; Pàmies, D.; Marques-Bonet, T.; Balloux, F.; van Dorp, L.; Lalueza-Fox, C. *Salmonella enterica* from a soldier from the 1652 siege of Barcelona (Spain) supports historical transatlantic epidemic contacts. *iScience* **2021**, *24*, 103021, doi:10.1016/j.isci.2021.103021.
71. Warinner, C.; García, N.R.; Spores, R.; Tuross, N. Disease, demography, and diet in early colonial New Spain: Investigation of a sixteenth-century Mixtec cemetery at Teposcolula Yucundaa. *Latin American Antiquity* **2012**, *23*, 467-489.
72. Vågane, Å, J.; Herbig, A.; Campana, M.G.; Robles García, N.M.; Warinner, C.; Sabin, S.; Spyrou, M.A.; Andrades Valtueña, A.; Huson, D.; Tuross, N., et al. *Salmonella enterica* genomes from victims of a major sixteenth-century epidemic in Mexico. *Nat Ecol Evol* **2018**, *2*, 520-528, doi:10.1038/s41559-017-0446-6.
73. Haller, M.; Callan, K.; Susat, J.; Flux, A.L.; Immel, A.; Franke, A.; Herbig, A.; Krause, J.; Kupczok, A.; Fouquet, G., et al. Mass burial genomics reveals outbreak of enteric paratyphoid fever in the Late Medieval trade city Lübeck. *iScience* **2021**, *24*, 102419, doi:10.1016/j.isci.2021.102419.
74. Wu, X.; Ning, C.; Key, F.M.; Andrades Valtueña, A.; Lankapalli, A.K.; Gao, S.; Yang, X.; Zhang, F.; Liu, L.; Nie, Z., et al. A 3,000-year-old, basal *S. enterica* lineage from Bronze Age Xinjiang suggests spread along the Proto-Silk Road. *PLoS pathogens* **2021**, *17*, e1009886, doi:10.1371/journal.ppat.1009886.
75. Zhou, Z.; Lundstrøm, I.; Tran-Dien, A.; Duchêne, S.; Alikhan, N.F.; Sergeant, M.J.; Langridge, G.; Fotakis, A.K.; Nair, S.; Stenøien, H.K., et al. Pan-genome analysis of ancient and modern *Salmonella enterica* demonstrates genomic stability of the invasive Para C lineage for millennia. *Curr Biol* **2018**, *28*, 2420-2428.e2410, doi:10.1016/j.cub.2018.05.058.
76. Biagini, P.; Thèves, C.; Balaesque, P.; Gérard, A.; Cannet, C.; Keyser, C.; Nikolaeva, D.; Gérard, P.; Duchesne, S.; Orlando, L., et al. Variola virus in a 300-year-old Siberian mummy. *The New England journal of medicine* **2012**, *367*, 2057-2059, doi:10.1056/NEJMc1208124.

77. Schuenemann, V.J.; Kumar Lankapalli, A.; Barquera, R.; Nelson, E.A.; Iraíz Hernández, D.; Acuña Alonzo, V.; Bos, K.I.; Márquez Morfín, L.; Herbig, A.; Krause, J. Historic *Treponema pallidum* genomes from Colonial Mexico retrieved from archaeological remains. *PLoS Negl Trop Dis* **2018**, *12*, e0006447, doi:10.1371/journal.pntd.0006447.
78. Majander, K.; Pfrengle, S.; Kocher, A.; Neukamm, J.; du Plessis, L.; Pla-Díaz, M.; Arora, N.; Akgül, G.; Salo, K.; Schats, R., et al. Ancient bacterial genomes reveal a high diversity of *Treponema pallidum* strains in early modern Europe. *Curr Biol* **2020**, *30*, 3788-3803.e3710, doi:10.1016/j.cub.2020.07.058.
79. Knell, R.J. Syphilis in renaissance Europe: rapid evolution of an introduced sexually transmitted disease? *Proceedings. Biological sciences* **2004**, *271 Suppl 4*, S174-176, doi:10.1098/rsbl.2003.0131.
80. Tampa, M.; Sarbu, I.; Matei, C.; Benea, V.; Georgescu, S.R. Brief history of syphilis. *J Med Life* **2014**, *7*, 4-10.
81. Beale, M.A.; Lukehart, S.A. Archaeogenetics: What can ancient genomes tell us about the origin of syphilis? *Curr Biol* **2020**, *30*, R1092-r1095, doi:10.1016/j.cub.2020.08.022.
82. Mühlemann, B.; Vinner, L.; Margaryan, A.; Wilhelmson, H.; de la Fuente Castro, C.; Allentoft, M.E.; de Barros Damgaard, P.; Hansen, A.J.; Holtsmark Nielsen, S.; Strand, L.M., et al. Diverse variola virus (smallpox) strains were widespread in northern Europe in the Viking Age. *Science (New York, N.Y.)* **2020**, *369*, doi:10.1126/science.aaw8977.
83. Toppinen, M.; Perdomo, M.F.; Palo, J.U.; Simmonds, P.; Lycett, S.J.; Söderlund-Venermo, M.; Sajantila, A.; Hedman, K. Bones hold the key to DNA virus history and epidemiology. *Sci Rep* **2015**, *5*, 17226, doi:10.1038/srep17226.
84. Mühlemann, B.; Margaryan, A.; Damgaard, P.B.; Allentoft, M.E.; Vinner, L.; Hansen, A.J.; Weber, A.; Bazaliiskii, V.I.; Molak, M.; Arneborg, J., et al. Ancient human parvovirus B19 in Eurasia reveals its long-term association with humans. *Proceedings of the National Academy of Sciences of the United States of America* **2018**, *115*, 7557-7562, doi:10.1073/pnas.1804921115.
85. Guzmán-Solís, A.A.; Villa-Islas, V.; Bravo-López, M.J.; Sandoval-Velasco, M.; Wesp, J.K.; Gómez-Valdés, J.A.; Moreno-Cabrera, M.L.; Meraz, A.; Solís-Pichardo, G.; Schaaf, P., et al. Ancient viral genomes reveal introduction of human pathogenic viruses into Mexico during the transatlantic slave trade. *eLife* **2021**, *10*, doi:10.7554/eLife.68612.
86. Loreille, O.; Roumat, E.; Verneau, O.; Bouchet, F.; Hänni, C. Ancient DNA from *Ascaris*: extraction amplification and sequences from eggs collected in coprolites. *Int J Parasitol* **2001**, *31*, 1101-1106, doi:10.1016/s0020-7519(01)00214-4.
87. Oh, C.S.; Seo, M.; Hong, J.H.; Chai, J.Y.; Oh, S.W.; Park, J.B.; Shin, D.H. Ancient mitochondrial DNA analyses of ascaris eggs discovered in coprolites from Joseon tomb. *Korean J Parasitol* **2015**, *53*, 237-242, doi:10.3347/kjp.2015.53.2.237.
88. Sørensen, M.J.; Nejsum, P.; Fredensborg, B.L.; Kapel, C.M. DNA typing of ancient parasite eggs from environmental samples identifies human and animal worm infections in Viking-age settlement. *J Parasitol* **2015**, *101*, 57-63, doi:10.1645/14-650.1.
89. Roche, K.; Nicolas, C.; Elsa, P.; Paolo, L.; Pasquino, P.; Raffaella, B.; Matthieu, L.B. Gastrointestinal parasite burden in 4th-5th c. CE Florence highlighted by microscopy and paleogenetics. *Infect Genet Evol* **2021**, *90*, 104713, doi:10.1016/j.meegid.2021.104713.
90. Flammer, P.G.; Dellicour, S.; Preston, S.G.; Rieger, D.; Warren, S.; Tan, C.K.W.; Nicholson, R.; Přichystalová, R.; Bleicher, N.; Wahl, J., et al. Molecular archaeoparasitology identifies cultural changes in the Medieval Hanseatic trading centre of Lübeck. *Proceedings. Biological sciences* **2018**, *285*, doi:10.1098/rspb.2018.0991.

91. Mitchell, P.D.; Yeh, H.Y.; Appleby, J.; Buckley, R. The intestinal parasites of King Richard III. *Lancet (London, England)* **2013**, *382*, 888, doi:10.1016/s0140-6736(13)61757-2.
92. Flammer, P.G.; Ryan, H.; Preston, S.G.; Warren, S.; Přichystalová, R.; Weiss, R.; Palmowski, V.; Boschert, S.; Fellgiebel, K.; Jasch-Boley, I., et al. Epidemiological insights from a large-scale investigation of intestinal helminths in Medieval Europe. *PLoS Negl Trop Dis* **2020**, *14*, e0008600, doi:10.1371/journal.pntd.0008600.
93. Ryan, H.; Flammer, P.G.; Nicholson, R.; Loe, L.; Reeves, B.; Allison, E.; Guy, C.; Doriga, I.L.; Waldron, T.; Walker, D., et al. Reconstructing the history of helminth prevalence in the UK. *PLoS Negl Trop Dis* **2022**, *16*, e0010312, doi:10.1371/journal.pntd.0010312.
94. Appelt, S.; Fancello, L.; Le Bailly, M.; Raoult, D.; Drancourt, M.; Desnues, C. Viruses in a 14th-century coprolite. *Appl Environ Microbiol* **2014**, *80*, 2648-2655, doi:10.1128/aem.03242-13.
95. Warinner, C.; Rodrigues, J.F.; Vyas, R.; Trachsel, C.; Shved, N.; Grossmann, J.; Radini, A.; Hancock, Y.; Tito, R.Y.; Fiddyment, S., et al. Pathogens and host immunity in the ancient human oral cavity. *Nat Genet* **2014**, *46*, 336-344, doi:10.1038/ng.2906.
96. Warinner, C.; Speller, C.; Collins, M.J.; Lewis, C.M., Jr. Ancient human microbiomes. *J Hum Evol* **2015**, *79*, 125-136, doi:10.1016/j.jhevol.2014.10.016.
97. Warinner, C.; Speller, C.; Collins, M.J. A new era in palaeomicrobiology: prospects for ancient dental calculus as a long-term record of the human oral microbiome. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2015**, *370*, 20130376, doi:10.1098/rstb.2013.0376.
98. Santiago-Rodriguez, T.M.; Fornaciari, G.; Luciani, S.; Dowd, S.E.; Toranzos, G.A.; Marota, I.; Cano, R.J. Gut microbiome of an 11th century A.D. pre-Columbian Andean mummy. *PloS one* **2015**, *10*, e0138135, doi:10.1371/journal.pone.0138135.
99. Santiago-Rodriguez, T.M.; Fornaciari, G.; Luciani, S.; Dowd, S.E.; Toranzos, G.A.; Marota, I.; Cano, R.J. Natural mummification of the human gut preserves bacteriophage DNA. *FEMS Microbiol Lett* **2016**, *363*, fnv219, doi:10.1093/femsle/fnv219.
100. Neukamm, J.; Pfrengle, S.; Molak, M.; Seitz, A.; Francken, M.; Eppenberger, P.; Avanzi, C.; Reiter, E.; Urban, C.; Welte, B., et al. 2000-year-old pathogen genomes reconstructed from metagenomic analysis of Egyptian mummified individuals. *BMC Biol* **2020**, *18*, 108, doi:10.1186/s12915-020-00839-8.
101. Wibowo, M.C.; Yang, Z.; Borry, M.; Hübner, A.; Huang, K.D.; Tierney, B.T.; Zimmerman, S.; Barajas-Olmos, F.; Contreras-Cubas, C.; García-Ortiz, H., et al. Reconstruction of ancient microbial genomes from the human gut. *Nature* **2021**, *594*, 234-239, doi:10.1038/s41586-021-03532-0.
102. Senti, S.; Habicht, M.E.; Rayo, E.; Eppenberger, P.E.; Rühli, F.J.; Galassi, F.M. Egyptian canopic jars at the crossroad of medicine and archaeology: Overview of 100 years of research and future scientific expectations. *Pathobiology* **2018**, *85*, 267-275, doi:10.1159/000490797.
103. Ruffer, M.A. Note on the presence of "Bilharzia haematobia" in Egyptian mummies of the Twentieth Dynasty [1250-1000 B.C.]. *Br Med J* **1910**, *1*, 16, doi:10.1136/bmj.1.2557.16-a.
104. Susat, J.; Lübke, H.; Immel, A.; Brinker, U.; Macãne, A.; Meadows, J.; Steer, B.; Tholey, A.; Zagorska, I.; Gerhards, G., et al. A 5,000-year-old hunter-gatherer already plagued by *Yersinia pestis*. *Cell Rep* **2021**, *35*, 109278, doi:10.1016/j.celrep.2021.109278.
105. Devault, A.M.; Golding, G.B.; Waglechner, N.; Enk, J.M.; Kuch, M.; Tien, J.H.; Shi, M.; Fisman, D.N.; Dhody, A.N.; Forrest, S., et al. Second-pandemic strain of *Vibrio cholerae* from the Philadelphia cholera outbreak of 1849. *The New England journal of medicine* **2014**, *370*, 334-340, doi:10.1056/NEJMoa1308663.
106. Zhu, T.; Korber, B.T.; Nahmias, A.J.; Hooper, E.; Sharp, P.M.; Ho, D.D. An African HIV-1 sequence from 1959 and implications for the origin of the epidemic. *Nature* **1998**, *391*, 594-597, doi:10.1038/35400.

107. Worobey, M.; Gemmel, M.; Teuwen, D.E.; Haselkorn, T.; Kunstman, K.; Bunce, M.; Muyembe, J.J.; Kabongo, J.M.; Kalengayi, R.M.; Van Marck, E., et al. Direct evidence of extensive diversity of HIV-1 in Kinshasa by 1960. *Nature* **2008**, *455*, 661-664, doi:10.1038/nature07390.
108. Taubenberger, J.K.; Reid, A.H.; Krafft, A.E.; Bijwaard, K.E.; Fanning, T.G. Initial genetic characterization of the 1918 "Spanish" influenza virus. *Science (New York, N.Y.)* **1997**, *275*, 1793-1796, doi:10.1126/science.275.5307.1793.
109. Reid, A.H.; Fanning, T.G.; Hultin, J.V.; Taubenberger, J.K. Origin and evolution of the 1918 "Spanish" influenza virus hemagglutinin gene. *Proceedings of the National Academy of Sciences of the United States of America* **1999**, *96*, 1651-1656, doi:10.1073/pnas.96.4.1651.
110. Worobey, M.; Han, G.Z.; Rambaut, A. Genesis and pathogenesis of the 1918 pandemic H1N1 influenza A virus. *Proceedings of the National Academy of Sciences of the United States of America* **2014**, *111*, 8107-8112, doi:10.1073/pnas.1324197111.
111. Patrono, L.V.; Vrancken, B.; Budt, M.; D ux, A.; Lequime, S.; Boral, S.; Gilbert, M.T.P.; Gogarten, J.F.; Hoffmann, L.; Horst, D., et al. Archival influenza virus genomes from Europe reveal genomic variability during the 1918 pandemic. *Nature communications* **2022**, *13*, 2314, doi:10.1038/s41467-022-29614-9.
112. Fanning, T.G.; Slemons, R.D.; Reid, A.H.; Janczewski, T.A.; Dean, J.; Taubenberger, J.K. 1917 avian influenza virus sequences suggest that the 1918 pandemic virus did not acquire its hemagglutinin directly from birds. *Journal of virology* **2002**, *76*, 7860-7862, doi:10.1128/jvi.76.15.7860-7862.2002.
113. Rekand, T.; Male, R.; Myking, A.O.; Nygaard, S.J.; Aarli, J.A.; Haarr, L.; Langeland, N. Detection of viral sequences in archival spinal cords from fatal cases of poliomyelitis in 1951-1952. *Journal of virological methods* **2003**, *114*, 91-96, doi:10.1016/j.jviromet.2003.08.013.
114. D ux, A.; Lequime, S.; Patrono, L.V.; Vrancken, B.; Boral, S.; Gogarten, J.F.; Hilbig, A.; Horst, D.; Merkel, K.; Prepoint, B., et al. Measles virus and rinderpest virus divergence dated to the sixth century BCE. *Science (New York, N.Y.)* **2020**, *368*, 1367-1370, doi:10.1126/science.aba9411.
115. Xiao, Y.L.; Kash, J.C.; Beres, S.B.; Sheng, Z.M.; Musser, J.M.; Taubenberger, J.K. High-throughput RNA sequencing of a formalin-fixed, paraffin-embedded autopsy lung tissue sample from the 1918 influenza pandemic. *The Journal of pathology* **2013**, *229*, 535-545, doi:10.1002/path.4145.
116. Carrick, D.M.; Mehaffey, M.G.; Sachs, M.C.; Altekruse, S.; Camalier, C.; Chuaqui, R.; Cozen, W.; Das, B.; Hernandez, B.Y.; Lih, C.J., et al. Robustness of Next Generation Sequencing on older formalin-fixed paraffin-embedded tissue. *PloS one* **2015**, *10*, e0127353, doi:10.1371/journal.pone.0127353.
117. Hykin, S.M.; Bi, K.; McGuire, J.A. Fixing formalin: A method to recover genomic-scale DNA sequence data from formalin-fixed museum specimens using high-throughput sequencing. *PloS one* **2015**, *10*, e0141579, doi:10.1371/journal.pone.0141579.
118. Xiao, Y.; Sheng, Z.M.; Taubenberger, J.K. Isolating viral and host RNA sequences from archival material and production of cDNA libraries for high-throughput DNA sequencing. *Curr Protoc Microbiol* **2015**, *37*, 1e.8.1-16, doi:10.1002/9780471729259.mc01e08s37.
119. Kocjan, B.J.; Ho snjak, L.; Poljak, M. Detection of alpha human papillomaviruses in archival formalin-fixed, paraffin-embedded (FFPE) tissue specimens. *Journal of clinical virology : the official publication of the Pan American Society for Clinical Virology* **2016**, *76 Suppl 1*, S88-s97, doi:10.1016/j.jcv.2015.10.007.
120. Van Doorslaer, K.; Chen, Z.; McBride, A.A. Detection and genotyping of human papillomaviruses from archival formalin-fixed tissue samples. *Curr Protoc Microbiol* **2016**, *43*, 14b.19.11-14b.19.20, doi:10.1002/cpmc.16.

121. Derkarabetian, S.; Benavides, L.R.; Giribet, G. Sequence capture phylogenomics of historical ethanol-preserved museum specimens: Unlocking the rest of the vault. *Mol Ecol Resour* **2019**, *19*, 1531-1544, doi:10.1111/1755-0998.13072.
122. Montiel, R.; Solórzano, E.; Díaz, N.; Álvarez-Sandoval, B.A.; González-Ruiz, M.; Cañadas, M.P.; Simões, N.; Isidro, A.; Malgosa, A. Neonate human remains: a window of opportunity to the molecular study of ancient syphilis. *PloS one* **2012**, *7*, e36371, doi:10.1371/journal.pone.0036371.
123. Schuenemann, V.J.; Bos, K.; DeWitte, S.; Schmedes, S.; Jamieson, J.; Mittnik, A.; Forrest, S.; Coombes, B.K.; Wood, J.W.; Earn, D.J., et al. Targeted enrichment of ancient pathogens yielding the pPCP1 plasmid of *Yersinia pestis* from victims of the Black Death. *Proceedings of the National Academy of Sciences of the United States of America* **2011**, *108*, E746-752, doi:10.1073/pnas.1105107108.
124. Castillo-Rojas, G.; Cerbón, M.A.; López-Vidal, Y. Presence of *Helicobacter pylori* in a Mexican Pre-Columbian Mummy. *BMC Microbiol* **2008**, *8*, 119, doi:10.1186/1471-2180-8-119.
125. Shin, D.H.; Oh, C.S.; Hong, J.H.; Lee, H.; Lee, S.D.; Lee, E. *Helicobacter pylori* DNA obtained from the stomach specimens of two 17(th) century Korean mummies. *Anthropol Anz* **2018**, *75*, 75-87, doi:10.1127/anthranz/2018/0780.
126. Fornaciari, G.; Zavaglia, K.; Giusti, L.; Vultaggio, C.; Ciranni, R. Human papillomavirus in a 16th century mummy. *Lancet (London, England)* **2003**, *362*, 1160, doi:10.1016/s0140-6736(03)14487-x.
127. Chan, J.Z.; Sergeant, M.J.; Lee, O.Y.; Minnikin, D.E.; Besra, G.S.; Pap, I.; Spigelman, M.; Donoghue, H.D.; Pallen, M.J. Metagenomic analysis of tuberculosis in a mummy. *The New England journal of medicine* **2013**, *369*, 289-290, doi:10.1056/NEJMc1302295.
128. Kim, Y.S.; Lee, I.S.; Oh, C.S.; Kim, M.J.; Cha, S.C.; Shin, D.H. Calcified pulmonary nodules identified in a 350-year-old-Joseon mummy: the first report on ancient pulmonary tuberculosis from archaeologically obtained pre-modern Korean samples. *J Korean Med Sci* **2016**, *31*, 147-151, doi:10.3346/jkms.2016.31.1.147.
129. Duggan, A.T.; Perdomo, M.F.; Piombino-Mascalì, D.; Marciniak, S.; Poinar, D.; Emery, M.V.; Buchmann, J.P.; Duchêne, S.; Jankauskas, R.; Humphreys, M., et al. 17(th) Century variola virus reveals the recent history of smallpox. *Curr Biol* **2016**, *26*, 3407-3412, doi:10.1016/j.cub.2016.10.061.
130. Ferrari, G.; Neukamm, J.; Baalsrud, H.T.; Breidenstein, A.M.; Ravinet, M.; Phillips, C.; Rühli, F.; Bouwman, A.; Schuenemann, V.J. Variola virus genome sequenced from an eighteenth-century museum specimen supports the recent origin of smallpox. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2020**, *375*, 20190572, doi:10.1098/rstb.2019.0572.
131. Gray, R.R.; Tanaka, Y.; Takebe, Y.; Magiorkinis, G.; Buskell, Z.; Seeff, L.; Alter, H.J.; Pybus, O.G. Evolutionary analysis of hepatitis C virus gene sequences from 1953. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2013**, *368*, 20130168, doi:10.1098/rstb.2013.0168.
132. Swanston, T.; Haakensen, M.; Deneer, H.; Walker, E.G. The characterization of *Helicobacter pylori* DNA associated with ancient human remains recovered from a Canadian glacier. *PloS one* **2011**, *6*, e16864, doi:10.1371/journal.pone.0016864.
133. Maixner, F.; Krause-Kyora, B.; Turaev, D.; Herbig, A.; Hoopmann, M.R.; Hallows, J.L.; Kusebauch, U.; Vigl, E.E.; Malfertheiner, P.; Megraud, F., et al. The 5300-year-old *Helicobacter pylori* genome of the Iceman. *Science (New York, N.Y.)* **2016**, *351*, 162-165, doi:10.1126/science.aad2545.
134. Drancourt, M.; Tran-Hung, L.; Courtin, J.; Lumley, H.; Raoult, D. *Bartonella quintana* in a 4000-year-old human tooth. *The Journal of infectious diseases* **2005**, *191*, 607-611, doi:10.1086/427041.

135. Mai, B.H.; Barbieri, R.; Chenal, T.; Castex, D.; Jonvel, R.; Tanasi, D.; Georges-Zimmermann, P.; Dutour, O.; Peressinotto, D.; Demangeot, C., et al. Five millennia of Bartonella quintana bacteraemia. *PloS one* **2020**, *15*, e0239526, doi:10.1371/journal.pone.0239526.
136. Thèves, C.; Senescau, A.; Vanin, S.; Keyser, C.; Ricaut, F.X.; Alekseev, A.N.; Dabernat, H.; Ludes, B.; Fabre, R.; Crubézy, E. Molecular identification of bacteria by total sequence screening: determining the cause of death in ancient human subjects. *PloS one* **2011**, *6*, e21733, doi:10.1371/journal.pone.0021733.
137. Guellil, M.; Keller, M.; Dittmar, J.M.; Inskip, S.A.; Cessford, C.; Solnik, A.; Kivisild, T.; Metspalu, M.; Robb, J.E.; Scheib, C.L. An invasive Haemophilus influenzae serotype b infection in an Anglo-Saxon plague victim. *Genome biology* **2022**, *23*, 22, doi:10.1186/s13059-021-02580-z.
138. Wagner, D.M.; Klunk, J.; Harbeck, M.; Devault, A.; Waglechner, N.; Sahl, J.W.; Enk, J.; Birdsell, D.N.; Kuch, M.; Lumibao, C., et al. Yersinia pestis and the plague of Justinian 541-543 AD: a genomic analysis. *The Lancet. Infectious diseases* **2014**, *14*, 319-326, doi:10.1016/s1473-3099(13)70323-2.
139. Fotakis, A.K.; Denham, S.D.; Mackie, M.; Orbegozo, M.I.; Mylopotamitaki, D.; Gopalakrishnan, S.; Sicheritz-Pontén, T.; Olsen, J.V.; Cappellini, E.; Zhang, G., et al. Multi-omic detection of Mycobacterium leprae in archaeological human dental calculus. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2020**, *375*, 20190584, doi:10.1098/rstb.2019.0584.
140. Jacobson, D.K.; Honap, T.P.; Monroe, C.; Lund, J.; Houk, B.A.; Novotny, A.C.; Robin, C.; Marini, E.; Lewis, C.M., Jr. Functional diversity of microbial ecologies estimated from ancient human coprolites and dental calculus. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2020**, *375*, 20190586, doi:10.1098/rstb.2019.0586.
141. Willmann, C.; Mata, X.; Hanghoej, K.; Tonasso, L.; Tisseyre, L.; Jeziorski, C.; Cabot, E.; Chevet, P.; Crubézy, E.; Orlando, L., et al. Oral health status in historic population: Macroscopic and metagenomic evidence. *PloS one* **2018**, *13*, e0196482, doi:10.1371/journal.pone.0196482.
142. Bravo-Lopez, M.; Villa-Islas, V.; Rocha Arriaga, C.; Villaseñor-Altamirano, A.B.; Guzmán-Solís, A.; Sandoval-Velasco, M.; Wesp, J.K.; Alcantara, K.; López-Corral, A.; Gómez-Valdés, J., et al. Paleogenomic insights into the red complex bacteria Tannerella forsythia in Pre-Hispanic and Colonial individuals from Mexico. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2020**, *375*, 20190580, doi:10.1098/rstb.2019.0580.
143. Philips, A.; Stolarek, I.; Handschuh, L.; Nowis, K.; Juras, A.; Trzciński, D.; Nowaczewska, W.; Wrzesińska, A.; Potempa, J.; Figlerowicz, M. Analysis of oral microbiome from fossil human remains revealed the significant differences in virulence factors of modern and ancient Tannerella forsythia. *BMC Genomics* **2020**, *21*, 402, doi:10.1186/s12864-020-06810-9.
144. Ubaldi, M.; Luciani, S.; Marota, I.; Fornaciari, G.; Cano, R.J.; Rollo, F. Sequence analysis of bacterial DNA in the colon of an Andean mummy. *Am J Phys Anthropol* **1998**, *107*, 285-295, doi:10.1002/(sici)1096-8644(199811)107:3<285::Aid-ajpa5>3.0.Co;2-u.
145. Cano, R.J.; Tiefenbrunner, F.; Ubaldi, M.; Del Cueto, C.; Luciani, S.; Cox, T.; Orkand, P.; Künzel, K.H.; Rollo, F. Sequence analysis of bacterial DNA in the colon and stomach of the Tyrolean Iceman. *Am J Phys Anthropol* **2000**, *112*, 297-309, doi:10.1002/1096-8644(200007)112:3<297.
146. Rifkin, R.F.; Vikram, S.; Ramond, J.B.; Rey-Iglesia, A.; Brand, T.B.; Porraz, G.; Val, A.; Hall, G.; Woodborne, S.; Le Bailly, M., et al. Multi-proxy analyses of a mid-15th century Middle Iron Age Bantu-speaker palaeo-faecal specimen elucidates the configuration of the 'ancestral' sub-Saharan African intestinal microbiome. *Microbiome* **2020**, *8*, 62, doi:10.1186/s40168-020-00832-x.

147. Santiago-Rodriguez, T.M.; Fornaciari, G.; Luciani, S.; Dowd, S.E.; Toranzos, G.A.; Marota, I.; Cano, R.J. Taxonomic and predicted metabolic profiles of the human gut microbiome in pre-Columbian mummies. *FEMS Microbiol Ecol* **2016**, *92*, doi:10.1093/femsec/fiw182.
148. Jensen, T.Z.T.; Niemann, J.; Iversen, K.H.; Fotakis, A.K.; Gopalakrishnan, S.; Vågene, Å, J.; Pedersen, M.W.; Sinding, M.S.; Ellegaard, M.R.; Allentoft, M.E., et al. A 5700 year-old human genome and oral microbiome from chewed birch pitch. *Nature communications* **2019**, *10*, 5520, doi:10.1038/s41467-019-13549-9.
149. Devault, A.M.; Mortimer, T.D.; Kitchen, A.; Kieseewetter, H.; Enk, J.M.; Golding, G.B.; Southon, J.; Kuch, M.; Duggan, A.T.; Aylward, W., et al. A molecular portrait of maternal sepsis from Byzantine Troy. *eLife* **2017**, *6*, doi:10.7554/eLife.20983.
150. Kay, G.L.; Sergeant, M.J.; Giuffra, V.; Bandiera, P.; Milanese, M.; Bramanti, B.; Bianucci, R.; Pallen, M.J. Recovery of a medieval *Brucella melitensis* genome using shotgun metagenomics. *mBio* **2014**, *5*, e01337-01314, doi:10.1128/mBio.01337-14.
151. Grunenwald, A.; Keyser, C.; Sautereau, A.M.; Crubézy, E.; Ludes, B.; Drouet, C. Novel contribution on the diagenetic physicochemical features of bone and teeth minerals, as substrates for ancient DNA typing. *Anal Bioanal Chem* **2014**, *406*, 4691-4704, doi:10.1007/s00216-014-7863-z.
152. Little, L.K. Life and afterlife of the first plague pandemic. In *Plague and the end of antiquity. The pandemic of 541–750*, Little, L.K., Ed. Cambridge University Press: Cambridge, UK, 2007; pp. 3-33.
153. Habicht, M.E.; Eppenberger, P.E.; Rühli, F. A critical assessment of proposed outbreaks of plague and other epidemic diseases in Ancient Egypt. *Int J Infect Dis* **2021**, *103*, 217-219, doi:10.1016/j.ijid.2020.11.160.
154. Vitiello, L.; Ilari, S.; Sansone, L.; Belli, M.; Cristina, M.; Marcolongo, F.; Tomino, C.; Gatta, L.; Mollace, V.; Bonassi, S., et al. Preventive measures against pandemics from the beginning of civilization to nowadays-How everything has remained the same over the millennia. *J Clin Med* **2022**, *11*, doi:10.3390/jcm11071960.
155. Rodríguez-Frías, F.; Quer, J.; Taberner, D.; Cortese, M.F.; Garcia-Garcia, S.; Rando-Segura, A.; Pumarola, T. Microorganisms as shapers of human civilization, from pandemics to even our genomes: Villains or friends? A historical approach. *Microorganisms* **2021**, *9*, doi:10.3390/microorganisms9122518.
156. Bodner, K. Mouse trap: A text-critical problem with rodents in the ark narrative. *The Journal of Theological Studies* **2008**, *59*, 634–649.
157. Asensi, V.; Fierer, J. Of rats and men: Poussin's plague at Ashdod. *Emerging infectious diseases* **2018**, *24*, 186-187, doi:10.3201/eid2401.AC2401.
158. Shapiro, B.; Rambaut, A.; Gilbert, M.T. No proof that typhoid caused the Plague of Athens (a reply to Papagrigorakis et al.). *Int J Infect Dis* **2006**, *10*, 334-335; author reply 335-336, doi:10.1016/j.ijid.2006.02.006.
159. Derat, M.-L. Du lexique aux talismans: occurrences de la peste dans la Corne de l'Afrique du XIIIe au XVe siècle. In *Afriques*, 2018; Vol. 09.
160. Connor, B.A.; Schwartz, E. Typhoid and paratyphoid fever in travellers. *The Lancet. Infectious diseases* **2005**, *5*, 623-628, doi:10.1016/s1473-3099(05)70239-5.
161. Manesh, A.; Meltzer, E.; Jin, C.; Britto, C.; Deodhar, D.; Radha, S.; Schwartz, E.; Rupali, P. Typhoid and paratyphoid fever: a clinical seminar. *J Travel Med* **2021**, *28*, doi:10.1093/jtm/taab012.
162. westbrabantsarchieff.nl/collectie/bladeren-in-bronnen/registers, Bergen op Zoom, Begraaflijsten ziekenbezoeker 1701-1804, map 5 (1709-1716, 1746-1747, 1797, 1802-1804).
163. van der Kuyl, A.C. *The Dutch cholera epidemic of 1832 as seen through 19th century medical publications*; Eburin Academic Publishers: Utrecht, the Netherlands, 2021.
164. Arntzenius, D.J.A. *Bijdragen tot de kennis en behandeling van den Aziatischen braakloop in Nederland*; Van der Vinne: Amsterdam, the Netherlands, 1832.

165. Heyman, P.; Cochez, C.; Hukić, M. The English sweating sickness: out of sight, out of mind? *Acta Med Acad* **2018**, *47*, 102-116, doi:10.5644/ama2006-124.221.
166. Cheshire, W.P.; van Gerpen, J.A.; Sejvar, J.J. Sudor Anglicus: an epidemic targeting the autonomic nervous system. *Clin Auton Res* **2020**, *30*, 317-323, doi:10.1007/s10286-020-00698-x.
167. Black, J. A boke or counseill against the disease commonly called the Sweate or the Sweating Sickness. *Bmj* **2007**, *335*, 1159, doi:10.1136/bmj.39385.490023.FA.
168. Buchner, W.F. *Aanteekeningen en opmerkingen betrekkelijk den Aziatischen braakloop te Gouda*; R. J. Bernthrop: Amsterdam, the Netherlands, 1833.
169. Thwaites, G.; Taviner, M.; Gant, V. The English sweating sickness, 1485 to 1551. *The New England journal of medicine* **1997**, *336*, 580-582, doi:10.1056/nejm199702203360812.
170. Taviner, M.; Thwaites, G.; Gant, V. The English sweating sickness, 1485-1551: a viral pulmonary disease? *Med Hist* **1998**, *42*, 96-98, doi:10.1017/s0025727300063365.
171. Heyman, P.; Simons, L.; Cochez, C. Were the English sweating sickness and the Picardy sweat caused by hantaviruses? *Viruses* **2014**, *6*, 151-171, doi:10.3390/v6010151.
172. Umapathi, T.; Poh, M.Q.W.; Fan, B.E.; Li, K.F.C.; George, J.; Tan, J.Y.L. Acute hyperhidrosis and postural tachycardia in a COVID-19 patient. *Clin Auton Res* **2020**, *30*, 571-573, doi:10.1007/s10286-020-00733-x.
173. Scala, I.; Rizzo, P.A.; Bellavia, S.; Brunetti, V.; Colò, F.; Broccolini, A.; Della Marca, G.; Calabresi, P.; Luigetti, M.; Frisullo, G. Autonomic dysfunction during acute SARS-CoV-2 infection: A systematic review. *J Clin Med* **2022**, *11*, doi:10.3390/jcm11133883.
174. Nandi, S.; Negi, B.S. Bovine ephemeral fever: a review. *Comparative immunology, microbiology and infectious diseases* **1999**, *22*, 81-91, doi:10.1016/s0147-9571(98)00027-7.
175. Burgess, G.W.; Spradbrow, P.B. Studies on the pathogenesis of bovine ephemeral fever. *Aust Vet J* **1977**, *53*, 363-368, doi:10.1111/j.1751-0813.1977.tb07952.x.
176. Kuhn, J.H.; Pān, H.; Chiu, C.Y.; Stremlau, M. Human tibroviruses: commensals or lethal pathogens? *Viruses* **2020**, *12*, doi:10.3390/v12030252.
177. Edridge, A.W.D.; Abd-Elfarag, G.; Deijs, M.; Jebbink, M.F.; Boele van Hensbroek, M.; van der Hoek, L. Divergent rhabdovirus discovered in a patient with new-onset Nodding Syndrome. *Viruses* **2022**, *14*, doi:10.3390/v14020210.
178. Stremlau, M.H.; Andersen, K.G.; Folarin, O.A.; Grove, J.N.; Odiā, I.; Ehiane, P.E.; Omoniwa, O.; Omoregie, O.; Jiang, P.P.; Yozwiak, N.L., et al. Discovery of novel rhabdoviruses in the blood of healthy individuals from West Africa. *PLoS Negl Trop Dis* **2015**, *9*, e0003631, doi:10.1371/journal.pntd.0003631.
179. Guy, P.L. Prospects for analyzing ancient RNA in preserved materials. *Wiley Interdiscip Rev RNA* **2014**, *5*, 87-94, doi:10.1002/wrna.1199.
180. Mahmoudi, A.; Kryštufek, B.; Sludsky, A.; Schmid, B.V.; AMP, D.E.A.; Lei, X.; Ramasindrazana, B.; Bertherat, E.; Yeszhanov, A.; Stenseth, N.C., et al. Plague reservoir species throughout the world. *Integr Zool* **2021**, *16*, 820-833, doi:10.1111/1749-4877.12511.
181. Key, F.M.; Posth, C.; Esquivel-Gomez, L.R.; Hübler, R.; Spyrou, M.A.; Neumann, G.U.; Furtwängler, A.; Sabin, S.; Burri, M.; Wissgott, A., et al. Emergence of human-adapted *Salmonella enterica* is linked to the Neolithization process. *Nat Ecol Evol* **2020**, *4*, 324-333, doi:10.1038/s41559-020-1106-9.
182. Dunstan, S.J.; Hue, N.T.; Han, B.; Li, Z.; Tram, T.T.; Sim, K.S.; Parry, C.M.; Chinh, N.T.; Vinh, H.; Lan, N.P., et al. Variation at HLA-DRB1 is associated with resistance to enteric fever. *Nat Genet* **2014**, *46*, 1333-1336, doi:10.1038/ng.3143.

183. Haller, M.; Bonczarowska, J.H.; Rieger, D.; Lenz, T.L.; Nebel, A.; Krause-Kyora, B. Ancient DNA study in medieval Europeans shows an association between HLA-DRB1*03 and paratyphoid fever. *Frontiers in immunology* **2021**, *12*, 691475, doi:10.3389/fimmu.2021.691475.
184. Raoult, D.; Ndiokubwayo, J.B.; Tissot-Dupont, H.; Roux, V.; Faugere, B.; Abegbinni, R.; Birtles, R.J. Outbreak of epidemic typhus associated with trench fever in Burundi. *Lancet (London, England)* **1998**, *352*, 353-358, doi:10.1016/s0140-6736(97)12433-3.
185. Tarasevich, I.; Rydkina, E.; Raoult, D. Outbreak of epidemic typhus in Russia. *Lancet (London, England)* **1998**, *352*, 1151, doi:10.1016/s0140-6736(05)79799-3.
186. McCormick, D.W.; Rowan, S.E.; Pappert, R.; Yockey, B.; Dietrich, E.A.; Petersen, J.M.; Hinckley, A.F.; Marx, G.E. Bartonella seroreactivity among persons experiencing homelessness during an outbreak of Bartonella quintana in Denver, Colorado, 2020. *Open Forum Infect Dis* **2021**, *8*, ofab230, doi:10.1093/ofid/ofab230.
187. Smith, O.; Clapham, A.; Rose, P.; Liu, Y.; Wang, J.; Allaby, R.G. A complete ancient RNA genome: identification, reconstruction and evolutionary history of archaeological Barley Stripe Mosaic Virus. *Sci Rep* **2014**, *4*, 4003, doi:10.1038/srep04003.
188. Peyambari, M.; Warner, S.; Stoler, N.; Rainer, D.; Roossinck, M.J. A 1,000-Year-Old RNA Virus. *Journal of virology* **2019**, *93*, doi:10.1128/jvi.01188-18.
189. Smith, O.; Dunshea, G.; Sinding, M.S.; Fedorov, S.; Germonpre, M.; Bocherens, H.; Gilbert, M.T.P. Ancient RNA from Late Pleistocene permafrost and historical canids shows tissue-specific transcriptome survival. *PLoS biology* **2019**, *17*, e3000166, doi:10.1371/journal.pbio.3000166.
190. Enard, D.; Petrov, D.A. Ancient RNA virus epidemics through the lens of recent adaptation in human genomes. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **2020**, *375*, 20190575, doi:10.1098/rstb.2019.0575.
191. van der Kuyl, A.C.; Berkhout, B. Viruses in the reproductive tract: On their way to the germ line? *Virus research* **2020**, *286*, 198101, doi:10.1016/j.virusres.2020.198101.
192. Horie, M.; Honda, T.; Suzuki, Y.; Kobayashi, Y.; Daito, T.; Oshida, T.; Ikuta, K.; Jern, P.; Gojobori, T.; Coffin, J.M., et al. Endogenous non-retroviral RNA virus elements in mammalian genomes. *Nature* **2010**, *463*, 84-87, doi:10.1038/nature08695.
193. Chase, A.F.; Chase, D.Z.; Fisher, C.T.; Leisz, S.J.; Weishampel, J.F. Geospatial revolution and remote sensing LiDAR in Mesoamerican archaeology. *Proceedings of the National Academy of Sciences* **2012**, *109*, 12916-12921, doi:doi:10.1073/pnas.1205198109.
194. Prümers, H.; Betancourt, C.J.; Iriarte, J.; Robinson, M.; Schaich, M. Lidar reveals pre-Hispanic low-density urbanism in the Bolivian Amazon. *Nature* **2022**, *606*, 325-328, doi:doi.org/10.1038/s41586-022-04780-4.
195. Albitar, O.; Ballouze, R.; Ooi, J.P.; Sheikh Ghadzi, S.M. Risk factors for mortality among COVID-19 patients. *Diabetes Res Clin Pract* **2020**, *166*, 108293, doi:10.1016/j.diabres.2020.108293.
196. Dessie, Z.G.; Zewotir, T. Mortality-related risk factors of COVID-19: a systematic review and meta-analysis of 42 studies and 423,117 patients. *BMC infectious diseases* **2021**, *21*, 855, doi:10.1186/s12879-021-06536-3.
197. Callaway, E. Divided by DNA: The uneasy relationship between archaeology and ancient genomics. *Nature* **2018**, *555*, 573-576, doi:10.1038/d41586-018-03773-6.
198. Downes, S.M. The role of ancient DNA research in archaeology. *Topoi* **2021**, *40*, 285-293, doi:10.1007/s11245-019-09665-2.
199. Rayo, E.; Ferrari, G.; Neukamm, J.; Akgül, G.; Breidenstein, A.M.; Cooke, M.; Phillips, C.; Bouwman, A.S.; Rühli, F.J.; Schuenemann, V.J. Non-destructive extraction of DNA from preserved tissues in medical collections. *Biotechniques* **2022**, *72*, 60-64, doi:10.2144/btn-2021-0014.

