

Supplementary materials

Efficient purification of *cowpea chlorotic mottle virus* by a novel peptide aptamer

Georg Tscheuschner ¹, Marco Ponader ¹, Christopher Raab ¹, Prisca S. Weider ¹, Reni Hartfiel ¹, Jan Ole Kaufmann ^{1,2,3}, Jule L. Völzke ¹, Gaby Bosc-Bierne ¹, Carsten Prinz ¹, Timm Schwaar ⁴, Paul Andrie ¹, Henriette Bäßler ¹, Khoa Nguyen ¹, Yanchen Zhu ⁵, Antonia S. J. S. Mey ⁵, Amr Mostafa ⁶, Ilko Bald ⁶, Michael G. Weller ^{1,*}

- ¹ Federal Institute for Materials Research and Testing (BAM), Richard-Willstätter-Straße 11, 12489 Berlin, Germany; georg.tscheuschner@bam.de (G.T.); marco.ponader@bam.de (M.P.); christopher.raab@bam.de (C.R.); prisca.weider@tum.de (P.S.W.); reni.hartfiel@outlook.com (R.H.); jan-ole.kaufmann@bam.de (J.O.K.); jule.voelzke@bam.de (J.L.V.); gaby.bosc-bierne@bam.de (G.B.); carsten.prinz@bam.de (C.P.); paul.andrie@bam.de (P.A.); henriette.baessler@uni-rostock.de (H.B.); khoa.nguyen@bam.de (K.N.); michael.weller@bam.de (M.G.W.);
 - ² Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Charitéplatz 1, 10117 Berlin, Germany; jan-ole.kaufmann@charite.de (J.O.K.)
 - ³ Technical University of Munich, Department of Diagnostic and Interventional Radiology, Ismaninger Str.22, 81675 Munich, Germany; jan-ole.kaufmann@tum.de (J.O.K.)
 - ⁴ SAFIA Technologies GmbH, Richard-Willstätter-Straße 11, 12489 Berlin, Germany; schwaar@safia.tech (T.S.);
 - ⁵ EaStCHEM School of Chemistry, University of Edinburgh, David Brewster Road, Edinburgh EH9 3FJ, United Kingdom; y.zhu-105@sms.ed.ac.uk (Y.Z.); antonia.mey@ed.ac.uk (A.S.J.S.M.)
 - ⁶ University of Potsdam, Institute of Chemistry - Physical Chemistry, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany; amrmostafa@uni-potsdam.de (A.M.); bald@uni-potsdam.de (I.B.)
- * Correspondence: michael.weller@bam.de; Tel.: +49-30-8104-1150

1. Peptide modeling with additional information from the Markov State Model

Figure S1 illustrates the implied timescales from the 5 μ s trajectory of MD simulations. The plot was used to determine the 1ns lagtime for the building of the MSM and the coarse-graining into two dominant states, as there is one slow time scale observed in the system (blue line). Figure S2 shows the projection onto two dominant independent components from the TICA analysis.




Figure S1. The convergence test of the implied timescales associated with the five slowest processes. Solid lines refer to the timescales estimated from the maximum likelihood MSM. Dashed lines refer to the ensemble means computed from Bayesian sampling, with shaded areas indicating the 95 % confidence intervals. The black line refers to the equality of the timescale and lag time. The lag time was chosen to be 1 ns for estimating MSM, as most implied timescales converge at this lag time. The relatively large gap between the slowest (blue) and the second slowest (red) timescales suggests two metastable states.




Figure S2. Top: the free energy surface shown on the first two independent components (IC1 and IC2) with microstates colored according to the metastable distribution of metastable state 1 (left) and state 2 (right). A dark color indicates a high probability of metastable assignment. Red stars refer to the centers of metastable states. The populations of metastable states 1 and 2 are 3.4 % and 96.6 %, respectively, computed as the sum of their microstate populations. Bottom: the ensembles of 20 conformations sampled from metastable states 1 (left) and 2 (right). The linker is shown in purple, and the aptamer is shown in green. More details of MSM estimation and analysis can be found at: https://github.com/meyresearch/CCMV_apptamer_msm

2. Degradation of CCMV due to aging leaves




Figure S3. Silver-stained SDS-PAGE (left) of partially degraded CCMV and Western-Blot (right) with peroxidase-conjugated mAb BAM-CCMV-29-81. The Western Blot shows that the bands seen in the SDS-PAGE gel all belong to the CCMV capsid protein. The older the leaves from the purified CCMV (1-3 in ascending age: 3, 6 and 9 weeks), the more fragmentation of the capsid protein takes place. The CCMV coat protein has a mass of ~20 kDa. Its degradation products are visible in the bands corresponding to masses lower than that.





Figure S4. MALDI-TOF mass spectra of CCMV with varying degrees of proteolytic degradation depending on the time that the virus was allowed to propagate in the leaves until harvested. After 9 weeks of being infected, leaves yielded significantly higher amounts of degraded capsid proteins, as can be seen from multiple fragments smaller than 20,300 Da and their respective multiply charged species.

3. Determination of CCMV purity by reversed-phase chromatography (HPLC) and detection at 220 nm



No.	Retention Time min	Area mAU*min	Height mAU	Relative Area %	Relative Height %	Peak Width min
1	10,407	0,210	1,156	0,23	0,17	0,24
2	11,854	0,208	1,955	0,23	0,28	0,21
3	12,101	0,570	2,608	0,62	0,38	0,56
4	12,667	0,151	1,104	0,16	0,16	0,30
5	14,881	0,334	1,530	0,36	0,22	0,34
6	20,341	90,282	679,853	98,39	98,79	0,19
Total:		91,756	688,207	100,00	100,00	

Figure S5. Summary of the report from the chromatographic analysis software Chromeleon (version 7.2.10.23925) for the determination of CCMV purity with detection at 220 nm. For peak picking and integration during gradient separation from 3 to 23 minutes, default settings of the software used. The software finds 6 peaks in total and gives a relative peak area of 98.39% for CCMV in this sample.

4. Determination of particle integrity




Figure S6. Absorption spectrum of CCMV after affinity purification with peptide aptamer. The absorbance ratio of 260/280 nm gives 1.6.




Figure S7. Dynamic light scattering measurements show a narrow distribution of particles with a mean size of 32.7 nm, which is in good agreement with TEM and AFM data. Note that DLS measurements include the hydration sphere of the particles.

5. Titer of anti-CCMV rabbit serum




Figure S8. Titer determination of rabbit antiserum against CCMV. Microplates were coated with CCMV. Subsequently, CCMV antiserum (red) and pre-immune serum (blue) were added at different dilutions. Peroxidase-conjugated anti-rabbit antibodies were used for detection. The resulting titer is at least 1:100,000,000 as signals are still distinguishable from blank measurements with pre-immune serum at this dilution.

6. Peptide Mass Fingerprints of BAM-CCMV-29-81 for identity confirmation




Figure S9. Peptide Mass Fingerprint of BAM-CCMV-29-81 with MALDI-TOF-MS after partial acidic hydrolysis (top) and tryptic digestion (bottom). Antibody cleavage was performed according to Tscheuschner et.al. (2022). The raw data of the fingerprint spectra (see electronic supplement) may be used for future comparison for identity confirmation using the open-source software ABID 2.0 (<https://bam.de/ABID>).

7. Estimation of LOD and LOQ of CCMV immunoassay




Figure S10. Linear regression of the lowest five test points of the immunoassay calibration curve. LOD and LOQ were determined as $y_b + 3S_b$ and $y_b + 10S_b$ respectively, where y_b is the average absorption of 26 blanks and S_b its standard deviation. LOQ and LOQ absorption values (y) were translated to concentration values (x) with the linear calibration function above using the following fit: $x = (y - 0.0823 \pm 0.00065) / (0.029 \pm 0.00037)$. For the LOD, 0.25 $\mu\text{g/L}$, and for the LOQ, 0.79 $\mu\text{g/L}$ were determined.