

Article

Not peer-reviewed version

---

# Effect of Slaughter and Drying Temperatures on Protein Hydrolysis of Black Soldier Fly Larvae Meal

---

[María Rodríguez-Rodríguez](#) , [María José Sánchez-Muros](#) , [María del Carmen Vargas-García](#) \* ,  
Agnes Timea Varga , [Dmitri Fabrikov](#) , Fernando García Barroso

Posted Date: 8 May 2024

doi: 10.20944/preprints202405.0465.v1

Keywords: insect processing; insect meal; digestibility; hydrolysis degree; *Hermetia illucens*



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# Effect of Slaughter and Drying Temperatures on Protein Hydrolysis of Black Soldier Fly Larvae Meal

María Rodríguez-Rodríguez <sup>1</sup>, María José Sánchez-Muros <sup>2</sup>, María del Carmen Vargas-García <sup>2,\*</sup>, Ágnes Timea Varga <sup>2</sup>, Dmitri Fabrikov <sup>2</sup> and Fernando García Barroso <sup>1</sup>

<sup>1</sup> Department of Biology and Geology, CECOUAL, University of Almería, Carretera de Sacramento s/n, 04120, Almería, Spain; fbarroso@ual.es

<sup>2</sup> Department of Biology and Geology, CEIMAR, University of Almería, Carretera de Sacramento s/n, 04120, Almería, Spain; mjmuros@ual.es (M.J.S.-M.); avarga@ual.es (A.T.V.); df091@ual.es (D.F.)

\* Correspondence: María del Carmen Vargas García Phone: +34950015892 e-mail: mcvargas@ual.es

**Simple Summary:** Insects, because of their protein content and environmental advantages, have been considered a promising alternative to other traditional protein sources. Black soldier fly (BSFL) is one of the most studied insects as an alternative feed for farmed animals (mammals, poultry, and fish). This paper studies the effect on protein digestibility, proximal composition, and organic matter digestibility of BSFL meal according to the method of slaughter (frozen, blanched, Melacide<sup>®</sup>+frozen, liquid nitrogen) and drying temperature (50, 70 and 90 °C). The results showed that the best slaughter methods were liquid nitrogen and blanched. While 70 °C was the drying temperature that showed the best digestibility results while maintaining good hygienic-sanitary conditions.

**Abstract:** In recent years, the potential of insects has been explored as a sustainable protein alternative to sustain the growing world population. Differences in the way insects are processed can affect their proximate composition and digestibility. This work studies the effects of the combination of different types of slaughter methods and drying temperatures on the proximate composition, organic matter digestibility (OMd), hydrolysis degree (DH/NH<sub>2</sub> and DH/100 g DM), total hydrolysis (TH) and hygienic sanitary study of BSFL (black soldier fly larvae). Four types of slaughter methods were used: frozen (F), blanched + frozen (B), Melacide<sup>®</sup> + frozen (M) and liquid nitrogen (N), each with three drying temperatures (50, 70 and 90 °C). A negative correlation between acid detergent fibre (ADF) and protein digestibility parameters was obtained. 70 °C was the most suitable drying temperature, as it reported higher values of protein digestibility (DH, TH), resulting in hygienic and sanitary conditions suitable for food use. Slaughtering with liquid nitrogen and blanching were more convenient than traditional freezing or the use of Melacide<sup>®</sup> to achieve higher protein digestibility results.

**Keywords:** insect processing; insect meal; digestibility; hydrolysis degree; *Hermetia illucens*

## 1. Introduction

The increase in population in recent years has resulted in a search for more sustainable protein sources to cope with the increase in protein demand [1]. Insects have been reported as a promising alternative due to the possibility of being raised with organic waste, low water consumption, high efficiency in their feed conversion, low emissions of greenhouse gases and ammonia, few animal welfare problems, and low risk of transmission of zoonotic infections [2]. In addition, they have a high protein content and a good amino acid composition [3].

One of the most studied and mass-produced species is *Hermetia illucens* (Linneo, 1758) (Black Soldier Fly Larvae) [4]. Both industry and research show interest in this specie due to its potential to contribute to the circular economy through waste management as they consume 25-500 mg of fresh matter/larva/day, in a wide range of decomposing organic materials (food or agricultural waste), as well as being a source of protein [5–7]. It is a dipteran with a life cycle that includes the next stages:

egg, larva, prepupa, pupa, and adult. However, the prepupal and larval stages are the most widely used in research as an alternative feed because it is a good source of protein (40-56.9 % in dry matter), lipids (35-40 % in DM) and minerals [8,9]. For that reason, its inclusion in the diet of terrestrial and aquatic animals has been studied [8–11]. This is the case for farm animals such as piglets, chickens, ducks and quail, and farmed fish such as seabass, turbot, jian carp, rainbow trout, and shrimp [2,9,12,13]. The optimal level of inclusion of BSFL seems to vary by species [11], with full inclusion achieved for Atlantic salmon without compromising growth or sensory qualities [14].

If new protein sources are used as food and feed, their digestibility should be assessed, as protein quality is determined by amino acid composition and digestibility. In a recent review, Rodríguez-Rodríguez et al. [15] noted that BSFL shows very good results for crude protein digestibility (CPd), with a maximum of 89.7 % [16] and lower data for prepupae, with 50 % [12] using commercial enzymes simulating dog and duck digestion, respectively. These results have been obtained for raw insects, but if the aim is to market insect meal for human consumption, it is vitally important to investigate the most appropriate processing method. Kinyuru et al. [17] state that processing methods can affect the nutritional composition of insects, especially the digestibility of their proteins.

The processing methods aim to improve the texture, palatability, nutrient bioavailability, or food safety of insect meals [15]. Although in recent years, several researchers have studied the effect of treatments such as killing methods, drying methods, defatting or extrusion on the nutritional value of BSFL [18–21], results on the impact of different processing techniques, or combinations thereof, on their nutritional value or digestibility are still limited. As a result, a standard procedure for processing BSFL meal has not yet been established.

Currently, the method most companies use to slaughter different insect species is slow freezing, which can lead to enzymatic browning of insects [20]. This browning is caused by the action of endogenous phenol oxidases (POs), which oxidize monophenols and diphenols to form *o*-quinones [19], which act during slow freezing. In living insects, this browning is a defense reaction or sclerotization [22]. In addition to the color change, this browning could affect both protein quality and digestibility [20]. This browning can be prevented by chemical inhibitors (such as ascorbic acid, sulfite, or Melacide®) or physical treatments (boiling, blanching, or sonication) [23].

Another critical process is heat treatment during drying prior to grinding, as protein-rich insects can show non-enzymatic browning (Maillard reaction), which can affect the nutritional value of BSFL meal [24,25]. Depending on temperature, heat treatment could improve protein digestibility by denaturing them, increasing enzyme susceptibility, or inactivating possible inhibitors of specific enzymes [26]. However, if the temperatures are too high, reactions between amino acids can occur, which prevents hydrolysis [27].

Therefore, it is necessary to investigate the optimal processing methods for each insect species. The aim of this work is to determine the effects of different types of slaughter methods and drying temperatures on the nutritional value and protein digestibility, in addition to the evaluation of the hygienic-sanitary quality of the BSFL meals as a feed source for farm animals.

## 2. Materials and Methods

BSFL were purchased from Entomo AgroIndustrial (Cehegín, Murcia, Spain). The insects were divided into different lots and subjected to different types of slaughter: Frozen: without any additional treatment, live larvae were placed in zipper bags in the freezer at -18 °C. Blanched: larvae were placed in boiling water at 100 °C for 1 minute. They were removed from the water with a strainer and placed in zipper bags in the freezer at -18 °C. Melacide®: the larvae were mixed with Melacide® (15 grams per kilo of BSFL). They were placed in zipper bags in the freezer at -18 °C. Liquid nitrogen: was poured into a container with the larvae. The larvae were placed in zipper bags in the freezer at -18 °C. Then each lot was subdivided into three different lots, each of which was dried at different temperatures (50, 70 and 90 °C) and ground (Table 1).

**Table 1.** Treatments according to the slaughter method and drying temperature.

<b>Slaughtering</b>	<b>Frozen</b>			<b>Blanched+Frozen</b>			<b>Melacide®+Frozen</b>			<b>Liquid Nitrogen</b>		
T <sup>a</sup> dried (°C)	50	70	90	50	70	90	50	70	90	50	70	90
Treatment	F50	F70	F90	B50	B70	B90	M50	M70	M90	N50	N70	N90

In addition, soybean meal (SBM), albumin (A), and whey protein (W) were used as comparisons. Albumin and whey protein are pure proteins, they can serve as a control for the hydrolysis process. Soya, on the other hand, is a vegetable protein source traditionally used in animal feed.

### 2.1. Proximate Composition

The proximate composition was determined according to the Association of Official Analytical Chemists [28]. Dry matter and ash were determined by gravimetry after drying at 105 °C (AOAC #934.01) in a conventional oven and combustion at 500 °C in a muffle furnace (AOAC #942.05). The crude protein (CP) was determined by the Kjeldahl method (AOAC #954.01). The CP data of the BSFL pretreatments were calculated using the conversion factor of 4.76. This coefficient has been used following the recommendation of Janssen et al. [29] to avoid the overestimation of protein in insects. For the remainder of the materials analyzed (albumin, whey protein, and soybean meal), the usual conversion coefficient of 6.25 was used. The contents of acid detergent fiber (ADF) were determined according to the Van Soest et al. [30] method using an Ankom<sup>200</sup> fiber analyzer. All analyzes were performed in triplicate.

### 2.2. In Vitro Protein Hydrolysis

The protocol described by Huang et al. [18] was followed with some modifications, 100 g of BSFL meal was mixed with 4 mL NaCl (2 g/L) + HCl (7 mL/L), pH 2. The mixture was incubated at 38 °C for 15 min in a shaking water bath. The gastric phase was carried out using porcine pepsin (P7000 Sigma–Aldrich, St. Louis, MO) for 240 min under constant agitation at 38 °C. After gastric digestion, the pH was adjusted to 7 with 0.2 M NaOH solution and the intestinal phase was carried out using porcine pancreatin (P1750 Sigma–Aldrich) for 240 min under constant agitation at 38 °C.

Before adding enzymes from the gastric and intestinal phases, samples were taken to be used as blanks. In this way, the data from both phases are not affected by the amino acids present at time 0. As the intestinal phase occurs after the gastric phase, we consider that intestinal digestibility includes the sum of the gastric + intestinal phase.

The reaction of the collected samples was stopped with an equal volume of 20 % trichloroacetic acid (TCA). Three replicates per sample were analyzed.

### 2.3. Organic Matter Digestibility (OMd)

After in vitro hydrolysis, the samples were centrifuged. They were washed with 80 ml of distilled water and again centrifuged. The residue was dried at 100 °C for 24 h to calculate the dry matter and then at 500 °C to obtain the residue ash. The following formula was used to calculate the digestibility of organic matter:

$$\% \text{ OM Digestibility} = \frac{(\text{DM}_i - \text{Ash}_i) - (\text{DM}_f - \text{Ash}_f)}{(\text{DM}_i - \text{Ash}_i)} \times 100 \quad (1)$$

where DM<sub>i</sub> is the initial dry matter, DM<sub>f</sub> is the final dry matter expressed in grams, Ash<sub>i</sub> is the initial ash of the BSFL meal, and Ash<sub>f</sub> is the final ash of the BSFL after hydrolysis.

### 2.4. O-Phthaldialdehyde (OPA) Method

The OPA methodology described by Church et al. [31] is based on the reaction of OPA and 2-mercaptoethanol with amino groups. It was used for the calculation of α-amino groups, using L-leucine as a standard. α-Amino groups are found in most amino acids that make up proteins. This method measures the α-amino groups released after proteolysis.

The OPA results were expressed as % of the free amino groups in regard to total amino groups (DH/NH<sub>2</sub>) and % of the free amino groups in regard to dry matter (DH/DM) according to the following calculation.

The degree of hydrolysis (DH/100 NH<sub>2</sub>) relates the number of peptide bonds broken during hydrolysis (h) to the total number of peptide bonds present in the sample (h<sub>tot</sub>).

$$\text{DH/NH}_2(\%) = \frac{h}{h_{\text{tot}}} \times 100 \quad (2)$$

where h was calculated by measuring the free amino groups by OPA at the end time of hydrolysis (480 min), and h<sub>tot</sub> is the total amino groups in the sample [32]. These were determined by hydrolyzing 50 mg of the initial sample with 2.5 ml of 6 M HCl for 24 h at 100 °C.

DH/100 g of DM refers to the number of peptide bonds broken during protein hydrolysis (h) per 100 grams of dry matter.

$$\text{DH/DM}(\%) = \frac{h_{(0,240,480)}}{\text{DM}_i} \times 100 \quad (3)$$

where h was calculated by measuring the free amino groups with OPA at the beginning of hydrolysis (h<sub>0</sub>) and after the end of the gastric (h<sub>240</sub>) and intestinal (h<sub>480</sub>) phases, and DM<sub>i</sub> is the organic matter introduced at the start of protein hydrolysis.

Time 0 was used as a blank for the gastric and intestinal phases, while the intestinal phase is the sum of the gastric and intestinal phases.

The reason for expressing the results in dry matter is no other than to measure the yield of the sample, since the purpose of these meals is to be used in feed as a substitute for other proteins.

### 2.5. Total Hydrolysis (TH)

The degree of hydrolysis only measures the broken peptide bonds at the end of in vitro hydrolysis (480 min). However, the small peptides present that are susceptible to end hydrolysis are not considered. Therefore, to calculate the total hydrolysis, the final hydrolysis supernatant was completely hydrolyzed with the same amount of 12 M HCl for 24 h at 100 °C and neutralized with 6 M NaOH.

$$\text{Total hydrolysis}(\%) = \frac{\text{supernatant completely hydrolyzed}}{h_{\text{tot}}} \times 100 \quad (4)$$

### 2.6. Hygienic-Sanitary Quality Study: Microbial Contamination Indicators

To establish the level of hygienic-sanitary safety provided by each of the slaughter and drying temperature combinations tested, the following microbial groups, usually considered indicators of contamination, were estimated. Sulfite-reducing clostridia (*Clostridium perfringens*) using SPS (Sulfite Polymyxin Sulfadiazine) agar as medium and incubated at 37 °C for 48 h. The potential presence of the *Salmonella* spp. pathogen was investigated according to the following protocol: pre-enrichment stage (incubation of the sample at 37 °C for 18 h, in buffered peptone water where the meal to nutrient medium ratio was 1:9); enrichment stage (incubation in Rappaport-Vassiliadis broth, a selective medium, at 41.5 °C for 24 h, extendable to 48 h in case of initial negative results); selective isolation stage (seeding from the presumptive positive selection medium grown on Hektoen Agar incubated at 37 °C for 24 h); and confirmation stage (study of suspected colonies by growth characteristics on Kligler Agar, incubated at 37 °C for 24 h). The presence of Enterobacteria was investigated in VRBG agar plates grown for 24 h at 37 °C. The presence of total aerobic mesophilic bacteria (TAMB) was investigated in PCA agar plates grown for 24 h at 37 °C.

### 2.7. Statistical Analyses

The experimental results were expressed as means ± SD. Statistical differences in insect digestibility between slaughtering and drying temperature were analyzed with the ANOVA multivariate test, followed by a comparison of means (Tukey test). Proximate composition, OMD, DH/NH<sub>2</sub>, DH/100 g DM, TH and hygienic-sanitary results were analyzed using simple ANOVA followed by a multiple comparison of means (Tukey test). The correlations were analyzed using multivariate pairwise correlation analysis. Normality test was performed. The data did not follow a normal distribution, so a normalization of the data was performed. Data were normalized using a

two-step transformation (first, the variable was transformed into a percentile rank and then the inverse-normal transformation) [33]. The normal distribution of the residuals and of the transformed data was confirmed. The tests were carried out with IBM SPSS Statistics software (29.0.1.0). For graphic representation, all data in tables and figures were kept in raw form (prior to normalization) for clarity of presentation.

### 3. Results

#### 3.1. Proximate Composition

Regarding the proximate composition (Tables 2 and 3), no significant differences were found in the protein data from the different slaughtering and drying temperatures of BSFL. Slaughter methods did not show significant differences in ADF (Table 2). On the other hand, it was affected by drying temperature, showing the highest data for 50 and the lowest for 70 °C. The ash content was higher for the M treatment, while no significant differences were found with the other slaughter methods. Drying at 50 °C showed the highest ash data.

**Table 2.** MANOVA and ANOVA fit statistics CP, ADF, Ash in BSFL treatments, soybean meal, albumin, and whey protein. CP: Crude protein; ADF: Acid detergent fiber.

	Factor	F	df	p-value	Test
CP	Temperature	0.029	2	0.971	ANOVA multivariate
	Sacrifice	0.234	3	0.872	
	T x S	0.767	6	0.604	
ADF	Temperature	32.662	2	<0.001	
	Sacrifice	0.717	3	0.552	
	T x S	6.952	6	<0.001	
Ash	Temperature	48.620	2	<0.001	
	Sacrifice	28.390	3	<0.001	
	T x S	3.905	6	<0.001	
CP	-	4.772	14	<0.001	
ADF	-	25.780	14	<0.001	
Ash	-	31.305	14	<0.001	

**Table 3.** Proximate composition (g/100 g dry matter) of BSFL treatments, soybean meal, albumin, whey protein and relative standard deviation ( $\pm$ SD). Significant differences in the Tukey test ( $p < 0.05$ ) are represented by different letters. A: Albumin; W: Whey protein; SBM: Soybean meal; M: Melacide®; F: Frozen; B: Blanched; N: Nitrogen; 50, 70, 90: dried at 50 °C, 70 °C, 90 °C; CP: Crude protein; ADF: Acid detergent fiber; nsd: no significative differences.

		CP	ADF	Ash	Test	
Slaughtering	Melacide®	nsd	nsd	16.2 $\pm$ 1.8 <sup>a</sup>	ANOVA multivariate	
	Frozen	nsd	nsd	13.8 $\pm$ 1.7 <sup>b</sup>		
	Blanched	nsd	nsd	14.0 $\pm$ 1.6 <sup>b</sup>		
	Nitrogen	nsd	nsd	13.6 $\pm$ 0.3 <sup>b</sup>		
Temperature	50 °C	nsd	5.8 $\pm$ 0.8 <sup>a</sup>	16.0 $\pm$ 1.9 <sup>a</sup>		
	70 °C	nsd	4.6 $\pm$ 0.3 <sup>c</sup>	13.6 $\pm$ 1.3 <sup>b</sup>		
	90 °C	nsd	5.1 $\pm$ 0.5 <sup>b</sup>	13.6 $\pm$ 0.7 <sup>b</sup>		
Sample	A	98.9 $\pm$ 0.9 <sup>a</sup>	0.0 $\pm$ 0.0 <sup>g</sup>	1.1 $\pm$ 0.0 <sup>h</sup>		ANOVA simple
	W	90.5 $\pm$ 1.3 <sup>b</sup>	0.0 $\pm$ 0.0 <sup>g</sup>	3.2 $\pm$ 0.1 <sup>gh</sup>		
	SBM	50.0 $\pm$ 0.5 <sup>c</sup>	8.4 $\pm$ 0.8 <sup>a</sup>	6.9 $\pm$ 0.0 <sup>fgh</sup>		
	M50	37.2 $\pm$ 3.5 <sup>d</sup>	6.1 $\pm$ 0.3 <sup>bc</sup>	18.6 $\pm$ 0.3 <sup>a</sup>		
	M70	34.7 $\pm$ 3.3 <sup>d</sup>	4.3 $\pm$ 0.2 <sup>fg</sup>	15.5 $\pm$ 0.8 <sup>bc</sup>		
	M90	34.0 $\pm$ 0.5 <sup>d</sup>	5.2 $\pm$ 0.5 <sup>def</sup>	14.7 $\pm$ 0.0 <sup>bcd</sup>		

F50	34.9 ± 1.7 <sup>d</sup>	6.6 ± 0.3 <sup>b</sup>	15.7 ± 1.5 <sup>b</sup>
F70	34.6 ± 1.3 <sup>d</sup>	4.5 ± 0.3 <sup>fg</sup>	12.6 ± 0.8 <sup>efgh</sup>
F90	33.9 ± 0.4 <sup>d</sup>	4.9 ± 0.5 <sup>def</sup>	13.2 ± 0.3 <sup>defg</sup>
B50	34.2 ± 1.5 <sup>d</sup>	5.9 ± 0.3 <sup>bcd</sup>	16.0 ± 0.2 <sup>b</sup>
B70	34.6 ± 0.4 <sup>d</sup>	4.9 ± 0.5 <sup>def</sup>	12.7 ± 0.6 <sup>efgh</sup>
B90	35.8 ± 1.8 <sup>d</sup>	4.9 ± 0.4 <sup>def</sup>	13.3 ± 0.2 <sup>defg</sup>
N50	34.8 ± 1.3 <sup>d</sup>	4.6 ± 0.1 <sup>efg</sup>	13.8 ± 0.1 <sup>bcde</sup>
N70	34.5 ± 0.2 <sup>d</sup>	4.7 ± 0.1 <sup>ef</sup>	13.5 ± 0.5 <sup>cdef</sup>
N90	34.8 ± 0.6 <sup>d</sup>	5.5 ± 0.4 <sup>cde</sup>	13.3 ± 0.0 <sup>defg</sup>

### 3.2. Organic Matter Digestibility (OMd)

The OMd results, according to multivariate statistics (Tables 4 and 5), showed that there were significant differences in terms of slaughter, and frozen had the lowest data. Significant differences were also found in drying temperature, with the highest results obtained at 70 °C.


**Table 4.** MANOVA fit statistics Organic Matter Digestibility in BSFL treatments. ANOVA fit statistics Organic Matter Digestibility in BSFL treatments, albumin, whey protein and soybean meal. OMd: Organic Matter digestibility.

	Factor	F	df	p-value	Test
OMd	Temperature	10.635	2	<0.001	ANOVA multivariate
	Sacrifice	10.705	3	<0.001	
	T x S	10.423	6	<0.001	
	-	19.395	14	<0.001	ANOVA simple

**Table 5.** Organic matter digestibility of BSFL treatments, and relative standard deviation (±SD). Significant differences in the Tukey test (p<0.05) are represented by different letters. OMd: Organic Matter digestibility.

		OMd	Test
Slaughtering	Melacide®	88.3 ± 6.9 <sup>a</sup>	ANOVA multivariate
	Frozen	77.8 ± 9.0 <sup>b</sup>	
	Blanched	86.3 ± 3.2 <sup>a</sup>	
	Nitrogen	90.0 ± 6.9 <sup>a</sup>	
Temperature	50 °C	83.6 ± 10.7 <sup>b</sup>	
	70 °C	89.9 ± 5.7 <sup>a</sup>	
	90 °C	83.3 ± 5.4 <sup>b</sup>	

When the combined effect was analysed, the simple ANOVA (Figure 1), showed high digestibility for M50 and N70, which achieved similar data to albumin and whey protein.



**Figure 1.** Organic matter digestibility (%) of BSFL pretreatments, soybean meal, albumin and whey protein and relative standard deviation ( $\pm$ SD). Significant differences in the Tukey test are represented by different letters ( $p < 0.05$ ). A: Albumin; W: Whey protein; SBM: Soybean meal; M: Melacide®; F: Frozen; B: Blanched; N: Nitrogen; 50, 70, 90: dried at 50 °C, 70 °C, 90 °C.

### 3.3. Hydrolysis Degree (DH/NH<sub>2</sub>) and Total Hydrolysis (TH)

Regarding slaughter methods, there were differences between DH/NH<sub>2</sub> and TH (Tables 6 and 7). Blanched and Nitrogen were the methods showing the highest DH/NH<sub>2</sub>, however, the Nitrogen method showed the lowest TH. Therefore, if both parameters are considered, Blanched is the slaughter with the best results. Furthermore, when analyzing the effect of drying temperature, the highest values for DH/NH<sub>2</sub> was obtained at 70 °C. However, TH was not affected by temperature (Table 6).

**Table 6.** MANOVA fit statistics Hydrolysis degree and Total hydrolysis in BSFL treatments. ANOVA fit statistics Hydrolysis degree and Total hydrolysis in BSFL treatments, albumin, whey protein and soybean meal.


	Factor	F	df	p-value	Test
DH	Temperature	11.781	2	<0.001	ANOVA multivariate
	Sacrifice	32.258	3	<0.001	
	T x S	8.391	6	<0.001	
TH	Temperature	3.273	2	0.057	
	Sacrifice	6.224	3	0.003	
	T x S	1.341	6	0.281	
DH	-	26.423	14	<0.001	ANOVA simple
TH	-	8.875	14	<0.001	

**Table 7.** Hydrolysis degree and Total hydrolysis between different slaughtering methods and drying temperatures. Significant differences in the Tukey test ( $p < 0.05$ ) are represented by different letters. DH: Hydrolysis degree; TH: Total hydrolysis.

		DH	TH	Test
Slaughtering	Melacide®	22.4 $\pm$ 3.9 <sup>b</sup>	60.4 $\pm$ 2.3 <sup>ab</sup>	ANOVA multivariate
	Frozen	19.5 $\pm$ 3.7 <sup>c</sup>	62.6 $\pm$ 1.9 <sup>a</sup>	
	Blanched	26.9 $\pm$ 3.8 <sup>a</sup>	62.0 $\pm$ 1.5 <sup>a</sup>	
	Nitrogen	27.5 $\pm$ 2.8 <sup>a</sup>	57.6 $\pm$ 3.2 <sup>b</sup>	
Temperature	50 °C	24.1 $\pm$ 3.3 <sup>b</sup>	nsd	

70 °C	26.2 ± 3.6 <sup>a</sup>	nsd
90 °C	21.9 ± 6.2 <sup>b</sup>	nsd

Analyzing the combination of treatments (Figure 2), the highest DH values were shown for B90, and N70, which outperformed SBM, while the worst were M90 and F90. However, when analyzing TH, these differences are clearly dampened, the results between treatments being equal, and only N50 showed lower hydrolysis.



**Figure 2.** DH (% of total amino groups) and Total hydrolysis (% of total amino groups) of BSFL pretreatments, fishmeal, soybean meal, albumin, and whey protein and relative standard deviation ( $\pm$ SD). Significant differences in the Tukey test are represented by different letters ( $p < 0.05$ ). A: Albumin; W: Whey protein; SBM: Soybean meal; M: Melacide®; F: Frozen; B: Blanched; N: Nitrogen; 50, 70, 90: dried at 50 °C, 70 °C, 90 °C; DH: Hydrolysis Degree; TH: Total Hydrolysis.

### 3.4. Gastric and Intestinal Digestion

Amino groups were observed prior to the onset of hydrolysis (Tables 8 and 9).

**Table 8.** MANOVA fit statistics Amino groups (g / 100 g DM) in BSFL treatments. ANOVA fit statistics Amino groups (g / 100 g DM) in BSFL treatments, albumin, whey protein and soybean meal.

	Factor	F	df	p-value	Test
Initial (0')	Temperature	47.222	2	<0.001	ANOVA multivariate
	Sacrifice	15.712	3	<0.001	
	T x S	4.721	6	0.003	
Gastric (240')	Temperature	0.300	2	0.744	
	Sacrifice	23.388	3	<0.001	
	T x S	7.631	6	<0.001	
Intestinal (480')	Temperature	12.982	2	<0.001	
	Sacrifice	42.667	3	<0.001	
	T x S	8.487	6	<0.001	
Initial (0')	-	19.021	14	<0.001	ANOVA simple
Gastric (240')	-	24.055	14	<0.001	
Intestinal (480')	-	26.855	14	<0.001	

**Table 9.** Amino groups (g / 100 g DM) of BSFL treatments, soybean meal, albumin, and whey protein at the start and end of the gastric and intestinal phases of protein hydrolysis and relative standard deviation ( $\pm$ SD). Significant differences in the Tukey test ( $p < 0.05$ ) obtained in the different ANOVAS are represented by different letters. A: Albumin; W: Whey protein; SBM: Soybean meal; M: Melacide®; F: Frozen; B: Blanched; N: Nitrogen; 50, 70, 90: dried at 50 °C, 70 °C, 90 °C; nsd: no significative differences.

		Initial (0')	Gastric (240')	Intestinal (480')	Test	
Slaughtering	Melacide	4.5 $\pm$ 0.9 <sup>a</sup>	6.1 $\pm$ 2.0 <sup>b</sup>	8.4 $\pm$ 1.5 <sup>c</sup>	ANOVA multivariate	
	Frozen	4.2 $\pm$ 0.4 <sup>b</sup>	4.6 $\pm$ 0.3 <sup>c</sup>	7.2 $\pm$ 1.4 <sup>d</sup>		
	Blanched	3.3 $\pm$ 0.9 <sup>c</sup>	8.3 $\pm$ 1.1 <sup>a</sup>	9.8 $\pm$ 1.3 <sup>b</sup>		
	Nitrogen	4.2 $\pm$ 0.9 <sup>b</sup>	6.5 $\pm$ 1.0 <sup>b</sup>	11.2 $\pm$ 1.1 <sup>a</sup>		
Temperature	50 °C	4.9 $\pm$ 0.4 <sup>a</sup>	nsd	9.2 $\pm$ 1.6 <sup>ab</sup>		
	70 °C	4.1 $\pm$ 0.8 <sup>b</sup>	nsd	9.9 $\pm$ 1.6 <sup>a</sup>		
	90 °C	3.2 $\pm$ 0.5 <sup>c</sup>	nsd	8.3 $\pm$ 2.4 <sup>b</sup>		
Sample	A	4.0 $\pm$ 0.1 <sup>cdef</sup>	18.8 $\pm$ 0.5 <sup>ab</sup>	39.3 $\pm$ 0.9 <sup>ab</sup>		ANOVA simple
	W	4.7 $\pm$ 0.1 <sup>abc</sup>	18.5 $\pm$ 0.8 <sup>a</sup>	46.0 $\pm$ 1.5 <sup>a</sup>		
	SBM	1.8 $\pm$ 0.0 <sup>g</sup>	9.4 $\pm$ 0.2 <sup>abc</sup>	11.2 $\pm$ 1.6 <sup>cd</sup>		
	M50	5.1 $\pm$ 0.1 <sup>ab</sup>	8.6 $\pm$ 0.7 <sup>cde</sup>	9.4 $\pm$ 0.5 <sup>def</sup>		
	M70	5.1 $\pm$ 0.4 <sup>a</sup>	4.6 $\pm$ 0.3 <sup>g</sup>	9.3 $\pm$ 1.0 <sup>def</sup>		
	M90	3.4 $\pm$ 0.1 <sup>defg</sup>	5.0 $\pm$ 0.6 <sup>fg</sup>	6.6 $\pm$ 0.6 <sup>fg</sup>		
	F50	4.7 $\pm$ 0.1 <sup>abc</sup>	4.5 $\pm$ 0.0 <sup>g</sup>	7.6 $\pm$ 0.7 <sup>efg</sup>		
	F70	4.2 $\pm$ 0.2 <sup>cde</sup>	4.6 $\pm$ 0.4 <sup>g</sup>	8.3 $\pm$ 1.0 <sup>efg</sup>		
	F90	3.7 $\pm$ 0.1 <sup>cdefg</sup>	4.7 $\pm$ 0.4 <sup>fg</sup>	5.6 $\pm$ 0.5 <sup>g</sup>		
	B50	4.4 $\pm$ 0.4 <sup>bcd</sup>	6.8 $\pm$ 0.2 <sup>defg</sup>	8.2 $\pm$ 0.5 <sup>efg</sup>		
	B70	3.1 $\pm$ 0.2 <sup>efg</sup>	8.9 $\pm$ 0.2 <sup>cde</sup>	10.2 $\pm$ 0.8 <sup>cde</sup>		
	B90	2.4 $\pm$ 0.1 <sup>fg</sup>	9.1 $\pm$ 0.1 <sup>bcd</sup>	10.8 $\pm$ 0.8 <sup>cd</sup>		
	N50	5.3 $\pm$ 0.3 <sup>a</sup>	6.4 $\pm$ 0.4 <sup>defg</sup>	11.5 $\pm$ 0.6 <sup>bcd</sup>		
N70	3.9 $\pm$ 0.2 <sup>cdef</sup>	7.3 $\pm$ 0.3 <sup>cdef</sup>	12.0 $\pm$ 0.9 <sup>bc</sup>			
N90	3.3 $\pm$ 0.3 <sup>defg</sup>	5.6 $\pm$ 1.2 <sup>efg</sup>	10.0 $\pm$ 0.9 <sup>cde</sup>			

Considering the drying temperatures data at the end of the hydrolysis, the lowest results were obtained at 90 °C. Regarding the method of slaughter, the highest values of hydrolysis during gastric digestion were obtained for Blanched, while in the intestinal digestion, the highest values of hydrolysis correspond to the simples treated with nitrogen.

When taking into account the combination of treatments, the samples matched the SBM results at the end of the hydrolysis, with the exception of B50, M (F50, F70, F90), and F (F50, F70, F90), which showed a significantly lower amount of free amino groups.

### 3.5. Correlations

As shown in Table 10, Omd is positively correlated with DH and TH. Furthermore, there was also a positive correlation between CP and the different protein digestibility measurements. Additionally, protein digestibility variables were positively correlated ( $r=0.456$ ).

**Table 10.** Correlations between proximal composition, organic matter digestibility, and protein digestibility (DH and TH). The correlation is significant at the 0.05 level. CP: Crude Protein; ADF: Acid Detergent Fibre; Omd: Organic Matter digestibility; DH: Hydrolysis Degree; TH: Total Hydrolysis; p: p-value.

	TH	p	DH	p	Omd	p	Ash	p	ADF	CP
CP	0.329	0.031	0.479	0.001	0.607	<0.001	-0.408	0.006	nsd	-
ADF	-0.385	0.011	-0.489	<0.001	-0.383	0.011	0.450	0.002	-	-
Ash	-0.302	0.047	-0.374	0.012	nsd	-	-	-	-	-

OMd	0.454	0.002	0.636	<0.001	-	-	-	-	-	-
DH	0.456	0.002	-	-	-	-	-	-	-	-

Furthermore, we obtained a negative correlation between ADF-DH ( $r=-0.489$ ) and ADF-TH ( $r=-0.385$ ).

### 3.6. Hygienic-Sanitary Quality Study: Microbial Contamination Indicators

The results obtained (Table 11) show a clear influence of the treatments applied on the levels of microbial groups indicative of contamination present in insects. The clearest case was that of the Enterobacteriaceae group, as all the combinations tested led to the disappearance of this type of bacteria, regardless of the method of slaughter or the drying temperature. The latter generated the lowest degree of Thermophilic Acidophilic Bacteria (TAMB) reduction due to temperature action. The presence of *Salmonella* and enterobacteria, was also safely controlled by the drying temperature, regardless of the slaughter method applied. However, in this case, the thermal value required was somewhat higher, as treatments performed at 50 °C were not sufficient to promote the disappearance of this pathogen. Finally, and in relation to SRCs, the only problematic batch was that of insects slaughtered with liquid nitrogen.

**Table 11.** Hygienic-sanitary quality, according to the levels of microbial contamination indicators, of insects after processing according to the different treatments. Values are expressed as Log<sub>10</sub> CFU mL<sup>-1</sup>. Mean values ± SD are shown. Significant differences in the Tukey test are represented by different letters ( $p<0.05$ ). A: Albumin; W: Whey protein; SBM: Soybean meal; M: Melacide®; F: Frozen; B: Blanched; N: Nitrogen; 50, 70, 90: dried at 50 °C, 70 °C, 90 °C. TAMB: Thermophilic Acidophilic Bacteria; SRC: sulfite-reducing Clostridia. \*Detection limit: 1 log CFU mL<sup>-1</sup>.

	TAMB	Enterobacteria	SRC*	Salmonella
Control	5.92 ± 0.35 <sup>ab</sup>	5.32 ± 0.13 <sup>a</sup>	Negative	Positive
M50	7.25 ± 0.33 <sup>a</sup>	< 1 <sup>b</sup>	Negative	Positive
M70	2.92 ± 2.53 <sup>cde</sup>	< 1 <sup>b</sup>	Negative	Negative
M90	1.65 ± 1.43 <sup>ef</sup>	< 1 <sup>b</sup>	Negative	Negative
F50	6.14 ± 0.25 <sup>ab</sup>	< 1 <sup>b</sup>	Negative	Positive
F70	4.84 ± 0.50 <sup>bc</sup>	< 1 <sup>b</sup>	Negative	Negative
F90	2.11 ± 1.92 <sup>de</sup>	< 1 <sup>b</sup>	Negative	Negative
B50	6.97 ± 0.13 <sup>a</sup>	< 1 <sup>b</sup>	Negative	Positive
B70	3.54 ± 0.92 <sup>cd</sup>	< 1 <sup>b</sup>	Negative	Negative
B90	< 1 <sup>f</sup>	< 1 <sup>b</sup>	Negative	Negative
N50	1.71 ± 0.06 <sup>ef</sup>	< 1 <sup>b</sup>	Positive	Negative
N70	< 1 <sup>f</sup>	< 1 <sup>b</sup>	Positive	Negative
N90	< 1 <sup>f</sup>	< 1 <sup>b</sup>	Positive	Negative

## 4. Discussion

To realize the full potential of insects as a novel and sustainable food in the near future, the insect mass-rearing industry needs to develop and optimize insect meal processing systems to maximize their nutritional potential and food safety. The high variability observed in the nutritional composition of the same insect species shows that, in addition to other factors, the slaughter and drying method can clearly affect it. All this makes it necessary to investigate the effects of common processing, slaughtering, and drying conditions on the nutritional characteristics and digestive proteolysis of BSLF meal.

It appears clear that the slaughter method and the drying temperature do not affect the protein content of BSLF (Table 3). However, drying at 70 °C showed a lower proportion of fibre and ash, and as pointed out by several studies [34,35], higher levels of fibre could be indicative of worse digestibility.

OMd has traditionally been used to assess the digestibility of food [36]. Based on our OMD results (Figure 1), Frozen is the worst slaughter method and 70 °C is the best drying temperature. In general, our data when combining both processes were similar to those obtained by Bosch et al. [37] who obtained 84.3 % for BSFL, and poultry meat meal (85.8 %). However, except for N70 (98.1%), they were less than SBM (93.8 %).

The DH data obtained (Table 7) showed higher results for the Nitrogen and Blanched slaughter. Frozen and Nitrogen are both freezing slaughter treatments, and their difference is in the freezing temperature and therefore the speed of slaughter, as nitrogen slaughter is instantaneous. This could explain the differences in their results. As it is a fast slaughter, insects do not have time to develop the strategies they carry out when stressed, such as enzymatic browning, which can hinder protein hydrolysis. Browning can affect protein digestibility and quality, and the melanization process seems to be related to protein aggregation due to insect stress [20]. According to Janssen et al. [29], five enzymes may play an important role in enzymatic browning in insects: phenol oxidase, laccase, tyrosine hydroxylase, DOPA decarboxylase, and peroxidase. These are related to defense mechanisms and react quickly to stress. They are also involved in the sclerotization of the exoskeleton. Wessels et al. [38] note that, of these, phenol oxidase is the key in browning and is likely to remain active after freezing. This is because there are anti-freeze proteins in insects that ensure that part of the water does not finish freezing, and that enzymatic reactions that favor the formation of aggregates and melanization continue to take place. This could explain the higher DH data in N with respect to F, but also the higher results for B (Table 7). Some studies confirm the improvement in digestibility using blanching, such as that of Leni et al. [20], who obtained higher DH results for BSFL in blanched (32 %) than frozen (16.5 %). In the industry, blanching has been used to inactivate enzymes and preserve food quality. This is because blanching blocks melanization by inactivating phenol oxidase [20]. Different studies have shown less browning with blanching than with traditional freezing slaughter (-18 °C) [20,30,39]. Other studies found no improvement in digestibility with blanching [27,40,41]. Even in Janssen et al. [19], DH data for *Alphitobius diaperinus* were the same in raw and blanched but did not improve digestibility in BSFL or *Tenebrio molitor*.

As indicated in the Materials and Methods section, in this experiment, we used Melacide®. It is an antioxidant, antimelanosis, color, and texture stabilizer additive used for frozen crustaceans and mollusks. No studies to date using Melacide® in insects have been found in the literature. This treatment was carried out precisely to try to reduce the possible melanization of the insects during slaughter. DH data showed higher results with Melacide® (M) than without Melacide® (F) (Table 7); therefore, it is a desirable treatment if slaughter is to be carried out by freezing.

Regarding drying temperatures, the highest DH were obtained at 70 °C and the lowest when increasing the temperature to 90 °C (Table 7). These data follow the same line as those obtained by Mancini et al. [40], in which they obtained a better digestibility for TM in a traditional oven at 70 °C than at 150 °C. High heat treatments appear to make proteins more compact by polymerizing them [18]. Janssen et al. [19] observed that this denaturation and aggregation reduced the accessibility of digestive enzymes, decreasing protein digestibility.

When the treatments were compared with SBM, all showed similar or even higher DH (B70, B90, N50, and N70) (Figure 2).

However, when these data are referred to as TH, which considers DH plus other small peptides that have been solubilized after hydrolysis and are expected to be digestible, the statistical differences disappear, showing similar digestibility among almost all treatments. TH was higher in the treatments with lower DH (Figure 2). This could indicate a slower velocity of digestion for these samples, maybe because they have a higher content and/or a less digestible structure of the scleroprotein in their exoskeleton. However, the TH data indicate that these differences are due to small peptides, not accounted for by DH, which are likely to be absorbed. This may be relevant if we want to value these insect meals as feed. Protein digestibility may be underestimated if only DH is taken into account. Differences were found between slaughter with N and the rest of the treatments. Regarding the drying temperature, again the best results were obtained at 70 °C (Table 7).

When evaluating free amino groups per g of sample (in DM), high initial values were observed (Table 9). This may indicate that the endogenous enzymes of the insect are active from the beginning. Therefore, as these enzymes are inactivated by blanching (B), this method of slaughter releases the fewest amino groups. According to Leni et al. [42] it was observed partial solubilization of insect protein under hydrolysis conditions without enzymatic action below 50 %. As stated above, these data were higher than those obtained by Hall et al. [43] (20 % for cricket) and similar to those of a previous study by Caligiani et al. [8] (43 % for BSFL). They suggested that these differences may depend on temperature, time, pH, and enzyme conditions. After the gastric and intestinal phases, it seems clear that the slaughter method with the lowest release of amino groups was F, and the methods with the highest release were B and N (Table 9).

The decrease in amino group content with increasing drying temperature (Table 9) could be due to an aggregation of proteins that hinders their breakage and therefore the release of their amino groups. It is known that increasing or decreasing temperature destabilizes proteins and can promote their aggregation [44]. Temperatures above 60-70 °C can destroy hydrogen bonds and electrostatic and hydrophobic interactions of proteins, and disorganize their structure, thus favoring protein denaturation and aggregation [45]. Santiago et al. [46], studying protein aggregation in *Gryllus assimilis*, observed that aggregation progressively increased from practically non-existent at 65 °C to reaching its maximum at 90-95 °C. Janssen et al. [19] linked protein denaturation and aggregation with worsening digestibility, as aggregates hinder the action of enzymes for protein degradation.

After the gastric and intestinal phases, there was a higher percentage of amino groups released in A and W as expected. In general, the BSFL meals had similar protein yields for the same amount of dry matter as SBM, and only a few BSFL treatments (M90, F50, F70, F90 y B50) showed significantly lower data (Table 9).

The diversity in the way authors approach protein digestibility makes it difficult to compare results. Therefore, we consider it of interest to study the potential correlation between the different parameters. A positive correlation was found between CP and the different protein digestibility measurements (Table 10). This could indicate that the higher the protein content, the higher the protein digestibility. According to Finke [34], the ADF of insects consists of chitin and associated cuticular proteins, and since the determination of chitin is complex, the ADF can be used to estimate digestibility. Protein digestibility is affected because chitin is neither degraded nor absorbed in the digestive tract [47]. Similar to us, Marono et al. [35] found a negative correlation between ADF and digestibility for BSFL. Therefore, although it is not a parameter that measures digestibility, when deep studies are not possible, determining ADF can be a rough predictive tool for the degree of protein digestibility of the insect. In general, ash and ADF showed a negative correlation with the different protein digestibility parameters analysed; nevertheless, this would require further research. The correlation between CP and TH or DH showed that both parameters can be used interchangeably.

The presence of certain microbial species in final products made from insects and destined for the food market is of great importance from a hygienic-sanitary perspective. In this sense, guaranteeing the suitability of the processing protocols applied to ensure their microbial quality gives these protocols additional value in the food sector. The efficiency in the progressive disappearance of the Enterobacteriaceae group seems to be mainly a consequence of the application of drying temperatures, as noted in other studies [48], which describe the high efficiency of thermal protocols, even with low potential, to control the presence of this type of bacteria. In the case of TAMB, the observed impact was not as considerable, although the reductions in the counts obtained were more significant as the drying temperature increased (Table 11). Thus, the important role of temperature in relation to the microbiological quality of the insect seems to be confirmed. However, the method of slaughter also contributed to the observed differences, with nitrogen treatment being the most favorable, compared to freezing. Treatments performed at 50 °C were not sufficient to ensure the avoidance of *Salmonella* (Table 11). Therefore, it is important to ensure the correct application of the treatment; otherwise, the resistance of this bacterium tends to increase [49]. Although there are few studies that combine the study of microbial quality with slaughter techniques based on the use of

liquid nitrogen, the presence of SRC in insects processed by this type of methodology has been described [50]. Taking into account what was mentioned, and from a microbial perspective, it seems logical to conclude that the temperature applied during the drying stage is more decisive than the method of slaughtering. Temperatures of approximately 70 °C should ensure the hygienic-sanitary quality of insects in terms of their use in the food sector.

## 5. Conclusions

ADF and DH/100 g DM are correlated with DH/NH<sub>2</sub> and TH. As these parameters are faster and simpler, they could be useful in cases where a very deep study on digestibility is not necessary or where there are time or economic limitations.

Regarding slaughter methods, Blanched and liquid Nitrogen showed higher results than Frozen and Melacide®, both in terms of protein digestibility and hygienic quality. However, in terms of economy and easy accessibility, blanching is more advisable.

The drying temperature plays an important role. Generally, 70 °C resulted in better digestibility data than 50 °C. At 90 °C the lowest results were obtained because at higher temperatures there is a risk of protein aggregation processes that would hinder digestibility.

**Author Contributions:** Conceptualization, María José Sánchez-Muros and Fernando G. Barroso; Data curation, María Rodríguez Rodríguez, Agnes Timea Varga and Dmitri Fabrikov; Formal analysis, María Rodríguez Rodríguez; Funding acquisition, María José Sánchez-Muros and Fernando G. Barroso; Methodology, María Rodríguez Rodríguez, María del Carmen Vargas-García and Agnes Timea Varga; Project administration, María José Sánchez-Muros and Fernando G. Barroso; Resources, María José Sánchez-Muros, María del Carmen Vargas-García and Fernando G. Barroso; Software, María Rodríguez Rodríguez and María del Carmen Vargas-García; Supervision, María José Sánchez-Muros and Fernando G. Barroso; Validation, María José Sánchez-Muros, María del Carmen Vargas-García and Dmitri Fabrikov; Writing – original draft, María Rodríguez Rodríguez; Writing – review & editing, María Rodríguez Rodríguez, María José Sánchez-Muros, María del Carmen Vargas-García, Agnes Timea Varga, Dmitri Fabrikov and Fernando G. Barroso.

**Data Availability Statement:** Data available in a publicly accessible repository. The data presented in this study are openly available in Repositorio Institucional de la Universidad de Almería at <http://hdl.handle.net/10835/14827>.

**Acknowledgments:** To the project “Determination of the best insect meal processing techniques to optimize their nutritional value, acceptability, assimilation and food safety” (UAL2020-AGR-B1882). Project co-financed by funds from the Operational Program Feder Andalucía 2014-2020 and by the Consejería de Universidad, Investigación e Innovación (Junta de Andalucía).

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Ojha, S.; Bekhit, A. E.-D.; Grune, T.; Schlüter, O. K. Bioavailability of nutrients from edible insects. *Current Opinion in Food Science* 2021, 41, 240–248. <https://doi.org/10.1016/j.cofs.2021.08.003>
2. Magalhães, R.; Sánchez-López, A.; Leal, R.S.; Martínez-Llorens, S.; Oliva-Teles, A.; Peres, H. Black soldier fly (*Hermetia illucens*) pre-pupae meal as a fish meal replacement in diets for European seabass (*Dicentrarchus labrax*). *Aquaculture* 2017, 476, 79-85. <https://doi.org/10.1016/j.aquaculture.2017.04.021>
3. Barroso, F.G.; de Haro, C.; Sánchez-Muros, M.-J.; Venegas, E.; Martínez-Sánchez, A.; Pérez-Bañón, C. The potential of various insect species for use as food for fish. *Aquaculture* 2014, 422-423, 193-201. <https://doi.org/10.1016/j.aquaculture.2013.12.024>
4. Eke, M.; Tougeron, K.; Hamidovic, A.; Tinkeu, L. S. N.; Hance, T.; Renoz, F. Deciphering the functional diversity of the gut microbiota of the black soldier fly (*Hermetia illucens*): recent advances and future challenges. *Animal Microbiome* 2023, 5(1). <https://doi.org/10.1186/s42523-023-00261-9>
5. Barroso, F. G., Sánchez-Muros, M.-J., Segura, M., Morote, E., Torres, A., Ramos, R., & Guil, J.-L. (2017). Insects as food: Enrichment of larvae of *Hermetia illucens* with omega 3 fatty acids by means of dietary modifications. *Journal of Food Composition and Analysis: An Official Publication of the United Nations University, International Network of Food Data Systems*, 62, 8–13. <https://doi.org/10.1016/j.jfca.2017.04.008>
6. Cohn, Z.; Latty, T.; Abbas, A. Understanding dietary carbohydrates in black soldier fly larvae treatment of organic waste in the circular economy. *Waste Management (New York, N.Y.)* 2022, 137, 9–19. <https://doi.org/10.1016/j.wasman.2021.10.013>

7. Walter, A.; Klammsteiner, T.; Gassner, M.; Heussler, C. D.; Kapelari, S.; Schermer, M.; Insam, H. Black soldier fly school workshops as means to promote circular economy and environmental awareness. *Sustainability* 2020, 12(22), 9574. <https://doi.org/10.3390/su12229574>
8. Caligiani, A.; Marseglia, A.; Leni, G.; Baldassarre, S.; Maistrello, L.; Dossena, A.; Sforza, S. Composition of black soldier fly prepupae and systematic approaches for extraction and fractionation of proteins, lipids and chitin. *Food Research International* 2018, 105, 812-820. <https://doi.org/10.1016/j.foodres.2017.12.012>
9. Li, S.; Ji, H.; Zhang, B.; Zhou, J.; Yu, H. Defatted black soldier fly (*Hermetia illucens*) larvae meal in diets for juvenile Jian carp (*Cyprinus carpio* var. Jian): Growth performance, antioxidant enzyme activities, digestive enzyme activities, intestine and hepatopancreas histological structure. *Aquaculture* 2017, 477, 62-70. <https://doi.org/10.1016/j.aquaculture.2017.04.015>
10. Dumas, A.; Raggi, T.; Barkhouse, J.; Lewis, E.; Weltzien, E. The oil fraction and partially defatted meal of black soldier fly larvae (*Hermetia illucens*) affect differently growth performance, feed efficiency, nutrient deposition, blood glucose and lipid digestibility of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 2018, 492, 24-34. <https://doi.org/10.1016/j.aquaculture.2018.03.038>
11. Fabrikov, D.; Sánchez-Muros, M. J.; Barroso, F. G.; Tomás-Almenar, C.; Melenchón, F.; Hidalgo, M. C.; Morales, A. E.; Rodríguez-Rodríguez, M.; Montes-Lopez, J. Comparative study of growth performance and amino acid catabolism in *Oncorhynchus mykiss*, *Tinca tinca* and *Sparus aurata* and the catabolic changes in response to insect meal inclusion in the diet. *Aquaculture* 2020, 529, 735731. <https://doi.org/10.1016/j.aquaculture.2020.735731>
12. Kovitvadh, A.; Chundang, P.; Thongprajukaew, K.; Tirawattanawanich, C.; Srikachar, S.; Chotimanothum, B. Potential of insect meals as protein sources for meat-type ducks based on in vitro digestibility. *Animals* 2019, 9(4), 155. <https://doi.org/10.3390/ani9040155>
13. Kovitvadh, A.; Chundang, P.; Pliantiangtam, N.; Thongprajukaew, K.; Tirawattanawanich, C.; Suwanasopee, T.; Koonawootrittriron, S. Screening of in vitro nutrient digestibility coefficients of selected insect meals in broiler chickens, black-meat chickens and quails. *Journal of Animal Physiology and Animal Nutrition* 2021, 105, 305-315. <https://doi.org/10.1111/jpn.13451>
14. Li, Y.; Kortner, T.M.; Chikwati, E.M.; Munang'andu, H.M.; Lock, E.-J.; Krogdahl, Å. Gut health and vaccination response in pre-smolt Atlantic salmon (*Salmo salar*) fed black soldier fly (*Hermetia illucens*) larvae meal. *Fish & Shellfish Immunology* 2019, 86, 1106-1113. <https://doi.org/10.1016/j.fsi.2018.12.057>
15. Rodríguez-Rodríguez, M.; Barroso, F.G.; Fabrikov, D.; Sánchez-Muros, M.J. In vitro crude protein digestibility of insects: A review. *Insects* 2022, 13(8), 682. <https://doi.org/10.3390/insects13080682>
16. Bosch, G.; Vervoort, J.J.M.; Hendriks, W.H. In vitro digestibility and fermentability of selected insects for dog foods. *Animal Feed Science and Technology* 2016, 221, 174-184. <https://doi.org/10.1016/j.anifeedsci.2016.08.018>
17. Kinyuru, J.N.; Kenji, G.M.; Njoroge, S.M.; Ayieko, M. Effect of processing methods on the in vitro protein digestibility and vitamin content of edible winged termite (*Macrotermes subhylanus*) and grasshopper (*Ruspolia differens*). *Food and Bioprocess Technology* 2010, 3, 778-782. <https://doi.org/10.1007/s11947-009-0264-1>
18. Huang, C.; Feng, W.; Xiong, J.; Wang, T.; Wang, W.; Wang, C.; Yang, F. Impact of drying method on the nutritional value of the edible insect protein from black soldier fly (*Hermetia illucens* L.) larvae: amino acid composition, nutritional value evaluation, in vitro digestibility, and thermal properties. *European Food Research and Technology* 2019, 245, 11-21. <https://doi.org/10.1007/s00217-018-3136-y>
19. Janssen, R.H.; Vincken, J.-P.; Arts, N.J.G.; Fogliano, V.; Lakemond, C.M.M. Effect of endogenous phenoloxidase on protein solubility and digestibility after processing of *Tenebrio molitor*, *Alphitobius diaperinus* and *Hermetia illucens*. *Food Research International* 2019, 121, 684-690. <https://doi.org/10.1016/j.foodres.2018.12.038>
20. Leni, G.; Caligiani, A.; Sforza, S. Killing method affects the browning and the quality of the protein fraction of Black Soldier Fly (*Hermetia illucens*) prepupae: a metabolomics and proteomic insight. *Food Research International* 2019, 115, 116-125. <https://doi.org/10.1016/j.foodres.2018.08.021>
21. Ottoboni, M.; Spranghers, T.; Pinotti, L.; Baldi, A.; De Jaeghere, W.; Eeckhout, M. Inclusion of *Hermetia Illucens* larvae or prepupae in an experimental extruded feed: process optimisation and impact on in vitro digestibility. *Italian Journal of Animal Science* 2018, 17(2), 418-427. <https://doi.org/10.1080/1828051x.2017.1372698>
22. Terwilliger, N.B. Hemolymph proteins and molting in crustaceans and insects. *American Zoologist* 1999, 39(3), 589-599. <https://doi.org/10.1093/icb/39.3.589>
23. Özoğlu, H.; Bayındırlı, A. Inhibition of enzymic browning in cloudy apple juice with selected antibrowning agents. *Food Control* 2022, 13(4-5), 213-221. [https://doi.org/10.1016/s0956-7135\(02\)00011-7](https://doi.org/10.1016/s0956-7135(02)00011-7)
24. Oliver, C.M.; Melton, L.D.; Stanley, R.A. Creating proteins with novel functionality via the Maillard reaction: a review. *Critical Reviews in Food Science and Nutrition* 2006, 46(4), 337-350. <https://doi.org/10.1080/10408690590957250>

25. Zhen, Y.; Chundang, P.; Zhang, Y.; Wang, M.; Vongsangnak, W.; Pruksakorn, C.; Kovitvadhi, A. Impacts of killing process on the nutrient content, product stability and in vitro digestibility of black soldier fly (*Hermetia illucens*) larvae meals. *Applied Sciences (Basel, Switzerland)* 2020, 10(17), 6099. <https://doi.org/10.3390/app10176099>
26. Opstvedt, J.; Nygård, E.; Samuelson, T.A.; Venturini, G.; Luzzana, U.; Mundheim, H. Effect on protein digestibility of different processing conditions in the production of fish meal and fish feed: Processing effects on protein quality of extruded fish feed. *Journal of the Science of Food and Agriculture* 2003, 83(8), 775-782. <https://doi.org/10.1002/jsfa.1396>
27. Caparros Megido, R.; Poelaert, C.; Ernens, M.; Liotta, M.; Blecker, C.; Danthine, S.; Tyteca, E.; Haubruge, É.; Alabi, T.; Bindelle, J.; Francis, F. Effect of household cooking techniques on the microbiological load and the nutritional quality of mealworms (*Tenebrio molitor* L. 1758). *Food Research International* 2018, 106, 503-508. <https://doi.org/10.1016/j.foodres.2018.01.002>
28. AOAC. Official Methods of Analysis of the Association of Official Analytical Chemists International 2005, 18th edn. AOAC International, Gaithersburg, USA.
29. Janssen, R.H.; Lakemond, C.M.M.; Fogliano, V.; Renzone, G.; Scaloni, A.; Vincken, J.-P. Involvement of phenoloxidase in browning during grinding of *Tenebrio molitor* larvae. *PloS One* 2017, 13(1), e0192015. <https://doi.org/10.1371/journal.pone.0192015>
30. Van Soest, P.J.; Robertson, J.B.; Lewis, B.A. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 1991, 74(10), 3583-3597. [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)
31. Church, F.C.; Swaisgood, H.E.; Porter, D.H.; Catignani, G. Spectrophotometric assay using o-phthaldehyde for determination of proteolysis in milk proteins. *Journal of Dairy Science* 1983, 66, 1219-1227. [http://dx.doi.org/10.3168/jds.S0022-0302\(83\)81926-2](http://dx.doi.org/10.3168/jds.S0022-0302(83)81926-2)
32. Adler-Nissen, J. Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzenesulfonic acid. *Journal of Agricultural and Food Chemistry* 1979, 27(6), 1256-1262. <https://doi.org/10.1021/jf60226a042>
33. Templeton, G. F. (2011). A two-step approach for transforming continuous variables to normal: Implications and recommendations for IS research. *Communications of the Association for Information Systems*, 28. <https://doi.org/10.17705/1cais.02804>
34. Finke, M.D. Estimate of chitin in raw whole insects. *Zoo Biology* 2007, 26, 105-115. <https://doi.org/10.1002/zoo.20123>
35. Marono, S.; Piccolo, G.; Loponte, R.; Di Meo, C.; Attia, Y.A.; Nizza, A.; Bovera, F. in vitro crude protein digestibility of *Tenebrio Molitor* and *Hermetia illucens* insect meals and its correlation with chemical composition traits. *Italian Journal of Animal Science* 2015, 14(3), 3889. <https://doi.org/10.4081/ijas.2015.3889>
36. Sumczynski, D.; Bubelová, Z.; Fišera, M. Determination of chemical, insoluble dietary fibre, neutral-detergent fibre and in vitro digestibility in rice types commercialized in Czech markets. *Journal of Food Composition and Analysis International Network of Food Data Systems* 2015, 40, 8-13. <https://doi.org/10.1016/j.jfca.2014.12.007>
37. Bosch, G.; Zhang, S.; Oonincx, D.G.A.B.; Hendriks, W.H. Protein quality of insects as potential ingredients for dog and cat foods. *Journal of Nutritional Science* 2014, 3, e29. <https://doi.org/10.1017/jns.2014.23>
38. Wessels, M.L.J.; Azzollini, D.; Fogliano, V. Frozen storage of lesser mealworm larvae (*Alphitobius diaperinus*) changes chemical properties and functionalities of the derived ingredients. *Food Chemistry* 2020, 320, 126649. <https://doi.org/10.1016/j.foodchem.2020.126649>
39. Tonnejck-Srpová, L.; Venturini, E.; Humblet-Hua, K.N.P.; Bruins, M.E. Impact of processing on enzymatic browning and texturization of yellow mealworms. *Journal of Insects as Food and Feed* 2019, 5(4), 267-277. <https://doi.org/10.3920/jiff2018.0025>
40. Mancini, S.; Mattioli, S.; Paolucci, S.; Fratini, F.; Dal Bosco, A.; Tuccinardi, T.; Paci, G. Effect of cooking techniques on the in vitro protein digestibility, fatty acid profile, and oxidative status of mealworms (*Tenebrio molitor*). *Frontiers in Veterinary Science* 2021, 8, 675572. <https://doi.org/10.3389/fvets.2021.675572>
41. Manditsera, F.A.; Luning, P.A.; Fogliano, V.; Lakemond, C.M.M. Effect of domestic cooking methods on protein digestibility and mineral bioaccessibility of wild harvested adult edible insects. *Food Research International* 2019, 121, 404-411. <https://doi.org/10.1016/j.foodres.2019.03.052>
42. Leni, G.; Soetemans, L.; Jacobs, J.; Depraetere, S.; Gianotten, N.; Bastiaens, L.; Caligiani, A.; Sforza, S. Protein hydrolysates from *Alphitobius diaperinus* and *Hermetia illucens* larvae treated with commercial proteases. *Journal of Insects as Food and Feed* 2020, 6(4), 393-404. <https://doi.org/10.3920/jiff2019.0037>
43. Hall, F.G.; Jones, O.G.; O'Haire, M.E.; Liceaga, A.M. Functional properties of tropical banded cricket (*Grylloides sigillatus*) protein hydrolysates. *Food Chemistry* 2017, 224, 414-422. <https://doi.org/10.1016/j.foodchem.2016.11.138>
44. Wang, W.; Nema, S.; Teagarden, D. Protein aggregation--pathways and influencing factors. *International Journal of Pharmaceutics* 2010, 390(2), 89-99. <https://doi.org/10.1016/j.ijpharm.2010.02.025>

45. Mishyna, M.; Martinez, J.-J.I.; Chen, J.; Davidovich-Pinhas, M.; Benjamin, O. Heat-induced aggregation and gelation of proteins from edible honey bee brood (*Apis mellifera*) as a function of temperature and pH. *Food Hydrocolloids* 2019, 91, 117-126. <https://doi.org/10.1016/j.foodhyd.2019.01.017>
46. Santiago, L.A.; Fadel, O.M.; Tavares, G.M. How does the thermal-aggregation behavior of black cricket protein isolate affect its foaming and gelling properties? *Food Hydrocolloids* 2021, 110, 106169. <https://doi.org/10.1016/j.foodhyd.2020.106169>
47. Vidanarachchi, J.; Kurukulasuriya, M.; Kim, S.-K. Chitin, chitosan, and their oligosaccharides in food industry. In *Chitin, Chitosan, Oligosaccharides and Their Derivatives*. CRC Press 2010, 543-560. <https://doi.org/10.1201/EBK1439816035-c38>
48. Klunder, H.C.; Wolkers-Rooijackers, J.; Korpela, J.M.; Nout, M.J.R. Microbiological aspects of processing and storage of edible insects. *Food Control* 2013, 26, 628-631. <https://doi.org/10.1016/j.foodcont.2012.02.013>
49. Gruzdev, N.; Pinto, R.; Sela, S. Effect of desiccation on tolerance of *Salmonella enterica* to multiple stresses. *Applied and Environmental Microbiology* 2011, 77, 1667-1773, <https://doi.org/10.1128/AEM.02156-10>
50. Larouche, J.; Deschamps, M.H.; Saucier, L.; Lebeuf, Y.; Doyen, A.; Vanderberg, G.W. Effects of killing methods on lipid oxidation, colour and microbial load of black soldier fly (*Hermetia illucens*) larvae. *Animals* 2019, 9(4), 182. <https://doi.org/10.3390/ani9040182>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.