

Article

Not peer-reviewed version

Temperatures Lower than Preferred Ones Maintain DNA Integrity and Sperm Quality of *Lepidophyma gaigeae* (SQUAMATA: XANTUSIIDAE)

[Daniel Uriostegui Escoto](#)*, Fausto Roberto Méndez de la Cruz, Mario García Lorenzana, Yolanda López Ramírez, [Edith Arenas Ríos](#)*

Posted Date: 11 April 2025

doi: 10.20944/preprints202504.0999.v1

Keywords: Sperm; Lizards; Behavior; Thermal effect; Reproduction; DNA fragmentation



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Temperatures Lower Than Preferred Ones Maintain DNA Integrity and Sperm Quality of *Lepidophyma gaigeae* (SQUAMATA: XANTUSIIDAE)

Daniel Uriostegui Escoto ^{1,*}, Fausto Roberto Méndez de la Cruz ², Mario García Lorenzana ³, Yolanda López Ramírez ¹ and Edith Arenas Ríos ^{3,*}

¹ Doctorado en Ciencias Biológicas y de la Salud. Universidad Autónoma Metropolitana, Iztapalapa, Ciudad de México, C.P. 09340, México

² Departamento de Zoología. Universidad Nacional Autónoma de México. Instituto de Biología, Universidad Nacional Autónoma de México, Coyoacán, A.P. 70515, C.P. 04510, México

³ Departamento de Biología de la Reproducción. Universidad Autónoma Metropolitana. Unidad Iztapalapa. Universidad Autónoma Metropolitana, Iztapalapa, Ciudad de México, C.P. 09340, México

* Correspondence: danuriostegui@gmail.com (D.U.E.); editharenas2000@yahoo.com.mx (E.A.R.)

Simple Summary: The imminent increase in global temperature has led to changes in ectothermic species, from their development and behavior to their reproduction. Although individuals such as lizards require a defined temperature range (preferred temperature) to carry out their physiological and metabolic activities, it is believed that in reproductive terms this range should be lower, as it could lead to problems in sperm maturation, as has been reported in other species. Given the above, in this work we seek to describe the effect of temperatures that may favor or affect sperm parameters in the lizard *Lepidophyma gaigeae*, finding that the thermal range that favors them is that which is below the preferred temperature. These results could emulate what would happen if the temperature increase in their environment continues, as well as explain how temperature affects reproductive environments, which could increase the risk of extinction in some lizard species.

Abstract: The global increase of temperature has directly impacted the environment and reproduction of ectothermic organisms, leading to defects in processes that intervene in sperm quality and maturation. The present study evaluates the effect of temperature on sperm quality of the lizard *Lepidophyma gaigeae*, in the surroundings of Landa de Matamoros, Querétaro. As a species under special protection, only 12 male individuals were captured during December 2020 and exposed at 1) preferred temperature (T_{pref}) at 24 °C; 2) Low Temperature (LT) at 21 °C; 3) individuals without treatment (Control) to compare the effect on viability, concentration and integrity of DNA, as well as if the presence of the cytoplasmic droplet (CD). Sperm viability decreased by approximately 50% in organisms exposed at T_{pref}, accompanied by a significant reduction in concentration. Likewise, the CD evaluation showed a retention percentage of 30% at the cauda level, while the LT and Control groups showed 5% in this terminal area of the epididymis. The percentage of sperm with intact DNA was also decreased in the T_{pref} group compared to the LT and Control groups. These findings highlight how thermal environments impact reptilian reproduction, reinforcing not only what has already been found in these species in recent years, but also the repercussions that it presents in the medium- or long-term exposure in these organisms at an ecological level, as well as the consequences with the potential damage and even infertility, in case of prolonged exposition to preferred temperatures.

Keywords: sperm; lizards; behavior; thermal effect; reproduction; DNA fragmentation

1. Introduction

Under the thermal dependence of physiological processes, ectotherms must regulate their body temperature close to their preferred temperature range (T_{pref}/T_{sel}) for the optimal performance of various physiological functions, such as metabolic rate, locomotor performance, digestive efficiency, growth rate, egg production, spermatogenesis and sperm maturation [1–5]. To this thermal dependence, reports on the susceptibility of germ cell damage to extreme temperature changes in lizards date back as far as the 1940 showed that high temperatures cause sterility [6] and subsequently, asserting that there is a greater sensitivity of germ cells concerning somatic cells under conditions of temperature stress [7].

A study in *Sceloporus siniferus* [8] revealed the possibility of a negative effect of temperature on epididymal sperm maturation due to CD retention since the collection of males between 2015 and 2016 shows a percentage of sperm with significantly different CD. It is observed that the sperm percentage with CD in the epididymis in 2015 was approximately 10%; in 2016, the percentage remained around 40%. Moreover, it was observed that the specimens collected in 2015 show a cloacal temperature of 28 °C, while those collected in 2016 presented a temperature of 34 °C

Studies in *Sceloporus aeneus* detail the positive and negative repercussions of subjecting specimens to preferred temperatures, highlighting locomotor performance and sperm quality. In the first case, animals that are kept in a preferred temperature environment favor motility interval, sprint speed, and foraging, however, some sperm parameters are decreased, such as concentration, viability, and motility, as well as the increase of spermatozoa with CD [3]. These aspects had been considered by Méndez-De La Cruz et al. (2014) [2], who proposed that low temperatures, also given by the inactivity schedules of the organisms, may be the ones in charge of favoring such reproductive parameters.

Recently, in *Sceloporus megalepidurus*, these aspects were shown, where temperatures below the preferred temperature favor viability and epididymal sperm maturation, contrary to individuals subjected to higher temperatures, where the cases of sperm DNA fragmentation and the presence of CD increased [5].

Even more, in *Lepidophyma pajapanense*, it was found that copulation occurs in the coldest months and the ovulation at the beginning of the warmest season; perhaps that asynchrony in reproduction is due to the favorable temperature for males' mismatch with the ones that are favorable for the fecundation. Moreover, a significant percentage of infertile eggs was found, perhaps due to low sperm quality or to the long sperm retention in females, or both [9]. Therefore, the interaction between sperm quality and temperature is an important issue to resolve.

The study of lizard thermal ecology is relevant to understanding their distribution, life history, ecology, and reproductive aspects. It also reveals how species can achieve physiologically active body temperatures, evaluating the risk of extinction in the face of climate change and its general consequences.

Also known as Gaiger's lizard [10], *Lepidophyma gaigeae* is an endemic species of Mexico, whose habitat includes pine and oak forests, grasslands and xerophytic scrub preferably in outcrops and between the crevices of limestone rocks [11], at altitude of 1500 to 2500 m, in states such as Hidalgo and Querétaro [12]. It is a small-sized viviparous lizard, between 47 and 54 mm (snout-vent length), one of the smallest species of the genus [13]. It presents a diurnal activity period [14], although some nocturnal individuals have been found [13]. In areas of thermal ecology, Arenas-Moreno et al. (2018) [13] state that *L. gaigeae* has a preferred temperature range of around 24 °C, a range similar to that reported in other species of its genera [15,16].

The present study aims to determine if the preferred temperature could be the cause of the alteration in some sperm parameters in the testis and/or epididymis, as well as the migration of CD and the affectation of DNA integrity in spermatozoa of the lizard *L. gaigeae*.

2. Materials and Methods

As a species under special protection according to Mexican Official Standard NOM-059-ECOL-1994 and so as not to alter availability of adult males in the area and consider their distribution in the future, only 12 male individuals were captured during December 2020 in the surroundings of Landa de Matamoros, Querétaro, in the area known as "El Lobo" (21°17'32.5 "N 99°07'13.1 "W), with a collection schedule from 9 am to approximately 3 pm, which coincides with their diurnal usual activity schedule.

For the selection, male lizards with snout-vent length (SVL) greater than 47 mm and a visible bulging in the ventral region due to the presence of hemipenes inside the cloaca, characteristics described by Goldberg & Camarillo (2003) [17].

The specimens were captured by direct capture methods, either manually or with the use of a fishing rod, depending on the location of the specimens and the availability of space for free collection of the specimens.

The captured animals were housed in 20 x 15cm blanket bags, with a maximum of 4 specimens each, to be later contained in plastic racks to facilitate their transfer and to be transported to the vicinity of the Laboratory of Morphophysiology and Biochemistry of Spermatozoa, Universidad Autónoma Metropolitana-Iztapalapa, to proceed with the acclimatization procedures. The transport vehicle maintained a temperature between 21 °C and 24 °C according to the preferred temperature range of the species (24 °C) [13], avoiding stretches that generate sudden movements during their arrival at the facilities.

2.1. Inclusion of Specimens

The lizards were separated into 3 groups (4 individuals per group): control group, analyzed without treatment; preferred temperature (Tpref) (24 °C), and low temperature (LT) (21 °C), following the considered observations of [2,3,5,17]), where a range between the lower limits of the Tpref and without reaching the upper limit of the critical minimum temperature of the species is considered. Treatments were maintained during the activity time of the lizards (9 am to approximately 3 pm), placed in incubators "Hova Bator 1602N" of thermal radiation, with the required temperature, including the temperature during scotophase [13]. They were checked daily and maintained with water and live food, ad libitum during the 14 days of treatment.

2.2. Collection of Biological Material

The control specimens were sacrificed by intraperitoneal injection with 0.10 mL of sodium pentobarbital (Sedalpharma, from Pet's Pharma de México, S.A. de C. V) 36 hours after the arrival of the specimens at the University facilities for the Control group and after 14 days of treatment for the remaining groups, with adherence to the guide for the care and use of laboratory animals (Institute of Laboratory Animal Resources, National Research Council, 1996), as well as the Mexican Official Standard NOM-033-SAG/ZOO-2014, Methods for killing domestic and wild animals. Both testes and epididymides were extracted, measured (diameters and lengths) with a digital caliper (precision=±0.5mm) and weighed with a METTLER TOLEDO AB204-S balance (precision =±0.1mg), of which, those on the left side were used for sperm parameters, although those on the right side were fixed in 4% paraformaldehyde (PAF) (Thermo Scientific) and subsequently sagittally sectioned for further study.

Each epididymis was sectioned into three regions with similar proportions: caput, corpus, and cauda.

To obtain sperm parameters, the organs were placed in a four-well Petri dish with 250 µL of physiological Ringer's solution (NaCl, KCl, KH₂PO₄, CaCl₂ 2H₂O from Baker Laboratories) to obtain as many spermatozoa as possible, then the biological material was transferred to a 1.5mL Eppendorf tube [18]. Subsequently, the material was centrifuged at 250 G for 5 minutes, the supernatant was

removed, and the cell button was disaggregated with 250 μ L of Ringer's solution twice. Thus, the resultant will be "washed spermatozoa".

The spermatozoa obtained were evaluated according to the World Health Organization specifications for the evaluation of human semen (WHO, 2010) with necessary adjustments for lizard cells obtained from the different organs. The variables and methods to obtaining them were as follows:

Total number of spermatozoa per organ (n spz $\times 10^6$ / organ): obtained by counting the spermatozoa with the Neubauer chamber, making a dilution depending on the density of spermatozoa identified at a field of 40x, filling both chambers with 10 to 15 μ L of the dilution.

To obtain the sperm concentration, the formula was used: $N/n \times (1/20) \times \text{dilution factor}$, where N is the total number of spermatozoa counted in both chambers and n is the number of squares counted.

To calculate the viability percentage and CD, a 5 μ L aliquot of the sperm suspension was placed on a slide and mixed with an aliquot of the same volume of Spermatovit supravital stain and then smeared and observed under brightfield microscopy for counting.

Percentage of DNA integrity: A 10 μ L aliquot of the sperm suspension was placed on a slide to make a smear and allowed to dry at room temperature, then included in Carnoy's solution (ethanol from Meyer Laboratories, chloroform and acetic acid from Baker Laboratories) for approximately 24 hrs in a cool, dark place. Afterwards, the fixative was removed and the slides were included in acridine orange solution for 5 minutes, then, the excess dye was removed and left to dry isolated from light, finally, the slides were analyzed with the aid of fluorescence microscopy at 40x.

The handling of chemical-biological waste generated in the experiment was instructed by the Mexican Official Standard NOM-033-STPS-2015, safety conditions for working in confined spaces.

2.3. Statistical Analysis

It was used to identify the existence of significant differences among groups, performing parametric analysis using an analysis of variance (ANOVA) for the evaluation of the three treatments, as well as to evaluate variability and random errors so that possible environmental and temporal effects are distributed equally among the treatments, followed by a Tukey post-hoc test and a Kruskal-Wallis non-parametric analysis for significantly different means. The data were previously analyzed to meet the assumptions of the parametric test used (normality and homoscedasticity), considering the data from the different organisms analyzed as independent.

3. Results

According to the criteria [13], as shown in Table 1, adult male individuals were collected. Presenting a SVL greater than 50 mm and a similar weight to the reported adult size.

After the period determined for the corresponding study, the organs of interest were obtained. It should be noted that all specimens presented spermatozoa in the testis and epididymis, however, those organs of the individuals that were subjected to some temperature treatment (24 °C and 21 °C) presented lower testicular weight and length.

Table 1. Length and weight records of 12 male *L. gaigeae* collected. The average length is shown to verify the status of an adult male.

Organisms	SVL (mm)	Body weight (g)
<i>L. gaigeae</i>	51.60 \pm 2.17	4.37 \pm 0.33

3.1. Gonadal Size and Weight

The records obtained show a lower testicular weight in the groups that were subjected to some temperature treatment (24 °C and 21 °C) (Figure 2), either preferred or low (0.010 \pm 0.003 and 0.009 \pm

0.001, respectively), as well as their length (0.43 ± 0.07 and 0.45 ± 0.04 , respectively) (Figure 3), compared to the control group. Despite this, the data obtained did not involve significant differences in the epididymis.

Although testicular morphology was similar between the groups, it was found that one of the individuals treated at low temperature showed an apparent decrease in the concentration of spermatozoa in the caput area (Figure 4b) in conjunction with two subjects who appeared to have a greater amount of blood vessels between the two organs of interest (Figure 4d).




Figure 1. Position and regionalization of male gonads of *L. gaigeae* in reproductive stage. In **A**: Location of the gonadal position. Arrows show the testis (T) and epididymis (Ep) of the specimen. In **B**: Testis and epididymis with regionalization to identify the area of the caput, corpus and cauda. Bar: 0.5 cm. In **C**: Photomicrographs in sequence showing the different histomorphology of the three epididymal regions in longitudinal section, obtained with H&E at 5x. Bar at 50 μ m.




Figure 2. Weight (g) of testis and epididymis of *L. gaigeae* specimens with and without treatment at different temperatures. The bars indicate the distribution of the mean value of the data obtained from each treatment and region. Different letters indicate significant differences between the comparison of groups, according to the analysis of variance (ANOVA) and the Tukey and Kruskal-Wallis post hoc test given with a p-value < 0.05.




Figure 3. Testicular and epididymal lengths of *L. gaigeae* specimens with and without treatment at different temperatures. The bars indicate the distribution of the mean value of the data obtained from each treatment and region. Different letters indicate significant differences between the comparison of groups, according to the analysis of variance (ANOVA) and the Tukey and Kruskal-Wallis post hoc test given with a p-value < 0.05.




Figure 4. Physical variations of testis and epididymis found in some individuals of *L. gaigeae* subjected to different temperatures. Images a and c correspond to organs collected in the preferred temperature treatment (Tpref); Images b and d correspond to organs collected in the low temperature treatment (LT). G: Fat body. T: Testis. Bars at 0.5 mm.

3.2. Sperm Parameters

3.2.1. Sperm Concentration

There are significant differences between groups: gonadal size and weight, sperm concentration ratified the decrease between the regions analyzed and between groups; both in testis and epididymis in the specimens with temperature treatment (24 °C and 21 °C).

The lower sperm concentration presented in the Tpref (24 °C) and LT (21 °C) treatments is manifested from the testicular region (25.3 ± 12.74 and 30.6 ± 6.60 , respectively) and continues in the same way in the epididymis (Table 3). It should be noted that, as described in Figure 4b, a specimen treated at LT (21 °C) presented a lower sperm concentration at the caput level (10.6 ± 0.81), which could be corroborated in the quantitative analysis and could provide a significant difference in the comparison between groups.

On the other hand, the groups treated at Tpref (24 °C) and LT (21 °C) showed similarities in corpus (19 ± 4 and 15.6 ± 4.27 respectively) and cauda (21.6 ± 3.05 and 23.6 ± 6.13 respectively) epididymis, however, having significant differences in comparison with the control group (38 ± 7.93 in corpus and 39 ± 9.53 in cauda).

It should be noted that the present study of sperm concentration is the first to show a quantitative analysis of the species in its reproductive stage, which could be very close to what would be found in natural conditions, as well as, to the impact that temperature would have on its activity period.

Table 3. Sperm concentration of *L. gaigae* with and without treatment at different temperatures. Data represent n106 spz/mL. Asterisks indicate significant differences between group comparison, according to analysis of variance (ANOVA) and Tukey's and Kruskal-Wallis post hoc test given at p-value < 0.05.

	Testis	Caput	Corpus	Cauda
Control	80 ± 8.88	18 ± 2	38 ± 7.93	39 ± 9.53
Tpref	* 25.3 ± 12.74	14.3 ± 3.51	* 19 ± 4	* 21.6 ± 3.05
LT	* 30.6 ± 6.60	* 10.6 ± 0.81	* 15.6 ± 4.27	* 23.6 ± 6.13

3.2.2. Sperm Viability

The sperm viability analysis (Figure 5), obtained by the eosin-nigrosin staining method (Figure 6), shows the impact of temperature treatments on these parameters. Contrary to the analysis of sperm concentration, significant differences between groups were found in one of them, especially at the epididymis level. The results stand out in different points, highlighting at least three important aspects to be considered.

Firstly, it should be noted that the percentage of sperm viability in the control group and LT (21 °C) is above 60% at the cauda level. In this case, the percentage of viable spermatozoa in the control group ranges from 50.66% at the testicular level to 65.6% in the cauda.

Secondly, it is important to mention that the group treated at LT (21 °C) showed a viability percentage ranging from 73% to 71%, even though, at the caput level, the count is 67.6%. No differences were found between the Control and Tpref (24 °C) groups. It is worth mentioning that it was in the Tpref group (24 °C) where the individuals showed a lower sperm concentration, mainly in the caput of one of them. A maximum of 50.3% in caput, but lower in the remaining regions up to 47.3% in cauda. Even though the impact may not be drastic compared to that found in other species [3,5]. The comparison with the concentration and other parameters corroborates the effect at the reproductive cellular level.




Figure 5. Percentage of sperm viability of *L. gaigeae* with and without treatment at different temperatures. The bars indicate the distribution of the mean value of the data obtained from each treatment and region. It is evident the maintenance of the viability percentage in the groups with low temperature treatment (21°C) and the increase of sperm viability in the control group compared to the remaining group, whose values decrease notably and maintain a viability of 50%. Different letters indicate significant differences between the group comparisons, according to the analysis of variance (ANOVA) and the Tukey and Kruskal-Wallis post hoc test given with a p-value < 0.05.



Figure 6. Photomicrographs of *L. gaigeae* spermatozoa with and without treatment at different temperatures, stained with eosin-nigrosin for viability assessment with the plasma membrane dye penetration method, obtained by brightfield microscopy showing A: viable spermatozoa. B: non-viable spermatozoa. X400. Bars corresponding to 10 μm.

3.2.3. Cytoplasmic Droplet Percentage

The evaluation of CD in the spermatozoa obtained from testis and epididymis (Figure 7) was carried out considering only the cells that presented the shape of a fully formed spermatozoa and without excess of cytoplasmic residues in the head (Figure 8). Only the spermatozoa counted were found with the CD in the neck of the cell. Thus, the control group, like the LT (21 °C) group, showed a significantly lower percentage of spermatozoa with CD compared to the group subjected to Tpref (24 °C). These values ranged from 54% and 57% in the testis, respectively, to around 5% of sperm with CD in both groups at the cauda level. Although the Tpref (24 °C) group also presented a lower percentage of CD retention, this remained above 30% in the last epididymal region, despite not showing significant differences at the testicular level compared with the two groups mentioned.




Figure 7. Percentage of spermatozoa with cytoplasmic droplet (CD) of *L. gaigeae* with and without treatment at different temperatures. The bars indicate the distribution of the mean value of the data obtained from each treatment and region. Evidence of CD removal at the testicular level is shown in the control group and with low temperature treatment (21°C). All groups showed a decrease in CD as they moved to the caput zone, with a difference in the percentage of spermatozoa found with this characteristic. Different letters indicate significant differences between the comparison of groups, according to the analysis of variance (ANOVA) and the Tukey and Kruskal-Wallis post hoc test given with a p-value < 0.05.



Figure 8. Photomicrographs of *L. gaigeae* spermatozoa stained with eosin-nigrosin (A and B) and unstained (C and D) for evaluation of the presence of cytoplasmic droplet (CD) by brightfield microscopy. A and C: Spermatozoa with CD. B and D: Spermatozoa without CD. X400. Bars indicate the dimensions of the photomicrographs, corresponding to 10 μm.

3.2.4. DNA Integrity

The projected staining after treatment with acridine orange, where spermatozoa with intact DNA are cataloged if they manifest a green hue at the level of the cell head, while those presenting an orange hue were counted as cells with fragmented DNA (Figure 10), allowed demonstrating the differences in terms of variation between groups and regions (Figure 9). In the first case, the Control group remained at a relatively constant interval, with nearly 80% found in practically all regions, with the caudal region of the epididymis being the area where the lowest percentage of spermatozoa with intact DNA was found. Despite this, the average recorded was 78.3%.

On the contrary, the group treated at Tpref (24 °C) exhibits a lower percentage of spermatozoa with intact DNA from the testicular level (63.3%), whereas in caput presents its lowest percentage value (54.6%) which subsequently increases in the corpus and cauda (58.6% and 64.3%, respectively).

This finding contrasts directly with the LT group (21 °C), whose case is similar to that found in the Control group, being its percentage in the caudal region even higher than in the last mentioned (82%).




Figure 9. Percentage of DNA integrity of *L. gaigeae* spermatozoa with and without treatment at different temperatures. The bars indicate the distribution of the mean value of the data obtained from each treatment and region. The percentage of spermatozoa with intact DNA is maintained, except for the group with treatment at the preferred temperature (Tpref) in all the areas evaluated, which seems to be increasing, shortening the difference between the remaining groups.




Figure 10. Photomicrographs of *L. gaigeae* spermatozoa stained with acridine orange for DNA integrity evaluation by fluorescence microscopy. **A:** Spermatozoa with intact DNA. **B:** Spermatozoa with fragmented DNA. **C:** Indicates the bipolarity of the staining found in the cell samples. X400. Bars at 10 μm.

4. Discussion

The formation of spermatozoa takes place in the seminiferous epithelium of the testis, through a process known as spermatogenesis, a process by which, from diploid round cells (spermatogonia) located adjacent to the basement membrane of the seminiferous tubules, they differentiate until forming spermatozoa, advancing towards the tubule lumen [19,20]. However, even when the morphological differentiation of the spermatozoa culminates in the testis when the spermatozoa are released from it, they have not yet acquired the capacity to fertilize the oocyte, for this reason they need to pass through the epididymis, where they acquire this capacity, a process known as epididymal sperm maturation [21,22], where the following stand out: loss of cytoplasmic droplet (CD) and chromatin hypercompaction about the maintenance of DNA integrity [8,23,24].

Climate change may cause the extinction of species, as well as a shift in their distribution in the coming decades [2,25]. Since 1975, 12% of local lizard populations have gone extinct, as well as 4% around the world. By 2080, extinction is projected to approach 39% globally and species extinction by 20%. If the maximum temperature derived from global warming continues unchanged, it is estimated that, in Mexico, in the year 2050, 56% of the viviparous species will become extinct and 66% by 2080. The causes of extinction were attributed to low thermal quality that impairs the lizards' ability to perform basic biological activities and increases the restriction hours [25]. Nevertheless, the temperatures may decrease the thermal quality of the sperm, and the males become infertile. The results of this study show that the sperm is labile (in quality and morphology) when lizards are overexposed to the preferred temperatures and is even worse at higher temperatures.

For example, as occurred with the difference in testicular size, previous studies with *Sceloporus megalepidurus* [5] showed that individuals who remained in treatment at the preferred temperature for 14 days presented smaller testis and epididymal, being one of the first characteristics described in reptiles affected by temperature changes. It is known that the volume of reproductive organs can be an indirect indicator of sperm concentration [26]. In the present study, we found that this difference in volume could be attributed to the decrease of spermatozoa in the group exposed to preferred temperature [3,21,27]. In the *Lepidophyma pajanapense*, a high percentage of infertile eggs was found [26]. That low percentage could be attributed to the long sperm retention or the low quality of the sperm cells, even when copula occurs during the coldest months of the year, perhaps due to the inability to manage the warm temperatures during the ovulation months.

It is known that chronic hyperthermia stress can cause alterations in spermatogenesis [28], as has also been shown in mammalian species such as rats [29], and sheep [30]. In addition, these thermal conditions tend to influence reproductive conditions from the hypothalamus-pituitary-gonad axis through the increase of glucocorticoids and the repression of gonadotropin-releasing hormone (GnRH), fundamental for the promotion of spermatogenesis and testosterone production [31], also causing a reduction in the number of Leydig cells as well as spermatogonia in the seminiferous tubule [32].

The integrity of the plasmatic membrane is one of the most important characteristics to consider for the evaluation of viable spermatozoa [33], therefore the use of the eosin-nigrosin dye allowed demonstrating the decrease of this characteristic in individuals maintained at preferred temperatures, bringing it below 50%, a value that resembles that shown in other reptile species [3,5]. On the other hand, knowing that sperm viability in each of the epididymal regions may be different according to the functionality of its cellular structure and the achievement of non-functional sperm absorption [34], the coincidence in the percentage, even in the four regions evaluated allows us to question whether the problem may even come from the concentration of androgens. Given that the epididymis is a hormone-dependent organ, mainly dihydrotestosterone [35] different analyses have coincided in that the lack of these contributes in the decrease of epididymal weight, as well as in the regulation of secretions that reach the luminal region, such as proteins and transcription factors important for sperm viability and flagellar motility mainly [13,36–42]. In addition, reduction of epididymal epithelium volume may also occur as a consequence of the effect of temperature, with increased apoptosis of constituent cells, such as principal cells [43].

Although once again the effect that CD causes in lizard spermatozoa has not been analyzed in depth, it is clear with the present study, as well as those reported [3,5,8], that thermal modifications tend to influence its presence along the different regions of the epididymis. This coincides with numerous reports in mammals that document the increase in CD under conditions of testicular hyperthermia, being indicative of immature sperm, given by changes in ambient osmolarity or decreased antioxidant system and increased reactive oxygen species (ROS) and plasma membrane lipid peroxidation [22,44–46].

However, some studies have raised the possibility of a protective function in the face of unfavorable conditions for spermatozoa. [47] showing that CD spermatozoa may have a higher tolerance to heat stress. In addition to this, it is known that CD has the presence of aquaporins, which

can collaborate with osmotic adaptation, migration or sperm storage [48], besides being enzymatically active [49] and being an important source of energy for the different maturational aspects of the cells, which is why they can remain in the cell throughout the epididymis [50].

The relationship that CD has had around the exposure of organisms to high temperatures is given in many cases by its relationship with lipid droplets (LD), which, although in some cases, supplements the function of the latter to provide energy [50], can also cause lipoperoxidation of the plasma membrane by increasing ROS given mainly by the decompensation of antioxidant enzymes. In hyperthermia treatments, GL has been found along Leydig cells, important for testosterone production and sperm maintenance [46], but mostly involved in the breakdown of disulfide bridges involved in DNA compaction [51]. The relationship of increased ROS with sperm viability and morphology also derives from the cell death by apoptosis, where studies show the activation of caspase 3 in seminal samples with high DNA fragmentation index [52].

Lizards around the world are facing problems to survive; one of the most important is global warming. This study determines that warming temperatures affect not only the hours of activity but also the fertility of males. It was found that rising temperatures arrested female fertility during the El Niño Southern Oscillation event in *Sceloporus mucronatus* [53] (Rodríguez Romero and Méndez de la Cruz, 2004). This study determines that sperm quality could be severely affected by environmental temperatures and may be the cause of that infertility. Therefore, the declining of populations may also be due to the infertility of males exposed to warmer temperatures. Furthermore, it may promote dissociation in reproductive cycles, as in *L. pajapanense*, where males copulate during the colder months while females ovulate in the warmer ones [9]. This study shows that the conditions for fully fertile males are an issue that demands more studies as there is a connection between physiology and environmental temperatures.

5. Conclusions

The present study has shown the effect of temperature on *Lepidophyma gaigeae* spermatozoa, given by the controlled thermal maintenance that emulates different environmental conditions in the organism, showing that the temperature at which the individual tends to perform better in terms of ethological, physiological and digestive parameters tends to be detrimental in reproductive terms, causing a decrease in gonadal size, concentration, viability and probably spermatogenic maturation. Furthermore, we demonstrate that temperatures below the preferred temperature maintain not only sperm quality but also DNA integrity.

Author Contributions: Conceptualization: Daniel Uriostegui Escoto, Edith Arenas Ríos, Mario García Lorenzana, Fausto R. Méndez de la Cruz; Field work: Daniel Uriostegui Escoto, Fausto R. Méndez de la Cruz; Laboratory work: Daniel Uriostegui Escoto, Yolanda López Ramírez; Resources: Mario García Lorenzana, Fausto R. Méndez de la Cruz, Edith Arenas Ríos; Data analysis: Daniel Uriostegui Escoto; Writing – Review & Editing: Daniel Uriostegui Escoto, Edith Arenas Ríos, Mario García Lorenzana, Fausto R. Méndez de la Cruz.

Funding: Funding for the work was provided by the Consejo Nacional de Humanidades Ciencia y Tecnología (PhD scholarship number: 932666) and Universidad Nacional Autónoma de México (PAPIIT IN-212724).

Institutional Review Board Statement:



Casa abierta al tiempo

UNIVERSIDAD AUTÓNOMA METROPOLITANA-Iztapalapa
COMISIÓN ACADÉMICA DE ÉTICA DE LA DIVISIÓN
DE CIENCIAS BIOLÓGICAS Y DE LA SALUD

Ciudad de México a 29 de octubre de 2021
Dictamen: CECBS21-04

A quien corresponda:


La Comisión Académica de Ética de la División de Ciencias Biológicas y de la Salud, conforme a sus competencias revisó el protocolo de Investigación de la alumna **Daniel Uriostegui Escoto** titulado: **"EFECTO DE LA TEMPERATURA EN LA ORGANIZACIÓN HISTOLÓGICA DE TESTÍCULO Y EPIDÍDIMO Y EN LA PRESENCIA DE INDICADORES DE MADURACIÓN ESPERMÁTICA DE LA LAGARTIJA *Lepidophyma gaigeae*"**, dirigido por la doctora: **Edith Arenas Rios**.

Encontrando que dicho proyecto, **SI CUMPLEN CON LOS LINEAMIENTOS PARA LA CONDUCCIÓN ÉTICA DE LA INVESTIGACIÓN, LA DOCENCIA Y LA DIFUSIÓN EN LA DIVISIÓN DE CIENCIAS BIOLÓGICAS Y DE LA SALUD DE ESTA INSTITUCIÓN.**

Cabe recordar que dichos lineamientos derivan de la legislación mexicana que atañe a la parte bioética de nuestro quehacer.

Se extiende la presente para los fines que convengan al interesado

ATENTAMENTE
Casa Abierta al Tiempo


DRA. MARÍA DEL ROSARIO TARRAGÓ CASTELLANOS
Presidenta

c.c.p. archivo

UNIDAD IZTAPALAPA Av. San Rafael Atlixco 186 Col. Vicentina. C.P. 09340, Ciudad de México

Data Availability Statement: We hereby declare the open accessibility of data from our research for transparency purposes. Due to time and technical status, our data is not yet available on Figshare to share DOI, however, we are committed to uploading it as soon as possible. For now, we can share them, if you wish, via email.

Acknowledgments: This work could not be possible without the joint work of the Sperm Morphophysiology and Biochemistry Laboratory, the Neuroimmunoendocrine Tissue Biology and Engineering Laboratory, and the Herpetology Laboratory of the Metropolitan Autonomous University and the National Autonomous University of Mexico. We want to thank all the colleagues who have supported us in this project, especially Mirna Crizel Vera Chávez, Misael Garrido Casas, Isela Quintero Pérez and Diego Arenas Moreno for their unconditional support in field work.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

CD: Cytoplasmic droplet

LT: Low temperature

T_{pref}: Preferred temperature

ROS: Reactive oxygen species

SVL: Snout-vent length

LD: Lipid droplet

References

1. Meek R, "Reptiles, termorregulation and the enviroment. TESTUDO. Vol. 4:(2), pp. 56-78. 1994.
2. Méndez-De La Cruz F, Manríquez N, Arenas-Ríos E, Ibargüengoytia N, "Male Reproductive Cycles in Lizards", Reproductive Biology and Phylogeny, vol. 10, pp: 303-339, 2014.
3. Quintero R, Méndez -de la Cruz R, Miles D, Vera M, López Ramírez Y, Arenas-Moreno D, Arenas-Ríos E. "Trade-off between thermal preference and sperm maturation in a montane lizard", Journal of Thermal Biology. Vol. 113: 103526. 2023.
4. Rangel-Patiño C, García-Morales C, Mastachi-Loza C, Eifler D, Ruiz-Gomez M., "When things get hot: Thermoregulation behavior in the lizard *Sceloporus aeneus* at different thermal conditions". Journal of Thermal Biology, 89, 102572. 2020.
5. Uriostegui-Escoto D., "Efecto de la temperatura en la calidad espermática de la lagartija *Sceloporus megalepidurus*". Tesis de Maestría. Universidad Autónoma Metropolitana. 2021.
6. Cowles R, Burleson G, "The Sterilizing Effect of High Temperature on The Male Germ-Plasm of The Yucca Night Lizard, *Xantusia vigilis*", The American Naturalist, vol. 79, pp. 417-435, 1945.
7. Licht, P., "The relation between preferred body temperatures and testicular heat sensitivity in lizards", Copeia, pp. 428-436, 1965.
8. Retana F., "Determinación de indicadores de maduración epididimaria en espermatozoides de la lagartija *Sceloporus siniferus* (Sauria: Phrynosomatidae)", Molecular Reproduction and Development, vol. 1, pp. 40-7, 2016.
9. Méndez-De La Cruz F, Villagrán M, Hernández-Gallegos O, Manríquez-Morán N, Rodríguez-Romero F. "Reproductive cycle of the tropical night lizard, *Lepidophyma pajapanensis* (Sauria: Xantusiidae)". Journal of Herpetology, vol. 33: pp. 340-343, 1999.
10. Liner E, Andreu C. Nombres estándar en español en inglés y nombres científicos de los anfibios y reptiles de México. Society for the Study of Amphibians & Reptiles; 2nd edition, 2008.
11. Bezy R, "Systematics of xantusiid lizards of the genus *Lepidophyma* in northeastern Mexico". Contributions in Science. Vol. 349, pp.1-16, 1984.
12. Bezy R, Camarillo R, "Systematics of xantusiid lizards of the genus *Lepidohpyma*". Contributions in science. Vol. 493, pp. 1-41, 2002.
13. Arenas-Moreno D, Santos-Bibiano R, Muñoz-Nolasco F, Charruau P, Méndez-de la Cruz F, "Thermal ecology and activity patterns of six species of tropical night lizards (Squamata: Xantusiidae: *Lepidophyma*) from Mexico". Journal of Thermal Biology, S0306456517304837. 2018.
14. Leavitt D, Hibbits T., "Lepidophyma gaigeae Mosauer Gaige's Tropical Night Lizard", Catalogue of American Amphibians and Reptiles, pp. 896, 2012.
15. Grigg J, Buckley L. "Conservatism of lizard thermal tolerances and body temperatures across evolutionary history and geography". Biol. Lett. Vol.9: pp. 20121056-20121056, 2013.
16. Clusella-Trullas S, Chown S, "Lizard thermal trait variation at multiple scales: a review". Journal of Comparative Physiology B, vol. 184: pp. 5-21, 2014.
17. Goldberg S, Camarillo-Rangel J. "Lepidophyma gaigeae (Gaige's tropical night lizard) reproduction". Herpetological Review. Vol 34: pp. 246, 2003.
18. Cervantes, M, Arenas-Ríos E, León-Galván M, López-Wilchis R, Ambriz D, Rosado A. "Spermatozoa epididymal maturation in the Mexican big-eared bat (*Corynorhinus mexicanus*)", Systems Biology in Reproductive Medicine, vol. 54:4-5, pp. 196-204, 2008.
19. Fink S., Cookson B. "Apoptosis, Pyroptosis, and Necrosis: Mechanistic description of dead and dying eukaryotic cells. Infection and Immunity, vol. 73, pp. 1907-1916, 2005.
20. Dacheux, J.L.; Castella, S.; Gatti, L.J. & Dacheux, F. "Epididymal cell secretory activities and the role of the proteins in boar sperm epididymis". Theriogenology, Vol. 63, No. 2, pp. 319-341, 2005.

21. Arrotéia K., Vianna P., Ferreira M., Lopes M. y Violin L., "The Epididymis: Embryology, Structure, Function and Its Role in Fertilization and Infertility", Embryology - Updates and Highlights on Classic Topics, Prof. Luis Violin Pereira, Eds. InTech. pp. 41-66, 2012.
22. Dacheux J., Dacheux F., "New insights Into Epididymal Functions in Relation to Sperm Maturation", Reproduction, vol. 147, pp. 27-42, 2014.
23. Ashok A., Tamer M., "Sperm chromatin assessment", Textbook of assisted reproductive techniques, pp. 65–87, 2004.
24. Cooper T., Yeung C., "Sperm maturation in the human epididymis", The Sperm Cell. Chapter 4, pp. 72-107, 2010.
25. Sinervo B, Mendez-de-la-Cruz F, Miles D, Heulin, B, Bastiaans E, Villagran-Santa Cruz M, Lara-Resendiz R, Martinez-Mendez N, Calderon-Espinosa M, Meza-Lazaro R. "Erosion of Lizard Diversity by Climate Change and Altered Thermal Niches". Science, vol. 328:(5980), pp. 894–899. 2010.
26. Adriansjah R, Kusumajaya C, Wijayanti Z, Successful testicular sperm extraction in infertile male with non-obstructive azoospermia presented with bilateral atrophic testis: A case report", Andrology and Fertility, vol. 33, 101300, 2020.
27. Fink S., Cookson B. "Apoptosis, Pyroptosis, and Necrosis: Mechanistic description of dead and dying eukaryotic cells. Infection and Immunity, vol. 73, pp. 1907-1916, 2005.
28. Cooper, J., "Wildlife Forensic Investigation, Principles and Practice". Capítulo 8. Working with Live Animals. Ed. Taylor & Francis Group. EEUU. pp 229, 2013.
29. Setchel B, Plöen, Ritzen E, "Reduction of long-term effects of local heating of the testis by treatment of rats with a GnRH agonist and an anti-androgen", Reproduction, vol. 122, pp. 255-263. 2001.
30. Mieusset P, Quintana L, Sanchez-Partida S, Sowerbutts J, Zupp L, Setchell B, "The Effects of Moderate Heating of the Testes and Epididymides of Rams by Scrotal Insulation on Body Temperature, Respiratory Rate, Spermatozoa Output and Motility, and on Fertility and Embryonic Survival in Ewes Inseminated with Frozen Semen", vol. 637 (1), pp 445-458, 1991.
31. Gao HB, Tong MH, Hu YQ, Guo QS, Ge R, Hardy M. "Glucocorticoid induces apoptosis in rat leydig cells". Endocrinology, vol. 143(1): pp. 130-138, 2002.
32. Whirledge S, Cidlowski J. "Glucocorticoids, Stress, and Fertility". Minerva Endocrinology. Vol. 35(2): pp. 109-125, 2010.
33. Simón-Salvador P, Álvarez T, Fuentes-Mascorro G. "Choque hipoosmótico en espermatozoides de víbora de cascabel", Revista Iberoamericana de Ciencias, vol. 3(3), pp. 52-57. 2016.
34. Kempinas W, Klinefelter G. "Interpreting histopathology in the epididymis". Spermatogenesis. Vol. 4:2, e979114. 2014.
35. Ramos-Ibeas, P., Pericuesta, E., Fernandez-Gonzalez, R., Ramirez, M. A., & Gutierrez-Adan, A. "Most regions of mouse epididymis are able to phagocytose immature germ cells". Reproduction, 146(5), 481–489, 2013.
36. Hinton B, Lan Z, Rudolph D, Labus J, Lye R, "Testicular Regulation of Epididymal Gene Expression", Journal of Reproduction and Fertility. Supplements, Vol. 53:47–57, 1998.
37. Ezer N, Robaire B, "Androgenic Regulation of The Structure and Functions of the Epididymis", In: Robaire B, Hinton BT, editors. The Epididymis: From Molecules to Clinical Practice. New York: Kluwer Academic/Plenum Publishers, pp. 297–316, 2002.
38. Avram C, Yeung CH, Nieschlag E, Cooper T, "Regulation of the Initial Segment of the Murine Epididymis by Dihydrotestosterone and Testicular Exocrine Secretions Studied by Expression of Specific Proteins and Gene Expression. Cell Tissue Research. Vol. 317:13–22, 2004.
39. Maclean J, Chen M, Wayne C, Bruce S, Rao M, Meistrich M, Macleod C, Wilkinson M, "Rhox: a new homeobox gene cluster". Cell, vol. 120:369–382, 2005.
40. Sipila P, Pujiato D, Shariatmadari R, Nikkila J, Lehtoranta M, Huhtaniemi I, Poutanen M, "Differential Endocrine Regulation of Genes Enriched In Initial Segment and Distal Caput of the Mouse Epididymis as Revealed by Genome-wide Expression Profiling", Biology of Reproduction, vol. 75:240–251, 2006.

41. Turner T, Bang H, Attipoe S, Johnston D, Tomsig J, "Sonic Hedgehog Pathway Inhibition Alters Epididymal Function as Assessed by the Development of Sperm Motility, *Journal of Andrology*, vol. 27:225–232, 2006.
42. Olesen I, Joensen U, Petersen J, Almstrup K, Rajpert-De Meyts E, Carlsen E, McLachlan R, Juul A, Jorgensen N, "Decrease in Semen Quality and Leydig Cell Function in Infertile Men: A Longitudinal Study", *Human Reproduction*, vol. 33: 1963-1974, 2018.
43. Ozturk U, Kefeli M, Asci R, Akpolat I, BuyukalPELLI R, Sarikaya S., "The Effects of Experimental Left Varicocele on the Epididymis". *Systems Biology in Reproductive Medicine*, 54 (4-5), 177-184, 2008.
44. Brouwers J, Silva P, Gadella B, "New Assays for Detection and Localization of Endogenous Lipid Peroxidation Products in Living Boar Sperm After BTS Dilution or After Freeze–thawing", *Theriogenology*, vol. 63: 458–469, 2005.
45. Cooper T., "The epididymis, cytoplasmic droplets and male fertility" *Asian Journal of Andrology*, vol. 13, pp. 130-138, 2011.
46. Durairajanayagam D, Agarwal A., Ong C., "Causes, Effects and Molecular Mechanisms of Testicular Heat Stress", *Reproductive Healthcare Ltd.* pp. 1472-6483, 2014.
47. Schulze M, Waberski D, "Compensability of Enhanced Cytoplasmic Droplet Rates in Boar Semen: Insights of a Retrospective Field Study", *Animals (Basel)*, vol. 12 (20): 2892, 2022.
48. Chen Q, Peng H, Lei L, Zhang Y, Kuang H, Cao Y, Shi Q, Ma T, Duan E, "Aquaporin3 is a Sperm Water Channel Essential for Postcopulatory Sperm Osmoadaptation and Migration", *Cell Research*, vol. 21 (6): 922-933, 2011.
49. Dott H, Dingle J, "Distribution of Lysosomal Enzymes in the Spermatozoa and Cytoplasmic Droplets of Bull and Ram", *Experimental Cell Research*, vol. 52: 523-540, 1968.
50. Zhang L, Yang P, Xunguang B, Zhang Q, "Modification of Sperm Morphology During Long-Term Sperm Storage in the Reproductive Tract of the Chinese Soft-shelled Turtle, *Pelodiscus sinensis*", *Scientific Reports*, vol. 5, 16096, 2015.
51. Xu D, Shen H, Zhu Q, Chua L, Wang Q, Taha S, Sayed S, Ghandour N, Mahran A, Saleh M, Amin M, Shamloul R. "Correlation between seminal lead and cadmium and seminal parameters in idiopathic oligoasthenozoospermic males". *Central European Journal of Urology*. Vol. 1 (66):84-92, 2013.
52. Manente L, Pecoraro S, Picillo E, Gargiulo U, Gargiulo P, De Luca A & Politano L, "Molecular evidence of apoptotic pathway activation in semen samples with high DNA fragmentation", *In Vivo*, vol 29, pp. 289–294. 2015.
53. Rodríguez-Romero F, Méndez-de la Cruz.F. "Reproductive Arrest in *Sceloporus mucronatus* (Lacertilia: Phrynosomatidae) Correlated with "El Niño Southern Oscillation". *Herpetological Review*, vol. 35 (2): 121-123, 2004.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.