

1 **Cancer and non-cancer risk of Santiago Island (Cape Verde) population due to**
2 **potential toxic elements exposure from soils**

3
4 Marina M.S. Cabral Pinto^{1*}, Maria M. V. Silva², Eduardo. A. Ferreira da Silva³, A. Paula. Reis³

5
6 ¹ Geobiotec Research Centre and CNC; Department of Earth Sciences, University of Coimbra,
7 3030-790 Coimbra, Portugal

8 ² CEMUC; Department of Earth Sciences, University of Coimbra, 3030-790 Coimbra, Portugal

9 ³ Geobiotec Research Centre, University of Aveiro, Department of Geosciences, 3810-193 Aveiro,
10 Portugal

11
12 * Corresponding author: mcp@uc.pt

13
14
15 **Abstract**

16 The hazard and the carcinogenic risks due to the exposure to some potentially toxic
17 elements by the Santiago Island (Cape Verde) population where calculated, considering
18 soil ingestion, inhalation and dermal contact as exposure pathways. The topsoil of
19 Santiago Island is enriched in Co, Cr, Cu, Ni, V, Zn, Mn and Cd to upper crust values.
20 Hazard indices (HI) were calculated for these metals and As exposures, of Santiago
21 Island population and the calculations were performed for children and adults. For
22 children HI are higher than 1 for Co, Cr and Mn. So there is indication of potential non-
23 carcinogenic risk for children, due to the high Co (HI=2.995), Cr (HI=1.329) and Mn
24 (HI=1.126), values in soils. For the other elements and for adults there is no potential
25 non-carcinogenic risk. Cancer risk was calculated for As, Cd, Cr and Ni exposures, for
26 adults and children and the results are always lower than the carcinogenic target risk of
27 1×10^{-6} , for As, Cd, and Ni. However, cancer risk are higher than the carcinogenic target
28 risk for Cr, for adults. Regarding As, for children the fraction due to Risk_{ingestion}
29 represents 51.6%, while Risk_{inhalation} represents 48.0% and Risk_{dermalcontact} represents only
30 0.4% of total risk. For adults Risk_{inhalation} represents 81.3%, Risk_{ingestion} represents 16.6%
31 and Risk_{dermal contact} represents 2.1%. These results reflect the higher daily ingestion dose

1 for children and the higher inhalation rate and higher dermal contact surface for adults.
2 For the other elements and for adults the cancer risk due to Cr, Ni and Cd inhalation is
3 always higher than for children, reflecting the higher inhalation rate for adults.

4

5 Keywords: cancer risk, risk assessment, volcanic soils, Santiago Island

6

7 **1. Introduction**

8 Soils are natural resources, formed at the Earth surface by weathering of underlying
9 rocks by the action of physical, chemical and biologic factors. They are the support of
10 agriculture, the main carbon reservoir of the terrestrial carbon cycle, act as sink for
11 pollutants, protecting groundwater from pollution and are also used as construction
12 material and support. So they have very important social, environmental and economic
13 functions (Vrščaj et al., 2008; Cabral Pinto et al., 2017a).

14 However, if contaminated or polluted they can transfer potentially toxic elements (PTE)
15 to groundwater, seepage waters and rivers and also to crops and vegetables which are
16 used by humans and animals, and consequently affect human health. The soil
17 contamination may be natural due to rock composition. Some elements can accumulate
18 on topsoil, to concentrations that are toxic to the plant, to the animal feeding on it and to
19 humans. Air quality may also be affect by contaminated soils, by the generation of
20 airborne particles and dust (Gray et al., 2003). In deeper soils due to changes in pH and
21 Eh, the PTE may be released to groundwater resulting in its contamination (Camobreco
22 et al., 1996; Pinto et al., 2004; Mirlean et al., 2007; Neiva et al., 2014).

23 The chemical composition of dust, soils and groundwater may cause metabolic changes
24 which may favor the occurrence of endemic diseases in humans (Salgueiro et al., 2010;
25 Cabral Pinto et al., 2014; Candeias et al., 2014; Patinha et al., 2015; Wu et al., 2015;
26 Centeno et al., 2016; Cabral Pinto et al. 2017b). The role of F, I, Se and As
27 concentrations in the health of human populations is well documented in the scientific
28 literature (Shamberger, 1983; Rayman, 2000; Kinniburgh and Smedley, 2001; Centeno
29 et al., 2002; Tchounwo et al., 2003; Charlet and Polya, 2006; Dissanayake and
30 Chandrajith, 2009; Tisserand et al., 2014; Centeno et al., 2016).

31 The high concentrations of PTE on topsoil can threaten human health via (a) soil

1 ingestion by geophagism, rare in adults, but quite common in children by hand-to-
2 mouth intake; (b) by inhalation of dust particles and (c) by dermal contact (Sun et al.,
3 2010; Xie et al., 2011), specially by farmers and construction workers and (d) indirectly
4 also by ingestion of contaminated groundwater.

5 Geochemistry of major, trace and rare earth elements (REEs) of soils of Santiago Island
6 (Cape Verde) has been done in order to characterize soils developed on volcanic rocks
7 and Quaternary sediments, contributing to the establishment of a geochemical atlas of
8 the island (Marques et al., 2012; Cabral Pinto et al., 2012; 2014; 2015).

9 Cabral Pinto et al. (2015) presented the Mapping of Estimated Background Values
10 (EBV), the agricultural and residential Environmental Risk Index (ERI) for each
11 element and the agricultural and residential Multi-Element (ME-ERI), which are the
12 average of the ERIs taken over all elements, for some harmful elements in soils of
13 Santiago Island (Cape Verde Archipelago). The present work follows the precedent
14 study in order to better understand the relationships between environmental
15 geochemistry and public health in a volcanic island which still preserve many pristine
16 geochemical characteristics, where the anthropogenic action is not yet too strong. We
17 present the hazard risk and the carcinogenic risk due to the exposure to some potentially
18 toxic elements by the Santiago Island population, according to the Exposure Factors
19 Handbook (USEPA, 2011). We consider soil ingestion, inhalation and dermal contact as
20 exposure pathways, because most of the population of the Island is rural and the island
21 is affected by strong winds (defined as "bruma seca") which mobilize significant
22 amounts of dust particles from soil (Almeida-Silva et al., 2013).


23

24 **2. Geographic, geologic and climate settings and soil types**

25 Cape Verde is formed by 10 islands, located at the African western shore (Fig. 1). The
26 country capital is located in Santiago Island, which is the biggest island. It is a
27 mountainous island with a maximum altitude of 1394 m. It has 215 km² of arable area
28 and estimated water resources of $56.6 \times 100 \text{ m}^3/\text{year}$ at the surface, and 42.4×100
29 m^3/year underground (PNUD, 1993).

30 The climate is semi-arid, with strong winds during the dry season, and a mean annual
31 precipitation of 321 mm, mainly due to torrential rains, in the wet season (INGM,

1 2005). According to the CCKP (2015) in the 1900-2012 years period the mean
 2 historical monthly rainfall attained 347.7 mm and the highest value (109.2 mm) was
 3 recorded in September. For the same years period the mean historical monthly
 4 temperature varied from 20.4 °C in February to 25.5 °C in September.




5
 6 Figure 1 - Cape Verde archipelago and its location.

7 .

8

9 The islands are volcanic, intraplate, located over a submarine plateau, the Cape Verde
 10 Rise, relatively stable within the African Plate and the volcanism resulted from the
 11 interaction of a mantle plume with the fractured lithosphere (Williams et al., 1990;

1 Millet et al., 2008). Santiago Island is a shield volcano, with periods of intense volcanic
 2 activity, with the emission of alkaline basaltic lava flows and subaerial pyroclastic
 3 materials, separated by erosion and sedimentations periods (Serralheiro, 1976). A brief
 4 description of the lithostratigraphic formations of Santiago is presented in Figure 2 (a)
 5 and Table 1.



6
 7 **Figure 1(a).** Geological cartography of the island of Santiago, Cape Verde, adapted from Serralheiro
 8 (1976); **(b).** Adapted soil cartography of the island of Santiago according to FAO/UNESCO Classification
 9 (1974), adapted from Hernandez (2008).

10


11 Table 1 -Brief description of each geological formation of soils of Santiago Island.

| Geological Formation | Rock type | Composition |
|--|---|--|
| CA - Ancient Internal Eruptive Complex | Subaerial and submarine lava flows and pyroclastic deposits; dykes and intrusive rocks | Basalts-basanites, phonolites-trachytes and carbonatites |
| FL – Flamengos formation | Submarine lava flows with subordinated breccias and tuffs | Basanites |
| CB – Orgãos Formation | Volcano-sedimentary deposits; rare lava flows | Diverse |
| PA – Pico da Antónia Eruptive Complex | Subaerial and submarine lava flows, dykes and pyroclastic material; intercalated sedimentary deposits | Basalts-basanites and phonolites-trachytes |
| AS – Assomada Formation | Subaerial lava flows and some pyroclasts | Basanites |
| MV- Monte das Vacas Formation | Subaerial pyroclasts and small subordinated lava flows | Basanites |
| CC – Recent sedimentary formations | Alluvial, aeolian and marine | Diverse |

deposits

1

2 Figure 2 (b) presents the adapted soil cartography of the Santiago Island, based on
 3 studies by Faria (1970), Diniz and Matos (1986), Hernandez (2008) and Cabral Pinto et
 4 al. (2012), according to the FAO/UNESCO classification and USA Soil Taxonomy.
 5 Table 2 shows a brief description of the soil cartography. The main soils are LT, RG, X
 6 and CM. Kastanozemes occur mainly in association with LV, which are the soil group
 7 typically used for agriculture in Cape Verde (Figure 2b and Table 2).



8

9

10 Table 2 -Brief description of each group of soils of Santiago Island.

| Pedological Formation | Development characteristics | Texture | Outcrop |
|-----------------------|--|---|---|
| LT - Lithosols | Immature incipient mineral soils with no or little differentiation (< 20 cm thickness). | Low clay and organic matter contents and high proportion of coarse-grained fractions. | North and centre |
| RG – Regosols | Incipient mineral and undifferentiated soils. No vegetation. | Mainly sandy and gravel materials. Low clay and organic matter contents. | In rare narrow and discontinuous coastal ridges. |
| FV – Fluvisols | Undifferentiated or show little differentiation. Developed on alluvial deposits on the banks of temporary or torrential streams. | Mainly sand and coarse particles. | In alluvial deposits on the banks of temporary or torrential streams. |

| | | | |
|-----------------|---|---|--|
| CM – Cambisols | Immature (profile AC), non-climate (20 - 30 cm thickness). | Mainly coarse-to-fine sand with high proportion of slightly weathered rock fragments. | Western part of the island, in the center and south. |
| K – Castanozems | Developed soils, but with moderately or poorly differentiated profiles and relatively rich in organic matter. | Fine-grained, mostly consisting of clay materials. | Center and in the southeast region of the island (small patches in the northern) |
| X – Xerosols | Sub-arid soils, with surface decalcification horizons and with some organic matter (0.8–1.8%). | Mainly coarse-to-fine material. | Southeastern coast. |
| VR - Vertisols | Non-lytic soils. Developed soils (ABC profile). | Fine-grained, up to 30% clay content. | South and east coastal areas. |
| LV – Luvisols | Developed soils (ABC profile). | High proportion of fine-grained particles (mainly clay) | Very small patches in the central and interior. |

1

2 **3. Methodologies**

3 **3.1. Sampling, chemical and statistical analysis**

4 The sampling, analytical and statistical methodologies were fully described in Cabral
5 Pinto et al. (2015), so only a brief description will be referred in the present paper.
6 Collection of 249 topsoil composite samples, away from potential anthropogenic
7 influence, was performed at a spatial resolution of 0.3 sites/km² (identified by GPS).
8 Duplicate field samples were collected at every 10 sites. The <2 mm fraction was
9 pulverized to < 75 µm, digested with *aqua regia* and analyzed by ICP-MS at the ACME
10 Analytical Laboratories, Canada. Lab-duplicate samples were taken at every 30 samples
11 to calculate the analytical precision (which was better than 10%) and certified standard
12 materials were analyzed to determine accuracy.

13 Variance analysis was performed to test the reliability of the data to be used in the
14 statistical analysis. The estimated background values (EBV-S) of the analyzed elements
15 was estimated as the median of the data limited by the Tukey Range. Principal
16 Component analysis (PCA) was performed to determine the associations of metals, with
17 Matlab 10 software.

1

2 **3.2. Risk assessment**

3 The environmental risk index (ERI) was calculated for PTE by Cabral Pinto et al.
4 (2015) using Canadian (Ministry of the Environment 2011) and Dutch (VROM, 2000),
5 legislations for soils. For each element $ERI=C(s)/P$, where computed. C(s) is the
6 element concentration at sampling site s and P is the permissive level of that element,
7 according to the legislations.

8 Non-cancer risk is represented in terms of hazard index (HI) for multiple substances
9 and/or exposure pathways (USEPA, 2011). HI is the sum of the hazard quotient (HQ),
10 for each element and each pathway and if it $HI < 1$ there is a very low chance for a non-
11 carcinogenic risk. $HQ = ADD/RfD$ where ADD is the average daily dose of an element
12 that a person is exposed and RfD is the reference dose (USEPA, 2011), below which the
13 non-cancer risk is negligible.

14 Hazard indices (HI) were calculated for Co, Cr, Cu, Ni, V, As, Zn, Mn and Cd exposure
15 of Santiago Island population, according to the Exposure Factors Handbook (USEPA,
16 2011). The equation used to calculate the average daily dose (ADD) for each pathway
17 are those presented in USEPA (2011).

18

$$19 \quad ADD_{\text{ingestion}} = \frac{C_{\text{soil}} \times \text{IngR} \times \text{EF} \times \text{ED}}{\text{BW} \times \text{AT}} \quad \text{Eq.- 1}$$

$$20 \quad ADD_{\text{dermal}} = \frac{C_{\text{soil}} \times \text{SA} \times \text{SAF} \times \text{DA} \times \text{EF} \times \text{ED}}{\text{BW} \times \text{AT}} \quad \text{Eq.- 2}$$

$$21 \quad ADD_{\text{inhalation}} = \frac{C_{\text{soil}} \times \text{InhR} \times \text{EF} \times \text{ED}}{\text{BW} \times \text{AT} \times \text{PEF}} \quad \text{Eq.- 3}$$

22

23 C_{soil} is the concentration of the element in the soil (mg.kg^{-1}) and we used the 95
24 percentile of the soil distribution values, presented in Cabral Pinto et al. (2015). IngR is
25 the soil ingestion rate and we used 200 mg.d^{-1} for children and 100 mg.d^{-1} for adults
26 (USEPA, 2011). EF is the exposure frequency and we used 365 days, considering the
27 inhabitants of the island. ED is the exposure duration and we assumed that an inhabitant
28 spend half of his life exposed, so we assumed 6 years for children and 35 years for adults

1 and AT is ED expressed in days for non-carcinogenic. For body weight (BW) we used
2 15 kg for children and 70 kg for adults and a life expectancy of 70 years.

3 The exposed skin area (SA) was taken as 2372 and 60,132 cm² for children and adults,
4 respectively. The used skin adherence factor (SAF) was 0.2 and 0.07 mg.cm⁻² for
5 children and adults respectively and the dermal absorption factor (DA) was 0.003 for As
6 and 0.001 for other elements (USEPA, 2001; 2011; USDE, 2013). InhR is the inhalation
7 rate taken as 7.6 and 20 m³.d⁻¹ for children and adults, respectively and the particle
8 emission factor (PEF) is 1.36x10⁹ m³.kg⁻¹.

9 The carcinogenic risks (Risk) were calculated for As, Cd, Cr and Ni exposure of
10 Santiago Island population, according to the Exposure Factors Handbook (USEPA,
11 2011) and using the RfD values according to USDE (2013).

12

13 4. Results and Discussion

14 The topsoil of Santiago Island is enriched in Cd, Ni, Co, Cu, V, Mn, Cr and Zn to upper
15 crust values (UCC) of Rudnick and Gao (2003), considering both the Estimated
16 Background Values (EBV) or the 95th percentile value calculated from the soil
17 distribution values (Table 3).

Table 3 - Estimated background values, 95 percentiles of Santiago soils (mg.kg⁻¹) and ratios of these values to upper continental crust (UCC) and published guidelines values (Gdl).

| | EBV | P95(s) | UCC | Guidelines* | EBV/UCC | P95(s)/UCC | EBV/Gdl | P95(s)/Gdl |
|----|-------|--------|-------|-------------|-------------|------------|------------|------------|
| As | 0.6 | 1.92 | 4.8 | 11 | 0.1 | 0.4 | 0.1 | 0.2 |
| Cd | 0.20 | 0.4 | 0.09 | 0.8 | 2.22 | 4.4 | 1.1 | 0.5 |
| Co | 46.4 | 67.11 | 17.3 | 9 | 2.7 | 3.9 | 5.2 | 7.5 |
| Cr | 118 | 298 | 92 | 67 | 1.3 | 3.2 | 1.8 | 4.4 |
| Cu | 50.8 | 81.8 | 28 | 36 | 1.8 | 2.9 | 1.4 | 2.3 |
| Mn | 1293 | 1954 | 774.5 | - | 1.7 | 2.6 | - | - |
| Ni | 136.1 | 267.9 | 47 | 36 | 2.9 | 5.7 | 3.8 | 7.4 |
| V | 169 | 260 | 97 | 86 | 1.7 | 2.7 | 2.0 | 3.0 |
| Zn | 79 | 120.2 | 67 | 140 | 1.2 | 1.8 | 0.6 | 0.9 |

EBV: estimated background values¹; P95(s) 95 percentile in Santiago soils¹; Gdl: guidelines;

UCC: Upper continental crust²; *: Canadian and Dutch Guidelines³.

¹Cabral Pinto et al. (2015); ²Rudnick and Gao (2003); ³Ministry of the Environment (2011) and (VROM, 2000)

18

19

20 In Santiago Island the soil geochemistry is mainly controlled by lithology, although

1 some elements may have an anthropogenic influence, such as As, Hg, Cd, Zn and Pb
 2 (Cabral Pinto et al., 2015). Using the permissible values for agricultural soils in the
 3 calculations, their results shown that soil of the entire island has environmental risk
 4 index (ERI) above 1 for Co, Ni, Cr, V, Cu considering the 95th percentile (Table 3).
 5 The Hazard Quotients (HQ) for ingestion, dermal contact and inhalation routes and
 6 Hazard Indices (HI) were calculated (Table 4), for the metals which show enrichment to
 7 UCC and also for As, which is a toxic and carcinogenic element, Zn and Mn.

Table 4- HQ values for various pathways and elements and HI for elements from Santiago Island

| | HQ ingestion | | HQ dermal | | HQ inhalation | | HI | |
|----|--------------|--------|-----------|--------|---------------|--------|----------|--------|
| | children | adult | children | adult | children | adult | children | adult |
| Co | 2.9827 | 0.3196 | 0.0084 | 0.0013 | 0.0042 | 0.0023 | 2.9952 | 0.3232 |
| Cr | 1.3244 | 0.1419 | 0.0037 | 0.0006 | 0.0011 | 0.0006 | 1.3293 | 0.1431 |
| V | 0.6878 | 0.0737 | 0.0019 | 0.0003 | 0.0010 | 0.0005 | 0.6907 | 0.0745 |
| Ni | 0.1786 | 0.0191 | 0.0005 | 0.0001 | 0.0011 | 0.0006 | 0.1802 | 0.0198 |
| Cu | 0.0273 | 0.0029 | 0.0001 | 0.0000 | | | | |
| As | 0.0853 | 0.0091 | 0.0007 | 0.0001 | 0.0000 | 0.0000 | 0.0861 | 0.0093 |
| Zn | 0.0053 | 0.0006 | 0.0000 | 0.0000 | | | | |
| Cd | 0.0053 | 0.0006 | 0.0006 | 0.0001 | 0.0000 | 0.0000 | 0.0059 | 0.0007 |
| Mn | 1.1111 | 0.1190 | | | 0.0149 | 0.0084 | 1.1260 | 0.1275 |

| Cancer risk | | | |
|-------------|----------|---------|---|
| | children | adult | |
| As | 6.4E-09 | 1.2E-08 | for ingestion, dermal contact and inhalation pathways |
| Cr | 3.9E-07 | 1.3E-06 | |
| Ni | 7.2E-09 | 2.4E-08 | for inhalation |
| Cd | 7.8E-11 | 2.6E-10 | |


8

9 The selected elements are potentially toxic elements and some (As, Cd, Cr and Ni) are
 10 also carcinogenic (USDE, 2013). The pathways chosen were ingestion, inhalation and
 11 dermal contact and the calculations were performed for children and adults. Major
 12 neurodegenerative disorders, including Alzheimer's and Parkinson's disease, are
 13 characterized by elevate tissue metals, such as Fe, Cu, Mn and Zn (Barnham and Bush
 14 2008). Environmental exposure to Mn can induce parkinsonism and although the long-
 15 term medical significance of this finding is unclear, the data are troubling and point to
 16 the need for further investigation of manganese's health risks (Ahlskog, 2016).

1 The non-carcinogenic hazard indices (HI) for all nine elements are given in Table 4. For
 2 adults the HI are always smaller than 1, whereas for children they are higher than 1 for
 3 Co, Cr, and Mn. The HI value of these elements is mainly controlled by the $HQ_{\text{ingestion}}$,
 4 which are also higher than 1 for these 3 elements. The $HQ_{\text{ingestion}}$ is always the highest
 5 for all the elements, while the $HQ_{\text{inhalation}}$ is the lowest (Figure 3).

6

7 Figure 3- Hazard index (HI) and hazard quotient (HQ) for potentially toxic elements from Santiago
 8 Island. Symbols: HI-crosses; HQ for ingestion-diamonds; HQ for dermal contact – squares; HQ for
 9 inhalation- triangles.



10

11 So there is indication of potential non-carcinogenic risk for children, due to the high Co
 12 (HI=2.995), Cr (HI=1.329) and Mn (HI=1.126), values in soils. For the other elements
 13 and for adults there is no potential non-carcinogenic risk. For both children and adults
 14 the HI is $Co > Cr > Mn$. Compared to adult, children health index is larger, and their
 15 cumulative effect is also of concern for children.

16 The evaluation of cancer risk was performed only with those elements which are
 17 potentially carcinogenic (USDE, 2013). For As, ingestion, inhalation and dermal contact
 18 exposure pathways were considered, but for the other carcinogenic elements (Cd, Cr
 19 and Ni) only the inhalation risk were computed because the the Risk Assessment
 20 Information System (USDE, 2013) does not present slope factors for the other
 21 exposures.

1 For children the As cancer risk ($risk_{total}$) is 6.38×10^{-9} (Table 4) and of this, the fraction
2 due to $Risk_{ingestion}$ represents 51.6%, while $Risk_{inhalation}$ represents 48.0% and
3 $Risk_{dermalcontact}$ represents only 0.4% of total risk. For adults $risk_{total}$ for As is 1.24×10^{-8}
4 (Table 4) where $Risk_{inhalation}$ represents 81.3%, $Risk_{ingestion}$ represents 16.6% and
5 $Risk_{dermal contact}$ represents 2.1%. These results reflect the higher daily ingestion dose for
6 children and the higher inhalation rate for adults. For adults the cancer risk due to Cr, Ni
7 and Cd inhalation is always higher than for children, reflecting the higher inhalation rate
8 for adults.

9 The results for cancer risk are higher than the carcinogenic target risk of 1×10^{-6}
10 (USEPA, 2011) only for Cr, for adults, but these results underestimate the risk. The
11 other pathways were not considered and they can be important for Cr, for children
12 which present a cancer risk of 1.3×10^{-6} , very close to the target risk. The lack of RfD for
13 some elements disables a more complete evaluation of the cancer-risk.

14 Santiago Island still has an almost pristine surface environment and the topsoil
15 composition is mainly determined by the composition of the underlying basic rock.
16 These rocks are rich in siderophile elements promoting a natural contamination of soils
17 in Co, Cr, Ni, Cu, V, Zn, Cd and Mn. Of these, Co, Cr and Mn present a potential non-
18 carcinogenic risk for children, which are vulnerable people. On the other way the soil
19 composition affects groundwater composition, so there is a flux of the natural
20 contamination from soils to groundwater which deserves to be evaluated. The
21 inhabitants of Santiago Island depend on groundwater for consumption and for
22 agriculture and the flux water-vegetables-men also deserve evaluation, because endemic
23 diseases can be controlled with proper measures, if its cause is well constrained.

24

25 **5. Conclusions**

26 The topsoil of Santiago Island, Cabo Verde, have a geochemical composition mainly
27 controlled by the type of underlying rock, as most of the elements in the topsoil have
28 mainly a geogenic origin.

29 The environmental risk index (ERI) calculations shown that the Santiago Island topsoil
30 are naturally contaminated in Co, Cr, Cu, Ni and V, because these elements have
31 contents well above those allowed by Canadian and Dutch legislations for agricultural

1 soils.

2 The non-carcinogenic hazard indices (HI) were calculated for 9 potentially toxic
3 elements and they are always smaller than 1 for adults, considering that the soil
4 contaminants enter the human body by soil ingestion, dermal contact and inhalation of
5 dust particles. For children the non-carcinogenic hazard indices are 2.9952 for Co,
6 1.3293 for Cr and 1.1111 for Mn. For the other elements they are lower than 1.

7 The cancer risk is higher than the carcinogenic target risk of 1×10^{-6} for Cr, for adults.
8 But these results may be underestimated as only the inhalation risk was calculated for
9 Cr, Ni and Cd. Also soil contaminants may be indirectly ingested by groundwater and
10 by crop and vegetables consumption. The elements flux soil-groundwater-crops and
11 vegetables may increase the hazard and cancer risk.

12 There is need of evaluation of the risks associated with groundwater consumption and
13 diet in Santiago Island.

14

15 **6. References**

16 Ahlskog, J.E., 2016. The new Parkinson's Disease. Oxford university press, New Yourk,
17 2° edition. 544 p.

18 Almeida-Silva, M., Almeida, S.M., Freitas, M.C., Pio, C.A., Nunes, T., Cardoso, J.,
19 2013. Impact of Sahara dust transport on Cape Verde atmospheric element
20 particles. *Journal of Toxicology and Environmental Health* 76(4-5), 240-251.

21 Barnham, K.J., Bush, A.I., 2008. Metals in Alzheimer's and Parkinson's
22 diseases. *Current opinion in chemical biology* 12(2), 222-228.

23 Cabral Pinto, M. M., Marinho-Reis, A. P., Almeida, A., Ordens, C. M., Silva, M. M.,
24 Freitas, S., ... & da Silva, E. A. F., 2017b. Human predisposition to cognitive
25 impairment and its relation with environmental exposure to potentially toxic
26 elements. *Environmental Geochemistry and Health*, 1-18.
27 <https://link.springer.com/article/10.1007/s10653-017-9928-3>

28 Cabral Pinto, M. M., Silva, M. M., da Silva, E. A. F., Dinis, P. A., & Rocha, F., 2017.
29 Transfer processes of potentially toxic elements (PTE) from rocks to soils and the
30 origin of PTE in soils: A case study on the island of Santiago (Cape

- 1 Verde). Journal of Geochemical Exploration.
2 <https://doi.org/10.1016/j.gexplo.2017.06.004>
- 3 Cabral Pinto, M.M.S., Ferreira da Silva E.A., Silva M.M.V.G., Melo-Gonçalves P.,
4 2015. Heavy metals of Santiago Island (Cape Verde) top soils: Estimated
5 Background Value maps and environmental risk assessment. Journal of African
6 Earth Sciences 101, 162–176.
- 7 Cabral Pinto, M.M.S., Ferreira da Siva, E.A., Silva, M.M.V.G., Melo-Gonçalves, P.,
8 Hernandez, R. Marinho, A.P., Inácio, M., Rocha, F., 2012. Mapping of Estimated
9 Geochemical Background Values of some harmful metals in Soils of Santiago
10 Island (Cape Verde archipelago), 10, 101-122. In 'Progress on Heavy Metals in the
11 Environment. Ref.:Ms. No. MARALTEBOOKS-D-11-00053R1. J., Nriagu, P.,
12 Szefer, J., Pacyna, B., Markert & S., Wünschmann Eds. DOI:10.5645/b.2.10
- 13 Cabral Pinto, M.M.S., Silva, E. A., Silva, M.M., Melo-Gonçalves, P., Candeias, C.,
14 2014. Environmental Risk Assessment Based on High-Resolution Spatial Maps of
15 Potentially Toxic Elements Sampled on Stream Sediments of Santiago, Cape
16 Verde. Geosciences 4, 297-315.
- 17 Camobreco, V.J., Richards B.K., Steenhuis T.S., Peverly J.H., McBride M.B., 1996.
18 Movement of heavy metals through undisturbed and homogenized soil columns.
19 Soil Science 161, 740-750.
- 20 Candeias, C., Silva, E.F., Ávila, P.F., Teixeira, J.P., 2014. Identifying sources and
21 assessing potential risk of exposure to heavy metals and hazardous materials in
22 mining areas: The case study of Panasqueira mine (Central Portugal) as an
23 example. Geosciences 4, 240-268.
- 24 CCKP: Climate Change Knowledge Portal, 2015.
25 [http://sdwebx.worldbank.org/climateportal/index.cfm?page=country_historical_cl](http://sdwebx.worldbank.org/climateportal/index.cfm?page=country_historical_climate)
26 [imate](http://sdwebx.worldbank.org/climateportal/index.cfm?page=country_historical_climate) ThisRegion=Africa&ThisCCCode=CPV.
- 27 Centeno, J. A., Finkelman, R. B., Selinus, O., 2016. Medical geology: Impacts of the
28 natural environment on public health. Geosciences 6(1), 8.
- 29 Centeno, J.A., Mullick, F.G., Martinez, L., Page, N. P., Gibb, H., Longfellow, D.,
30 Thompson, D., Ladich, E.R., 2002. Pathology Related to Chronic Arsenic
31 Exposure. Environmental Health Perspectives 110(5), 883-886.

- 1 Charlet, L., Polya, D.A., 2006. Arsenic in shallow, reducing groundwater in southern
2 Asia: An environmental health disaster. *Elements* 2, 91–96.
- 3 Diniz, A., Matos, G., 1986. Carta de zonagem agro-ecológica e de vegetação de Cabo
4 Verde. I Ilha de Santiago, Cabo Verde. *Garcia da Orta* 83, 982 p.
- 5 Dissanayake, C.B., Chandrajith, R., 2009. Introduction to Medical Geology Focus on
6 tropical environments. Springer, 297 p.
- 7 Faria, X., 1970. Os Solos da ilha de Santiago. *Est Ens Doc, Junta de Investigações do*
8 *Ultramar, Lisboa.*
- 9 Gray, C., McLaren, R., Roberts, A., 2003. Atmospheric accessions of heavy metals to
10 some New Zealand pastoral soils. *Science of the Total Environment* 305, 105-115.
- 11 Hernandez, R.V.A., 2008. Caracterização dos solos da ilha de Santiago (Cabo Verde)
12 numa perspectiva de sustentabilidade ambiental. Unpublished MSc thesis. Aveiro
13 University, 344 p.
- 14 Kinniburgh, D.G., Smedley, P.L., (eds.). 2001. Arsenic concentrations of groundwater in
15 Bangladesh. British Geological Survey Technical Report WC/00/19, 1, 14p.
- 16 Marques, R., Prudêncio, M. I., Rocha, F., Cabral Pinto, M.M.S., Silva, M.M.V., Ferreira
17 da Silva, E.A., 2012. REE and other trace and major elements in the topsoil layer
18 of Santiago Island, Cape Verde. *Journal of African Earth Sciences* 64, 20-33.
- 19 Millet, M.-A., Doucelance, R., Schiano, P., David, C., Bosp, C., 2008. Mantle plume
20 heterogeneity versus shallow-level interactions: a case study, the São Nicolau
21 Island, Cape Verde archipelago. *J. Volcanol Geotherm Res.* 176, 265–276.
- 22 Ministry of the Environment, 2011. Soil, Groundwater and Sediment Standards for Use
23 Under Part XV.1 of the Environmental Protection Act. Ministry of the
24 Environment.
- 25 Mirlean, N., Roisenberg, A., Chies, J.O., 2007. Metal contamination of vineyard soils in
26 wet subtropics (southern Brazil). *Environmental Pollution* 149 (1), 10-17.
- 27 Neiva, A.M.R., Carvalho, P.C.S., Antunes, I.M.H.R., Silva, M.M.V.G., Santos, A.C.T.
28 Cabral Pinto, M.M.S., Cunha, P.P., 2014. Contaminated water, stream sediments
29 and soils close to the abandoned Pinhal do Souto uranium mine, central Portugal.
30 *Journal of Geochemical Exploration* 136, 102–117.

- 1 Patinha, C., Reis, A.P., Dias, A.C., Abduljelil, A.A., Noack, Y., Robert, S., ... Silva,
2 E.F., 2015. The mobility and human oral bioaccessibility of Zn and Pb in urban
3 dusts of Estarreja (N Portugal). *Environmental geochemistry and health* 37(1),
4 115-131.
- 5 Pinto, M. M. S. C., Silva, M. M. V. G., & Neiva, A. M. R., 2004. Pollution of water and
6 stream sediments associated with the Vale De Abrutiga Uranium Mine, Central
7 Portugal. *Mine Water and the Environment*, 23(2), 66-75.
- 8 PNUD, 1993. United Nations Development Program for Cape Verde, PNUD. New
9 York, NY, USA.
- 10 Rayman M.P., 2000. The importance of selenium to human health. *The Lancet* 356, 233-
11 241.
- 12 Rudnick, R.L., Gao, S., 2003. Composition of the Continental Crust. In Holland, H.D.,
13 Turekian, K.K. (Eds.), *Treatise on Geochemistry*, vol. 3. Elsevier, New York, 1-
14 64.
- 15 Salgueiro, A. R., Ávila, P. H., Pereira, H. G., & da Silva, E. F. (2010). Assessment of the
16 impact of pollution by arsenic in the vicinity of Panasqueira Mine (Portugal).
17 In *geoENV VII—Geostatistics for Environmental Applications*, 151-159. Springer
18 Netherlands.
- 19 Serralheiro, A., 1976. A Geologia da ilha de Santiago (Cabo Verde). *Boletim Museu*
20 *Laboratório Mineralógico Geológico Faculdade de Ciências de Lisboa* 14, (2).
- 21 Shamberger, R.J., 1983. Selenium and health. Pp 167-181. In *Trace elements in health:*
22 *a review of current issues*, (Ed) J. Rose, Butterworths.
- 23 Sun, Y., Zhou, Q., Xie, X., Liu, R., 2010. Spatial, sources and risk assessment of heavy
24 metal contamination of urban soils in typical regions of Shenyang, China, *Journal of*
25 *hazardous materials* 174(1-3), 455-462.
- 26 Tchounwou, P.B., Patlolla, A.K., Centeno, J.A., 2003. Carcinogenic and systemic health
27 effects associated with arsenic exposure – a critical review. *Toxicologic Pathology*
28 31, 575-588.
- 29 Tisserand, D., Pili, E., Hellmann, R., Boullier, A-M., Charlet, L., 2014. Geogenic
30 arsenic in groundwater in the western Alps. *Journal of Hydrology* 518, 317–325.

- 1 USDE. 2013. U.S. Department of Energy. The risk assessment Information System
2 (RAIS); U.S. Department of Energy's Oak Ridge Operations Office: Oak Ridge,
3 USA. Consulted in April 2016 in: <https://rais.ornl.gov/>
- 4 USEPA. 2001. United States Environmental Protection Agency. Risk Assessment
5 Guidance for Superfund: Volume III–Part A, Process for Conducting Probabilistic
6 Risk Assessment; EPA 540-R-02-002.
- 7 USEPA. 2011. United States Environmental Protection Agency. Exposure factors
8 handbook 2011 edition (Final). Consulted in April 2016 in:
9 <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=236252>.
- 10 VROM. 2000. Circular on target values and intervention values for soil remediation.
11 Netherlands Government Gazette, 39. Ministry of Housing, Spatial Planning and
12 Environment, Directorate General for Environmental Protection, Department of
13 Soil Protection. The Hague Widerlund A. Shcherbakova.
- 14 Vrščaj B., Poggio, L., Marsan, F.A., 2008. A method for soil environmental quality
15 evaluation for management and planning in urban areas. *Landscape and Urban*
16 *Planning* 88 (2-4), 81-94.
- 17 Williams, C., Hill, I., Young, R., White, R.S., 1990. Fracture zones across the Cape
18 Verde Rise, NE Atlantic. *J Geol. Soc. London* 147, 851–857.
- 19 Wu, S., Peng, S., Zhang, X., Wu, D., Luo, W., Zhang, T.,..., Wu, L., 2015. Levels and
20 health risk assessments of heavy metals in urban soils in Dongguan,
21 China. *Journal of Geochemical Exploration* 148, 71-78.
- 22 Xie, Y., Chen, T. B., Lei, M., Yang, J., Guo, Q.J., Song, B., 2011. Spatial distribution of
23 soil heavy metal pollution estimated by different interpolation methods: accuracy and
24 uncertainty analysis. *Chemosphere* 82(3), 468-476.

25

26 **Acknowledgements**

27 The authors would like to thank the funding provided by the Portuguese Foundation for
28 Science and Technology (SFRH/BPD/71030/2010 and UID/GEO/04035/2013 projects),
29 Geobiotec Centre from Aveiro University, and CNC, CEMUC and Geosciences Centre
30 from Coimbra University.