

Proteostatic Signaling & Control of Protein Synthesis

AJ Keefe

Department of Biology, Big Bend Community College, Moses Lake, Washington, 98837
Email: AJK@BigBend.edu

Keywords:

Proteostasis, translation, neurodegeneration, mtUPR, UPR, stress granules, ATF4

Pages: 15

Body Word Count: 6289

Character Count Excluding Spaces: 39166

Figures: 2

Citations: 121

The author has declared no conflicts of interest

The author received no funding for article preparation

Figure Captions

Figure 1. The UPR Signaling Network

Protein misfolding in the ER activates PERK and Ire1 by competing BiP away from their luminal domains and promoting multimerization. PERK phosphorylates eIF2a resulting in attenuated translation and ATF4 expression. PERK also activates NRF2, and in concert with ATF4, these transcription factors promote cytoprotective gene expression. Prolonged ATF4 transcription promotes cell death through the expression of CHOP and PUMA. Ire1 activates both proapoptotic and antiapoptotic pathways through RIDD and XBP1, respectively.

Figure 2. mTORC & mtUPR Signaling Network

Proteostatic networks activated by mtUPR and chaperone sequestration. mTORC inhibition, derived from JNK or chaperone depletion, results in downregulated translation. The mtUPR activates a myriad of pathways that converge of eIF2a phosphorylation, ATF4 expression, and mTORC inhibition. Both events lead to stress granule assembly which may also attenuate translation.

Highlights:

- The Integrated Stress Response (ISR) functions to maintain proteostasis by stalling translation.
- The mitochondrial UPR (mtUPR) operates through the ISR.
- mTORC1 mediated translation is lost during proteostasis
- Stress granule formation can also cause translation stalling

ABBREVIATIONS

UPR, Ubiquitin Proteasome System; UPR, Unfolded Protein Response; mtUPR, Mitochondrial Unfolded Response; ER, Endoplasmic Reticulum; ISR, Integrated Stress Response; eIF2a, Eukaryotic Initiation Factor 2 Alpha; uORF, Upstream Open Reading Frame; ATF4, Activating Transcription Factor 4; PKR, Protein Kinase R; PERK, protein kinase R (PKR)-like endoplasmic reticulum kinase; ARE, Antioxidant Response Elements; CRT, Calreticulin; TOP, Terminal Oligopyrimidine Tract; RIDD, Regulation Ire1 Dependent Decay; HSR, Heat Shock Response; HSF1, Heat Shock Factor 1; MAMs, Mitochondrial Associated Membranes.

ABSTRACT

The tremendous diversity and complexity of proteins invariably results in protein misfolding, to which cells have evolved numerous mechanisms of mitigating. Degrading misfolded proteins is perhaps the most intuitive strategy, but also critical to managing proteostasis are the elaborate mechanisms of translational control. Attenuated rates of translation ameliorate protein misfolding by downregulating the flux of new protein and conserving ATP. Loss of translational control, particularly in neurons, constitutes a major proteostatic dysfunction capable of causing or exacerbating neurodegeneration, while interventions aimed at downregulating protein synthesis are generally neuroprotective. In this review, I examine the critical neuronal signaling networks employed to control translation with an emphasis on current research. This includes the Unfolded Protein Response (UPR), the mitochondrial UPR (mtUPR), mTORC1 signaling, and stress granule formation.

INTRODUCTION

Neuronal proteostasis relies on highly regulated coordination between protein translation, chaperone-assisted folding, and protein degradation. This "central dogma" of protein homeostasis encompasses a wide range of interconnected pathways such as the Ubiquitin-Proteasome System (UPS), the UPR, and autophagy. Although a great deal of research has interrogated mechanisms of degrading proteotoxic protein, an underappreciated cellular response to protein misfolding is simply preventing or controlling protein synthesis until proteostatic conditions are restored.

Neurons maintain stringent and selective control of translation during basal steady-state conditions and is typically only derepressed during neuronal stimulation [1]. Neurons are thus sensitive and primed to respond to proteotoxic stress by utilizing these same translational regulations. Uncoupling translational repression in response to altered proteostasis can aggravate protein misfolding and compromise cell viability [2, 3, 4]. In contrast, genetic, dietary, and pharmacological means of downregulating translation is sufficient to extend lifespan and protect against protein misfolding diseases by improving proteostasis [6, 7, 8, 9, 10, 11, 12, 13]. The cytoprotection offered by attenuated translation is attributed to three principles. First, slowing the flux of new protein affords chaperones and degradative machinery more time to restore proteostasis [14]. Temporarily halting translation is also an effective strategy of maintaining ATP concentrations since translation is energetically expensive [15]. Calculations have estimated that translation is responsible for consuming 50-70% of a cell's pool of ATP [16, 17]. Lastly, translation is an error-prone process and requires chaperones that, during proteostatic disruption, could be preoccupied or sequestered in aggregates resulting in protein misfolding during translation. The inaccuracy of translational machinery is estimated to result in point mutations every 10^3 to 10^4 codons, affecting about 18% of all proteins [18, 19]. And of these mutations, roughly 10-50% of them are loss of folding mutations [20]. Stress-induced deficiencies in translation fidelity increases the rate of protein misfolding, underscoring the importance of chaperone surveillance during translation [121, 123]. Many proteins, and protein complexes such as mTORC1, also require an available pool of chaperones in order to simply adopt correctly folded conformations, regardless of the presence of mutations. Indeed, up to 30% of newly synthesized proteins emerge from the ribosome misfolded, and, in total, about 12-15% of them are promptly degraded in a process termed cotranslational ubiquitination [21]. Thus deregulated translation during cellular stress challenges metabolic homeostasis by depleting ATP, while additionally straining proteostasis due to the unsupervised

production of misfolded protein that may exceed the capacity of refolding and degradative networks.

Although transient translational stalling can ameliorate protein misfolding, it is vital that a neuron restores proteostasis expeditiously, since sustained translation inhibition due to unresolved protein misfolding leads to neuronal dysfunction and, eventually, cell death [22, 23]. The physiological consequence of persistent translation inhibition is exemplified in Vanishing White Matter Syndrome, caused by mutations in the eIF2B translation initiation factor, resulting in significant defects in translation and early onset neurodegeneration [24]. There is also evidence that conventional protein misfolding disorders, like Alzheimer's Disease (AD), Parkinson's Disease, and polyglutamine repeat disorders, are deleteriously affected by chronic translation inhibition [124]. Translation inhibition becomes unsustainable as the proteome becomes increasingly carbonylated from ROS signaling, depleted of critical proteins from degradation, and eventually the induction of cell death from the accumulation of apoptotic post-translational modifications. Proteostatic control of protein translation is likely beneficial during the early stages of disease while proteotoxic stress is manageable, but contributes to cellular dysfunction when the cell becomes incorrigibly overwhelmed by misfolded protein. Thus dynamic monitoring and control of translation, along with efficient proteome turnover, are both critical facets to healthy cellular proteostasis.

Protein misfolding and aggregate formation is nearly ubiquitous across neurodegenerative diseases and, not surprisingly, significant perturbations in the major proteostatic signaling networks (UPS, autophagy, UPR, mTORC, and mtUPR) are common. Additionally, many genetic manipulations of these networks in mouse models, or SNPs in humans, cause or increase susceptibility to neurodegenerative disease [27]. Neuronal vulnerability to proteotoxic stressors stems primarily from their status as post-mitotic cells, which are unable to dilute aggregates through cell division. The accumulation of aggregates may also be uniquely neurotoxic since neurons must facilitate vesicular transport through axons that can become physically obstructed by aggregated protein. Additionally, neurons are metabolically demanding, and even transient lapses in ATP production lead to cell death, as illustrated by their susceptibility to oxygen deprivation or oxidative phosphorylation failure. Neurodegenerative diseases are fundamentally disorders of altered proteostasis, thus a general understanding of how proteostatic signaling networks are activated, regulated, and communicated within the neuron may yield important discoveries and therapeutic opportunities.

In this review, I describe how proteostatic dysfunction is communicated within the neuron to achieve translational control and appropriate gene expression in order to ameliorate

proteotoxic stress. Specifically, I will examine how protein misfolding in the endoplasmic reticulum (ER) and the mitochondria initiate proteostatic measures that regulate translation through the Integrated Stress Response (ISR), how the inactivation of mTORC1 complements this process, and how stress granules sequester latent mRNA and protein during translational shutdown.

The ISR

The ISR is an evolutionarily conserved cellular program that converges on the phosphorylation of ribosomal initiation complex subunit eIF2a (eukaryotic initiation factor 2 alpha). Phosphorylation of eIF2a at serine 51 blocks the formation of the ribosomal initiation complex on mRNA resulting in attenuated rates of translation and ATP conservation. Paradoxically, p-eIF2a actually promotes the translation of certain transcripts bearing alternative upstream open reading frames (uORFs). Activating Transcription Factor 4 (ATF4) is the most well-understood p-eIF2a activated uORF and has earned a reputation as a stress-induced transcription factor capable of inducing proteostatic, autophagic, and antioxidant gene expression. By utilizing a basic leucine zipper domain, which requires dimerization for transcriptional activation, ATF4's transcriptional outcome can be fine-tuned by a diverse set of co-transcription factors. Although ATF4 typically homodimerizes, it also heterodimerizes with many other transcription factors such as AP-1, ATF3, or CHOP, and these interactions vastly expand ATF4's transcriptional versatility. ATF4/CHOP dimers, for example, are responsible for shifting the ISR's gene expression from cytoprotective to proapoptotic [28]. By utilizing stringent stress-activated translation through p-eIF2a, and heterodimerization with a diverse set of transcription factors, ATF4 has become a central player in the transduction of multiple stress pathways.

The critical checkpoint dictating this pathway is eIF2a phosphorylation, which is orchestrated by only four kinases that are each expressed in the brain - GCN2, HRI, PKR, and PERK. GCN2 is canonically activated by amino acid starvation, HRI by heme deficiency, PKR by double-stranded RNA, and PERK by ER protein misfolding. The activation of each of these kinases is derived from unique stimuli, thereby granting cells the ability to integrate diverse sources of stress into a single, but highly malleable, cytoprotective transcriptional response. Each of the following pathways discussed in this review involves, at least in part, this Integrated Stress Response, and thus the ISR has emerged as a recurring theme in proteostasis.

The UPR

Misfolded proteins are capable of triggering the ISR, and thus translational stalling and eIF2a induced ATF4 expression, through the PERK arm of the UPR. PERK, an ER membrane-embedded kinase activated by dimerization, is constitutively bound and inhibited by ER folding chaperone BiP. Misfolded proteins, however, compete BiP away from PERK enabling its homodimerization, activation, and phosphorylation of eIF2a. ER calcium depletion, a phenomenon observed in various neurodegenerative diseases like ALS, HD, and PD, has also been shown to activate PERK [29]. Since many ER folding chaperones, including BiP itself, require high calcium concentrations in order to function, protein misfolding could be exacerbated by chronic calcium depletion. Regardless, PERK-mediated phosphorylation of eIF2a triggers a rapid decrease in protein translation (~80% decrease), and the noncanonical translation of ATF4, which subsequently translocates to the nucleus and upregulates cytoprotective gene expression. Prolonged ATF4 transcriptional activity leads to the upregulation of CHOP, a secondary transcription factor that promotes apoptotic gene expression and accounts for ATF4's duration dependent toxicity. In neurons, ATF4/CHOP dimers trigger ER stress-induced apoptosis through the BH3 protein PUMA [30]. CHOP also upregulates GADD34, a stress-induced subunit of the PP1A phosphatase that redirects its substrate specificity to P-eIF2a and restores normal translation [31]. Premature restoration of protein synthesis, through both GADD34 dependent and independent PP1A activation, has been shown to promote cell death by depleting ATP and increasing ROS generation [31, 32].

Although PERK is recognized most often for its role in connecting the UPR with the ISR and ATF4, the kinase is also important for initiating an antioxidant and autophagic response through the activation of NRF2. PERK-mediated phosphorylation of NRF2 displaces its negative regulator KEAP1, preventing NRF2 ubiquitination and unveiling a nuclear localization signal [33, 34]. After activation by PERK, NRF2 translocates to the nucleus and binds Antioxidant Response Elements (ARE) promoting cytoprotective gene expression. NRF2 promotes the expression of ER folding chaperones, antioxidant proteins, and various autophagic proteins like LAMP2A, for chaperone-mediated autophagy, and p62, for macroautophagy [35, 36, 37]. Interestingly, p62 is capable of reciprocally activating NRF2 by binding and displacing KEAP1, potentially fueling a positive feedback loop during NRF2 dependent autophagy. Like NRF2, ATF4 is also responsible for activating autophagic gene expression including genes such as ATG5, p62, and Beclin1 [38, 39]. A study by B'Chir et al has even suggested that p-eIF2a/ATF4 is required for stress-induced autophagy [40]. In addition to the transcriptional upregulation of autophagic genes, PERK/eIF2a phosphorylation may also be critical to the processing of the

autophagic cargo adaptor receptor LC3 into LC3-II following expanded polyglutamine expression, but the mechanism is not understood [41]. The activation of autophagy is not only a consequence of translation inhibition but can also sustain it by promoting the degradation of ribosomes (ribophagy) [42]. Thus the influence ATF4 and NRF2 have on autophagy could, in theory, function to prolong translation inhibition. Indeed, ATF4 activation results in decreased ribosomal subunit protein expression [43]. There is also evidence that ATF4 and NRF2 physically interact, and do so in such a way to prevent ATF4 from binding and initiating transcription at the promoter of proapoptotic CHOP [44]. In addition to preventing apoptotic gene expression, ATF4/NRF2 cotranscription potentiates antioxidant gene expression [45]. The significant overlap between the cytoprotective transcriptional outcomes of NRF2 and ATF4, and their common mechanism of activation through PERK, suggests that the UPR likely relies on both transcription factors for proteostatic gene expression in response to ER protein misfolding

Loss of BiP due to the presence of misfolded protein can also activate the ER stress sensor Ire1, resulting in its multimerization and formation of an atypical endoribonuclease. UPR induced Ire1 oligomerization communicates to the nucleus via the removal of an inhibitory mRNA motif in the 5' UTR of the XBP1 transcription factor. Following Ire1 dependent splicing, XBP1 is translated and undergoes nuclear translocation to trigger proteostatic gene expression, including folding chaperones and machinery involved in ER-associated degradation. Dimerized Ire1 can also participate directly in translational control by a process termed RIDD, Regulated Ire1 Dependent Decay, whereby mRNA, miRNA, and even rRNA critical to translation, are cleaved at consensus sites that share homology to XBP1 resulting in their degradation. RIDD has been demonstrated to target and degrade CReP mRNA, a regulatory subunit of eIF2a that contains constitutive phosphatase activity and maintains a low basal level of p-eIF2a [46]. Ire1 deletion nearly doubles the expression of CReP and, as a result, greatly decreases p-eIF2a in response to ER stress. By facilitating the degradation of CReP, RIDD functions to amplify PERK/eIF2a/ATF4 signaling. Despite their similarities, RIDD mediated RNA decay and XBP1 cleavage appear to be distinct and pharmacologically separable mechanisms. Generally, dimerized Ire1 is associated with RIDD and cell death, while oligomerized Ire1 (four or more) is associated with XBP1 splicing and cell survival [47]. Interestingly, the proapoptotic membrane pore-forming proteins Bak and Bax, which localize to the ER membrane in addition to the mitochondria, bind to Ire1 and promote cytoprotective oligomerization. During chronic ER stress, however, Ire1's capacity to bind and sequester Bak/Bax diminishes leading to pore formation and

cytotoxic release of ER contents, an event partially dependent on the expression of the BH3 protein Bnip3 [48]. ER membrane permeabilization, like mitochondrial membrane permeabilization, promotes apoptosis by facilitating the release of cytotoxic molecules such as calcium, ROS, and apoptotic proteins like Calreticulin (CRT). CRT can be repurposed and relocated from the ER, where it functions as a critical folding chaperone, to the plasma membrane, where it serves as an immunogenic "eat me" signal, and has been shown to promote the phagocytosis of live viable neurons by microglia [49, 50]. Although the details are complex, current models suggest Bak/Bax interactions facilitate Ire1 clustering during proteotoxic stress and promotes cytoprotective XBP1 splicing, but prolonged proteotoxic stress, and increased BH3 protein expression, exhausts Ire1's capacity to inhibit Bak/Bax resulting in Ire1 declustering, a shift from XBP1 splicing to RIDD, and apoptotic ER membrane permeability [51]. It is not clear what declusters Ire1, and thus releases Bak/Bax, but one possibility is that XBP1 mediated gene expression upregulates folding chaperones that monomerize Ire1, and possibly facilitates the release of Bak/Bax. ER co-chaperone ERdj4, for example, is upregulated by XBP1 and functions to disaggregate Ire1 clusters by reattaching BiP, which could, in the process of Ire1 monomerization, enhance Bak/Bax oligomerization [52]. Besides its endoribonuclease activity, oligomerized Ire1 serves as a signaling platform that recruits ASK1 and TRAF2, which collectively lead to the recruitment and activation of the stress signaling kinase JNK, discussed more in later sections [53]. In sum, Ire1 dimerization or oligomerization is induced following the loss of BiP to misfolded protein, leading to several distinct processes including RIDD, XBP1 cleavage, Bak/Bax induced pore formation, and JNK activation.

The mtUPR

The ISR is also activated by the mitochondrial unfolded protein response, mtUPR. Despite its name, and the suggestion of a regulated response to misfolded protein, studies examining tissues from patients with bona fide mitochondrial protein misfolding diseases have failed to find patterns of gene expression marked by the induction of mitochondrial chaperones or proteases, as the classic mtUPR program was envisioned [55]. Instead, recent studies into mitochondrial protein misfolding have suggested the phenomenon presents as a broader pattern of gene expression including diverse transcripts such as PPAR γ , promoting mitochondrial biogenesis, genes required for glucose, lipid, and folate driven one-carbon cycle metabolism, oxidative stress-responsive genes, and many others. The inability to clearly define the mtUPR probably results from the interconnectedness of mitochondrial dysfunctions. For example, mitochondrial

defects, such as import inhibition, oxidative phosphorylation (OXPHOS) deficiency, or mitochondrial or nuclear DNA mutations, can all result in misfolded proteins, and thus the mtUPR. Likewise, the accumulation of misfolded proteins in the mitochondria can cause import inhibition, OXPHOS deficiency, and mtDNA mutations. The term mtUPR is thus used to describe any mitonuclear signaling event occurring during mitochondrial dysfunction that functions to restore mitochondrial homeostasis.

The mtUPR triggers a host of cytoprotective genes that, like the ISR, appear to rely primarily on p-eIF2a mediated ATF4 gene expression. A recent high throughput evaluation of the mitochondrial stress response in HeLa cells by Quiros et al, found many conventional mitochondrial toxins, including uncoupling agent FCCP, OXPHOS inhibitors, mitochondrial ribosomal inhibitor doxycycline, or a mitochondrial import inhibitor, each triggered broad cytoprotective gene expression that was largely dependent on p-eIF2a induced ATF4 activation [43]. Interestingly, none of the conventional eIF2a kinases were implicated, suggesting a unique mechanism of ISR activation (discussed below). Analysis of KEGG pathways affected by these toxins found mitochondrial ribosomal biogenesis to be the most significantly down-regulated pathway, while mitochondrial folding chaperones (the classic mtUPR) were neither upregulated or downregulated. In mouse skeletal muscle the mtUPR, caused by a genetic autophagy deficiency that results in the accumulation of dysfunctional mitochondria, upregulates p-eIF2a and ATF4 dependent expression of Fgf21 [56]. Fgf21 is a critical stress induced mitokine that promotes β -oxidation and glucose uptake. Similarly, the metabolic reprogramming of mitochondria towards folate driven one-carbon metabolism in order to promote redox homeostasis is upregulated in patients with mitochondrial diseases [57]. Many of these upregulated genes contain AAREs, Amino Acid Response Elements, in their promoter which are universally recognized by ATF transcription factors, including ATF3, ATF4, and ATF5. In *Drosophila*, the expression of rate-limiting genes in 1C metabolism following mitochondrial toxin treatment is dependent on PERK/eIF2a/ATF4 signaling [58]. The evolutionary conservation of this pathway has recently been demonstrated in mouse models of mitochondrial myopathy, caused by mutations in the mitochondrial helicase Twinkle, which requires ATF4 expression for the upregulation of rate limiting genes in one-carbon metabolism [59]. In a separate study examining mitochondrial stress signaling, authors Michel et al found mtDNA depletion triggers the ISR and, interestingly, discovered this process to be dependent on GCN2 expression [60]. In a *Drosophila* model of mtUPR induced dendritic degeneration, researchers Tsuyama et al discovered eIF2a phosphorylation by PERK was essential in mediating dendritic retraction [61]. Interfering with the ISR

prevented dendritic retraction in response to mitochondrial dysfunction or protein misfolding. This suggests that dendritic degeneration in response to mitochondrial dysfunction, a degenerative but stress adaptive response aimed at preserving ATP, may function through the ISR signaling network. In addition to PERK and GCN2, PKR has also been implicated in the transduction of mitochondrial stress. The mtUPR induced by mutant OTC, a protein that misfolds during mitochondrial import, activated a transcriptional response in intestinal cells that was contingent on PKR mediated eIF2a phosphorylation [62]. The convergence of various mitochondrial stressors onto eIF2a phosphorylation suggest global translational inhibition, and downstream ATF4 transcription, are critical factors orchestrating the mtUPR.

Besides the traditional eIF2a kinases, a recent study by Khan et al provided evidence that mTORC1 can activate ATF4 translation following mitochondrial dysfunction [63]. Despite this seemingly paradoxical relationship (mTORC1 promotes translation while eIF2a/ATF4 inhibits translation), this study provides compelling evidence that mTORC1 can, at least in some circumstances, function upstream of ATF4 activation during mitochondrial stress. Indeed, the Quiros et al study discussed above did find significant upregulation of mTORC signaling components following the mtUPR. The implication of each eIF2a kinase, or none at all, in mediating the mtUPR suggests there are multiple mechanisms by which mitochondrial dysfunction is relayed to the ISR.

In addition to ATF4, the structurally and functionally related ATF5 transcription factor has also been implicated in the mtUPR. Research by Fioerese et al discovered that ATF5, a transcription factor also induced by eIF2a phosphorylation, is essential to the transduction of the mtUPR in *C. elegans* [64]. ATF5 is an intriguing candidate due to its partial homology with ATFS-1, a previously validated mtUPR transcription factor in *C. elegans*. Mammalian ATF5, but not ATF4, can functionally replace ATFS-1 in *C. elegans* and ameliorate mitochondrial dysfunction. In mammalian cells, ATF5 expression can be induced by proteostatic stress derived from proteasome inhibition and mitochondrial depolarization, both of which trigger its expression through eIF2a phosphorylation. Following translation, ATF5 translocates to the nucleus where it upregulates various mitochondrial folding chaperones and ultimately restores mitochondrial proteostasis. Since ATF4 and CHOP have both been confirmed to bind the promoter of ATF5 and activate its transcription, it is plausible that ATF5 functions secondary to ATF4 activation [65].

The mechanistic link between mitochondrial stress and eIF2a phosphorylation, and thus ATF4 and ATF5 translation, has not been identified, but several candidate mechanisms exist. Since the mtUPR is marked by membrane depolarization and import deficiency, it is possible that the

well-characterized PINK1/PARKIN signaling cascade could be involved. In this pathway, mitochondrial membrane depolarization prevents the PINK1 kinase from translocating into the mitochondria and being degraded and instead results in PINK1 accumulating on the outer mitochondrial membrane. PINK1 then phosphorylates and activates both PARKIN1, an E3 ubiquitin ligase, and its preferred species of ubiquitin, resulting in protein phosphoubiquitination and mitophagy. Induction of the mtUPR by supplying cells with mutant OTC does indeed induce mitophagy suggesting PINK1/PARKIN1 is activated in response to the mtUPR [66]. The stress-induced expression of PARKIN1 is also critically reliant on PERK/ATF4 mediated gene expression, but any mechanistic link between these processes has not been identified [67].

The loss of mitochondrial electrochemical potential prevents not just PINK1 from entering the mitochondria, but many other mitochondrial preproteins as well. In *C. elegans*, the cytosolic accumulation of mitochondrial preproteins results from depolarization induced deficits in preprotein import, or actively facilitated by mitochondrial matrix peptide exporter HAF1 that functions during the mtUPR to expel partially degraded peptides [68]. The release of these short mitochondrial peptides are required in some way for ATSF1 mediated gene expression and mtUPR resolution. Recent work in mice by Liu et al discovered mitochondrial import inhibition resulted in the toxic accumulation of metastable mitochondrial preproteins in the cytosol that formed LC3/ubiquitin positive aggregates and, in mouse models, caused neurodegeneration [69]. Perhaps physiological levels of preprotein accumulation during mitochondrial stress could somehow function to activate the ISR, and thus mediate the mtUPR. Mitochondrial preprotein accumulation in yeast does indeed downregulate cytosolic protein translation, in addition to stimulating proteasomal degradation, but a link to eIF2a has yet to be identified [70]. Interestingly, stress-induced mitochondrial import deficiencies have been linked to the activation of PERK, which can trigger the degradation of TIM17A, a critical subunit of the mitochondrial import receptor [71].

Although typically associated with transducing the UPR, PERK also influences mitochondrial dynamics at mitochondrial-associated membranes (MAMs), where it is functionally involved in tethering the two membranes, independent of its kinase activity [72]. Owing to the extensive communication that occurs at these MAMs, it is possible that the mtUPR could be launched through MAMs, thus linking PERK to the mtUPR. Since mitochondrial stress does indeed activate the UPR, it would be intriguing to explore whether MAMs are essential in mediating this process, and perhaps yield important insight into how the mtUPR triggers the ISR [73]. Another ISR kinase, GCN2, has been shown to be activated by ROS signaling in addition to uncharged tRNA

[74]. Although the exact mechanism is not understood, ROS production during mitochondrial dysfunction can facilitate eIF2a phosphorylation through GCN2.

Lastly, mtRNA has recently been shown to occupy and activate a large portion of endogenous PKR, which could be released as a result of mitochondrial rupture or facilitated export [75]. Supporting this notion, recent research has revealed that mitochondrial stress leads to the cytosolic accumulation of mtDNA [76, 76, 78]. Interestingly, PINK1/PARKIN1 mediated mitophagy in mice is required to suppress mtDNA release from stressed mitochondria, a phenomenon that subsequently leads to neurodegeneration through a cGAS/STING dependent proinflammatory response. Thus deficiencies in mitophagy may lead to the deleterious rupture of mitochondria, releasing mtDNA/mtRNA and potentially activating PKR and p-eIF2a. It is important to emphasize that none of these mechanisms are mutually exclusive, and it is highly likely that mitochondria use multiple signaling routes to activate the ISR. Ultimately there are dozens of pathways activated by mitochondrial stress, and any number of them could potentially activate the ISR and stall translation, but that discussion exceeds the scope of this review.

mTORC1

Loss of proteostasis can influence translation and metabolism through the modulation of mTORC1. mTORC1 is a serine/threonine kinase composed of three primary subunits; RAPTOR, critical for substrate recognition, mLST8, required for mTORC1's active site formation, and mTOR, the functional kinase. mTORC1's regulation is complex, and reviewed elsewhere, but can generally be summarized as a kinase activated by growth signaling and nutrient availability, and inactivated by nutrient deprivation and stress [79]. When active mTORC1 controls translation initiation by phosphorylating and inactivating 4E-BP, a protein that, when relieved from mTORC1's negative regulation, potently inhibits eIF4e of the translation initiation complex. mTORC1 upregulates translation elongation as well by phosphorylating and activating S6K1, a kinase that increases the processivity of the ribosome. mTORC1 also negatively regulates eIF2a phosphorylation by phosphorylating eIF2B resulting in the recruitment of NCK1, which mediates eIF2a dephosphorylation (possibly by recruiting PP1A) [80]. In addition, mTORC1 has been shown to enhance the rate of rRNA biogenesis by phosphorylating the critical RNA polymerase I transcription factor TIF-1A, and the inhibition of mTORC1 disrupts ribosomal assembly [81, 82]. mTORC1 thus serves to initiate translation through its inhibition of 4E-BP and p-eIF2a, upregulate the rate of translation elongation through S6K1 activation, and promote ribosome biogenesis by TIF-1a activation. Besides its modulation of translation, mTORC1 phosphorylates and

inactivates ULK1, the central kinase of the autophagic preinitiation complex. Autophagy is additionally downregulated at the transcriptional level by mTORC1 mediated phosphorylation of TFEB, a critical autophagic and lysosomal transcription factor, resulting in its negative regulation by binding inhibitory 14-3-3 class of phosphoserine-binding proteins [83]. Thus in order to restore proteostasis, it seems practical that mTORC1 would be inactivated during protein misfolding in order to simultaneously activate autophagy and inactivate translation, and indeed proteostatic disturbances consistently result in mTORC1 inhibition.

Considering mTORC1's pivotal role in modulating metabolic and degradative pathways, determining how proteostatic disruption affects mTORC1 signaling has become a central question. Generally, events that disrupt proteostasis are accompanied by the attenuation of mTORC1 signaling. For example, acute inhibition of the proteasome, a traditional method of disrupting proteostasis, rapidly deactivates mTORC1, even in the presence of translation inhibitors suggesting new protein synthesis is not required. Since certain species of misfolded protein have been shown to either inhibit the proteasome, or sequester it in aggregates, understanding how this deactivates mTORC1, and thus translation may provide critical insight into neurodegenerative processes. The loss of proteasomal function could result in the accumulation of DEPTOR, a protein that potently inhibits mTORC1 but is normally ubiquitinated by SCF^{BTrCP} ubiquitin ligases and degraded in proteasomes [84, 85]. mTORC1 activity may also be limited by its rate of formation, a process that is critically reliant on the HSP90 chaperone. The loss of HSP90 due to the presence of misfolded proteins, which can function as chaperone sinks, has been shown to limit the rate at which mTORC1 is constructed resulting in decreased mTORC1 signaling [86]. The activation of the UPR is also accompanied with decreased mTORC1 signaling, and thus translation, an effect that occurs independently of eIF2a phosphorylation [87]. mTORC1 inhibition accounts for the downregulation of translation in eIF2a knockout cells following ER stress (although this effect takes hours, as opposed to minutes in WT cells) suggesting the UPR can mediate mTORC inhibition independent of the ISR. Interestingly, mTORC1 mediated translation following UPR stress could be restored by transducing cells with ATF4, suggesting ATF4 transcription positively regulates mTORC1 (supported by Quiros et al above). In contrast, another study found ATF4 mediated transcription sustained mTORC1 inhibition by upregulating Sestrin [88]. Whether ATF4 transcriptional activity promotes or inhibits mTORC1 is not clear, but regardless there appears to important transcriptional crosstalk between these pathways.

How does UPR signaling result in mTORC1 suppression independent of eIF2a phosphorylation? Loss of

mTORC signaling during prolonged ER stress is associated with the loss of Akt signaling, an upstream activator of mTORC1, and activation of AMPK [89]. AMPK leads to the inactivation of mTORC1 by phosphorylation of its substrate recognition subunit RAPTOR, causing dissociation and binding to 14-3-3. Exactly how chronic ER stress results in AMPK activation and Akt inactivation is not clear. Alternatively, as discussed above, protein misfolding in the ER can lead to the activation of JNK, a stress signaling kinase that both associates with and phosphorylates mTORC1 (and also negatively regulates Akt). Recent research by Su et al found JNK mediated phosphorylation of mTORC1's RAPTOR domain induces the disassembly of the mTORC1 complex [90]. However, during normal proteostatic conditions JNK is bound and sequestered away from mTORC1 by HSF1. The induction of misfolded proteins through heat shock, proteasome inhibition, or HSP70 inhibition, displaces HSF1 from JNK resulting in the inhibitory phosphorylation of Raptor and mTORC1 disassembly. Knockdown of JNK prevented proteasome inhibition from resulting in mTORC1 inhibition, suggesting JNK is the essential missing link between proteasome inhibition and mTORC1 signaling. Additionally, HSF1 deletion in mice, which would, in theory, lead to constitutively active JNK and inhibited mTORC signaling, does indeed disrupt mTORC activity [91]. The potential contribution of Ire1/TRAFF/JNK in mediating this pathway has not been studied, but regardless, these experiments help link proteasome inhibition, HSF1 activation, and JNK signaling, to mTORC1 inactivation and translational control.

Mitochondrial insults, including mitochondrial translation inhibition and ETC disruption, have also been shown to deactivate mTORC1, thus promoting autophagy, stalling translation, and preserving ATP [92, 93]. Although it is clear mTORC1 is inactivated by mitochondrial stress, the mechanistic details bridging these two phenomena are poorly understood. It is reasonable that the loss of ATP production, and thus AMP accumulation, could activate AMPK and mediate mTORC1 inhibition. It was also reported that mitochondrial stress induced depolarization resulted in the activation of PARKIN1, discussed above, that polyubiquitinates mTORC1 triggering its proteasomal degradation [94]. mTORC1 is indeed associated with mitochondrial outer membranes, cofractionates with mitochondria, and is typically inactive while localized to mitochondria (mTORC1 requires lysosomal localization for activation) [95, 96]. Also associated with the mitochondrial outer membrane is JNK, which translocates to mitochondria during stress and is anchored by the protein Sab [97, 98]. It would be intriguing to test whether Sab mediated recruitment of JNK to the mitochondria facilitates mTORC1 phosphorylation and degradation. In sum, mTORC is inhibited by multiple proteostatic disturbances including the UPR,

proteasome inhibition, chaperone depletion, and the mtUPR, although more research is needed to determine the underlying signaling networks.

STRESS GRANULES

Stress granules are membraneless ~200 nm pseudo organelles that form in the cytoplasm during proteostatic or ribostatic stress, and function to temporarily sequester mRNA until translation resumes. Stress granules are dynamic structures that generally contain dense immobile cores and gel-like outer shells, the latter of which turns over on the order of seconds as measured by FRAP studies [99]. Although the majority of a cell's pool of mRNAs (about 90%) are targeted to stress granules during stress, only about 10% of mRNAs are consistently nucleated into aggregates (perhaps the population of mRNA locked in insoluble cores) [100]. The rest appear to only transiently associate with stress granules before being released, or funneled into P bodies for degradation. Only 185 genes have greater than 50% of their mRNA localized within stress granules, suggesting stress granules serve a more specialized function than previously thought, rather than broad sequestration of mRNA. These mRNAs prone to aggregation in stress granules appear to be related by extended 3' UTRs and larger than average transcript size, leading to poor translatability. These criteria would encompass all viral RNAs and result in their preferential accumulation in SGs, which is likely the evolutionary purpose of stress granules [101]. Indeed, many viruses employ transcripts specifically evolved to inhibit stress granule nucleating machinery. Larger proteins are also more prone to misfolding or causing translational errors, thus their preferential targeting to stress granules may aid in proteostatic recovery. There is also evidence that certain classes of mRNA, such as the HSP folding chaperones, are preferentially excluded from stress granules resulting in their upregulation during stress. The inability to sequester mRNA in stress granules during proteotoxic stress can accelerate neuronal aggregate formation and rate of cell death [102, 103, 104]. The loss of stress granule nucleation machinery alone is sufficient to induce neurodegeneration, particularly in the hippocampus [105, 106]. Similarly, the irreversible aggregation of stress granules also constitutes a proteostatic dysfunction that can promote protein misfolding and cell death [107]. Thus a careful balance of stress granule formation and dissolution in response to translation status is an important component of the proteostatic network.

Stress granules are a proteostatic response to translation inhibition derived from two primary physiological sources; mTORC inhibition and eIF2a phosphorylation. Since both of these events are fundamental to the proteostatic response, stress granules are nearly ubiquitous across protein

misfolding disorders. The RNA binding proteins responsible for nucleating stress granules, including most notably TIA1 and G3BP1, bind through prion-related protein interactions between intrinsically disordered domains promoting their aggregation [108, 109]. During healthy proteostatic conditions, continuous ATP dependent chaperone activity and the absence of ribosome-free mRNA precludes the formation of stress granules [110, 111]. It is thus the non-association of mRNA with ribosomes due to p-eIF2a or mTORC inhibition, and loss of enzymatically active chaperones like HSP70 (perhaps due to the presence of misfolded protein or ATP depletion), that induces the aggregation of mRNA into stress granules. During these conditions, RNA binding protein TIA1 is recruited to 3' TOP sequences of mRNA (Terminal Oligopyrimidine Tracts), and G3BP to stagnated 40s small ribosomal subunits, thus increasing their local concentration, promoting interactions between IDRs, and facilitating aggregation [112, 113].

Although stress granules nucleate around mRNA, certain proteins also become sequestered in stress granules, a regulated process that usually depends on the presence of an IDR (for interaction with TIA1/G3BP) or an RNA binding domain (for interaction with aggregated mRNA). For example, a cell's population of Ubiquilin-2 proteins, a protein linked to ALS, are nearly completely shuttled into SGs owing to an IDR [114]. Interestingly, the ubiquilin family of proteins are responsible for binding transmembrane mitochondrial outer membrane proteins that fail to be inserted into the mitochondrial membrane and delivers them to the proteasome [115]. Since the loss of ubiquilins results in the cytosolic accumulation and aggregation of mitochondrial outer membrane preproteins, it may be possible that stress granule formation, and sequestration of ubiquilin proteins, promotes the mtUPR. In addition to Ubiquilin-2, critical RNA binding proteins required for nucleocytoplasmic shuttling, splicing, and snRNA biogenesis, are also sequestered in stress granules [116]. By sequestering a host of RNA shuttling proteins, including Ran, Exportins, and Importins, stress granules significantly downregulate the flux of new mRNA into the cytoplasm, serving as an additional mechanism of proteostatic control of translation [106]. Similarly, the sequestration of TDP-43 in stress granules, a major ALS associated protein normally localized to Cajal bodies, could result in abnormally spliced transcripts and RNA processing [117]. Collectively, the aggregation of RNA binding proteins in stress granules could collapse protein assembly lines, which, like other inhibitors of translation, can be beneficial or detrimental depending on duration.


The importance of stress granules in attenuating translation has been challenged by recent research demonstrating that G3BP knockout cells, which cannot form stress granules following eIF2a phosphorylation, are not


defective in translational arrest [113]. Additionally, the majority of mRNA are not sequestered in stress granules, discovered following the high throughput analysis of stress granule transcriptomes [100]. Since stress granules are nucleated principally by eIF2a phosphorylation, they may simply be associated with stalled translation, and not causative. If not critical to translational attenuation, why are stress granules central to maintaining proteostasis? By nucleating mRNA, ribosomes, and translation initiation factors, stress granules may, paradoxically, be critical for the rapid reinitiation of translation by colocalizing translational machinery [118]. Perhaps the sequestration of mRNA is simply a consequence of colocalizing RNA binding proteins and ribosomes. Another possibility is that stress granules shield latent mRNA from ROS induced mutations thereby helping to mitigate subsequent translation errors. And lastly, stress granules may influence post-transcriptional modification dynamics. Storing mRNA in dense stress granules could serve as a mechanism for preventing potentially deleterious or otherwise unintended mRNA modifications. Regardless, the association of translational machinery, RNA binding proteins, and mRNA with stress granules strongly suggests they serve an important function in translation, although it may not be as straightforward as simply decreasing protein synthesis.

Once formed, a cell must efficiently disaggregate stress granules in order to resume proper translation, RNA processing, and recover from stress. The majority of stress granules in HeLa cells are cleared by the HSPB8/BAG3/HSP70 chaperone complex, although many other chaperones are also involved [111]. The efficient clearance of stress granules seems to require the recruitment of the 26S proteasome, a process conducted by ZFAND1/p97 [119]. As an ATPase segregase, p97 (also known as VCP) has been repeatedly implicated in the clearance of both stress granules and aggregates, and various mutations within p97 have been linked to ALS and FTD. In addition, p97's ATPase activity can remodel stress granules so as to expose ubiquitin moieties required for autophagic clearance (in yeast), although the importance of autophagic clearance of SGs in mammalian cells is disputed [120]. Regardless, since ZFAND1 is responsible for recruiting both p97 and the proteasome, it likely functions upstream of both clearance mechanisms. The current model of stress granule clearance involves the disaggregation of RNA binding proteins by chaperones like HSP70 and p97 (and many others) and degradation in the proteasome. In sum, stress granules aid proteostatic networks by sequestering mRNA, and proteins critical to mRNA splicing and nucleocytoplasmic shuttling, thereby downregulating protein synthesis.

Neurons have evolved complex mechanisms of translational control in order to manage the proteostatic and energetic challenges of neuronal function. However, in order to more completely integrate this mechanism with the neurodegenerative process, a better understanding of how translation is affected by ageing is needed. Ageing is the most significant risk factor for neurodegenerative disease, and any attempt to understand proteostasis must be examined through the lens of ageing. Old age is not only associated with aberrant protein accumulation, but also the accumulation of undegradable "biochemical garbage", such as lipofuscin in lysosomes, advanced glycation end products, or extracellular atherosclerotic cholesterol plaques. Why does cellular garbage accumulate as we age? The age-dependent accumulation of genetic mutations, known as the Somatic Mutation Theory of Ageing, could steadily escalate proteostatic pressures as protein function deteriorates and they become increasingly prone to misfold. Indeed, DNA damage syndromes have been linked to protein misfolding and proteostatic collapse [121]. Alternatively, a large body of evidence suggests the presence of various environmental toxins that accelerate aging. These "gerontogens" are diverse, cumulative in their effect, and are often related to the induction of DNA damage (a carcinogen). The age-dependent loss of immune function has also been linked to the reactivation of dormant viruses, such as HSV6/7 in the brain of Alzheimer's patients [122]. Many of the proteostatic pathways discussed in this paper have evolutionary roots in the cellular response to viral RNA and protein, and so the possibility that these proteostatic disturbances are caused by viruses is not implausible. Future research will benefit significantly from examining interactions between these mechanisms of ageing and translational control, and could yield critical insight into the neurodegenerative process.

CONCLUSION





1. Eom, T., Muslimov, I. A., Tsokas, P., Berardi, V., Zhong, J., Sacktor, T. C., & Tiedge, H. (2014). Neuronal BC RNAs cooperate with eIF4B to mediate activity-dependent translational control. *The Journal of Cell Biology*, 207(2), 237-252. doi:10.1083/jcb.201401005
2. Tain, L. S., Mortiboys, H., Tao, R. N., Ziviani, E., Bandmann, O., & Whitworth, A. J. (2009). Rapamycin activation of 4E-BP prevents parkinsonian dopaminergic neuron loss. *Nature Neuroscience*, 12(9), 1129-1135. doi:10.1038/nn.2372
3. D'Antonio, M., Musner, N., Scapin, C., Ungaro, D., Carro, U. D., Ron, D., . . . Wrabetz, L. (2013). Resetting translational homeostasis restores myelination in Charcot-Marie-Tooth disease type 1B mice. *The Journal of Cell Biology*, 201(2), 13-13. doi:10.1083/jcb20120ia3
4. Imai, Y., Gehrke, S., Wang, H., Takahashi, R., Hasegawa, K., Oota, E., & Lu, B. (2008). Phosphorylation of 4E-BP by LRRK2 affects the maintenance of dopaminergic neurons in *Drosophila*. *The EMBO Journal*, 27(18), 2432-2443. doi:10.1038/emboj.2008.163
5. Banko, J. L. (2005). The Translation Repressor 4E-BP2 Is Critical for eIF4F Complex Formation, Synaptic Plasticity, and Memory in the Hippocampus. *Journal of Neuroscience*, 25(42), 9581-9590. doi:10.1523/jneurosci.2423-05.2005
6. Tsaytler, P., Harding, H. P., Ron, D., & Bertolotti, A. (2011). Selective Inhibition of a Regulatory Subunit of Protein Phosphatase 1 Restores Proteostasis. *Science*, 332(6025), 91-94. doi:10.1126/science.1201396
7. Vellai T, Takacs-Vellai K, Zhang Y, Kovacs AL, Orosz L, Muller F. 2003. Genetics: influence of TOR kinase on lifespan in *C. elegans*. *Nature* 426: 620
8. Kapahi P, Zid BM, Harper T, Koslover D, Sapin V, Benzer S. 2004. Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr Biol* 14: 885-890
9. Hansen M, Taubert S, Crawford D, Libina N, Lee SJ, Kenyon C. 2007. Lifespan extension by conditions that inhibit translation in *Caenorhabditis elegans*. *Aging Cell* 6:95-110.
10. Selman C, Tullet JM, Wieser D, Irvine E, Lingard SJ, Choudhury AI, Claret M, Al-Qassab H, Carmignac D, Ramadani F, et al. 2009. Ribosomal protein S6 kinase 1 signaling regulates mammalian life span. *Science* 326: 140-144.
11. Liu, S., & Lu, B. (2010). Reduction of Protein Translation and Activation of Autophagy Protect against PINK1 Pathogenesis in *Drosophila melanogaster*. *PLoS Genetics*, 6(12). doi:10.1371/journal.pgen.1001237
12. Wang, X., Zuo, X., Kucejova, B., & Chen, X. J. (2008). Reduced cytosolic protein synthesis suppresses mitochondrial degeneration. *Nature Cell Biology*, 10(9), 1090-1097. doi:10.1038/ncb1769
13. Sherman, M. Y., & Qian, S. (2013). Less is more: Improving proteostasis by translation slow down. *Trends in Biochemical Sciences*, 38(12), 585-591. doi:10.1016/j.tibs.2013.09.003
14. Morimoto, R. I. (2008). Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes & Development*, 22(11), 1427-1438. doi:10.1101/gad.1657108
15. Zheng, X., Boyer, L., Jin, M., Kim, Y., Fan, W., Bardy, C., . . . Hunter, T. (2016). Alleviation of neuronal energy deficiency by mTOR inhibition as a treatment for mitochondria-related neurodegeneration. *ELife*, 5. doi:10.7554/elife.13378
16. Stouthamer, A. H. (1973). A theoretical study on the amount of ATP required for synthesis of microbial cell material. *Antonie Van Leeuwenhoek*, 39(1), 545-565. doi:10.1007/bf02578899
17. Lever, M. A., Rogers, K. L., Lloyd, K. G., Overmann, J., Schink, B., Thauer, R. K., . . . Jørgensen, B. B. (2015). Life under extreme energy limitation: A synthesis of laboratory- and field-based investigations. *FEMS Microbiology Reviews*, 39(5), 688-728. doi:10.1093/femsre/fuv020
18. E.B. Kramer, P.J. Farabaugh. (2007). The frequency of translational misreading errors in *E. coli* is largely determined by tRNA competition. *RNA*, 13, pp. 87-96
19. M. Ogle, V. Ramakrishnan. (2005) Structural insights into translational fidelity. *Annu. Rev. Biochem.*, 74, pp. 129-177
20. H.H. Guo, J. Choe, L.A. Loeb (2004). Protein tolerance to random amino acid change. *Proc. Natl. Acad. Sci. USA*, 101, pp. 9205-9210
21. Wang, F., Durfee, L., & Huijbregtse, J. (2013). A Cotranslational Ubiquitination Pathway for Quality Control of Misfolded Proteins. *Molecular Cell*, 50(3), 368-378. doi:10.1016/j.molcel.2013.03.009

REFERENCES

1. Eom, T., Muslimov, I. A., Tsokas, P., Berardi, V., Zhong, J., Sacktor, T. C., & Tiedge, H. (2014). Neuronal BC RNAs cooperate with eIF4B to mediate activity-dependent translational control. *The Journal of Cell Biology*, 207(2), 237-252. doi:10.1083/jcb.201401005
2. Tain, L. S., Mortiboys, H., Tao, R. N., Ziviani, E., Bandmann, O., & Whitworth, A. J. (2009). Rapamycin activation of 4E-BP prevents parkinsonian dopaminergic neuron loss. *Nature Neuroscience*, 12(9), 1129-1135. doi:10.1038/nn.2372
3. D'Antonio, M., Musner, N., Scapin, C., Ungaro, D., Carro, U. D., Ron, D., . . . Wrabetz, L. (2013). Resetting translational homeostasis restores myelination in Charcot-Marie-Tooth disease type 1B mice. *The Journal of Cell Biology*, 201(2), 13-13. doi:10.1083/jcb20120ia3
4. Imai, Y., Gehrke, S., Wang, H., Takahashi, R., Hasegawa, K., Oota, E., & Lu, B. (2008). Phosphorylation of 4E-BP by LRRK2 affects the maintenance of dopaminergic neurons in *Drosophila*. *The EMBO Journal*, 27(18), 2432-2443. doi:10.1038/emboj.2008.163
5. Banko, J. L. (2005). The Translation Repressor 4E-BP2 Is Critical for eIF4F Complex Formation, Synaptic Plasticity, and Memory in the Hippocampus. *Journal of Neuroscience*, 25(42), 9581-9590. doi:10.1523/jneurosci.2423-05.2005
6. Tsaytler, P., Harding, H. P., Ron, D., & Bertolotti, A. (2011). Selective Inhibition of a Regulatory Subunit of Protein Phosphatase 1 Restores Proteostasis. *Science*, 332(6025), 91-94. doi:10.1126/science.1201396
7. Vellai T, Takacs-Vellai K, Zhang Y, Kovacs AL, Orosz L, Muller F. 2003. Genetics: influence of TOR kinase on lifespan in *C. elegans*. *Nature* 426: 620
8. Kapahi P, Zid BM, Harper T, Koslover D, Sapin V, Benzer S. 2004. Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr Biol* 14: 885-890
9. Hansen M, Taubert S, Crawford D, Libina N, Lee SJ, Kenyon C. 2007. Lifespan extension by conditions that inhibit translation in *Caenorhabditis elegans*. *Aging Cell* 6:95-110.
10. Selman C, Tullet JM, Wieser D, Irvine E, Lingard SJ, Choudhury AI, Claret M, Al-Qassab H, Carmignac D, Ramadani F, et al. 2009. Ribosomal protein S6 kinase 1 signaling regulates mammalian life span. *Science* 326: 140-144.
11. Liu, S., & Lu, B. (2010). Reduction of Protein Translation and Activation of Autophagy Protect against PINK1 Pathogenesis in *Drosophila melanogaster*. *PLoS Genetics*, 6(12). doi:10.1371/journal.pgen.1001237
12. Wang, X., Zuo, X., Kucejova, B., & Chen, X. J. (2008). Reduced cytosolic protein synthesis suppresses mitochondrial degeneration. *Nature Cell Biology*, 10(9), 1090-1097. doi:10.1038/ncb1769
13. Sherman, M. Y., & Qian, S. (2013). Less is more: Improving proteostasis by translation slow down. *Trends in Biochemical Sciences*, 38(12), 585-591. doi:10.1016/j.tibs.2013.09.003
14. Morimoto, R. I. (2008). Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes & Development*, 22(11), 1427-1438. doi:10.1101/gad.1657108
15. Zheng, X., Boyer, L., Jin, M., Kim, Y., Fan, W., Bardy, C., . . . Hunter, T. (2016). Alleviation of neuronal energy deficiency by mTOR inhibition as a treatment for mitochondria-related neurodegeneration. *ELife*, 5. doi:10.7554/elife.13378
16. Stouthamer, A. H. (1973). A theoretical study on the amount of ATP required for synthesis of microbial cell material. *Antonie Van Leeuwenhoek*, 39(1), 545-565. doi:10.1007/bf02578899
17. Lever, M. A., Rogers, K. L., Lloyd, K. G., Overmann, J., Schink, B., Thauer, R. K., . . . Jørgensen, B. B. (2015). Life under extreme energy limitation: A synthesis of laboratory- and field-based investigations. *FEMS Microbiology Reviews*, 39(5), 688-728. doi:10.1093/femsre/fuv020
18. E.B. Kramer, P.J. Farabaugh. (2007). The frequency of translational misreading errors in *E. coli* is largely determined by tRNA competition. *RNA*, 13, pp. 87-96
19. M. Ogle, V. Ramakrishnan. (2005) Structural insights into translational fidelity. *Annu. Rev. Biochem.*, 74, pp. 129-177
20. H.H. Guo, J. Choe, L.A. Loeb (2004). Protein tolerance to random amino acid change. *Proc. Natl. Acad. Sci. USA*, 101, pp. 9205-9210
21. Wang, F., Durfee, L., & Huijbregtse, J. (2013). A Cotranslational Ubiquitination Pathway for Quality Control of Misfolded Proteins. *Molecular Cell*, 50(3), 368-378. doi:10.1016/j.molcel.2013.03.009

22. Radford, H., Moreno, J. A., Verity, N., Halliday, M., & Mallucci, G. R. (2015). PERK inhibition prevents tau-mediated neurodegeneration in a mouse model of frontotemporal dementia. *Acta Neuropathologica*, 130(5), 633-642. doi:10.1007/s00401-015-1487-z

23. Halliday, M., Radford, H., Sekine, Y., Moreno, J., Verity, N., Quesne, J. L., . . . Mallucci, G. R. (2015). Partial restoration of protein synthesis rates by the small molecule ISRib prevents neurodegeneration without pancreatic toxicity. *Cell Death and Disease*, 6(3). doi:10.1038/cddis.2015.49

24. Scheper, G. C., Knaap, M. S., & Proud, C. G. (2007). Translation matters: Protein synthesis defects in inherited disease. *Nature Reviews Genetics*, 8(9), 711-723. doi:10.1038/nrg2142

25. Caviston, J. P., Ross, J. L., Antony, S. M., Tokito, M., & Holzbaur, E. L. (2007). Huntingtin facilitates dynein/dynactin-mediated vesicle transport. *Proceedings of the National Academy of Sciences*, 104(24), 10045-10050. doi:10.1073/pnas.0610628104

26. Lu, X., Mattis, V. B., Wang, N., Al-Ramahi, I., Berg, N. V., Fratantoni, S. A., . . . Yang, X. W. (2014). Targeting ATM ameliorates mutant Huntingtin toxicity in cell and animal models of Huntington's disease. *Science Translational Medicine*, 6(268). doi:10.1126/scitranslmed.3010523

27. Kaushik, S., & Cuervo, A. M. (2015). Proteostasis and aging. *Nature Medicine*, 21(12), 1406-1415. doi:10.1038/nm.4001

28. Galehdar, Z., Swan, P., Fuerth, B., Callaghan, S. M., Park, D. S., & Cregan, S. P. (2010). Neuronal Apoptosis Induced by Endoplasmic Reticulum Stress Is Regulated by ATF4-CHOP-Mediated Induction of the Bcl-2 Homology 3-Only Member PUMA. *Journal of Neuroscience*, 30(50), 16938-16948. doi:10.1523/jneurosci.1598-10.2010

29. Mekahli, D., Bultynck, G., Parys, J. B., Smedt, H. D., & Missiaen, L. (2011). Endoplasmic-Reticulum Calcium Depletion and Disease. *Cold Spring Harbor Perspectives in Biology*, 3(6). doi:10.1101/cshperspect.a004317

30. Reid, D. W., Tay, A. S., Sundaram, J. R., Lee, I. C., Chen, Q., George, S. E., . . . Shenolikar, S. (2016). Complementary Roles of GADD34- and CReP-Containing Eukaryotic Initiation Factor 2 α Phosphatases during the Unfolded Protein Response. *Molecular and Cellular Biology*, 36(13), 1868-1880. doi:10.1128/mcb.00190-16

31. Marciniak, S. J. (2004). CHOP induces death by promoting protein synthesis and oxidation in the stressed endoplasmic reticulum. *Genes & Development*, 18(24), 3066-3077. doi:10.1101/gad.1250704

32. Han, J., Back, S. H., Hur, J., Lin, Y., Gildersleeve, R., Shan, J., . . . Kaufman, R. J. (2013). ER-stress-induced transcriptional regulation increases protein synthesis leading to cell death. *Nature Cell Biology*, 15(5), 481-490. doi:10.1038/ncb2738

33. Cullinan, S. B., & Diehl, J. A. (2004). PERK-dependent Activation of Nrf2 Contributes to Redox Homeostasis and Cell Survival following Endoplasmic Reticulum Stress. *Journal of Biological Chemistry*, 279(19), 20108-20117. doi:10.1074/jbc.m314219200

34. Xie, W., Pariolla, M., Wixted, W. E., Chitnis, N., Fornwald, J., Truong, M., . . . Li, H. (2015). Identification and Characterization of PERK Activators by Phenotypic Screening and Their Effects on NRF2 Activation. *Plos One*, 10(3). doi:10.1371/journal.pone.0119738

35. Pajares, M., Rojo, A. I., Arias, E., Díaz-Carretero, A., Cuervo, A. M., & Cuadrado, A. (2018). Transcription factor NFE2L2/NRF2 modulates chaperone-mediated autophagy through the regulation of LAMP2A. *Autophagy*, 14(8), 1310-1322. doi:10.1080/15548627.2018.1474992

36. Jain, A., Rusten, T. E., Katheder, N., Elvenes, J., Bruun, J., Sjøttem, E., . . . Johansen, T. (2015). P62/Sequestosome-1, Autophagy-related Gene 8, and Autophagy in Drosophila Are Regulated by Nuclear Factor Erythroid 2-related Factor 2 (NRF2), Independent of Transcription Factor TFEB. *Journal of Biological Chemistry*, 290(24), 14945-14962. doi:10.1074/jbc.m115.656116

37. Jiang, T., Harder, B., Vega, M. R., Wong, P. K., Chapman, E., & Zhang, D. D. (2015). P62 links autophagy and Nrf2 signaling. *Free Radical Biology and Medicine*, 88, 199-204. doi:10.1016/j.freeradbiomed.2015.06.014

38. Bretin, A., Carrière, J., Dalmasso, G., Bergougnoux, A., Bchir, W., Maurin, A., . . . Nguyen, H. T. (2016). Activation of the EIF2AK4-EIF2A/eIF2 α -ATF4 pathway triggers autophagy response to Crohn disease-associated adherent-invasive Escherichia coli infection. *Autophagy*, 12(5), 770-783. doi:10.1080/15548627.2016.1156823

39. Deegan, S., Koryga, I., Glynn, S. A., Gupta, S., Gorman, A. M., & Samali, A. (2015). A close connection between the PERK and IRE arms of the UPR and the transcriptional regulation of autophagy. *Biochemical and Biophysical Research Communications*, 456(1), 305-311. doi:10.1016/j.bbrc.2014.11.076

40. B'Chir, W., Maurin, A., Carraro, V., Averous, J., Jousse, C., Muranishi, Y., . . . Bruhat, A. (2013). The eIF2 α /ATF4 pathway is essential for stress-induced autophagy gene expression. *Nucleic Acids Research*, 41(16), 7683-7699. doi:10.1093/nar/gkt563

41. Kouraku, Y., Fujita, E., Tanida, I., Ueno, T., Isoai, A., Kumagai, H., . . . Momoi, T. (2006). ER stress (PERK/eIF2 α phosphorylation) mediates the polyglutamine-induced LC3 conversion, an essential step for autophagy formation. *Cell Death and Differentiation*, 14(2), 230-239. doi:10.1038/sj.cdd.4401984

42. Gretzmeier, C., Eiselein, S., Johnson, G. R., Engelke, R., Nowag, H., Zarei, M., . . . Dengjel, J. (2017). Degradation of protein translation machinery by amino acid starvation-induced macroautophagy. *Autophagy*, 13(6), 1064-1075. doi:10.1080/15548627.2016.1274485

43. Quirós, P. M., Prado, M. A., Zamboni, N., D'Amico, D., Williams, R. W., Finley, D., . . . Auwerx, J. (2017). Multi-omics analysis identifies ATF4 as a key regulator of the mitochondrial stress response in mammals. *The Journal of Cell Biology*, 216(7), 2027-2045. doi:10.1083/jcb.201702058

44. Zong, Z., Du, Z., Li, N., Li, C., Zhang, Q., Liu, B., . . . Wang, H. (2012). Implication of Nrf2 and ATF4 in differential induction of CHOP by proteasome inhibition in thyroid cancer cells. *Biochimica Et Biophysica Acta (BBA) - Molecular Cell Research*, 1823(8), 1395-1404. doi:10.1016/j.bbamcr.2012.06.001

45. He, C. H., Gong, P., Hu, B., Stewart, D., Choi, M. E., Choi, A. M., & Alam, J. (2001). Identification of Activating Transcription Factor 4 (ATF4) as an Nrf2-interacting Protein. *Journal of Biological Chemistry*, 276(24), 20858-20865. doi:10.1074/jbc.m101198200

46. So, J., Cho, S., Min, S., Kimball, S. R., & Lee, A. (2015). IRE1 α -Dependent Decay of CReP/Ppp1r15b mRNA Increases Eukaryotic Initiation Factor 2 α Phosphorylation and Suppresses Protein Synthesis. *Molecular and Cellular Biology*, 35(16), 2761-2770. doi:10.1128/mcb.00215-15

47. Tam, A., Koong, A., & Niwa, M. (2014). Ire1 Has Distinct Catalytic Mechanisms for XBP1/HAC1 Splicing and RIDD. *Cell Reports*, 9(3), 850-858. doi:10.1016/j.celrep.2014.09.016

48. Kanekura, K., Ma, X., Murphy, J. T., Zhu, L. J., Diwan, A., & Urano, F. (2015). IRE1 prevents endoplasmic reticulum membrane permeabilization and cell death under pathological conditions. *Science Signaling*, 8(382). doi:10.1126/scisignal.aaa0341

49. Fricker, M., Oliva-Martín, M. J., & Brown, G. C. (2012). Primary phagocytosis of viable neurons by microglia activated with LPS or A β is dependent on calreticulin/LRP phagocytic signalling. *Journal of Neuroinflammation*, 9(1). doi:10.1186/1742-2094-9-196

50. Panaretakis, T., Kepp, O., Brockmeier, U., Tesniere, A., Bjorklund, A., Chapman, D. C., . . . Kroemer, G. (2009). Mechanisms of pre-

apoptotic calreticulin exposure in immunogenic cell death. *The EMBO Journal*, 28(5), 578-590. doi:10.1038/emboj.2009.1

51. Wochlbier, U., & Hetz, C. (2011). Modulating stress responses by the UPRosome: A matter of life and death. *Trends in Biochemical Sciences*, 36(6), 329-337. doi:10.1016/j.tibs.2011.03.001

52. Amin-Wetzel, N., Saunders, R. A., Kamphuis, M. J., Rato, C., Preissler, S., Harding, H. P., & Ron, D. (2017). A J-Protein Co-chaperone Recruits BiP to Monomerize IRE1 and Repress the Unfolded Protein Response. *Cell*, 171(7). doi:10.1016/j.cell.2017.10.040

53. Zhu, X., Zhang, J., Sun, H., Jiang, C., Dong, Y., Shan, Q., . . . Liu, S. (2014). Ubiquitination of Inositol-requiring Enzyme 1 (IRE1) by the E3 Ligase CHIP Mediates the IRE1/TRAF2/JNK Pathway. *Journal of Biological Chemistry*, 289(44), 30567-30577. doi:10.1074/jbc.m114.562868

54. Verma, P., Pfister, J. A., Mallick, S., & Dmello, S. R. (2014). HSF1 Protects Neurons through a Novel Trimerization- and HSP-Independent Mechanism. *Journal of Neuroscience*, 34(5), 1599-1612. doi:10.1523/jneurosci.3039-13.2014

55. Suomalainen, A., & Battersby, B. J. (2017). Mitochondrial diseases: The contribution of organelle stress responses to pathology. *Nature Reviews Molecular Cell Biology*, 19(2), 77-92. doi:10.1038/nrm.2017.66

56. Kim, K. H., Jeong, Y. T., Oh, H., Kim, S. H., Cho, J. M., Kim, Y., . . . Lee, M. (2012). Autophagy deficiency leads to protection from obesity and insulin resistance by inducing Fgf21 as a mitokine. *Nature Medicine*, 19(1), 83-92. doi:10.1038/nm.3014

57. Bao, X. R., Ong, S., Goldberger, O., Peng, J., Sharma, R., Thompson, D. A., . . . Mootha, V. K. (2016). Mitochondrial dysfunction remodels one-carbon metabolism in human cells. *ELife*, 5. doi:10.7554/elife.10575

58. Celardo, I., Lehmann, S., Costa, A. C., Loh, S. H., & Martins, L. M. (2017). DATF4 regulation of mitochondrial folate-mediated one-carbon metabolism is neuroprotective. *Cell Death and Differentiation*, 24(4), 638-648. doi:10.1038/cdd.2016.158

59. Nikkanen, J., Forsström, S., Euro, L., Paetav, I., Kohnz, R., Wang, L., . . . Suomalainen, A. (2016). Mitochondrial DNA Replication Defects Disturb Cellular dNTP Pools and Remodel One-Carbon Metabolism. *Cell Metabolism*, 23(4), 635-648. doi:10.1016/j.cmet.2016.01.019

60. Michel, S., Canonne, M., Arnould, T., & Renard, P. (2015). Inhibition of mitochondrial genome expression triggers the activation of CHOP-10 by a cell signaling dependent on the integrated stress response but not the mitochondrial unfolded protein response. *Mitochondrion*, 21, 58-68. doi:10.1016/j.mito.2015.01.005

61. Tsuyama, T., Tsubouchi, A., Usui, T., Imamura, H., & Uemura, T. (2017). Mitochondrial dysfunction induces dendritic loss via eIF2 α phosphorylation. *The Journal of Cell Biology*, 216(3), 815-834. doi:10.1083/jcb.201604065

62. Rath, E., Berger, E., Messlik, A., Nunes, T., Liu, B., Kim, S. C., . . . Haller, D. (2011). Induction of dsRNA-activated protein kinase links mitochondrial unfolded protein response to the pathogenesis of intestinal inflammation. *Gut*, 61(9), 1269-1278. doi:10.1136/gutjnl-2011-300767

63. Khan, N. A., Nikkanen, J., Yatsuga, S., Jackson, C., Wang, L., Pradhan, S., . . . Suomalainen, A. (2017). MTORC1 Regulates Mitochondrial Integrated Stress Response and Mitochondrial Myopathy Progression. *Cell Metabolism*, 26(2). doi:10.1016/j.cmet.2017.07.007

64. Fiorese, C., Schulz, A., Lin, Y., Rosin, N., Pellegrino, M., & Haynes, C. (2016). The Transcription Factor ATF5 Mediates a Mammalian Mitochondrial UPR. *Current Biology*, 26(15), 2037-2043. doi:10.1016/j.cub.2016.06.002

65. Teske, B. F., Fusakio, M. E., Zhou, D., Shan, J., Mcclintick, J. N., Kilberg, M. S., & Wek, R. C. (2013). CHOP induces activating transcription factor 5 (ATF5) to trigger apoptosis in response to perturbations in protein homeostasis. *Molecular Biology of the Cell*, 24(15), 2477-2490. doi:10.1091/mbc.e13-01-0067

66. Jin, S. M., & Youle, R. J. (2013). The accumulation of misfolded proteins in the mitochondrial matrix is sensed by PINK1 to induce PARK2/Parkin-mediated mitophagy of polarized mitochondria. *Autophagy*, 9(11), 1750-1757. doi:10.4161/auto.26122

67. Bouman, L., Schlierf, A., Lutz, A. K., Shan, J., Deinlein, A., Kast, J., . . . Winklhofer, K. F. (2010). Parkin is transcriptionally regulated by ATF4: Evidence for an interconnection between mitochondrial stress and ER stress. *Cell Death & Differentiation*, 18(5), 769-782. doi:10.1038/cdd.2010.142

68. Haynes, C. M., Yang, Y., Blais, S. P., Neubert, T. A., & Ron, D. (2010). The Matrix Peptide Exporter HAF-1 Signals a Mitochondrial UPR by Activating the Transcription Factor ZC376.7 in *C. elegans*. *Molecular Cell*, 37(4), 529-540. doi:10.1016/j.molcel.2010.01.015

69. Liu, W., Duan, X., Fang, X., Shang, W., & Tong, C. (2018). Mitochondrial protein import regulates cytosolic protein homeostasis and neuronal integrity. *Autophagy*, 14(8), 1293-1309. doi:10.1080/15548627.2018.1474991

70. Wrobel, L., Topf, U., Bragoszewski, P., Wiese, S., Sztolszener, M. E., Oeljeklaus, S., . . . Chacinska, A. (2015). Mistargeted mitochondrial proteins activate a proteostatic response in the cytosol. *Nature*, 524(7566), 485-488. doi:10.1038/nature14951

71. Rainbolt, T., Atanassova, N., Genereux, J., & Wiseman, R. (2013). Stress-Regulated Translational Attenuation Adapts Mitochondrial Protein Import through Tim17A Degradation. *Cell Metabolism*, 18(6), 908-919. doi:10.1016/j.cmet.2013.11.006

72. Verfaillie, T., Rubio, N., Garg, A. D., Bultynck, G., Rizzuto, R., Decuyper, J., . . . Agostinis, P. (2012). PERK is required at the ER-mitochondrial contact sites to convey apoptosis after ROS-based ER stress. *Cell Death & Differentiation*, 19(11), 1880-1891. doi:10.1038/cdd.2012.74

73. Taniuchi, S., Miyake, M., Tsugawa, K., Oyadomari, M., & Oyadomari, S. (2016). Integrated stress response of vertebrates is regulated by four eIF2 α kinases. *Scientific Reports*, 6(1). doi:10.1038/srep32886

74. Baker, B. M., Nargund, A. M., Sun, T., & Haynes, C. M. (2012). Protective Coupling of Mitochondrial Function and Protein Synthesis via the eIF2 α Kinase GCN-2. *PLoS Genetics*, 8(6). doi:10.1371/journal.pgen.1002760

75. Kim, Y., Park, J., Kim, S., Kim, M. A., Kang, M., Kim, B., . . . Kim, V. N. (2018). PKR Senses Nuclear and Mitochondrial Signals by Interacting with Endogenous Double-Stranded RNAs. *SSRN Electronic Journal*. doi:10.2139/ssrn.3155545

76. Liu, S., Feng, M., & Guan, W. (2016). Mitochondrial DNA sensing by STING signaling participates in inflammation, cancer and beyond. *International Journal of Cancer*, 139(4), 736-741. doi:10.1002/ijc.30074

77. Sliter, D. A., Martinez, J., Hao, L., Chen, X., Sun, N., Fischer, T. D., . . . Youle, R. J. (2018). Parkin and PINK1 mitigate STING-induced inflammation. *Nature*. doi:10.1038/s41586-018-0448-9

78. Nakahira, K., Haspel, J. A., Rathinam, V. A., Lee, S., Lam, H. C., Rabinovitch, M., . . . Choi, A. M. (2011). Autophagy Proteins Regulate Innate Immune Response By Inhibiting NALP3 Inflammasome-Mediated Mitochondrial DAN Release. *A21. Innate Immunity*. doi:10.1164/ajrccm-conference.2011.183.1_meetingabstracts.a1077

79. Saxton, R. A., & Sabatini, D. M. (2017). MTOR Signaling in Growth, Metabolism, and Disease. *Cell*, 169(2), 361-371. doi:10.1016/j.cell.2017.03.035

80. Gandin, V., Masvidal, L., Cargnello, M., Gynis, L., McLaughlan, S., Cai, Y., . . . Topisirovic, I. (2016). MTORC1 and CK2 coordinate ternary and eIF4F complex assembly. *Nature Communications*, 7, 11127. doi:10.1038/ncomms11127

81. Iadevaia, V., Huo, Y., Zhang, Z., Foster, L., & Proud, C. (2012). Roles of the mammalian target of rapamycin, mTOR, in controlling ribosome biogenesis and protein synthesis: Figure 1. *Biochemical Society Transactions*, 40(1), 168-172. doi:10.1042/bst20110682

82. Mayer, C., Zhao, J., Yuan, X. and Grummt, I. (2004) mTOR-dependent activation of the transcription factor TIF-IA links rRNA synthesis to nutrient availability. *Genes Dev.* 18, 423–434

83. Martina, J. A., Chen, Y., Gucek, M., & Puertollano, R. (2012). MTORC1 functions as a transcriptional regulator of autophagy by preventing nuclear transport of TFEB. *Autophagy*, 8(6), 903-914. doi:10.4161/auto.19653

84. Gao, D., Inuzuka, H., Tan, M., Fukushima, H., Locasale, J., Liu, P., . . . Wei, W. (2011). MTOR Drives Its Own Activation via SCF β TrCP-Dependent Degradation of the mTOR Inhibitor DEPTOR. *Molecular Cell*, 44(2), 290-303. doi:10.1016/j.molcel.2011.08.030

85. Zhao, Y., Xiong, X., & Sun, Y. (2011). DEPTOR, an mTOR Inhibitor, Is a Physiological Substrate of SCF β TrCP E3 Ubiquitin Ligase and Regulates Survival and Autophagy. *Molecular Cell*, 44(2), 304-316. doi:10.1016/j.molcel.2011.08.029

86. Qian, S., Zhang, X., Sun, J., Bennink, J. R., Yewdell, J. W., & Patterson, C. (2010). MTORC1 Links Protein Quality and Quantity Control by Sensing Chaperone Availability. *Journal of Biological Chemistry*, 285(35), 27385-27395. doi:10.1074/jbc.m110.120295

87. Guan, B., Krokowski, D., Majumder, M., Schmotzer, C. L., Kimball, S. R., Merrick, W. C., . . . Hatzoglou, M. (2014). Translational Control during Endoplasmic Reticulum Stress beyond Phosphorylation of the Translation Initiation Factor eIF2a. *Journal of Biological Chemistry*, 289(18), 12593-12611. doi:10.1074/jbc.m113.543215

88. Ye, J., Palm, W., Peng, M., King, B., Lindsten, T., Li, M. O., . . . Thompson, C. B. (2015). GCN2 sustains mTORC1 suppression upon amino acid deprivation by inducing Sestrin2. *Genes & Development*, 29(22), 2331-2336. doi:10.1101/gad.269324.115

89. Preston, A. M., & Hendershot, L. M. (2013). Examination of a second node of translational control in the unfolded protein response. *Journal of Cell Science*, 126(18), 4253-4261. doi:10.1242/jcs.130336

90. Su, K., Cao, J., Tang, Z., Dai, S., He, Y., Sampson, S. B., . . . Dai, C. (2016). HSF1 critically attunes proteotoxic stress sensing by mTORC1 to combat stress and promote growth. *Nature Cell Biology*, 18(5), 527-539. doi:10.1038/ncb3335

91. Qiao, A., Jin, X., Pang, J., Moskophidis, D., & Mivechi, N. F. (2017). The transcriptional regulator of the chaperone response HSF1 controls hepatic bioenergetics and protein homeostasis. *The Journal of Cell Biology*, 216(3), 723-741. doi:10.1083/jcb.201607091

92. Brüning, A., Brem, G. J., Vogel, M., & Mylonas, I. (2014). Tetracyclines cause cell stress-dependent ATF4 activation and mTOR inhibition. *Experimental Cell Research*, 320(2), 281-289. doi:10.1016/j.yexcr.2013.11.012

93. Zhou, Q., Liu, C., Liu, W., Zhang, H., Zhang, R., Liu, J., . . . Chen, L. (2014). Rotenone Induction of Hydrogen Peroxide Inhibits mTOR-mediated S6K1 and 4E-BP1/eIF4E Pathways, Leading to Neuronal Apoptosis. *Toxicological Sciences*, 143(1), 81-96. doi:10.1093/toxsci/kfu211

94. Park, D., Lee, M. N., Jeong, H., Koh, A., Yang, Y. R., Suh, P., & Ryu, S. H. (2014). Parkin ubiquitinates mTOR to regulate mTORC1 activity under mitochondrial stress. *Cellular Signalling*, 26(10), 2122-2130. doi:10.1016/j.cellsig.2014.06.010

95. Betz, C., & Hall, M. N. (2013). Where is mTOR and what is it doing there? *The Journal of Cell Biology*, 203(4), 563-574. doi:10.1083/jcb.201306041

96. Desai, B. N., Myers, B. R., & Schreiber, S. L. (2002). FKBP12-rapamycin-associated protein associates with mitochondria and senses osmotic stress via mitochondrial dysfunction. *Proceedings of the National Academy of Sciences*, 99(7), 4319-4324. doi:10.1073/pnas.261702699

97. Win, S., Than, T. A., Fernandez-Checa, J. C., & Kaplowitz, N. (2014). JNK interaction with Sab mediates ER stress induced inhibition of mitochondrial respiration and cell death. *Cell Death & Disease*, 5(1). doi:10.1038/cddis.2013.522

98. Win, S., Than, T. A., Le, B. H., García-Ruiz, C., Fernandez-Checa, J. C., & Kaplowitz, N. (2015). Sab (Sh3bp5) dependence of JNK mediated inhibition of mitochondrial respiration in palmitic acid induced hepatocyte lipotoxicity. *Journal of Hepatology*, 62(6), 1367-1374. doi:10.1016/j.jhep.2015.01.032

99. Mollie, A., Temirov, J., Lee, J., Coughlin, M., Kanagaraj, A., Kim, H., . . . Taylor, J. (2015). Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and Drives Pathological Fibrillization. *Cell*, 163(1), 123-133. doi:10.1016/j.cell.2015.09.015

100. Khong, A., Matheny, T., Jain, S., Mitchell, S. F., Wheeler, J. R., & Parker, R. (2017). The Stress Granule Transcriptome Reveals Principles of mRNA Accumulation in Stress Granules. *Molecular Cell*, 68(4). doi:10.1016/j.molcel.2017.10.015

101. McCormick, C., & Khaperskyy, D. A. (2017). Translation inhibition and stress granules in the antiviral immune response. *Nature Reviews Immunology*, 17(10), 647-660. doi:10.1038/nri.2017.63

102. Arimoto-Matsuzaki, K., Saito, H., & Takekawa, M. (2016). TIA1 oxidation inhibits stress granule assembly and sensitizes cells to stress-induced apoptosis. *Nature Communications*, 7, 10252. doi:10.1038/ncomms10252

103. Maharjan, N., Künzli, C., Buthey, K., & Saxena, S. (2016). C9ORF72 Regulates Stress Granule Formation and Its Deficiency Impairs Stress Granule Assembly, Hypersensitizing Cells to Stress. *Molecular Neurobiology*, 54(4), 3062-3077. doi:10.1007/s12035-016-9850-1

104. Zou, T., Yang, X., Pan, D., Huang, J., Sahin, M., & Zhou, J. (2011). SMN Deficiency Reduces Cellular Ability to Form Stress Granules, Sensitizing Cells to Stress. *Cellular and Molecular Neurobiology*, 31(4), 541-550. doi:10.1007/s10571-011-9647-8

105. Martin, S., Zekri, L., Metz, A., Maurice, T., Chebli, K., Vignes, M., & Tazi, J. (2013). Deficiency of G3BP1, the stress granules assembly factor, results in abnormal synaptic plasticity and calcium homeostasis in neurons. *Journal of Neurochemistry*, 125(2), 175-184. doi:10.1111/jnc.12189

106. Zekri, L., Chebli, K., Tourriere, H., Nielsen, F. C., Hansen, T. V., Rami, A., & Tazi, J. (2005). Control of Fetal Growth and Neonatal Survival by the RasGAP-Associated Endoribonuclease G3BP. *Molecular and Cellular Biology*, 25(19), 8703-8716. doi:10.1128/mcb.25.19.8703-8716.2005

107. Shukla, S., & Parker, R. (2016). Hypo- and Hyper-Assembly Diseases of RNA-Protein Complexes. *Trends in Molecular Medicine*, 22(7), 615-628. doi:10.1016/j.molmed.2016.05.005

108. Gilks, N., Kedersha, N., Ayodele, M., Shen, L., Stoecklin, G., Dember, L. M., & Anderson, P. (2004). Stress Granule Assembly Is Mediated by Prion-like Aggregation of TIA-1. *Molecular Biology of the Cell*, 15(12), 5383-5398. doi:10.1091/mbc.e04-08-0715

109. Cherkasov, V., Hofmann, S., Druffel-Augustini, S., Mogk, A., Tyedmers, J., Stoecklin, G., & Bukau, B. (2013). Coordination of Translational Control and Protein Homeostasis during Severe Heat Stress. *Current Biology*, 23(24), 2452-2462. doi:10.1016/j.cub.2013.09.058

110. Walters, R. W., Muhlrad, D., Garcia, J., & Parker, R. (2015). Differential effects of Ydj1 and Sis1 on Hsp70-mediated clearance of stress granules in *Saccharomyces cerevisiae*. *Rna*, 21(9), 1660-1671. doi:10.1261/rna.053116.115

111. Ganassi, M., Mateju, D., Bigi, I., Mediani, L., Poser, I., Lee, H. O., . . . Carra, S. (2016). A Surveillance Function of the HSPB8-BAG3-HSP70 Chaperone Complex Ensures Stress Granule Integrity and Dynamism. *Molecular Cell*, 63(5), 796-810. doi:10.1016/j.molcel.2016.07.021

112. Damgaard, C. K., & Lykke-Andersen, J. (2011). Translational coregulation of STOP mRNAs by TIA-1 and TIAR. *Genes & Development*, 25(19), 2057-2068. doi:10.1101/gad.17355911

113. Kedersha, N., Panas, M. D., Achorn, C. A., Lyons, S., Tisdale, S., Hickman, T., . . . Anderson, P. (2016). G3BP-Caprin1-USP10 complexes mediate stress granule condensation and associate with 40S subunits. *The Journal of Cell Biology*, 212(7), 845-860. doi:10.1083/jcb.201508028

114. Dao, T. P., Kolaitis, R., Kim, H. J., O'Donovan, K., Martyniak, B., Colicino, E., . . . Castañeda, C. A. (2018). Ubiquitin Modulates Liquid-Liquid Phase Separation of UBQLN2 via Disruption of Multivalent Interactions. *Molecular Cell*, 69(6). doi:10.1016/j.molcel.2018.02.004

115. Itakura, E., Zavodszky, E., Shao, S., Wohlever, M., Keenan, R., & Hegde, R. (2016). Ubiquilins Chaperone and Triage Mitochondrial Membrane Proteins for Degradation. *Molecular Cell*, 63(1), 21-33. doi:10.1016/j.molcel.2016.05.020

116. Zhang, K., Daigle, J. G., Cunningham, K. M., Coyne, A. N., Ruan, K., Grima, J. C., . . . Lloyd, T. E. (2018). Stress Granule Assembly Disrupts Nucleocytoplasmic Transport. *Cell*, 173(4). doi:10.1016/j.cell.2018.03.025

117. Vanderweyde, T., Youmans, K., Liu-Yesucevitz, L., & Wolozin, B. (2013). Role of Stress Granules and RNA-Binding Proteins in Neurodegeneration: A Mini-Review. *Gerontology*, 59(6), 524-533. doi:10.1159/000354170

118. Protter, D. S., & Parker, R. (2016). Principles and Properties of Stress Granules. *Trends in Cell Biology*, 26(9), 668-679. doi:10.1016/j.tcb.2016.05.004

119. Turakhiya, A., Meyer, S. R., Marincola, G., Böhm, S., Vanselow, J. T., Schlosser, A., . . . Buchberger, A. (2018). ZFAND1 Recruits p97 and the 26S Proteasome to Promote the Clearance of Arsenite-Induced Stress Granules. *Molecular Cell*, 70(5). doi:10.1016/j.molcel.2018.04.021

120. Meyer, H., & Weihl, C. C. (2014). The VCP/p97 system at a glance: Connecting cellular function to disease pathogenesis. *Journal of Cell Science*, 127(18), 3877-3883. doi:10.1242/jcs.093831

121. Alupei, M. C., Maity, P., Esser, P. R., Krikki, I., Tuorto, F., Parlato, R., . . . Iben, S. (2018). Loss of Proteostasis Is a Pathomechanism in Cockayne Syndrome. *Cell Reports*, 23(6), 1612-1619. doi:10.1016/j.celrep.2018.04.041

122. Readhead, B., Haure-Mirande, J., Funk, C., Richards, M., Shannon, P., Haroutunian, V., . . . Dudley, J. (2018). Multiscale Analysis of Independent Alzheimer's Cohorts Finds Disruption of Molecular, Genetic, and Clinical Networks by Human Herpesvirus. *Neuron*, 99(1), 64-82. doi:10.1016/j.neuron.2018.05.023

123. Meriin, A. B., Mense, M., Colbert, J. D., Liang, F., Bihler, H., Zaarur, N., . . . Sherman, M. Y. (2012). A Novel Approach to Recovery of Function of Mutant Proteins by Slowing Down Translation. *Journal of Biological Chemistry*, 287(41), 34264-34272. doi:10.1074/jbc.m112.397307

124. Moon, S. L., Sonenberg, N., & Parker, R. (2018). Neuronal Regulation of eIF2 α Function in Health and Neurological Disorders. *Trends in Molecular Medicine*, 24(6), 575-589. doi:10.1016/j.molmed.2018.04.001