

Raising Concern About Cross-Infection Between Humans and Companion Animals by Evolutionary Analysis of Receptors for SARS-CoV-2

Huihao Tang^{1*}, Mayan Ali^{1*}, Lina Zhang^{1*}, Cheng Deng^{1#}

¹ Jiangsu Key Laboratory for Biodiversity and Biotechnology, College of Life Sciences, Nanjing Normal University, Nanjing, China

*These authors contributed equally to this work.

#Corresponding author. E-mail: 12065106@s.upc.edu.cn

Abstract

Since the COVID-19 caused by SARS-CoV-2 break out in Wuhan China from Dec. 2019, it has spread to hundreds of countries up to now. Scientists from all over the world have paid tremendous efforts to research and try to control the disease. Previous studies suggested that some of the wild animals could be intermediate hosts between humans and origination of SARS-CoV-2, and some companion animals of humans can be infected by SARS-CoV-2, which raised our curiosity about cross-infection of SARS-CoV-2 between animals and humans. Thus, we select some kinds of animals that might have contact with humans to estimate the susceptibility to SARS-CoV-2 in different animals by evolutionary analysis of their receptors for SARS-CoV-2. The results show that some companion animals of the Felidae family like the cat has a higher infection possibility while the species of the Rodent family like the rat and mouse having close contact with humans show an opposite result, which consist with recent animal experiments and researches. These should raise concerns about cross-infection between human and companion animals or animals to have close contact with humans which might grow into depositaries of the virus after control of SARS-CoV-2 spreading and cause second or more infection wave after social reopening. Another side of our results stands by the opinion that bioinformatic analysis can be consistent with practical experiments in some respects so that we can prevent unnecessary sacrifice of laboratory animals in future experiments.

Keywords: COVID-19, SARS-CoV-2, companion animal, cross-infection.

Result

The worldwide breaking out coronavirus disease 2019 (COVID-19) caused by SARS-CoV-2 has spread to hundreds of countries with over 3 million cases(1) and attracted more and more scientists' attention. The background of COVID-19 has been stated over and over again by other scientists(2, 3), so we need no repeat in this article.

Like a host of scientists who want to figure out which species transfer the coronavirus to humans from the most possible original virus carrier—bats(4), we initially interested in potential intermediate hosts of SARS-CoV-2. Since previous researches have already found that S-protein of SARS-CoV-2 interacts with ACE2 of human finally induce the infection of the virus(4-6), we analyze the ACE2 protein sequences (Supplemental Table. Sequences were downloaded from <https://www.ncbi.nlm.nih.gov>) of different species which partially selected from reported for selling in Wuhan Huanan Seafood market (Supplemental Figure) and others

are species from companion animals of humans or having possible contact with humans to find out the potential intermediate host/hosts of SARS-CoV-2. Finally, we find that some of the species we analyzed could be susceptible to SARS-CoV-2 especially some companion animals and domestic animals that have close contact with humans. Thus, we suggest that actions should be taken to prevent the second wave after reopening worldwide because of the cross-infection of these animals to humans.

We find that the phylogenetic tree (Figure 1. A) constructed by these selected species is not much as identical as their time tree (Figure 1. B) as we expected, especially when we focus on *Homo sapiens*. This inconsistency caused by different selective pressures, changes in evolutionary rates, or horizontal gene transfer has been mentioned by J. Castresana, L. S. Haggerty, M. Ruvolo, and Hervé Philippe *et al.* (7-10). But in this case, we are more interested in the phylogenetic tree rather than the time tree because the virus was reported interacting with ACE2, not other receptors(4-6). Generally, the phylogenetic tree can sufficiently represent the susceptibility of species to SARS-CoV-2 if it merely interacts with ACE2 in an organism without any other interference. Yet as we all know our body is a massive system with many unknown signaling pathways so that we cannot take the phylogenetic tree of ACE2 into account only. To further estimate the difference between two type trees of these species, we use the TN (time period plus the number of nodes) method to quantitate the evolutionary distance between ACE2 of *Homo sapiens* and of other species selected in the time tree (Supplemental Table. the smaller TN number the less evolutionary distance to *Homo*, the time tree of the species was construct in <http://www.timetree.org>(11)). Next, we measure the Pairwise distances between ACE2 of *Homo sapiens* and other species by MegaX Software(12) in phylogenetic tree (Supplemental Table). To compare the results of two type trees uniformly, we sort the species by TN distances (by descending order) and Pairwise distances (also by descending order) and then compare them by ranking number (Figure 2, Supplemental Table. sorted from smallest to largest by results of Pairwise distances minus TN distances).

We first speculate that the intermediate hosts may be wild animals that contact humans and transfer the virus to humans by chance. From all species selected we set the cut-off as -5 and 5 in our result (Figure 2), which means species scored lower than -5 have much possibility to be intermediate hosts of SARS-CoV-2 (they have less distance against human in the phylogenetic tree than in the time tree) while species scored higher than 5 have contrary results. As our results shown in (Figure 2), the species listed in the bottom of histogram should have higher susceptibility to SARS-CoV-2 according to our method. As previous studies reported *Manis javanica*(13) and *Bungarus multicinctu*(14) could be possible intermediate hosts. Then we find that *Manis javanica* has a low score -7 (Figure 2) which means it might be a possible intermediate host, just like T. T. Lam *et al.*(13) point out that *Manis javanica* has a higher virus carrier rate according to whole-genome analysis. *Manis javanica* could be a possible species who contact both bats and humans so that they can transfer the virus to us because there are still some people who eat *Manis javanica* in China despite the limitation of law. That is why we think it can be a candidate not like another species--snake(14), which is reported could be possible intermediate host locating at the middle of histogram results (Figure 2. *Thamnophis elegans*: 2) and we don't think it has the possibility to be an intermediate host according to our analysis.

Then, when we look through other species locating at the bottom or top of our results histogram.

We find that the Felidae family gathering at bottom of the histogram (Figure 2) with lower ranking scores (*Puma concolor*: -30, *Felis catus*: -29, *Lynx pardinus*: -29, *Lynx canadensis*: -26, *Acinonyx jubatus*: -19) than any other species, which means that the Felidae family should be more susceptibility than other species we selected. However, another Order of species the Rodent family show contrary results which locating at top of the histogram (Figure 2) with higher scores (*Rattus norvegicus*: 16, *Mus musculus*: 16), even though they are closer to humans than the Felidae family does in the time tree. Some subfamily of Rodents like Rats and Mice are considered as the closest model organism to our Primates in laboratories, while this time they lose their status in our analysis. Both of results show above about our analysis are confirmed in experiment recently: J. Shi *et al.* reported that SARS-CoV-2 replicates efficiently in cats(15) and some zoo tigers also reported testing positive for COVID-19, while P. Zhou *et al* show that nCoV-2019 is unable to use mouse ACE2 as an entry receptor in the ACE2-expressing cells(4). Previous study report that the Felidae family is divergence earlier than the Rodent family and Primates at chromosome level(16), while our results find that cats have more similar ACE2 sequences with human than rodents so that virus are more easily transfer to cats from humans. This can explain why some early researches find that SARS-CoV-2 antibody could be detected *in vivo* of cats in the pandemic spreading area(17), but cannot infect cells transfected with mouse ACE2 *in vitro*(4). Lynx is another subfamily of the Felidae with low score not reported getting COVID-19 yet and keep as pet in some country, but we believe they are susceptibility to SARS-CoV-2.

In addition to the species we mentioned above, there are still some of species with not much significant difference between two types of trees but also score upper 5 or lower than -5. Species get lower score like *Canis lupus familiaris*, *Vulpes vulpes*, and *Paguma larvata* (Figure 2) can be carriers after contact with humans but may slightly affect by SARS-CoV-2 or have no symptom. Early study report that SARS-CoV-2 could be poorly detected in dogs(15) but some press reported that there is a dog dying after testing positive for COVID-19 which we need further confirmation. One of the moderate scored species--*Paguma larvata* was reported as an intermediate host of another coronavirus SARS-COV (Severe Acute Respiratory Syndrome Coronavirus) in 2003 which has around 80% sequence identity between SARS-CoV-2(4) so *Paguma larvata* may have possibility infected by SARS-CoV-2. Other species got higher numbers above 5 (Figure 2) may have low susceptibility to SARS-CoV-2 like pig(15) according to our results. These species with not much significant difference may not easy to determine their susceptibility to SARS-CoV-2 because of their moderate scores.

To further investigate much more detail between SARS-CoV-2 and its receptor ACE2, we choose a particular alpha helix amino acids sequence of ACE2 that is binding with Receptor Binding Domain (RBD) of SARS-CoV-2(18) and treat this helix sequence with the same method above (Figure 3A). The helix is at the interface between SARS-CoV-2 and ACE2 compound and role as part of Ligand Binding Domain (LBD) of ACE2 (Figure 3A). Most results are consistent with full length sequence results of ACE2 especially in the Felidae family, they locating at the bottom of the histogram (Figure 3B) which means that their amino acids of alpha helix are similar with human thus they are closer to human than they should be in time tree distances. What's more we find that *Capra hircus*, *Bos Taurus*, and *Ovis aries* which belong to Suborder Ruminantia have even lower scores than the Felidae family (Figure 3B). So, we speculate that they might also have susceptibility to SARS-CoV-2 according to our results

because they have similar alpha helix sequence with human that in the interface between SARS-CoV-2 and ACE2.

The result of alpha helix is only as identical as part of ACE2 intact sequence analysis, this may because the interface of ACE2 and SARS-CoV-2 contains not only the helix we selected but also some residues from other second structure of ACE2 which we cannot analyses them as a serial sequence combining with the helix by our method. We believe these residues have their important roles through interaction with SARS-CoV-2, so just select a part of the whole protein may merely represent a partial respect of full fact. Nevertheless, the single alpha helix also reconfirm our analysis above reminding us about concern of cross-infection between humans and other animals that might cause the pandemic comeback.

Discussion

Our results reveal that in the worldwide pandemic situation not only should consider infection between humans and take social distancing guideline but also should pay attention to animals around us being possible virus depository that may cause next wave of infection in future. In this study, we find out that some species do have higher susceptibility than other species which might be the virus depository not on purpose to encourage them to kill their pets regardless the humanity but warning people to prevent these risks.

Method

Sequences:

All protein sequences were downloaded from <https://www.ncbi.nlm.nih.gov> and performed with Aliview software(19). All of the species are as follows: *Acinonyx jubatus*, *Anas platyrhynchos*, *Aotus nancymae*, *Balaenoptera acutorostrata scammoni*, *Bos taurus*, *Camelus dromedarius*, *Canis lupus familiaris*, *Capra hircus*, *Carlito syrichta*, *Chlorocebus aethiops*, *Cricetulus griseus*, *Crocota crocuta*, *Equus caballus*, *Erinaceus europaeus*, *Felis catus*, *Gallus gallus*, *Gorilla gorilla gorilla*, *Heterocephalus glaber*, *Homo sapiens*, *Ictidomys tridecemlineatus*, *Lynx canadensis*, *Lynx pardinus*, *Macaca fascicularis*, *Macaca mulatta*, *Macaca nemestrina*, *Manis javanica*, *Marmota flaviventris*, *Mesocricetus auratus*, *Microcebus murinus*, *Microtus ochrogaster*, *Mus musculus*, *Mus pahari*, *Myotis lucifugus*, *Nomascus leucogenys*, *Octodon degus*, *Oryctolagus cuniculus*, *Ovis aries*, *Paguma larvata*, *Pan paniscus*, *Pan troglodytes*, *Papio anubis*, *Physeter catodon*, *Ptilocolobus tephrosceles*, *Pongo abelii*, *Puma concolor*, *Rattus norvegicus*, *Rhinopithecus roxellana*, *Sapajus apella*, *Sus scrofa*, *Thamnomphs elegans*, *Theropithecus gelada*, *Urocitellus parryii*, *Ursus arctos horribilis*, *Vicugna pacos*, *Vulpes vulpes*.

Trees:

The time tree was constructed by online server (<http://www.timetree.org>) with species listed above. The phylogenetic tree (Neighbor-Joining tree) was constructed by MegaX software(12) (Poisson model, Uniform Rates, Pairwise deletion) using downloaded sequences. Both trees were drawing by iTOL online server (<https://itol.embl.de>)(20)


Histogram and Diagram:

Histograms of the results were performed using GraphPad Prism software and the diagram of the alpha helix structure was drawing by PyMOL software (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.)

References:

1. World Health Organization (2020) Coronavirus disease (COVID-19). in *Situation Report – 107* (World Health Organization).
2. V. Coronaviridae Study Group of the International Committee on Taxonomy of, The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536-544 (2020).
3. A. S. Gonzalez-Reiche *et al.*, Introductions and early spread of SARS-CoV-2 in the New York City area. *medRxiv* (2020).
4. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).
5. Q. Wang *et al.*, Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. *Cell* 10.1016/j.cell.2020.03.045 (2020).
6. M. Hoffmann *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278 (2020).
7. J. Castresana, Topological variation in single-gene phylogenetic trees. *Genome Biol* **8**, 216 (2007).
8. L. S. Haggerty, F. J. Martin, D. A. Fitzpatrick, J. O. McInerney, Gene and genome trees conflict at many levels. *Philos Trans R Soc Lond B Biol Sci* **364**, 2209-2219 (2009).
9. M. Ruvolo, Molecular phylogeny of the hominoids: inferences from multiple independent DNA sequence data sets. *Mol Biol Evol* **14**, 248-265 (1997).
10. H. Philippe, A. Chenuil, A. Adoutte, Can the Cambrian explosion be inferred through molecular phylogeny? *Development* **1994**, 15-25 (1994).
11. S. B. Hedges, J. Marin, M. Suleski, M. Paymer, S. Kumar, Tree of life reveals clock-like speciation and diversification. *Mol Biol Evol* **32**, 835-845 (2015).
12. S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549 (2018).
13. T. T. Lam *et al.*, Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. *Nature* 10.1038/s41586-020-2169-0 (2020).
14. W. Ji, W. Wang, X. Zhao, J. Zai, X. Li, Cross-species transmission of the newly identified coronavirus 2019-nCoV. *J Med Virol* **92**, 433-440 (2020).
15. J. Shi *et al.*, Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science* 10.1126/science.abb7015 (2020).
16. J. U. Pontius *et al.*, Initial sequence and comparative analysis of the cat genome. *Genome Res* **17**, 1675-1689 (2007).
17. Q. Zhang *et al.*, SARS-CoV-2 neutralizing serum antibodies in cats: a serological investigation. *BioRxiv* (2020).
18. R. Yan *et al.*, Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* **367**, 1444-1448 (2020).
19. A. Larsson, AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**, 3276-3278 (2014).
20. I. Letunic, P. Bork, Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and annotation. *Bioinformatics* **23**, 127-128 (2007).

A



B




Figure 1. Time Tree of selected species and Phylogenetic Tree (Neighbor-Joining Tree) of their ACE2. (A) Time tree was constructed by online server (<http://www.timetree.org>). The Internal tree scale represent the time scale of these species (unit: MYA/Millions of Years Ago). (B) Phylogenetic tree of ACE2 protein sequences. The branch lengths represent the percentage of amino acids mutation in different branches. The Felidae family is label by red leaf ranges.




Figure 2. Differences of distance against to human between Time Tree and Phylogenetic Tree. Red bracket represent the Felidae family locating at the bottom of the histogram. Cut-off is set at -5 and 5.




Figure 3. Differences of distance against to human between Time Tree and Phylogenetic Tree of the alpha helix. (A) Diagram of alpha helix in the interface between ACE2 and SARS-CoV-2 (R. Yan et al.)(18).(B) Red bracket represent the Felidae family locating at the bottom of the histogram.