

Risk perceptions of antibiotic usage and resistance: a cross-sectional survey of poultry farmers in Kwara State, Nigeria

Al-Mustapha A. Ibrahim ^{a,b,c}, Victoria O. Adetunji ^b and Annamari Heikinheimo ^c

^a Department of Veterinary Services, Kwara State Ministry of Agriculture and Rural Development, Ilorin, Nigeria.

^b Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary medicine, University of Ibadan, Nigeria.

^c Department of Food Hygiene and Environmental Health, Faculty of Veterinary Medicine, University of Helsinki, Finland.

*Corresponding author: Al-Mustapha, Ahmad Ibrahim ahmad.al-mustapha@helsinki.fi

Abstract

There are overwhelming empirical evidences highlighting the contribution of indiscriminate antibiotic usage (ABU) in food animals to the overall burden of antibiotic resistance (ABR) in humans, thus making antibiotic use the main selective pressure driving antibiotic resistance. Social and behavioral perspective on antibiotic use and resistance in poultry is limited. Our study therefore aimed at obtaining information on antibiotic usage, awareness of ABR, and the attitude and perceptions towards prudent antibiotic usage and ABR. A cross-sectional survey using a structured questionnaire was conducted in 125 poultry farms in Kwara state in December 2019. Most farmers (69.6%, n=87/125) were aware of ABR and had satisfactory knowledge about ABR with a mean knowledge score of 3.16 ± 1.47 . The age, gender, level of education of farmers, and their flock size were significantly associated with a satisfactory knowledge of ABR ($p < 0.05$). Tertiary education was significantly associated with ABR awareness (OR: 4.7; 95% CI: 0.0690, 0.654; $p = 0.007$) and the ABR knowledge level (OR: 7.8269; 95% CI: 3.2693, 18.7381; $p < 0.01$). Higher flock size was significantly associated with a satisfactory knowledge of ABR (OR: 9.4551; 95%CI: 3.7928, 23.5707; $p < 0.01$). Most of the poultry farmers (68%) had positive attitudes towards prudent antibiotic use with a mean score of 2.75 ± 0.89 . On the contrary, only 32.8% of poultry farmers had a good perception of ABR with a mean perception score of 4.95 ± 1.12 . The ABR knowledge level was significantly associated with the perceptions of farmers ($p < 0.05$) but not their attitudes toward ABU and ABR ($P = 0.083$). There was evidence of unprescribed use of antibiotics in poultry and a failure to observe antibiotic withdrawal periods. These constitute a risk of exposure to unacceptable levels of drug residues from poultry products and an increased risk of ABR. Improving education and communication on antibiotic stewardship programs are crucial to prevent the looming antibiotic apocalypse.

Keywords: Antibiotic usage; Antibiotic Resistance; Poultry; KAP; Kwara; Nigeria

1 Background

The illicit and uncontrolled access to and use of antibiotics in humans and animals is one of the major drivers of antibiotic resistance (ABR). It is a global health threat that is estimated to cause approximately 10 million deaths and over USD 100 trillion by 2050 if no global actions were

established (1; 2). It is estimated that the greatest impact of ABR will be in Sub-Saharan Africa and Asia due to the disproportionately high infectious disease burden, overwhelmed health care systems of most countries, poor livelihoods and living conditions, and poor healthcare infrastructures. Globally, ABR has gained global attention due to the increasing incidence of multidrug-resistant (MDR) organisms causing treatment failures, antibiotic residues in food, and the public health risks it poses (3). There is an increase in the incidence and dissemination of MDR organisms in humans, health facilities, animals, foods, and the environment (4; 5; 6; 7; 8; 9).

In Nigeria, the misuse of antibiotics, proliferation of unlicensed drug stores, sub-therapeutic use of antibiotics in food animals for prophylaxis and growth promotion, poor ABR awareness, and lack of stewardship programs have further frustrated efforts at controlling ABR in the human and animal health sectors (10). In Nigeria today, most antibiotics are available as over-the-counter (OTC) drugs. In humans, the rate of antibiotic prescription was 49.1% (11). A point prevalence study showed 80% of all hospitalized patients were on antibiotic course(12). In a 2015 global survey by the world health organization (WHO), Nigeria was the country with the highest number of respondents who reported having obtained antibiotics from a stall or hawker (13).

With a population of over 200 million people, there is extensive economic pressure on the poultry industry as a vital source of animal protein for the teeming population of Nigeria. The poultry industry in Nigeria has 180 million birds and produces 650,000 metric tonnes of egg and 300,000 metric tonnes of poultry meat annually (14). Globally, there are increasing empirical evidence and epidemiological studies highlighting the contribution of indiscriminate antibiotic usage (ABU) and ABRin animals to the overall burden of ABRin human (15; 16; 17). This interconnectedness of ABU and ABR in animals and human health requires a multi-sectoral one health approach (18).

There is a paucity of data on ABU and ABR from a social and behavioral change perspective which is essential to achieve attitudinal change necessary to control the imminent threat of ABR. Hence, improving the public awareness and knowledge on prudent use of antibiotics will play a significant role in reducing the illicit consumption of antibiotics, the main driver of ABR especially in the animal health sector in Nigeria. As such, we aimed to assess the knowledge, attitude, and perceptions of poultry farmers regarding ABU and ABR.

This paper is part of a broader project on animal health and disease demographic survey in Nigeria.

2. Materials and Methods

2.1. Ethical considerations

The Kwara State Ministry of Agriculture and Natural Resources, Ilorin, Nigeria (reference number: VKW/714/1/103) approved this study. Participation was anonymous and voluntary. Informed consent was sought from the respondents and participants could withdraw from the survey at any time in line with stipulations of the World Medical Association Declaration of Helsinki Ethical principles (19).

2.2. Study design, study participants, and sample size.

This study was conducted in December 2019as a cross-sectional survey of poultry farmers in Kwara State. A comprehensive list of poultry farms was obtained from the Ministry of Agriculture and the Poultry Association of Nigeria (n=197 farms). The targeted respondents were farm owners or managers. To calculate the sample size, we hypothesized that at a 95% confidence level, the assumed prevalence of antibiotic use was 50% of all farms. The total sample size was 131 farms. So, a random sampling of 131 farms was done to evaluate antibiotic usage and the farmer's perception of ABR.

2.3. Questionnaire design

A structured questionnaire was designed to conduct this study. The questionnaire was pre-validated by two independent reviewers, and a pilot study was conducted with 10 respondents. The responses from the pre-test were not included in the analyzed data. The questionnaire consisted of 4 parts: a) Demography of respondents, b) Antibiotic usage c) Knowledge of antibiotics and ABR d) Attitude and perceptions towards ABR. Some of the questionnaires were administered to farmers in their farms (n = 80) while others (n = 45) were administered in feed mills where farmers come to purchase poultry feeds.

2.4. Data analysis

Data were summarized using Microsoft Excel 2016 and analyzed using Minitab v.19.1.1. Descriptive statistics (frequency and proportions) were used to summarize the obtained data. To assess the knowledge, attitude, and perception levels of the poultry farmers, a numeric scoring system was used, and outcome variables – knowledge, attitude, and perception – were computed. These outcome variables were further categorized as binary (satisfactory or unsatisfactory) based on the cut-off (mean scores) marks. Chi-square test was used to test for association between independent variables (demographics) and outcome variables (knowledge, attitude, and perception) at a 95% confidence interval with significant variables ($p < 0.05$) subjected to a logistic regression model.

3. Results

3.1. Respondent demographics

The questionnaire was administered to 131 poultry farmers. However, only 125 responses were received as 6 farmers did not consent to participate in the study. Of these, female respondents accounted for most (56.8%, n = 71) of the responses. Most poultry farmers (72%, n = 90) had tertiary education and 63 (50.4%) of all farmers employed 1-2 workers (Table 1).

Table 1. Demographic structure of respondents (n=125).

Variables	No. of respondents (%)
Gender	
Female	71 (56.8)
Male	54 (43.2)
Age	
20-30	26 (20.8)
30-40	18 (14.4)
40-49	39 (31.2)
50-59	29 (23.2)
60-69	13 (10.4)
Level of Education	
Secondary education	35 (28)
Tertiary education	90 (72)
No of workers on the farm	
1	31 (24.8)
2	32 (25.6)
3	20 (16)
4	13 (10.4)
5	9 (7.2)
6	14 (11.2)
7	4 (3.2)
9	2 (1.6)

3.2. Numeric scoring system

Each graded question was allotted 1 point. A total score for each respondent was computed. Respondents that received scores greater than the mean scores for knowledge (3.16 ± 1.47), attitude (2.75 ± 0.89), and perception (4.95 ± 1.12) were deemed to be satisfactory responses and vice versa (Table 2).

Table 2. Description of scores obtained by respondents ($n = 125$) for knowledge, attitude, and perception towards ABR.

Outcome variables	Maximum obtainable scores	Scores received by respondents		Mean \pm SD	Satisfactory n (%)	Unsatisfactory n (%)
		Minimum score	Maximum score			
Knowledge	7	0	6	3.16 ± 1.47	87 (69.6)	38 (30.4)
Attitude	5	0	4	2.75 ± 0.89	85 (68)	40 (32)
Perception	8	3	7	4.95 ± 1.12	41 (32.8)	84 (67.8)

3.3. Antibiotic Usage in poultry

Most of the farms (89%, $n = 111$) had layers and 87% ($n = 109$) of all farms had less than 1,000 birds (small scale farmers). Similarly, 95.2% ($n = 119$) of all farms have completed their vaccination schedule against endemic poultry diseases. The majority of farmers (83.2%, $n = 104$) used antibiotics in the last 4 weeks but only one farm (0.8%) took samples for laboratory testing before the administration of antibiotics. During this survey, gentamicin based (68.8%, $n = 86$), sulfonamide (44%, $n = 55$) and quinolone-based antibiotics (30.4%, $n = 38$) were the most frequently administered antibiotics in poultry. Most of the farmers (92%, 115) purchased antibiotics from licensed drug stores. The majority of farmers (56%, $n = 70$) purchased antibiotics based on their previous experiences (Table 3).

Table 3. Descriptive statistics on poultry flocks and antibiotic usage in selected poultry farms in Kwara state ($n=125$).

Type of Birds	No. of respondents (%)
Broilers	12 (9.6)
Cockerel	2 (1.6)
Layers	111 (88.8)
Population of birds	
<500	37 (29.6)
500 – 1000	72 (57.6)
>1000	16 (12.8)
Ever sampled birds for lab testing?	
No	124 (99.2)
Yes	1 (0.8)
Vaccination status	
Complete vaccination schedule	119 (95.2)
Incomplete vaccination schedule	6 (4.8)
Was vaccination done by a vet?	
NO	104 (83.2)
YES	21 (16.8)
When last did you administer antibiotics to your birds?	
<4 weeks	104 (83.2)
>4 weeks	21 (16.8)
Which class of antibiotics did you use?	
Gentamicin based	86 (68.8)
Quinolones based	38 (30.4)
Sulphadimidine based	55 (44)
Oxytetracycline based	40 (32)
Sources of antibiotics	

Drug peddlers	7 (5.60)
Licensed store	115 (92)
Other sources	3 (2.40)
Do you purchase antibiotics based on other farmers' experiences?	
No	101 (80.8)
Yes	24 (19.2)
Do you purchase antibiotics based on your own previous experiences?	
No	55 (44)
Yes	70 (56)

3.3. Knowledge, attitude, and practices towards antibiotic resistance in Kwara State.

The majority of the poultry farmers (69.6%; n = 87) were aware of antibiotic resistance. The mean knowledge score was 3.16 ± 1.47 . Using the mean score as the cut-off; most of the poultry farmers (87; 69.6%) have satisfactory knowledge about ABR. Most farmers (72%, n = 90) knew that bacteria in poultry could become resistant to drugs but only 49 farmers (39.2%) agreed that ABR could make treatment difficult in birds. On the contrary, 39 (31.2%) of the farmers were unaware that ABR pathogens in birds can affect man. Also, 43 farmers (34.4%) were unaware that antibiotics cannot be used to treat viral, fungal, or parasitic infections in birds (Table S1). The age and the level of education of farmers were significantly associated with increased ABR awareness ($p < 0.05$). Poultry farmers with secondary education were $4.7 \times$ (95% CI: 0.0690, 0.654; $p = 0.007$) more likely to be aware of ABR than those with tertiary education (Table 5). Similarly, the age, gender, level of education of farmers, and their flock size were significantly associated with a satisfactory knowledge of ABR ($P < 0.05$) (Table S4). Male farmers were $8.51 \times$ (95% CI: 3.0339, 23.8732; $p < 0.01$) more likely to have satisfactory knowledge. An increase in flock size was significantly associated with a satisfactory knowledge of ABR (OR: 9.4551; 95%CI: 3.7928, 23.5707; $p < 0.01$) (Table 5). Respondents that have commercial poultry farms (but not full-term farmers) were $1.9765 \times$ (95% CI: 0.8544, 4.5722; $p = 0.111$) more likely to have satisfactory knowledge than full-term poultry farmers (Table S4).

Most of the poultry farmers (68%, n = 85) had positive attitudes towards prudent antibiotic use with a mean score of 2.75 ± 0.89 . Most farmers (88%, n = 110) did not believe that there was excessive antibiotic use in poultry. Only (48.8%, n = 61) farmers got their antibiotic prescription from a vet. While most farmers (89.6%, n = 112) claim to observe the withdrawal period of antibiotics as stipulated on each antibiotic sachet or vial, none of them discarded the eggs in the course of antibiotic therapy (Table S2). The farmer's age was significantly associated with a positive attitude towards prudent antibiotic usage ($p < 0.05$) (Table S5).

On the contrary, only 41 farmers (32.8%) had a good perception of ABR with a mean perception score of 4.95 ± 1.12 . Most farmers (85.6%, n = 107) did not believe that ABR is a major health threat that needs to be addressed in Nigeria. Similarly, 84 farmers (67.2%) did not believe that farmers must reduce the use of antibiotics in birds. Only 59 farmers (47.2%) felt that only a certified veterinarian should be allowed to prescribe antibiotics. Some farmers (48%, n = 60) of the farmers believe that proper routine vaccinations could reduce dependence on antibiotics. Most of the farmers (72%, n = 90) thought that the threat of ABR only affects farms that use antibiotics. While only 56 farmers (44.8%) thought there is nothing that they can do to reduce the emergence and transmission of MDR bacteria, most farmers (81.6%, n = 102) believed that frequent hand washing is important for poultry farmers after attending to their birds (Table S3). Farmer's education was significantly associated with a good perception of ABR (OR: 0.1304; 95% CI: 0.0547, 0.3108; $p < 0.01$) (Table S5).

The ABR knowledge level was significantly associated with the perceptions of farmers ($p < 0.05$) but not their attitudes toward ABU and ABR ($P = 0.083$) (Table 4). The flock size of the farmers was significantly associated with awareness rates as well as knowledge and perception ($p < 0.05$)

but not with the attitude of poultry farmers ($p = 0.468$) (Table S4; S5). Secondary education was the major predictor of higher ABR awareness and a good perception of ABR (Table 5).

Table 4. Association of knowledge level and the attitude and perceptions of poultry farmers on ABR.

		Attitude				
		Good (%)	Poor (%)	χ^2	DF	p - value
Knowledge	Satisfactory	55 (64.7)	32 (80)	3.01	1	0.083
	Unsatisfactory	30 (35.3)	8 (20)			
			Perception			
	Satisfactory	16 (39)	71 (84.5)	26.96	1	<0.01
	Unsatisfactory	25 (61)	13 (15.5)			

Table 5. Predictors of ABR awareness, satisfactory knowledge, and perceptions of ABR among poultry farmers in Kwara state.

	Variables	Univariate analysis		Multivariate analysis	
		OR (95% CI)	P-value	OR (95% CI)	P-value
Awareness of ABR	Secondary education	4.7058 (1.5274, 14.4927)	0.007	2.2471 (0.104, 5.2521)	< 0.01
	Tertiary education	7.8269 (3.2693, 18.7381)	< 0.01	2.1848 (0.7400, 6.4507)	< 0.01
Knowledge of ABR	Male Gender	8.5105 (3.0339, 23.8732)	< 0.01	4.1776 (1.1805, 14.7837)	0.027
	Flock size between 500-1000	9.4551 (3.7928, 23.5707)	< 0.01	4.5306 (1.5566, 13.1863)	0.021
Perceptions of ABR	Secondary education	7.6687 (3.2175, 18.2815)	< 0.01	3.5486 (1.2840, 9.8039)	0.015

OR- Odds ratio; 95%CI – 95% confidence interval.

4. Discussion

Frequent and sub-therapeutic doses of antibiotics create the ideal selective pressure for the emergence of resistant micro-organisms. The excessive use (or misuse) of antibiotics in animal production has severe consequences for public health and the environment (6; 9). MDR organisms have been isolated in poultry, poultry environment, and in poultry workers and this poses serious public health threats especially in LMICs like Nigeria (20; 21; 22). These MDR bacteria can be transmitted to humans via the food chain, the environment, water bodies, or by close contact with these animals (6; 23; 24).

In this study, the majority of farmers (83.2%) used antibiotics in their poultry during the last four weeks but only one (0.8%) farm took samples for laboratory testing before the administration of the antibiotics. For economic reasons, small-scale poultry farmers did not consult veterinarians before the administration of antibiotics and the majority of farmers (56%) purchased antibiotics based on their previous experiences (Table 3). In Nigeria, a cocktail of antibiotics with multivitamins and mineral elements is common. Gentamicin, sulfonamide, and quinolone-based antibiotics were the most frequently administered antibiotics in these birds. This is similar to reports by Adebawale et al., (25) and Ogunleye et al., (26) where gentamicin, tetracycline, quinolones, and sulfonamides were the most frequently used antibiotics in poultry in Ogun state. Their popularity amongst farmers might be because they are very cheap and readily available (27). Although 95.2% of all farms had completed their vaccination schedule against endemic poultry diseases in Nigeria, only some (48%) knew that vaccinations could prevent the occurrence of disease and this will reduce antibiotic consumption. The misuse, abuse, and resistance to

quinolones (ciprofloxacin) in animals are particularly worrisome because ciprofloxacin is on the essential medicines list for humans (28). Surprisingly, over the last two months (8 weeks), no farmer reported the use of any of the banned antibiotics in animals such as furazolidone and chloramphenicol.

While the majority of poultry farmers (69.6%) were aware of ABR, there were obvious gaps in their knowledge of ABR. Some poultry farmers (34.4%) thought antibiotics can be used to treat viral, parasitic, and fungal diseases. This might be due to the use of antibiotics to treat secondary bacterial infections associated with viral diseases such as Newcastle disease or fowl pox. Although most farmers were familiar with ABR as a term, they do not know what it means and the implications for human health. This is further evidenced by the fact that most farmers knew that bacteria in birds could become resistant to antibiotics but were unaware that these resistant bacteria could make treatment difficult in birds. Older farmers had significantly higher ABR awareness rates, knowledge levels, and attitudes than younger respondents. This might be due to hands-on experience acquired over the years. Farmers with secondary education were more aware, had better attitudes and perceptions of ABR than those with tertiary education. However, those with tertiary education had better knowledge of ABR (Table S5). Male farmers were more likely to have a better knowledge of ABR than females. Therefore, there is a need to improve ABR awareness among female farmers. This can be achieved by collaborations with women in agriculture groups throughout Nigeria. Farmers with higher flock sizes were more aware and had better knowledge of ABR. However, for economic reasons and production pressure, these variables do not influence the attitude of poultry farmers towards prudent antibiotic use.

The attitude of most poultry farmers towards ABR (68%) was positive with a mean score of 2.75 ± 0.89 . However, through enhanced behavioral changes communications, the attitude of poultry farmers needs to be improved with an emphasis on the observance of withdrawal periods for antibiotics, and the public health impact it has on human health. There are a plethora of studies that reported the non-adherence to antibiotics withdrawal periods as the major cause of antibiotic drug residues in foods of animal origins (3; 29; 30; 31; 32; 33; 34). As previously reported by Adebawale et al., (25) and Geidam et al., (35), some poultry farmers (38.4%) in this study stopped antibiotic therapy when they noticed improvements in their birds.

With a mean of 4.95 ± 1.12 , only 32.8% of poultry farmers had a satisfactory perception of ABR. Most poultry farmers neither believed they practice excessive antibiotic use nor perceived the imminent threat of ABR. Sequel to this, most farmers (67.2%) do not see the need to reduce antibiotic use in birds. The majority of poultry farmers did not believe that ABR is a major health threat, hence the national Action plan for antimicrobial resistance (NAP-AMR) should be focused on raising awareness of ABR in these farmers. This is even more important with the increasing number of poultry farmers in Nigeria. Farmers should also be educated that mechanical transmission of drug-resistant microorganisms is possible in farms that do not use antibiotics. Farmers could introduce MDR-organisms into their farms from feed mills, through feed sacks, transport vehicles, and farmworkers. To contribute to the fight against ABR and preserve the efficacy of drugs in humans and animals, farmers should practice good management practices, administer essential poultry vaccinations, tighten the biosecurity measures on their farms, and make use of prebiotics and probiotics.

We will like to promote the concept of antibiotic-free birds among consumers of poultry products. It is essential to assess and quantify antibiotic usage in other livestock production systems such as in aquaculture and dairy farms.

5. Conclusion

This study reports the extensive use of unprescribed essential antibiotics in poultry. The majority of antibiotics were received as OTC drugs and the withdrawal period of antibiotics were not followed. Farmers knew about ABR but were unaware of their harm for their animals and the potentials for human transmission. Therefore, interventions such as the full implementation of the

NAP-AMR, antibiotic stewardship programs, and behavioral change communications to livestock farmers should be instituted to prevent the looming antibiotic apocalypse. This apocalypse can only be solved by multi-sectoral collaborations using the one-health approach (36). Furthermore, a functional national ABR surveillance program in the livestock sector is long overdue.

Declarations

Availability of data and materials

The survey instrument and dataset are available as supplementary data.

Competing interests

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgments

We acknowledge Emmanuel Awosanya for validating the survey instrument.

Authors' contributions

AIA: VO, and AH contributed equally to the study. AIA drafted the manuscript. All co-authors did the overall review of the manuscript and approved the final study.

References

1. World Health Organization (WHO). New report calls for urgent action to avert antibiotic resistance crisis [Internet]. 2020 [cited 9 May 2020]. Available from: <https://www.who.int/news-room/detail/29-04-2019-new-report-calls-for-urgent-action-to-avert-antibiotic-resistance-crisis>
2. O'Neill J. Antibiotics in Agriculture and the Environment: Reducing unnecessary Use And Waste. 2015.
3. Annan-Prah A, Agbemafle E, Asare PT, Akorli SY. Antibiotic use, abuse and their public health implication: The contributory role of management flaws in the poultry industry in two Agro-ecological zones in Ghana. *J Vet Adv.* 2012; (2):199-208.
4. World Health Organization (WHO). New report calls for urgent action to avert antibiotic resistance crisis [Internet]. 2020 [cited 9 May 2020]. Available from: <https://www.who.int/news-room/detail/29-04-2019-new-report-calls-for-urgent-action-to-avert-antibiotic-resistance-crisis>
5. Kayode A, Okunrounmu P, Olagbende A, Adedokun O, Hassan A, Atilola G. High prevalence of multiple drug resistant enteric bacteria: Evidence from a teaching hospital in Southwest Nigeria. *Journal of Infection and Public Health.* 2020;13(4):651-656.
6. Caudell M, Dorado-Garcia A, Eckford S, Creese C, Byarugaba D, Afakye K, et al. Towards a bottom-up understanding of antibiotic use and resistance on the farm: A knowledge, attitudes, and practices survey across livestock systems in five African countries. *PLOS ONE.* 2020;15(1):e0220274.
7. Aworh M, Kwaga J, Okolocha E, Mba N, Thakur S. Prevalence and risk factors for multi-drug resistant *Escherichia coli* among poultry workers in the Federal Capital Territory, Abuja, Nigeria. *PLOS ONE.* 2019;14(11):e0225379.
8. Siwakoti S, Subedi A, Sharma A, Baral R, Bhattarai N, Khanal B. Incidence and outcomes of multidrug-resistant gram-negative bacteria infections in intensive care unit from Nepal- a prospective cohort study. *Antibiotic Resistance & Infection Control.* 2018;7(1).
9. Chandra B, Amit K, Rajeshwari S, Mouna N. Antibiotic Resistance in Poultry Environment: Spread of Resistance from Poultry Farm to Agricultural Field, Centre for Science and Environment. 2017. New Delhi
10. Federal Ministry of Agriculture, Environment, and Health. Antibiotic Use and Resistance in Nigeria; Situation Analysis and Recommendations. 2017.
11. Egwuenu A, Obasanya J, Okeke I, Aboderin O, Olayinka A, Kwange D et al. Antibiotic use and resistance in Nigeria: situation analysis and recommendations, 2017. *Pan African Medical Journal Conference Proceedings.* 2018;2.

12. Abubakar U. Antibiotic use among hospitalized patients in northern Nigeria: a multicenter point-prevalence survey. *BMC Infectious Diseases*. 2020;20(1).
13. World Health Organization. Antibiotic Resistance: Multi-Country Public Awareness Survey. Geneva; 2015.
14. Food and Agricultural Organization (FAO). Livestock and livelihoods spotlight Cattle and poultry sectors Nigeria. 2017.
15. Advice on impacts of using antibiotics in animals - European Medicines Agency [Internet].2020 [cited 31 May 2020]. Available from: <https://www.ema.europa.eu/en/veterinary-regulatory/overview/antibiotic-resistance/advice-impacts-using-antibiotics-animals>
16. Marshall B, Levy S. Food Animals and Antibiotics: Impacts on Human Health. *Clinical Microbiology Reviews*. 2011;24(4):718-733.
17. den Heijer C, Beerepoot M, Prins J, Geerlings S, Stobberingh E. Determinants of Antibiotic Resistance in *Escherichia coli* Strains Isolated from Faeces and Urine of Women with Recurrent Urinary Tract Infections. *PLoS ONE*. 2012;7(11):e49909.
18. Robinson T, Bu D, Carrique-Mas J, Fèvre E, Gilbert M, Grace D et al. Antibiotic resistance is the quintessential One Health issue. *Transactions of The Royal Society of Tropical Medicine and Hygiene*. 2016;110(7):377-380.
19. WMA. World Medical Association Declaration of Helsinki ethical principles for medical research involving human subjects. *JAMA*, 2013, 310, 2191–2194.
20. Aworh M, Kwaga J, Okolocha E, Harden L, Hull D, Hendriksen R et al. Extended-spectrum β -lactamase-producing *Escherichia coli* among humans, chickens and poultry environments in Abuja, Nigeria. *One Health Outlook*. 2020;2(1).
21. Ahmed A, Raji M, Mamman P, Kwanashie C, Raufu I, Aremu A et al. Salmonellosis: Serotypes, prevalence and multi-drug resistant profiles of *Salmonella enterica* in selected poultry farms, Kwara State, North Central Nigeria. *Onderstepoort Journal of Veterinary Research*. 2019;86(1).
22. Kabir J, Umoh V, Audu-okoh E, Umoh J, Kwaga J. Veterinary drug use in poultry farms and determination of antibiotic drug residues in commercial eggs and slaughtered chicken in Kaduna State, Nigeria. *Food Control*. 2004;15(2):99-105.
23. Lawal J, Jajere S, Geidam Y, Bello A, Wakil Y, Mustapha M. Antibiotic Residues in Edible Poultry Tissues and Products in Nigeria: A Potential Public Health Hazard. *International Journal of Animal and Veterinary Advances*. 2015;7(3):55-61.
24. Lutful-Kabir S. Avian Colibacillosis and Salmonellosis: A Closer Look at Epidemiology, Pathogenesis, Diagnosis, Control and Public Health Concerns. *International Journal of Environmental Research and Public Health*. 2010;7(1):89-114.
25. Adebawale O, Adeyemo O, Awoyomi O, Dada R, Adebawale O. Antibiotic use and practices in commercial poultry laying hens in Ogun State Nigeria. *Revue d'élevage et de médecine vétérinaire des pays tropicaux*. 2016;69(1):41.
26. Ogunleye AO, Oyekunle MA, Sonibare AO. Multidrug-resistant *Escherichia coli* isolates of poultry origin in Abeokuta, South Western Nigeria. *Vet. Arh.*, 2008;78(6):501-509.
27. Sirdar M, Picard J, Bisschop S, Jambalang A, Gummow B. A survey of antibiotic residues in table eggs in Khartoum State, Sudan, 2007–2008. *Onderstepoort J Vet Res*. 2012;79(1).
28. World Health Organization (WHO). The 2019 WHO AWaRe classification of antibiotics for evaluation and monitoring of use. Geneva: World Health Organization; 2019. (WHO/EMP/IAU/2019.11). License: [CC BY-NC-SA 3.0 IGO](https://creativecommons.org/licenses/by-nc-sa/3.0/igo/).
29. Beyene T. Veterinary Drug Residues in Food-animal Products: Its Risk Factors and Potential Effects on Public Health. *Journal of Veterinary Science & Technology*. 2015;07(01).
30. Omeiza GK, Ajayi IE, Ode OJ. Assessment of antibiotic drug residues in beef in Abuja, the Federal Capital Territory, Nigeria. *Vet Ital*. 2012;48(3):283-289.
31. Amosun, EA.,.Antibiotic resistance among commensals *Escherichia coli* from cattle faeces and beef in Ibadan, Nigeria. *African journal of biotechnology*. 2012;11(58).
32. Adesokan H, Akanbi I, Akanbi I, Obaweda R. Pattern of antibiotic usage in livestock animals in south-western Nigeria: The need for alternative plans. *Onderstepoort J Vet Res*. 2015;82(1).

33. Olatoye I, Ehinmowo A. Oxytetracycline residues in edible tissues of cattle slaughtered in Akure, Nigeria. *Nigerian Veterinary Journal*. 2011;31(2).
34. Adesokan H, Agada C, Adetunji V, Akanbi I. Oxytetracycline and penicillin-G residues in cattle slaughtered in south-western Nigeria: Implications for livestock disease management and public health. *Journal of the South African Veterinary Association*. 2013;84(1).
35. Geidam Y, Ibrahim U, Grema H, Sanda K, Suleiman A, Mohzo D. Patterns of Antibiotic Sales by Drug Stores and Usage in Poultry Farms: A Questionnaire-Based Survey in Maiduguri, Northeastern Nigeria. *Journal of Animal and Veterinary Advances*. 2012;11(16):2852-2855.
36. Zinsstag J, Schelling E, Waltner-Toews D, Tanner M. From “one medicine” to “one health” and systemic approaches to health and well-being. *Preventive Veterinary Medicine*. 2011;101(3-4):148-156.

Supplementary Data

Table S1. Frequency and proportion of respondents' knowledge of ABR (n=125).

Awareness of ABR	No. of respondents (%)
No	38 (30.4)
Yes	87 (69.6)
Can antibiotics be used to treat viral, fungal, or parasitic infections in birds?	
No	82 (65.6)
Yes	43 (34.4)
Can ABR pathogens in birds affect man?	
No	86 (68.8)
Yes	39 (31.2)
Can poultry be resistant to drugs?	
No	35 (28)
Yes	90 (72)
Does ABR make treatment difficult in birds?	
Don't know	62 (49.6)
No	14 (11.2)
Yes	49 (39.2)

Table S2. Frequency and proportion of respondents' attitude towards prudent antibiotic usage in poultry (n=125).

Do you believe there is excessive antibiotic usage in birds?	No. of respondents (%)
No	110 (88)
Yes	15 (12)
Do you stop treatment when your birds have shown improvements?	
No	77 (61.6)
Yes	48 (38.4)
Did you get antibiotic prescription from a vet?	

No	64 (51.2)
Yes	61 (48.8)
Do you observe withdrawal period of antibiotics?	
No	13 (10.4)
Yes	112 (89.6)
Do you discard eggs during antibiotic therapy?	
No	125 (100)
Yes	0 (0)
Do you	

Table S3. Frequency and proportion of respondents' perceptions on ABU and ABRin poultry (n=125).

Is ABR a major problem in Nigeria?	No. of respondents (%)
No	107 (85.6)
Yes	18 (14.4)
Only vets should be allowed to prescribe antibiotics	
1 (Strongly disagree)	17 (13.6)
2 (Disagree)	20 (16)
3 (Neither agree nor disagree)	29 (23.2)
4 (Agree)	40 (32)
5 (Strongly agree)	19 (15.2)
Farmers must reduce antibiotic use	
1 (Strongly disagree)	64 (51.2)
2 (Disagree)	20 (16)
3 (Neither agree nor disagree)	9 (7.2)
4 (Agree)	21 (16.8)
5 (Strongly agree)	11 (8.8)
Proper vaccination will reduce dependence on antibiotics	
1 (Strongly disagree)	29 (23.2)
2 (Disagree)	31 (24.8)
3 (Neither agree nor disagree)	5 (4)
4 (Agree)	5 (4)
5 (Strongly agree)	55 (44)
Antibiotic resistance can only affect farms that use antibiotics	
No	90 (72)
Yes	35 (28)
Antibiotics should only be prescribed when needed	
1 (Strongly disagree)	26 (20.8)
2 (Disagree)	6 (4.8)
3 (Neither agree nor disagree)	23 (18.4)

4 (Agree)	10 (8)
5 (Strongly agree)	60 (48)
There is nothing I can do to stop antibiotic resistance	
1 (Strongly disagree)	0 (0)
2 (Disagree)	7 (5.6)
3 (Neither agree nor disagree)	62 (49.6)
4 (Agree)	17 (13.6)
5 (Strongly agree)	39 (31.2)
Is hand hygiene important for poultry farmers?	
Don't know	23 (18.4)
No	0 (0)
Yes	102 (81.6)

Table S4. Analysis of demographic characteristics as factors influencing knowledge, attitude, and perception levels of poultry farmers in Kwara state.

Outcome variable	Variable	Referent	OR (95% CI)	p-value	OR (95% CI)	p-value			
Univariate analysis									
Awareness of ABR	Age	18 years	19 years and above	1.044 (1.0110, 1.0780)	0.009	1.1072 (1.0550, 1.1619) < 0.01			
	Level of Education	Secondary	Tertiary	0.2125 (0.0690, 0.6547)	0.007	0.0445 (0.0104, 0.1904) < 0.01			
	Gender	Female	Male	0.7843 (0.3644, 1.6880)	0.534	-			
	Population of birds	100-499	500-1000	1.5826 (0.6825, 3.6697)	0.494	-			
		>1000		1.8261 (0.4916, 6.7828)					
Multivariate analysis									
Knowledge level of ABR	Age	18 years	19 years and above	1.1295 (1.0799, 1.1814)	< 0.01	1.0903 (1.0381, 1.1452) 0.033			
	Level of Education	Secondary	Tertiary	7.8269 (3.2693, 18.7381)	< 0.01	2.1848 (0.7400, 6.4507) < 0.01			
	Gender	Female	Male	8.5105 (3.0339, 23.8732)	< 0.01	4.1776 (1.1805, 14.7837) 0.027			
	Population of birds	100-499	500-1000	9.4551 (3.7928, 23.5707)	< 0.01	4.5306 (1.5566, 13.1863) 0.021			
		>1000		1.19954E+07 (0.0000, *)		1.03325E+06 (0.0000, 2.08674E+297)			
Occupation									
Farmers									
Other professions									
No. of workers									
1									
2 and above									
1.1132 (0.9049, 1.3696)									
0.310									

Table S5. Analysis of demographic characteristics as factors influencing knowledge, attitude, and perception levels of poultry farmers in Kwara state.

Outcome variable	Variable	Referent	OR (95% CI)	p-value	OR (95% CI)	p-value
Univariate analysis						
Attitude towards ABR	Age	18 years	19 years and above	1.034 (0.9370, 0.9981)	0.038	-
	Level of Education	Secondary	Tertiary	0.5370 (0.2185, 1.3200)	0.175	-
	Gender	Female	Male	1.2987 (0.3632, 1.6479)	0.506	-
	Occupation	Farmers	Other professions	0.6004 (0.2780, 1.2965)	0.194	-
	No. of workers	1	2 and above	1.1129 (0.9149, 1.3781)	0.267	-
	Population of birds	100-499	500-1000	0.7407 (0.3086, 1.7778)	0.705	-
			>1000	0.6173 (0.1777, 2.1445)		
Perception of ABR	Age	18 years	19 years and above	0.8865 (0.8484, 0.9262)	< 0.01	0.9020 (0.8623, 0.9435)
	Level of Education	Secondary	Tertiary	0.1304 (0.0547, 0.3108)	< 0.01	0.2818 (0.1020, 0.7788)
	Population of birds	100-499	500 -1000	0.5176 (0.2284, 1.1732)	0.062	-
			>1000	0.1681 (0.0334, 0.8463)		
	Gender	Female	Male	1.2987 (0.3632, 1.6479)	0.506	-
	Occupation	Farmers	Other professions	0.5133 (0.2271, 1.1606)	0.109	-
	No. of workers	1	2 and above	0.9394 (0.7708, 1.1448)	0.535	-