
Review

Elevated CO₂ and Water Stress in Combination in Plants: Brothers in Arms or Partners in Crime?

Arun K. Shanker*, Deepika Gunnapaneni, Divya Bhanu, M. Vanaja, M. Jyothilakshmi, S.K. Yadav, M. Prabhakar and V.K. Singh

ICAR- Central Research Institute for Agriculture, Santoshnagar, Saidabad P.O, Hyderabad – 500059, India

*Corresponding author: arunshank@gmail.com

ORCID - <https://orcid.org/0000-0001-5635-878X>

Abstract: The changing dynamics in climate is the primary and important determinant of agriculture productivity. The effects of this changing climate on overall productivity in agriculture can be understood when we study the effects of individual components contributing to the changing climate on plants and crops. Elevated CO₂ (eCO₂) and drought due to high variability in rainfall is one of the important manifestations of the changing climate. There is a considerable amount of literature that addresses climate effects on plant systems from molecules to ecosystems. Of particular interest is the effect of increased CO₂ on plants in relation to drought and water stress. As it is known that one of the consistent effects of increased CO₂ in the atmosphere is increased photosynthesis, especially in C₃ plants, it will be interesting to know the effect of drought in relation to elevated CO₂. The potential of elevated CO₂ ameliorating the effects of water deficit stress is evident from literature suggesting that these two are agents are brothers in arms protecting the plant from stress rather than partnering in crime, specifically water deficit when in isolation. The possible mechanisms by which this occurs will be discussed in this minireview. Interpreting the effects of short-term and long-term exposure of plants to elevated CO₂ in the context of ameliorating the negative impacts of drought will show us the possible ways by which there can be effective adaption to crops in the changing climate scenario.

Keywords: elevated CO₂; drought; photosynthesis; transpiration rate; stomatal conductance; C₄ enzymes; malate; water deficit stress; abscisic acid

1. Introduction

Agriculture is one of the dominant drivers of change in the Anthropocene era, at present about 11 per cent which is about 1.5 billion ha of the total land surface area is used for the production of crops which is about 36 per cent of the land which is suitable for agriculture [1]. Agriculture, on one hand, is affected by the changing climate and paradoxically also is contributing to it. The changing dynamics in climate are the primary and important determinant of agriculture productivity. The effects of this changing climate on overall productivity in agriculture can be understood when we study the effects of individual components contributing to the changing climate on plants and crops.

There is a continuing need to feed the growing population and globally, the human population of 7.2 billion in mid-2013 is expected to increase to almost 8.1 billion in 2025, and to further grow to 9.6 billion in 2050 [2]. Food security in terms of food availability is imperative in such a scenario. Changing climate is a reality and slowly we are learning to adapt to it, also we are in the process of devising mitigation strategies so that we can put the brakes on the changes which in general are harmful. The competition for natural resources like land, water and energy will keep growing at a pace with which it would be difficult for us to catch up unless we have sound strategies in place to adapt to the harmful effects of changing climate and further mitigate the effects of changing climate.

The effects of changing climate on particular areas specifically agriculture are difficult to predict with a great degree of accuracy although the overall effects are known and understood. Reports indicate that global average temperatures have increased by about 1 °C since the pre-industrial era and that anthropogenic warming is adding around 0.2 °C to global average temperatures every decade [3]. The global CO₂ in the atmosphere reached 407 ppm in 2018 [4]. Given the current rate of generation of CO₂, it can be expected that it will exceed 600 ppm by the end of this century [5]. The levels of greenhouse gas (GHG) are changing rapidly, CO₂ concentration in the atmosphere can directly affect the growth and development of vegetation in general and it is indirectly affecting plant growth due to seasonality and variability in rainfall it causes.

Elevated CO₂ and drought due to low variability in rainfall are important manifestations of the changing climate. There is a considerable amount of literature that addresses these aspects in terms of effects on plant systems from molecules to ecosystems. Of particular interest is the effect of increased CO₂ on plants in relation to drought and water stress. Increases in the source of carbon can have favourable effects on plants in relation to their growth and development, this can be all the more pronounced in the presence of optimum to high levels of nutrients in the soil and increased water availability. These effects may be of short duration and can vary according to the photosynthetic metabolism of the plants like in C₃, C₄, CAM and C₃ - C₄ intermediate plants. In addition, there are studies which show that C₃ crops show increased growth and yield under eCO₂ when grown under both wet and dry growing conditions. C₄ crops show increased growth and yield only under dry growing conditions and drought leads to stomatal limitations of C₃ and C₄ crops and is alleviated by eCO₂.

The forecasts for the coming decades have projected varying changes in precipitation that can result from the increasing frequency of droughts and floods [6]. Drought is one of the important abiotic stresses in the present changing climate scenario and the study of the mechanism by which it affects plant metabolism, growth and development is of paramount importance. In the past decade, global losses in crop production due to drought totalled ~\$30 billion [7]. The loss in crop production due to drought in the past ten years has been close to about 30 billion and it is estimated that about 5 billion people will be in water-scarce regions of the world by 2050, emphasising the importance of studying all the facets of drought and plant growth. Interestingly there are studies where crops are grown under field conditions, the positive impact of elevated atmospheric CO₂ concentrations on productivity was found to be significantly stronger under soil water limitation than under potential growth conditions, as reported in [8] for cotton, [9] for wheat, [10] for alfalfa and also for temperate pasture species [11]. There are also evident interactive effects of elevated CO₂ and other environmental conditions which are indicative of changing climate like drought, heat and other stresses which invariably accompany elevated CO₂ conditions in the atmosphere. The importance of understanding this complex relationship is imperative in a high CO₂ atmosphere which is envisaged in future to counter the effects of changing climate. As it is known that one of the consistent effects of increased CO₂ in the atmosphere is increased photosynthesis, especially in C₃ plants, it will be interesting to know the effect of drought in relation to elevated CO₂. The possible mechanisms by which this occurs will be discussed in this minireview. Interpreting the effects of short-term and long-term exposure of plants to elevated CO₂ in the context of ameliorating the negative impacts of drought will show us the possible ways by which there can be effective adaption to crops in the changing climate scenario.

2. Water relations, transpiration and stomatal conductance

2.1. Stomatal Dynamics

Elevated CO₂ concentration is known to mitigate the effects of drought stress, in a study on *Populus* spp. and *Salix* spp. by [12] it was found that when these two species were grown in ambient (350 $\mu\text{mol mol}^{-1}$) or elevated (700 $\mu\text{mol mol}^{-1}$) predawn water potential reduced as water stress increased as against midday water potential which did not show

any changes. The changes observed were 0.1 MPa at predawn and 0.2 MPa at mid-day. Increased elasticity of the cell wall is usually observed when there are altered water relations. These cellular changes allow the plant to maintain higher turgor at lower water potentials and tissue water content. The mitigating effect of higher CO₂ was by increasing ψ_p at the same levels ψ_w which can result in osmotic adjustment. This mechanism of osmotic adjustment can improve plant metabolism or at least maintain plant metabolism at optimal levels resulting in acclimation to drought. Stomatal dynamics drive the carbon uptake during water deficit stress and when there is accompanying stress like short-term elevated CO₂, the role of stomatal limitation in the assimilation of carbon may reduce with a reduction in photorespiration and increase in the partitioning of soluble sugars and increase in water use efficiency.

eCO₂-mediated regulation of stomatal conduction and transpiration rate is mainly by regulating stomatal aperture as a short-duration response [13-15] and other long-duration morphological modifications like changes in stomatal density [16,17]. Varied crop-specific responses were also seen [17] in stomal density where eCO₂ increased the density of stomata in maize whereas the same decreased in Amarnath. It is interesting to note here the differences in dicot and monocot response of both the C₄ crops. The general response observed in both the C₄ crops is because under water deficit conditions C₄ crops are better performing under elevated CO₂ as they have a CO₂ concentrating mechanism, this mechanism favours optimum photosynthesis even under lower stomatal conductance, and they can close their stomata and still perform the dark reaction with an optimum amount of CO₂. On the other hand, the differences between dicot and monocot C₄ plants under elevated CO₂ may be due to the higher degree of suberization in the kranz anatomy specifically in the NADP – ME subtypes which are not seen in the dicots specifically in the NAD – ME subtype [17].

Studies on stomatal density have been indecisive in their outcomes as to what exactly is governing the decrease and increase in the density under stress conditions, although a large body of evidence says that it is one of the key morphological traits that regulates transpirational flux resistance in the leaf and conductance of stomata under eCO₂. The underlying mechanism is shifting the balance in favour of CO₂ uptake by increasing it under water loss conditions. On the other hand, a recent study has also suggested that stomatal density may be equally or more affected by temperature, specifically the large continental-scale geographical variations with an interplay of precipitation [18].

The mechanism of guard cell sensing of CO₂ especially in enriched conditions and this sensing playing a role in the turgor dynamics of the cells has gained much acceptance in recent times, the support for this comes from the fact the CO₂ itself is lipophilic and can easily diffuse across membranes and also move through mass flow across aquaporins. The mechanism is explained by the triggering of CO₂ of the efflux channels of K⁺ out which in turn increases the water potential inside the cell and this results in water moving out and in effect resulting in stomatal closure [19,20].

2.2. The ABA conundrum

Abscisic acid (ABA) is mainly involved in the regulation of many important physiological processes in the plant at the cellular level. ABA synthesis activates many types of countering mechanisms in plants under stress among which the main mechanism is the stomatal movement. Opening and closing are regulated in such a way that there is minimum loss of water during water deficit conditions [21,22]. The interplay of ABA and eCO₂ has been of interest to researchers as some of the mechanism by which they regulate stomatal dynamics seems to be the same.

We have seen that eCO₂ can mitigate drought-induced stress in plants through osmotic adjustment, changes in turgor pressure and also changes in root shoot ratio and the mechanism here is higher hydraulic conductance induced maintenance of higher relative water content (RWC). This is in addition to optimum water status being maintained by hydraulic conductance. On the other hand, when there is an interaction of eCO₂ with

drought, we are faced with the question as to what exactly is contributing to the stomatal dynamics, is it the eCO₂-induced changes in the stomata or is it the drought-induced ABA production that is instrumental or is it an action of both these agents in tandem.

The mechanism and the effect become complex when we see that both ABA and eCO₂ induce stomatal closure and in the case of ABA it is reasoned that closure is to prevent excessive loss of water during stress and in the case of eCO₂ the reason for the induced closure is debated. The complexity further increases when we see that under water stress conditions in eCO₂ there can be a combined action of both ABA and eCO₂. Further, taking the complexity to next level is the differential effects of eCO₂ seen in C₃ and C₄ plants where the responses are distinct and conserved within the photosynthetic types [19].

Two Tomato genotypes, one of them being a mutant deficient in ABA were tested for responses of hydraulic conductance at eCO₂ by [23], they found that a reduction in the transpiration rate and a concomitant increase in the water use efficiency (WUE) was seen in the wild type and not in the mutant, clearly indicating a role of ABA in this response. On the other hand, both in the mutant and the wild type, increased water use and osmotic adjustment was seen, this shows us that plant water consumption which also includes water transpired is not entirely controlled by ABA. This also shows that osmotic adjustment as a response to water stress can have several other metabolic players and can occur even in the absence of ABA. It is possible that the protective role of ABA under stress is regulated by a higher concentration of CO₂ and is manifested in higher WUE and reduced transpiration rate. It is generally thought that the eCO₂-mediated closure of stomata and opening of stomata is independent of the ABA pathway, on the other hand, some signalling components of the ABA pathway have been implicated to work in tandem, suggesting that some of the components of the regulatory mechanism are shared [24].

The response triggered by both these agents eCO₂ and water deficit is interconnected where ABA is shown to modulate and also regulate the effect of eCO₂. Water deficit stress is known to have a stronger effect on stomatal conductance as compared to eCO₂ and when in combination with water deficit stress causes a larger decrease in the stomatal conductance which could be an additive effect.

We see here that ROS is a necessary intermediate for ABA-mediated stomatal action in both eCO₂ and water deficit and while ROS is a well-known response under water stress. It is not so in eCO₂, so the condition of ROS being a necessary intermediate for stomatal dynamics under sole eCO₂ throws up some mechanistic challenges as to how this condition is satisfied or is there an alternative mechanism. This question to an extent justifies the certain degree of controversy that exists in the convergence of ABA and CO₂ signalling [25].

SLAC1 is a membrane protein that is multispanning and is mainly expressed in the guard cells and has an important role in the regulation of ion homeostasis in the cell and also is involved in S-type anion currents. It is a ubiquitous protein for effecting stomatal closure under various environmental signals like eCO₂, water deficit stress, ozone, light regimes and many more. Studies have shown that SLAC1 activity loss due to mutation continues to affect CO₂ responsiveness in stomatal closure and does not affect the same way under ABA, this suggests the presence of an ABA independent signalling network under eCO₂ conditions to cause stomatal closure. This adds to the intrigue in the signalling response and possible answers can be found when we can characterize the full complement of guard cell signalling sensors [26-30]. The role of guard cell chloroplasts in regulating CO₂ has also been extensively studied, they are not directly involved in the control of stomatal closure as induced by CO₂, it is controlled by the conversion of CO₂ to protons by carbonic anhydrases with HCO₃⁻ being the primary signalling molecules bringing about changes in the proton concentration and as a result, controlling the opening and closure of the stomata [31-34].

We already know that the lower the concentration of CO₂ the more the opening of stomata and as it goes higher the stomata start to close, CO₂-induced closure is mediated by Ca²⁺ and protein phosphorylation, and the specific phosphorylation events are set into motion by signal transduction by Calcium-dependent protein kinases (CPKs) and

calcineurin- B-like proteins (CBLs) with the secondary messenger being Ca^{2+} [35]. Ca^{2+} also has an ABA modulated and accelerated response, hence Ca^{2+} transporters and proteins may have a twin function connected to both eCO_2 and ABA [36-38]. Recent research has shown a role for both eCO_2 and ABA in stomatal closure.

The common pathway or overlap or sometimes called the convergence point in the mechanism of stomatal closure involves three different events. The first is the signal perception by SLAC1 of HCO_3^- where there is an involvement of several protein kinases and this signalling activates the SLAC1 anion channel. The signalling is downstream of the Open Stomata 1 and Sucrose non-fermenting related Kinase (1OST1/SnRK1) pathway [39,40,15]. The mechanistic differences in the eCO_2 -mediated stomatal closure and ABA-mediated stomatal closure is shown in Fig.1.




Figure 1. A simplified model of stomatal closure effected by eCO_2 (a) and ABA (b) with commonality and convergence in the mechanism shown in green. Several Aquaporins facilitate the entry of CO_2 in guard cells. Plasma membrane intrinsic protein (PIP2;1) aquaporin that facilitates water transport across the cell membrane and carbonic anhydrases (b CA4 and b CA 1) interact leading to the increased formation of Bicarbonate (HCO_3^-). The multidrug and toxic compound extrusion (MATE)-type transporter RESISTANT TO HIGH CARBON DIOXIDE 1 (RHC1) gene product senses HCO_3^- signalling. Carbon Dioxide and Bicarbonate together act as signal transduction molecules. Under eCO_2 the possible action would be the activation of MPK12 and MPK 4 resulting in the inhibition of expression of protein kinase HIGH LEAF TEMPERATURE1 (HT1). When ABA enters the guard cells, in the ABA-mediated closure PYRABACTIN RESISTANCE1 (PYR1)/PYR1-LIKE

(PYL)/REGULATORY COMPONENTS OF ABA RECEPTORS (RCAR) interacts with Type 2C protein phosphatases (PP2Cs) and inhibits them. The proton translocating ATPase is inhibited in the process by ABA and this prevents proton entry into the guard cells and regulated its pH. There is a subsequent release of Ca²⁺-independent protein kinases (SnRK2s). SLAC1 ion channel is phosphorylated by the activation of SnRK2 and calcium dependant protein kinases (CPK). The convergence step under both eCO₂ and ABA is the activation of Rapid type ion channel aluminium-activated malate transporter 12/quickly activating anion channel 1 (ALMT12/QUAC1) which leads to the turgor dynamics and K⁺ ion efflux and resultant stomatal closure.

2.3. Water relations

In a study with field experiments and process-based simulations [41] the authors have shown that CO₂ enrichment contributes to decreased water stress and also contributed to higher yields of maize under restricted water conditions. They showed from their studies that elevated CO₂ decreases transpiration without any effect on soil moisture at the same time it increases evaporation. Modelling has shown that water stress is reduced to an extent of 37 per cent under elevated CO₂, a simulated increase in stomatal resistance being the reason for this.

Some of the effects of water stress in combination with elevated CO₂ can be understood when we see the effects observed in Free Air CO₂ enrichment (FACE) experiments. In maize elevated CO₂ reduces transpiration and this, in turn, contributed to the increase in soil moisture and evaporation. In a simulated study [41] it was seen that transpiration was reduced by 22 per cent in the first year of the experiment. In another study[42] the authors showed that in a FACE experiment transpiration in maize was reduced significantly under 550 ppm CO₂ concentration. Daily sap flow and vapour pressure deficit (VPD) of maize were investigated [43], it was seen that whole-plant transpiration was reduced by 50 per cent in drought as compared to wet in ambient CO₂ concentrations and 37 per cent reduction was observed in elevated CO₂ concentration of 550 ppm. Enrichment of CO₂ did not affect sap flow under drought and a 20 per cent decrease was seen under wet conditions. Maize under elevated CO₂ had a higher transpiration rate which was due to lower sap flow in the preceding period when plant-available soil water was minimum, this shows that reduction in canopy transpiration by elevated CO₂ can delay the effects of water stress and can contribute to increased plant biomass production.

Another study [44] on the physiological response of two C₃ and C₄ mechanisms syndromes, Napier grass (*Pennisetum purpureum* Schumach × *Pennisetum glaucum* (L.) R. Br) and hydric common reed grass (*Phragmites australis* (Cav.) Trin. Ex Steud) under water stress and elevated CO₂ it was seen that there was a general response of increase in photosynthesis, reduced leaf water potential and increase in transpiration in both the grass species. A contrasting response was seen in the two types of grass to elevated CO₂ and water stress, the difference in the species response was due to the stomatal characteristics as evident by the changes in transpiration rate and osmotic adjustment. Water status adjustment by modification of xylem anatomy and hydromolytic properties is a mechanism found in many plants, its relationship with the observed effect of elevated CO₂ to increase plant water potential via reduced stomatal conductance and water loss has been studied [45]. One of the known adaptations to water stress by plants is to maintain high water potential and turgor pressure under water-deficient conditions. The authors saw in their study that water deficit significantly decreased xylem vessel diameter, conduit roundness and stem cross-section area, it was seen that these impacts of water deficit were relieved at elevated CO₂. In another study [46] where the adverse effects of the drought were studied on soyabean under elevated CO₂, the authors found that elevated CO₂ increased WUE contributing to countering drought, but they did not find any positive effects on osmotic adjustments.

The effects of Elevated CO₂ individually and in combination with a water deficit in soyabean were studied [47]. In instantaneous water stress treatment, elevated CO₂ reverted the expression of genes related to stress, transport and nutrient deficiency that was induced by water stress, the interaction of drought and elevated CO₂ affected the

expression of genes with physiological and transcriptomic analysis showing that elevated CO₂ can mitigate the negative effects of water stress in soyabean roots.

3. Dry matter production

3.1. Photosynthesis and growth

In addition to understanding the acclimation pattern of plants under a combination of water stress and elevated CO₂, future yield prediction can also be done under the changing climate scenario from precise data on the effects of elevated CO₂ and drought on biomass and soil water conditions. Growth modelling under these conditions has contributed to our knowledge of these effects. Under sufficient water supply, C₃ crops recorded increased yield under elevated CO₂ whereas C₄ crops did not show much change in the yield. A 10 – 15 per cent increase in biomass has been seen in C₃ crops under FACE experiments due to the CO₂ fertilizing effect [48,49], on the other hand, C₄ crops maize and sorghum did not respond similarly under water sufficient conditions [50,51]. In a study [52], it was seen that adverse effects of heat and drought were alleviated by improved water relations under elevated CO₂. The authors also saw that the mechanism of photosynthesis reduction under the combination of heat and drought was due to increased drying of soil and a decrease in stomatal conductance.

In C₃ plants, the most prominent effect is the increased photosynthesis due to the higher availability of CO₂ to rubisco and also due to reduction in photorespiration [53,54]. The effects of eCO₂ on C₃ plants have been widely studied and from a recent comprehensive meta-analysis [55] it is seen that under eCO₂ that leaf biomass per unit leaf area (LMA) increases with a slight decrease in leaf N content. The effect on photosynthesis is mainly due to higher concentrations of CO₂ in the vicinity of Rubisco and thus an increased affinity and reduced photorespiration. Some of the other effects were, an almost doubling of photosynthetic rate at higher concentrations of CO₂ (1000 ppm range) with a halving of stomatal conductance and transpiration per unit leaf area [55], this was seen when the stomatal density as such was not affected, implying that the reduced transpiration is due to closure of stomata (the mechanism of which is discussed in another section in this review). This increase in photosynthesis accompanied by reduced transpiration leads to higher WUE which is also a characteristic of C₄ plants under eCO₂ and it is one of the reasons for eCO₂ under water deficit conditions being less detrimental to the plant as against water-deficit stress in isolation. A schematic representation of the interconnected effects of both eCO₂ and water deficit stress in isolation and in combination in both C₃ and C₄ crops is given in Fig.2.




Figure 2. Schematic representation of individual and interactive effects of water deficit stress and eCO₂ in C3 and C4 plants.

In a study on Macauba palm [56] the authors investigated the effects of elevated CO₂ and drought on photosynthesis, they found that at elevated CO₂ the plants were capable of recovering more from water stress due to increased Rubisco carboxylation rate and electron transport rate thus preventing reduction in total dry matter production. The authors noted that drought and increased CO₂ affected stem length and total dry matter production, it was seen that at elevated CO₂ there was no reduction in stem length and total biomass due to drought.

In coffee, it was seen by authors [57] that at 723 ± 83 ppm concentration of CO₂ for a period of seven months s increased biomass accumulation even under water deficit treatments with reduced rates of photorespiration and oxidative pressure under drought. The plants under drought and elevated CO₂ showed high respiratory carbon flux which is high respiration rates and also an energy status that supported increased root growth under drought. These results show a new mitigating method of elevated CO₂ for the maintenance of photosynthetic performance under drought. Other studies have shown that in soyabean [58] drought effect on photosynthesis was not alleviated by elevated CO₂, the authors found that net photosynthetic rate and chlorophyll b content reduced under drought and elevated CO₂. In another study [48] evaluated biomass accumulation in long-term experiments under elevated CO₂ and drought and the autors reported that there was a multiple response pattern and they suggested long-term experiments to access the future impact of climate change. One of the ways eCO₂ increases the biomass is by increasing Leaf Area Index (LAI) at early developmental stages leading to increased utilization of incident radiation and in turn higher carbon fixation [59,60].

Among the growth parameters relative growth rate (RGR), net assimilation rate (NAR) and leaf area ratio (LAR) are the key parameters that have shown consistent increase due to eCO₂. Under water deficit, we see that all these parameters are negatively

affected and the positive effect of CO₂ is shown to protect the plants from adverse effects of drought in isolation. Under eCO₂ the higher NAR is compensated by reduced LAR which in turn is caused by the decrease in specific leaf area (SLA), this has been consistent with all the plant parts above ground and below ground [55,61,62].

The observed negative response or lack of acclimation of photosynthetic carbon fixation to long-term eCO₂ can be due to enrichment-induced disruption of RuBP and Pi regeneration in the leaves which can alter photosynthetic rates effectively lowering it down. This long-term effect has not been studied well for reasons mainly that most experiments with eCO₂ are for two crop seasons for 3 – 5 years which does not simulate the environmentally relevant concentration of CO₂ over a long term which constitutes the future scenario of climate change.

3.2. Malate Maelstrom

Some of the major difference in the response to eCO₂ and drought among plants is due to the operation of different types of photosynthetic pathways in plants namely the C₃ and C₄ pathways. The C₄ plants are adapted to low concentrations of CO₂ in the atmosphere as they have a special anatomy called Kranz anatomy which helps them concentrate CO₂ within the cells and reduce the oxygenase activity of RuBPcase thus reducing photorespiration and increasing photosynthesis. Consequently, the C₄ plants are not benefited from increased CO₂ concentrations in the atmosphere as are C₃ plants. On the other hand, the temperature optimum for C₄ plants is high and so under eCO₂ and drought conditions, this adaptability confers a degree of tolerance to these plants. The main enzymes of the C₄ plant that play an important role in photosynthesis are the decarboxylating enzymes NAD-dependent malic enzyme (NAD-ME) and NADP-dependent malic enzyme (NADP-ME). They are of diverse phylogenetic origin and they are present in mitochondrial isoforms in C₃ plant not playing a role in carbon fixation as they don't have the oxaloacetate decarboxylase (OAD) action [63-66]. In higher vascular plants the Malic enzymes are seen as a widely distributed isoforms in the plastids and mitochondria, they have multivariable functions ranging from deference, conferring tolerance to abiotic stresses, metabolic control and stomatal dynamics [65-68].

In the NAD-ME type plants, the enzyme has the main function of decarboxylation of malate to produce pyruvate, NADH and CO₂, this enzyme uses NAD⁺ and does not decarboxylate Oxalo Acetic Acid (OAA) and is present in mitochondria. Drought and C₄ enzymes specifically the NAD and NADP malic enzyme action dynamics have been well studied [67,69,70,14]. The possible mechanism can be that the entry of CO₂ is restricted under eCO₂ conditions due to reduction in stomatal density and also closure either due to eCO₂ or drought, the ME in C₄ can contribute to higher accumulation of CO₂ in the bundle sheath cells, thus increasing photosynthesis, this is in contrast with the commonly accepted view that C₄ plants don't have a distinct advantage under eCO₂ conditions. There is also evidence that ME has an important role in balancing out malic acid and also decreasing the Reactive Oxygen Species (ROS) generation and thereby offering protection against oxidative damage [71]. Considering all these aspects of Malic Enzymes, we did a sequence search, alignment and homology modelling of four different kinds of MEs to understand if there are possible structural differences in these different enzymes that can explain the functional diversity. It was seen that there were differences in the protein secondary structure (Table 1) and also distinct hairpin bend differences in the structure of the enzyme (Fig 3.).

Table 1. Secondary structure difference between selected representative NADP-ME and NAD-ME in different plants.

Enzyme	Sheets	Beta alpha beta units	Beta Hairpins	Beta bulge	Strands	Helices	Helix-helix interacs	Beta turns	Gamma turns
--------	--------	--------------------------------	------------------	---------------	---------	---------	-------------------------	---------------	----------------

NADP-ME Maize	4	5	1	0	14	33	47	36	3
NADP-ME Rice	4	5	2	0	15	33	44	33	4
NAD-ME Solanum	3	5	2	1	14	35	44	47	6
NAD-ME Amaranthus	4	5	2	1	14	36	45	39	5




Figure 3. Homology models of NADP-ME and NAD-ME enzymes with their beta hairpin bends shown in green.

The difference in the beta-hairpin motifs across these NAD-ME and NADP-ME can play an important role in the versatility of the structure and the enzyme action. Positional differences in the amino acids in the hairpins can explain substrate binding and affinity differences in these enzymes. Engineering these enzymes by CRISPR CAS genome editing techniques can be a way to change the activities of these enzymes in both C₃ and C₄ plants and it can play an important role under both eCO₂ and water deficit conditions.

4. Nutrients in a nutshell

Nutrition limitation under eCO₂ and water deficit conditions is one of the main reasons for lower dry matter production apart from the other stress-related responses. Adequate nutrient supply and an adaptive mode of transport into the plant system can alleviate these stresses in a big way. The long-term responses and short-term responses to both eCO₂ and water deficit could be very different and even opposite in their manifestation. Critical inquiry into the mechanistic process involved here will throw light on possible ways by which these stresses can be tackled. In a study on the possible adaptive response

of semi-dwarf durum wheat cultivars by physiological and molecular mechanisms [72] it was seen that elevated CO₂ and water stress increased d15N, which was cultivar dependent and the effect diminished as water stress increased. Shifts in N metabolism could reflect in decreased root-to-shoot translocation of N. The authors observed d13C increased under moderate stress irrespective of the CO₂ concentration indicative of higher water-use efficiency. Phosphoenol Pyruvate Carboxylase (PEPC) expression was increased under water stress and elevated CO₂ combination. Carbohydrates which are the substrates for PEPC increased under these stresses and this showed the role of PEPC in providing carbon skeleton for amino acid and lipid biosynthesis. It is seen that there is a transcript level coordination in C and N metabolism under a combination of water stress and elevated CO₂. The dehydrin genes DHN11 and DHN16 showed changes in expression under water stress and elevated CO₂ with genotype-dependent change in transcript levels, this shows that the interactive effects of both elevated CO₂ and water stress varies according to the genotype in wheat.

In systems where there is no limitation for nutrients and also where there is a sufficient amount of nutrients either in the natural ecosystem or managed system some species do not show any adaptation or acclimation to higher levels of CO₂ [73]. There has been ongoing research on the reasons for this lack of acclimation to elevated CO₂, especially on an ecosystem scale and the possible reasons are not clearly explained yet, although some explanations have been put forth among which is feedback inhibition in the source-sink relationship caused by increased carbohydrate loading in the leaves which can reduce photosynthesis [74].

The observed decrease in nitrate concentration in plants under eCO₂ especially in the leaves can be explained by the lower rates of transpiration also observed at high CO₂ which causes a reduced soil solute mass flow along the plant's above-ground parts starting from the roots in the soil plant atmosphere continuum [75,76]. The reduction could also be due to a down-regulation of nitrogen assimilation, differentially regulating N transporters which maybe dose-dependent [77,78].

A summary of effects of e CO₂ and drought in plants is given in Table 2

Table 2. Summary of effects of Elevated CO₂ and drought in plants.

Plant/Crop/Tree	CO ₂ concentration	Water stress imposition	Effect	Reference
Poplar (<i>Populus</i> spp.)	700 ± 50 μmol mol ⁻¹	Soil drying cycle by withholding water	Reduced Gas exchange, decreased leaf conductance, increased photosynthesis, increased transpiration efficiency	[12]
Wheat (<i>Triticum aestivum</i>)	400 μmol mol ⁻¹ , 790 μmol mol ⁻¹	Progressive restriction of water from 10 percent to 60 percent pot capacity	Reduced plant biomass, Stomatal conductance and carbon isotope signature indicated water saving strategy. PEPC expression increased	[72]
<i>Tabernaemontana divaricata</i>	1000 μmol mol ⁻¹ , 700 μmol mol ⁻¹	70 percent of field capacity (FC) for 4 days and 30 percent of FC for 4 days	Increase in stomatal conductance (g _s), plant height (PH) and plant girth (PG)	[82]
Maize (<i>Zea mays</i>)	550 μmol mol ⁻¹	Half water in water stress treatment compared to control	37 percent reduction in whole plant transpiration	[43]
Napier grass (<i>Pennisetum purpureum</i> Schumach × <i>Pennisetum glaucum</i> (L.) R. Br) and hydric common reed grass (<i>Phragmites australis</i> (Cav.) Trin. Ex Steud)	563 ± 6.7 μmol mol ⁻¹ , 541 ± 6.9 μmol mol ⁻¹ , 601 ± 9.1 μmol mol ⁻¹	Withdrawing irrigation	Increase in Photosynthesis, reduced leaf water potential and increase in transpiration	[44]
Maize (<i>Zea mays</i>)	700 μmol mol ⁻¹ , 900 μmol mol ⁻¹ , and 1,200 μmol mol ⁻¹	deficit irrigation	Decreases in stomatal conductance and reduced transpiration rate	[45]
Soyabean (<i>Glycine max</i>)	Ambient + 200 μmol mol ⁻¹	35- 45 percent of RWC	Elevated CO ₂ enhanced the resistance to drought by improving the capacity of photosynthesis and WUE in soybean leaves	[46]
<i>Pinus halepensis</i> (Aleppo pine)	867 ± 157 μmol mol ⁻¹	10 Percent Relative Substrate Water Content	Under drought, the effect of CO ₂ on WUE was pronounced, with intercellular CO ₂ being increased near stomatal closure,	[83]

Lemon (<i>Citrus limon</i>)	650 and 850 $\mu\text{mol mol}^{-1}$	leaf water potential of -3.5 MPa	Stomatal downregulation at elevated CO_2 reduced water-use but not photosynthesis.	[84]
Soybean (<i>Glycine max</i>)	800 $\mu\text{mol mol}^{-1}$	water deficit was applied by randomly removing plants out of the hydroponic solution exposing the roots to ambient- or elevated-air	Responses of soybean roots to short-term water deficit are buffered by Elevated CO_2	[47]
Cassava (<i>Manihot esculenta</i> Crantz)	750 $\mu\text{mol mol}^{-1}$	Stopping irrigation for 7 days	Elevated CO_2 reduced the negative effect of drought on transpiration, water use efficiency, all growth measures and harvest index.	[85]
Faba bean (<i>Vicia faba</i> L.)	550 $\mu\text{mol mol}^{-1}$	Water was withheld until nodulation and nodule density helped 30 percent FC	Elevated CO_2 -induced stimulation of	[86]
Andiroba (<i>Carapa surinamensis</i>)	700 $\mu\text{mol mol}^{-1}$	50 percent field capacity	Whole-plant water-use efficiency (WUE) improved under combination treatments	[87]
<i>Hymenaea stigonocarpa</i> Mart. ex Hayne, <i>Solanum lycocarpum</i> A. St.-Hil. and <i>Tabebuia aurea</i> (Silva Manso) Benth. and Hook. f. ex S. Moore	700 $\mu\text{mol mol}^{-1}$	Water stress was introduced three times during the experiment by halting irrigation 1 month before the fourth (360 days old), fifth (450 days old) and sixth (540 days old) morphophysiological surveys	Water stress decreased biomass production under high CO_2	[88]
Grapevines (<i>Vitis labrusca</i>)	800 $\mu\text{mol mol}^{-1}$	Stopping irrigation	Elevated CO_2 delayed drought effects on both net photosynthetic rate and Rubisco activity for four days, by reducing stomatal conductance, transpiration and stomatal density	[89]

<i>Brassica napus</i>	800 $\mu\text{mol mol}^{-1}$	Withholding water for 7 days	Elevated CO_2 diminished the adverse effect by improved water relations	[52]
Maize (<i>Zea mays</i>)	550, 700, and 900 $\mu\text{mol mol}^{-1}$	Deficit irrigation	Photosynthetic rate in elevated CO_2 concentrations was higher under Deficit irrigation than under regular irrigation.	[90]
<i>Acrocomia aculeata</i>	700 $\mu\text{mol mol}^{-1}$	Water withholding	Higher Rubisco carboxylation rate (V_c max) and electron transport rate (J max) contributed to recovery from drought	[56]
Cucumber (<i>Cucumis sativus</i> L.)	$800 \pm 20 \mu\text{mol mol}^{-1}$	$\psi_w = -0.05 \text{ MPa}$ and $\psi_w = -0.15$ with PEG 6000	Higher photosynthetic performance and increased grana thickness under moderate drought stress, increased palisade cells length and chloroplasts number per palisade cell under severe drought stress.	[91]

A proposed mechanistic explanation for this is also reduced photorespiration which can reduce N assimilation [79] where the authors show that chloroplast vesiculation (cv) gene expression which destabilizes photosynthesis when silenced maintained the N assimilation status in plants under e CO_2 . The key point to note here is that the total N in the plant is increased under e CO_2 because of the biomass increase seen in these conditions, this mechanism as such may not be a part of the mitigating factor when e CO_2 and water deficit are in combination as the reduction in the biomass induced by water stress is not fully compensated by e CO_2 although the combination is seen to reduce the reduction caused by water deficit stress.

Adequate Phosphorus nutrition is an ameliorative agent under water deficit stress in e CO_2 conditions. The mechanism by which P acts under e CO_2 conditions is by increasing the concentration of soluble sugars and Pi and maintaining relative water content in the leaves. In a study [80] in field peas, the authors found that stress tolerance index (STI) was higher in plants with adequate P under e CO_2 conditions under water deficit, this exemplifies the interactive role of P and e CO_2 as drought ameliorative combination. The role of P is to stabilize water relations, stomatal conductance and transpiration rate synergistically with e CO_2 . It is instrumental in decreasing stomatal conductance and concomitantly reduces water loss through transpiration thus increasing the transpiration efficiency under drought stress conditions. Total soluble sugars increase in leaves due to accumulation of leaf Pi under e CO_2 offers better disposition of plants to stave off drought-related stress than when in e CO_2 . The positive effects of Pi namely maintenance of high metabolic readiness to tackle low water status are enhanced under e CO_2 . A similar trend was seen in

Potassium (K) supplementation under water deficit conditions [81]. The authors found that K deficiency impeded the ameliorative effect of eCO₂ on drought. Nutrient sufficiency is an important prerequisite to realise the beneficial effects of eCO₂ and the drought mitigatory effect of eCO₂ is dependent on nutrient sufficiency.

5. Future perspectives

Deficit irrigation to economize water use and to induce acclimation by plant physiological adjustments is an approach that can be advocated to counter the adverse effects of changing climate, our mini-review here shows that this can be an important strategy in future agriculture under elevated CO₂ which effectively decrease the impact of low soil water on photosynthesis and in turn biomass accumulation and yield in crops. Plant water relations are mainly affected by gas exchange and stomatal physiology which in turn is affected by elevated CO₂ and drought and there is a complex manifestation when these stresses act in combination, these are the critical factors when the goal is to evolve climate ready cultivars. To device strategies for adaption in crops in agricultural systems, we have to understand and elucidate how these processes operate across a range from ecosystems to organismal level and from cellular, biochemical to the molecular level. Adaptation in agriculture to changing climate is occurring all over the world, the practices should now be based on the findings that e CO₂ under drought and water stress conditions can be effective in alleviating the effects of climate change. There is a consensus and better understanding of effects not that can be put to use for tackling climate-related effects on crop production.

One of the important facets that have come out of this mini-review is that most of the effects observed need to be looked into with a mechanistic perspective to arrive at correct inferences that can help us move ahead with the goal of evolving climate-ready cultivars. In many of the studies, the causal association are observed which needs further investigations we trust this mini-review will invigorate researchers.

Acknowledgments: The author wished to acknowledge the project of ICAR, National Innovation on Climate Resilient Agriculture for funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Alexandratos, N.; Bruinsma, J.; World agriculture towards 2030/2050: the 2012 revision. 2021.
2. United Nations. Working Paper No. ESA/P/WP.227. Department of Economic and Social Affairs, Population Division; 2013. World Population Prospects: The 2012 Revision, Key Findings and Advance Tables.
3. IPCC (Intergov. Panel Clim. Change); Global Warming of 1.5° C. An IPCC special report on the impacts of global warming of 1.5° C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty. 2018.
4. Friedlingstein, P.; Jones, M.; O'sullivan, M.; Andrew, R.; Hauck, J.; Peters, G.; Peters, W.; Pongratz, J.; Sitch, S.; Le Quéré, C.; DBakker, O.; Global carbon budget 2019. *Earth System Science Data*, 2019, 11(4), 1783-1838.
5. IPCC (2007) Climate Change 2007: Impacts, Adaptation and Vulnerability. Geneva: IPCC Secretariat.
6. Shanker, A.K.; Maheswari, M.; Yadav, S.K.; Desai, S.; Bhanu, D.; Attal, N.B.; Venkateswarlu, B.; Drought stress responses in crops. *Functional & integrative genomics*, 2014, 14(1), 11-22.
7. Gupta, A.; Rico-Medina, A.; Caño-Delgado, A.I.; The physiology of plant responses to drought. *Science*, 2020, 368(6488), 266-269.
8. Kimball, B.A.; LaMorte, R.L.; Seay, R.S.; Pinter Jr, P.J.; Rokey, R.R.; Hunsaker, D.J.; Dugas, W.A.; Heuer, M.L.; Mauney, J.R.; Hendrey, G.R.; Lewin, K.F.; Effects of free-air CO₂ enrichment on energy balance and evapotranspiration of cotton. *Agricultural and Forest Meteorology*, 1994, 70(1-4), 259-278.
9. Pinter Jr, P.J.; Kimball, B.A.; Garcia, R.L.; Wall, G.W.; Hunsaker, D.J.; LaMorte, R.L.; Free-air CO₂ enrichment: responses of cotton and wheat crops. In: Koch GW, Mooney HA, eds. Carbon Dioxide and terrestrial ecosystems San Diego, CA, USA. Academic Press, 1996, 215-264.
10. De Luis, I.; Irigoyen, J.J.; Sánchez-Díaz, M.; Elevated CO₂ enhances plant growth in droughted N₂-fixing alfalfa without improving water status. *Physiologia Plantarum*, 1999, 107(1), 84-89.

11. Clark, H.; Newton, P.C.D.; Barker, D.J.; Physiological and morphological responses to elevated CO₂ and a soil moisture deficit of temperate pasture species growing in an established plant community. *Journal of Experimental Botany*, 1999, 50(331), 233-242.

12. Johnson, J.D.; Tognetti, R.; Paris, P.; Water relations and gas exchange in poplar and willow under water stress and elevated atmospheric CO₂. *Physiologia Plantarum*, 2002, 115(1), 93-100.

13. Xu, Z.; Jiang, Y.; Jia, B.; Zhou, G.; Elevated-CO₂ response of stomata and its dependence on environmental factors. *Frontiers in plant science*, 2016, 7, 657.

14. Zhang, X.; Pu, P.; Tang, Y.; Zhang, L.; Lv, J.; C₄ photosynthetic enzymes play a key role in wheat spike bracts primary carbon metabolism response under water deficit. *Plant Physiology and Biochemistry*, 142, 2019, 163-172.

15. Habermann, E.; Dias de Oliveira, E.A.; Contin, D.R.; San Martin, J.A.; Curtarelli, L.; Gonzalez-Meler, M.A.; Martinez, C.A.; Stomatal development and conductance of a tropical forage legume are regulated by elevated [CO₂] under moderate warming. *Frontiers in plant science*, 2019, 10, 609.

16. Caldera, H.I.U.; De Costa, W.J.M.; Woodward, F.I.; Lake, J.A.; Ranwala, S.M.; Effects of elevated carbon dioxide on stomatal characteristics and carbon isotope ratio of *Arabidopsis thaliana* ecotypes originating from an altitudinal gradient. *Physiologia plantarum*, 2017, 159(1), 74-92.

17. Wei, Z.; Abdelhakim, L.O.A.; Fang, L.; Peng, X.; Liu, J.; Liu, F.; Elevated CO₂ effect on the response of stomatal control and water use efficiency in amaranth and maize plants to progressive drought stress. *Agricultural Water Management*, 2022, 266, 107609.

18. Marek, S.; Tomaszewski, D.; Źytkowiak, R.; Jasińska, A.; Zadworny, M.; Boratyńska, K.; Dering, M.; Danusevičius, D.; Oleksyn, J.; Wyka, T.P.; Stomatal density in *Pinus sylvestris* as an indicator of temperature rather than CO₂: Evidence from a pan-European transect. *Plant, Cell & Environment*, 2022.

19. Israel, W.K.; Watson-Lazowski, A.; Chen, Z.H.; Ghannoum, O.; High intrinsic water use efficiency is underpinned by high stomatal aperture and guard cell potassium flux in C₃ and C₄ grasses grown at glacial CO₂ and low light. *Journal of Experimental Botany*, 2022, 73(5), 1546-1565.

20. Du, B.; Ding, R.; Yang, X.; Du, T.; How are leaf carbon-and water-related traits coordinated acclimation to elevated CO₂ by its anatomy: A case study in tomato. *Environmental and Experimental Botany*, 2022, 104898.

21. Lim, C.W.; Baek, W.; Jung, J.; Kim, J.H.; Lee, S.C.; Function of ABA in stomatal defense against biotic and drought stresses. *International journal of molecular sciences*, 2015, 16(7), 15251-15270.

22. Bharath, P.; Gahir, S.; Raghavendra, A.S.; Abscisic acid-induced stomatal closure: An important component of plant defense against abiotic and biotic stress. *Frontiers in Plant Science*, 2021, 12, 324.

23. Li, S.; Fang, L.; Hegelund, J.N.; aLiu, F.; Elevated CO₂ modulates plant hydraulic conductance through regulation of PIPs under progressive soil drying in tomato plants. *Frontiers in plant science*, 2021, 12, 916.

24. Ma, X.; Bai, L.; Elevated CO₂ and reactive oxygen species in stomatal closure. *Plants*, 2021, 10(2), 410.

25. Murata, Y.; Mori, I.C.; Munemasa, S.; Diverse stomatal signaling and the signal integration mechanism. *Annual Review of Plant Biology*, 2015, 66, 369-392.

26. Assmann, S.M.; Jegla, T.; Guard cell sensory systems: recent insights on stomatal responses to light, abscisic acid, and CO₂. *Current opinion in plant biology*, 2016, 33, 157-167.

27. Negi, J.; Matsuda, O.; Nagasawa, T.; Oba, Y.; Takahashi, H.; Kawai-Yamada, M.; Uchimiya, H.; Hashimoto, M.; Iba, K.; CO₂ regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. *Nature*, 2008, 452(7186), 483-486.

28. Monda, K.; Mabuchi, A.; Negi, J.; Iba, K.; Cuticle permeability is an important parameter for the trade-off strategy between drought tolerance and CO₂ uptake in land plants. *Plant Signaling & Behavior*, 2021, 16(6), 1908692.

29. Dubeaux, G.; Hsu, P.K.; Ceciliato, P.H.; Swink, K.J.; Rappel, W.J.; Schroeder, J.I.; Deep dive into CO₂-dependent molecular mechanisms driving stomatal responses in plants. *Plant Physiology*, 2021, 187(4), 2032-2042.

30. Jalakas, P.; Nuhkat, M.; Vahisalu, T.; Merilo, E.; Brosché, M.; Kollist, H.; Combined action of guard cell plasma membrane rapid- and slow-type anion channels in stomatal regulation. *Plant physiology*, 2021, 187(4), 2126-2133.

31. Zhang, J.; De-oliveira-Ceciliato, P.; Takahashi, Y.; Schulze, S.; Dubeaux, G.; Hauser, F.; Azoulay-Shemer, T.; Töldsepp, K.; Kollist, H.; Rappel, W.J.; Schroeder, J.I.; Insights into the molecular mechanisms of CO₂-mediated regulation of stomatal movements. *Current Biology*, 2018, 28(23), 1356-1363.

32. Driesen, E.; Van den Ende, W.; De Proft, M.; Saeys, W.; Influence of environmental factors light, CO₂, temperature, and relative humidity on stomatal opening and development: A review. *Agronomy*, 2020, 10(12), 1975.

33. Ren, Z.; Suolang, B.; Fujiwara, T.; Yang, D.; Saito, Y.; Kinoshita, T.; Wang, Y.; Promotion and upregulation of a plasma membrane proton-ATPase strategy: principles and applications. *Frontiers in plant science*, 2021, 12.

34. Nobori, T.; Closing the gap: a plasma membrane H⁺-ATPase regulates stomatal closure. *The Plant Cell*, 2022, 1-2.

35. Schulze, S.; Dubeaux, G.; Ceciliato, P.H.; Munemasa, S.; Nuhkat, M.; Yarmolinsky, D.; Aguilar, J.; Diaz, R.; Azoulay-Shemer, T.; Steinhorst, L.; Offenborn, J.N.; A role for calcium-dependent protein kinases in differential CO₂-and ABA-controlled stomatal closing and low CO₂-induced stomatal opening in *Arabidopsis*. *New Phytologist*, 2021, 229(5), 2765-2779.

36. Jing, X.; Cai, C.; Fan, S.; Wang, L.; and Zeng, X.; Spatial and temporal calcium signaling and its physiological effects in Moso Bamboo under drought stress. *Forests*, 2019, 10(3), 224.

37. Kudla, J.; Becker, D.; Grill, E.; Hedrich, R.; Hippler, M.; Kummer, U.; Parniske, M.; Romeis, T.; Schumacher, K.; Advances and current challenges in calcium signaling. *New Phytologist*, 2018, 218(2), 414-431.

38. Richards, M.; Xu, W.; Mallozzi, A.; Errera, R.M.; Supan, J.; Production of calcium-binding proteins in *Crassostrea virginica* in response to increased environmental CO₂ concentration. *Frontiers in Marine Science*, 2018, 5, 203.

39. Hsu, P.K.; Takahashi, Y.; Munemasa, S.; Merilo, E.; Laanemets, K.; Waadt, R.; Pater, D.; Kollist, H.; Schroeder, J.I.; Abscisic acid-independent stomatal CO₂ signal transduction pathway and convergence of CO₂ and ABA signaling downstream of OST1 kinase. *Proceedings of the National Academy of Sciences*, 2018, 115(42), 9971-9980.

40. Engineer, C.B.; Hashimoto-Sugimoto, M.; Negi, J.; Israelsson-Nordström, M.; Azoulay-Shemer, T.; Rappel, W.J.; Iba, K.; Schroeder, J.I.; CO₂ sensing and CO₂ regulation of stomatal conductance: advances and open questions. *Trends in Plant Science*, 2016, 21(1), 16-30.

41. Kellner, J.; Houska, T.; Manderscheid, R.; Weigel, H.J.; Breuer, L.; Kraft, P.; Response of maize biomass and soil water fluxes on elevated CO₂ and drought—From field experiments to process-based simulations. *Global change biology*, 2019, 25(9), 2947-2957.

42. Hussain, M.Z.; VanLoocke, A.; Siebers, M.H.; Ruiz-Vera, U.M.; Cody Markelz, R.J.; Leakey, A.D.; Ort, D.R.; Bernacchi, C.J.; Future carbon dioxide concentration decreases canopy evapotranspiration and soil water depletion by field-grown maize. *Global Change Biology*, 2013, 19(5), 1572-1584.

43. Manderscheid, R.; Erbs, M.; Burkart, S.; Wittich, K.P.; Lopmeier, F.J.; Weigel, H.J.; Effects of free-air carbon dioxide enrichment on sap flow and canopy microclimate of maize grown under different water supply. *Journal of Agronomy and Crop Science*, 2016, 202(4), 255-268.

44. Mwendia, S.W.; Yunusa, I.A.M.; Sindel, B.M.; Whalley, R.D.B.; and Bruhl, J.J.; Osmotic adjustment, stomata morphology and function show contrasting responses to water stress in mesic and hydric grasses under elevated CO₂ concentration. *Photosynthetica*, 2019, 57(1), 121-131.

45. Liu, J.; Kang, S.; Davies, W.J.; Ding, R.; Elevated [CO₂] alleviates the impacts of water deficit on xylem anatomy and hydraulic properties of maize stems. *Plant, Cell & Environment*, 2020, 43(3), 563-578.

46. Wang, A.; Lam, S.K.; Hao, X.; Li, F.Y.; Zong, Y.; Wang, H.; Li, P.; Elevated CO₂ reduces the adverse effects of drought stress on a high-yielding soybean (*Glycine max* (L.) Merr.) cultivar by increasing water use efficiency. *Plant physiology and biochemistry*, 2018, 132, 660-665.

47. Bencke-Malato, M.; De Souza, A.P.; Ribeiro-Alves, M.; Schmitz, J.F.; Buckeridge, M.S.; Alves-Ferreira, M.; Short-term responses of soybean roots to individual and combinatorial effects of elevated [CO₂] and water deficit. *Plant Science*, 2019, 280, 283-296.

48. Andrensen, L.C.; Yuan, N.; Seibert, R.; Moser, G.; Kammann, C.I.; Luterbacher, J.; Erbs, M.; Muller, C.; Biomass responses in a temperate European grassland through 17 years of elevated CO₂. *Global Change Biology*, 2018, 24(9), 3875-3885.

49. Weigel, H.J.; Manderscheid, R.; Crop growth responses to free air CO₂ enrichment and nitrogen fertilization: Rotating barley, ryegrass, sugar beet and wheat. *European Journal of Agronomy*, 2012, 43, 97-107.

50. Leakey, A.D.; Uribelarrea, M.; Ainsworth, E.A.; Naidu, S.L.; Rogers, A.; Ort, D.R.; Long, S.P.; Photosynthesis, productivity, and yield of maize are not affected by open-air elevation of CO₂ concentration in the absence of drought. *Plant physiology*, 2006, 140(2), 779-790.

51. Manderscheid, R.; Erbs, M.; Weigel, H.J.; Interactive effects of free-air CO₂ enrichment and drought stress on maize growth. *European Journal of Agronomy*, 2014, 52, 11-21.

52. Diksaityte, A.; Virsile, A.; Zaltauskaite, J.; Januskaite, I.; Juozapaitiene, G.; Growth and photosynthetic responses in *Brassica napus* differ during stress and recovery periods when exposed to combined heat, drought and elevated CO₂. *Plant Physiology and Biochemistry*, 2019, 142, 59-72.

53. South, P.F.; Cavanagh, A.P.; Lopez-Calcagno, P.E.; Raines, C.A.; Ort, D.R.; Optimizing photorespiration for improved crop productivity. *Journal of Integrative Plant Biology*, 2018, 60(12), 1217-1230.

54. Dusenge, M.E.; Duarte, A.G.; Way, D.A.; Plant carbon metabolism and climate change: elevated CO₂ and temperature impacts on photosynthesis, photorespiration and respiration. *New Phytologist*, 2019, 221(1), 32-49.

55. Poorter, H.; Knopf, O.; Wright, I.J.; Temme, A.A.; Hogewoning, S.W.; Graf, A.; Cernusak, L.A.; Pons, T.L.; A meta-analysis of responses of C₃ plants to atmospheric CO₂: dose-response curves for 85 traits ranging from the molecular to the whole-plant level. *New Phytologist*, 2022, 233(4), 1560-1596.

56. Rosa, B.L.; Souza, J.P.; Pereira, E.G.; Increased atmospheric CO₂ changes the photosynthetic responses of *Acrocomia aculeata* (Arecaceae) to drought. *Acta Botanica Brasilica*, 2019, 33(3), 486-497.

57. Avila, R.T.; de Almeida, W.L.; Costa, L.C.; Machado, K.L.; Barbosa, M.L.; de Souza, R.P.; Martino P.B.; Juárez, M.A.; Marçal, D.M.; Martins, S.C.; Ramalho, J.D.; Elevated air [CO₂] improves photosynthetic performance and alters biomass accumulation and partitioning in drought-stressed coffee plants. *Environmental and Experimental Botany*, 2020, 104137.

58. Li, B.; Feng, Y.; Zong, Y.; Zhang, D.; Hao, X.; Li, P.; Elevated CO₂-induced changes in photosynthesis, antioxidant enzymes and signal transduction enzyme of soybean under drought stress. *Plant Physiology and Biochemistry*, 2020, 154, 105-114.

59. Ewert, F.; Modelling plant responses to elevated CO₂: how important is leaf area index?. *Annals of botany*, 2004, 93(6), 619-627.

60. Lemordant, L.; Gentile, P.; Swann, A.S.; Cook, B.I.; Scheff, J.; Critical impact of vegetation physiology on the continental hydrologic cycle in response to increasing CO₂. *Proceedings of the National Academy of Sciences*, 2018, 115(16), 4093-4098.

61. Qi, W.; Liu, J.; Leung, F.; A framework to quantify impacts of elevated CO₂ concentration, global warming and leaf area changes on seasonal variations of water resources on a river basin scale. *Journal of Hydrology*, 2019, 570, 508-522.

62. Jena, U.R.; Bhattacharya, S.; Swain, D.K.; Maiti, M.K.; Differential effect of elevated carbon dioxide on sucrose transport and accumulation in developing grains of three rice cultivars. *Plant Gene*, 2021, 28, 100337.

63. Rao, X.; Dixon, R.A.; The differences between NAD-ME and NADP-ME subtypes of C₄ photosynthesis: more than decarboxylating enzymes. *Frontiers in plant science*, 2016, 7, 1525.

64. Maier, A.; Zell, M.B.; Maurino, V.G.; Malate decarboxylases: evolution and roles of NAD (P)-ME isoforms in species performing C₄ and C₃ photosynthesis. *Journal of Experimental Botany*, 2011, 62(9), 3061-3069.

65. Tronconi, M.A.; Andreo, C.S.; Drincovich, M.F.; Chimeric structure of plant malic enzyme family: different evolutionary scenarios for NAD- and NADP-dependent isoforms. *Frontiers in plant science*, 2018, 9, 565.

66. Ludwig, M.; Busch, F.A.; Khoshravesh, R.; Covshoff, S.; Understanding C₄ evolution and function. *Frontiers in Plant Science*, 2021, 12.

67. Wen, Z.; Wang, Y.; Xia, C.; Zhang, Y.; Zhang, H.; Chloroplastic SaNADP-ME4 of C₃–C₄ woody desert species *Salsola laricifolia* confers drought and salt stress resistance to *Arabidopsis*. *Plants*, 2021, 10(9), 1827.

68. Begara-Morales, J.C.; Sánchez-Calvo, B.; Gómez-Rodríguez, M.V.; Chaki, M.; Valderrama, R.; Mata-Pérez, C.; López-Jaramillo, J.; Corpas, F.J.; Barroso, J.B.; Short-term low temperature induces nitro-oxidative stress that deregulates the NADP-malic enzyme function by tyrosine nitration in *Arabidopsis thaliana*. *Antioxidants*, 2019, 8(10), 448.

69. Chen, Q.; Wang, B.; Ding, H.; Zhang, J.; Li, S.; The role of NADP-malic enzyme in plants under stress. *Plant science*, 2019, 281, 206-212.

70. Correia, P.M.; da Silva, A.B.; Vaz, M.; Carmo-Silva, E.; da Silva, J.M.; Efficient regulation of CO₂ assimilation enables greater resilience to high temperature and drought in Maize. *Frontiers in plant science*, 2021, 12.

71. Sun, X.; Han, G.; Meng, Z.; Lin, L.; Sui, N.; Roles of malic enzymes in plant development and stress responses. *Plant signaling & behavior*, 2019, 14(10), 1644596.

72. Medina, S.; Vicente, R.; Amador, A.; and Araus, J.L.; Interactive effects of elevated [CO₂] and water stress on physiological traits and gene expression during vegetative growth in four durum wheat genotypes. *Frontiers in plant science*, 2016, 7, 1738.

73. Curtis, P.S.; Drake, B.G.; Whigham, D.F.; Nitrogen and carbon dynamics in C₃ and C₄ estuarine marsh plants grown under elevated CO₂ in situ. *Oecologia*, 1989, 78(3), 297-301.

74. Delucia, E.H.; Sasek, T.W.; Strain, B.R.; Photosynthetic inhibition after long-term exposure to elevated levels of atmospheric carbon dioxide. *Photosynthesis Research*, 1985, 7(2), 175-184.

75. Jin, J.; Armstrong, R.; Tang, C.; Impact of elevated CO₂ on grain nutrient concentration varies with crops and soils—A long-term FACE study. *Science of The Total Environment*, 2019, 651, 2641-2647.

76. Houshmandfar, A.; Fitzgerald, G.J.; O'Leary, G.; Tausz-Posch, S.; Fletcher, A.; Tausz, M.; The relationship between transpiration and nutrient uptake in wheat changes under elevated atmospheric CO₂. *Physiologia Plantarum*, 2018, 163(4), 516-529.

77. Adavi, S.B.; Sathee, L.; Elevated CO₂ alters tissue balance of nitrogen metabolism and downregulates nitrogen assimilation and signalling gene expression in wheat seedlings receiving high nitrate supply. *Protoplasma*, 2021, 258(1), 219-233.

78. Adavi, S.B.; Sathee, L.; Elevated CO₂ differentially regulates root nitrate transporter kinetics in a genotype and nitrate dose-dependent manner. *Plant Science*, 2021, 305, 110807.

79. Umnajitikorn, K.; Sade, N.; Rubio Wilhelm, M.D.M.; Gilbert, M.E.; Blumwald, E.; Silencing of OsCV (chloroplast vesiculation) maintained photorespiration and N assimilation in rice plants grown under elevated CO₂. *Plant, Cell & Environment*, 2020, 43(4), 920-933.

80. Jin, J.; Lauricella, D.; Armstrong, R.; Sale, P.; Tang, C.; Phosphorus application and elevated CO₂ enhance drought tolerance in field pea grown in a phosphorus-deficient vertisol. *Annals of botany*, 2015, 116(6), 975-985.

81. Asif, M.; Yilmaz, O.; Ozturk, L.; Potassium deficiency impedes elevated carbon dioxide-induced biomass enhancement in well watered or drought-stressed bread wheat. *Journal of Plant Nutrition and Soil Science*, 2017, 180(4), 474-481.

82. Thruppooyil, S.B.; Ksiksi, T.; Time-dependent stomatal conductance and growth responses of *Tabernaemontana divaricata* to short-term elevated CO₂ and water stress at higher than optimal growing temperature. *Current Plant Biology*, 2020, 22, 100127.

83. Birami, B.; Nägele, T.; Gattmann, M.; Preisler, Y.; Gast, A.; Arneth, A.; Ruehr, N.K.; Hot drought reduces the effects of elevated CO₂ on tree water-use efficiency and carbon metabolism. *New Phytologist*, 2020.

84. Paudel, I.; Halpern, M.; Wagner, Y.; Raveh, E.; Yermiyahu, U.; Hoch, G.; and Klein, T.; Elevated CO₂ compensates for drought effects in lemon saplings via stomatal downregulation, increased soil moisture, and increased wood carbon storage. *Environmental and Experimental Botany*, 2018, 148, 117-127.

85. Cruz, J.L.; LeCain, D.R.; Alves, A.A.; Coelho Filho, M.A.; Coelho, E.F.; Elevated CO₂ reduces whole transpiration and substantially improves root production of cassava grown under water deficit. *Archives of Agronomy and Soil Science*, 2018, 64(12), 1623-1634.

86. Parvin, S.; Uddin, S.; Tausz-Posch, S.; Fitzgerald, G.; Armstrong, R.; Tausz, M.; Elevated CO₂ improves yield and N₂ fixation but not grain N concentration of faba bean (*Vicia faba* L.) subjected to terminal drought. *Environmental and Experimental Botany*, 2019, 165, 161-173.

87. de Oliveira, M.F.; Marenco, R.A.; Gas exchange, biomass allocation and water-use efficiency in response to elevated CO₂ and drought in andiroba (*Carapa surinamensis*, Meliaceae). *iForest-Biogeosciences & Forestry*, 2019, 12(1).

88. Souza, J.P.; Melo, N.M.; Pereira, E.G.; Halfeld, A.D.; Gomes, I.N.; Prado, C.H.B.; Responses of woody *Cerrado* species to rising atmospheric CO₂ concentration and water stress: gains and losses. *Functional Plant Biology*, 2016, 43(12), 1183-1193.

89. da Silva, J.R.; Patterson, A.E.; Rodrigues, W.P.; Campostrini, E.; Griffin, K.L.; Photosynthetic acclimation to elevated CO₂ combined with partial rootzone drying results in improved water use efficiency, drought tolerance and leaf carbon balance of grapevines (*Vitis labrusca*). *Environmental and Experimental Botany*, 2017, 134, 82-95.

90. Li, X.; Kang, S.; Zhang, X.; Li, F.; Lu, H.; Deficit irrigation provokes more pronounced responses of maize photosynthesis and water productivity to elevated CO₂. *Agricultural Water Management*, 2018, 195, 71-83.

91. Liu, B.B.; Li, M.; Li, Q.M.; Cui, Q.Q.; Zhang, W.D.; Ai, X.Z.; Bi, H.G.; Combined effects of elevated CO₂ concentration and drought stress on photosynthetic performance and leaf structure of cucumber (*Cucumis sativus* L.) seedlings. *Photosynthetica*, 2018, 56(3), 942-952.