

Review

Bacteria DNA conjugation: from the cellular to the community level

Chloé Virolle^{1†}, Kelly Goldlust^{1†}, Sarah Djermoun^{1†}, Sarah Bigot¹ and Christian Lesterlin^{1*}

¹ Microbiologie Moléculaire et Biochimie Structurale (MMSB), Université Lyon 1, CNRS, Inserm, UMR5086, 69007, Lyon, France

[†] Co-first: Contributed equally

* Correspondence: Christian.lesterlin@ibcp.fr

Abstract: Bacterial conjugation, also referred to as bacterial sex, is a major horizontal gene transfer mechanism where the DNA is transferred from a donor to a recipient bacterium by direct contact. Conjugation is universally conserved among bacteria and occurs in a wide range of environments (soil, plant surfaces, water, sewage, biofilms and host-associated bacterial communities). Within these habitats, conjugation drives the rapid evolution and adaptation of bacterial strains by mediating the propagation of various metabolic properties, including symbiotic life-style, virulence, biofilm formation, or resistance to heavy metals and, most importantly, resistance to antibiotics. These properties make of conjugation a fundamentally important process at the center of extensive study. Here, we review the key steps of conjugation by following the life-cycle of the F plasmid during transfer from the donor to the recipient cell. We also discuss our current knowledge of the extent and impact of conjugation within an environmentally and clinically relevant bacterial habitat, bacterial biofilms.

Keywords: horizontal gene transfer, bacterial conjugation, phenotypic conversion, drug-resistance dissemination, bacterial biofilms, mobile plasmids, F plasmid

1. Introduction

Conjugation was first discovered in 1946 by Edward Tatum and Joshua Lederberg, who showed that bacteria can exchange genetic information by unidirectional transfer of DNA, mediated by a so-called F (Fertility) factor, [1]. It was later realized that the F factor is a replicative extra-chromosomal genetic element, for which they later coined term plasmid, that can be transferred across the cell membrane of the parental strains. Since this seminal discovery, the identification of a plethora of conjugative elements, including plasmids, conjugative transposons and integrative conjugative element (ICEs), has revealed that conjugation is a universally conserved DNA transfer mechanism among Gram-negative and Gram-positive bacteria [2,3]. Conjugation was also shown to be a ubiquitous process that occurs in environment bacterial communities present in the soil, on plant surfaces, in water and sewage, as well as in biofilms and bacterial communities associated with plant or animal hosts [4]. Within these niches, conjugation facilitates the adaptation of bacterial strains by mediating the propagation of advantageous metabolic properties, such as symbiotic life-style, virulence, or resistance to heavy metals and antimicrobials. Conjugation is therefore a major driver of the rapid evolution of bacterial genomes [5,6]. Such fundamental importance has made of conjugation the focus of extensively study over the last decades. Experimental approaches have provided a detailed understanding of the molecular mechanism of conjugational DNA transfer, while systematic sequencing has uncovered the extent of conjugation at the ecological scale.

Conjugative plasmids generally carry all the genes required for their maintenance during the vertical transfer from the mother to the daughter cells, as well as the genes necessary for horizontal transfer during conjugation from the donor to the recipient cell. These functions are encoded by different regions or modules that compose what is generally referred to as the plasmid backbone.

Isolation and sequence analysis of an increasing number of conjugative plasmids has revealed considerable diversity in terms of genetic properties and organisation. This diversity also indicates that different plasmids might use various regulations, molecular reactions and strategies to achieve productive conjugational transfer and maintenance.

In this article, we review the key steps of conjugation by following the life-cycle of the plasmid during transfer from the donor to the recipient cell (Figure 1). We focus on the first discovered and extensively described F plasmid, which we use as a paradigm to discuss other conjugative systems. The first chapter describes events occurring within the donor cell, i.e., the expression regulation of the plasmid *tra* genes required for conjugation; the processing of the plasmid by the relaxosome prior to transfer; the composition and function of the conjugative pilus in the mating pair formation process; the central role of the Type IV coupling proteins (T4CPs) and the transfer by the type IV secretion system (T4SS). The second chapter focuses on the dynamics of the newly acquired plasmid within the recipient cell, i.e., the plasmid establishment, which includes protection against host systems dedicated to foreign DNA elimination, early expression of leading genes and the conversion of the ssDNA plasmid into dsDNA; the plasmid maintenance, which includes plasmid replication and segregation; and the eventual phenotypic conversion of the transconjugant into a new donor cell with novel metabolic properties. In the third chapter, we review our current knowledge of the extent and impact of conjugation within an environmentally and clinically relevant bacterial habitat, bacterial biofilms.

2. WITHIN THE DONOR CELL

2.1. Transfer genes expression

2.1.1. Regulation of *tra* genes expression

The ability of the donor strain to perform conjugation requires the expression of the transfer genes clustered in the *tra* region of the plasmid. The transfer genes encode all the protein factors involved in the elaboration of the conjugative pilus and the T4SS required for formation of the mating pair, as well as the relaxosome components needed for the processing of the plasmid prior to transfer (Figure 1, step i). The expression of *tra* genes is regulated by several factors, including plasmid and host proteins, cell-cycle progression and environmental conditions. Most *tra* genes are gathered in one operon under the control of the P_Y promoter, while *traJ* and *traM* genes are located upstream and controlled by independent promoters (Figure 2) [7]. Transfer gene expression follows a specific regulation cascade that starts with the production of TraJ protein (Figure 2, Step1), which activates the P_Y promoter and the transcription of the *tra* operon (Figure 2, Step2). The first gene to be transcribed, *traY*, encodes the TraY regulator protein that activates the P_M promoter, resulting in the production of the relaxosome accessory protein TraM (Figure 2, Step3) [8]. Therefore, this regulation cascade results in the expression of all genes involved in the elaboration of the conjugative pilus, the T4SS system and the relaxosome composed of TraY, TraM, and TraI. It is observed that *tra* genes are normally repressed, most probably to prevent the fitness cost that would be associated with their constitutive expression [9]. It is interesting to remark that most regulation systems act by modulating cellular levels or activity of the primary activator TraJ. In most F-like plasmids (R100, R1, R6-5, ColB2-K77), the expression of *traJ*, and therefore that of other transfer genes, is repressed at the post-transcriptional level by the fertility inhibition system FinOP (Figure 2) [10,11]. FinP is an antisense RNA complementary to the stem-loop structures of the *traJ* mRNA. FinP binding hides the ribosome binding site and prevents TraJ translation [12,13]. FinO is an RNA chaperon that protects FinP for degradation by RNase E and stabilizes the formation of the duplex FinP-mRNA *traJ* [14–16]. Besides, *tra* gene expression is also regulated by chromosomal-encoded host factors (reviewed in [17]). One such regulation involves the silencing of P_Y , P_M , and P_J promoters by the chromosome encoded histone-like nucleoid structuring protein H-NS [18,19]. H-NS copy number per cell varies during growth [20], thus rendering F plasmid transfer rate growth-phase-dependent,

i.e., maximum in exponential, reduced in mid-exponential and mostly abolished in stationary phase [21,22]. Yet, during the exponential phase, H-NS repression activity is itself counteracted by the cooperative binding of TraJ and the host protein ArcA (Aerobic respiration control of anoxic redox control) to the P_γ promoter [23]. In the case of the virulence plasmid pSLT of *Salmonella enterica*, H-NS repression activity was also reported to depend on Dam (DNA adenine methylase) methylation of the DNA [24]. Other examples of host factor-mediated regulation of *tra* gene expression includes the repression by the RNA binding protein Hfq that destabilizes both *traJ* and *traM* transcripts [18], and by the GroEL chaperone proteins that directly activate plasmid R1 TraJ proteolysis during the cellular heat shock response [25].

For some conjugation systems, *tra* gene expression is regulated by quorum sensing (QS) mechanisms. This is the case for the Tumor inducing conjugative plasmid (pTi), which allows *Agrobacterium* to infect and disseminate within plant hosts. At high cellular density, *Agrobacterium* produces agrocinopines molecules that activate different operons, including the *arc* operon that encodes TraR (unrelated to the F TraR protein), a LuxR-like protein. Binding of TraR to the QS molecule 3-oxo-octanoylhomoserine lactone (OOHL) triggers the transcription of *trb* and *tra* operons, resulting in the production of the T4SS and the relaxosome proteins. The QS lactonase BlcC is also produced, allowing the degradation of OOHL molecules in the stationary phase or during carbon and nitrogen starvation associated with the host plant death. Such regulation provides coordination between the pTi conjugation proficiency and the host state or to the bacterial density within the plant tissues during infection [26]. *Pseudomonas aeruginosa* uses QS as a defense mechanism against interspecies conjugation through the production of the QS molecule N-acyl homoserine lactone (AHL), involved in *P. aeruginosa* populational regulation of mechanisms such as virulence, biofilm formation, and metabolism [27]. AHL produced by *P. aeruginosa* can bind to the *E. coli* LuxR-like transcriptional factor SdiA, which then represses *traI* gene expression and prevents the conjugation of the RP4 broad-host-range plasmid integrated into the chromosome of the *E. coli* donors.

Regulating the expression of transfer genes is the chief strategy to modulate the transfer efficiency of conjugative plasmids. The above examples illustrate that *tra* gene expression is controlled by complex regulatory circuits, which involve the combined activities of plasmids and chromosomal host factors. This tight regulation allows controlling the transfer efficiency in connection with the plasmid life-cycle and the host physiology, in response to environmental conditions and populational interactions.

2.1.2. Superspreader mutations

Over the years, several studies have revealed genetic modifications, so-called superspreader mutations, which dramatically enhanced the conjugation efficiency of conjugative plasmids belonging to diverse incompatibility groups. The first superspreader mutation was characterized on the F plasmid, which carries an IS3 insertion sequence into the *finO* gene. FinO inactivation destabilizes the FinP-mRNA *traJ* duplex, thus resulting in the up-regulation of *traJ* and the constitutive expression of *tra* genes [28]. This naturally occurring mutation accounts for the enhanced transfer efficiency of the F plasmid compared to the related IncF plasmids R100, R6-5, and R1, in which the FinOP regulatory system is still active [29]. More recently, genetically induced superspreader mutations of several resistance plasmids have been isolated in laboratory settings. In the IncI plasmid pESBL associated with extended-spectrum beta-lactamases production in *Enterobacteria*, inactivation of the Hft locus triggered the overexpression of conjugative pili and 20-fold enhancement of the transfer efficiency [30,31]. In the *Citrobacter freundii* IncM group plasmid pCTX-M3 that carries the *bla*_{CTX-M-3} gene, the deletion of two genes (*orf35* and *orf36*) resulted in the enhanced expression of the *tra* genes and increased plasmid transfer [32]. Another example was reported in the Gram-positive broad-host-range (Inc18) plasmid pIP501 involved in the propagation of vancomycin resistance from *Enterococci* to methicillin-resistant strains of *Staphylococcus aureus*. In

this case, the deletion of the *traN* gene encoding the small cytosolic protein TraN (unrelated to the F TraN protein) resulted in the upregulation of transfer factors and enhancement of the transfer efficiency [33].

Inducing the over-expression of plasmid transfer genes might not be the only way through which *superspreader* mutations increase the transfer efficiency of conjugative plasmids. It was shown that insertion of the Tn1999 transposon in the *tir* (transfer inhibition of RP4) gene of the IncL/M-type plasmid pOXA-48a, responsible for the dissemination of specific extended-spectrum β -lactamase genes in *Enterobacteriaceae*, increases the transfer efficiency by 50-100 fold without affecting *traM* expression levels [34]. The mechanism by which the inactivation of the Tir protein enhances transfer efficiency remains to be elucidated. These studies show that superspreader mutations can emerge by various mechanisms in different conjugative plasmids, and have the potential to aggravate the spread of drug-resistance plasmids among bacterial organisms.

2.2. Conjugative pilus, mating pair formation, and stabilization

2.2.1. F pilus structure and biosynthesis

Bacterial conjugation is a contact-dependent horizontal gene transfer mechanism that involves a conjugative pilus associated with a T4SS. Electron microscopy imaging was instrumental in analyzing the morphology of numerous conjugation pili encoded by plasmids belonging to different incompatibility groups [35–38]. These studies revealed that pili fall into two main morphological categories, thin flexible and thick rigid, which influence the ability to support conjugation in liquid or on a solid surface. F encodes a thin flexible pilus, which has a tubular structure $\sim 8 \mu\text{m}$ in diameter and up to $\sim 20 \mu\text{m}$ in length and which is constituted by the helicoidal arrangement of a unique protein subunit, the F pilin or TraA [39–44] (Figure 1, step i). The *traA* gene encodes a 121 amino acids pro-pilin peptide, that is subsequently matured into a 70 amino acids F Pilin [45,46]. The maturation process involves TraQ and TraX proteins [47]. TraQ chaperone-like protein binds transiently to TraA pro-pilin precursor, thus allowing its accumulation into the inner membrane by an ATP-dependent pathway [48] and giving it the right conformation for a signal peptide cleavage. Processing of pro-pilin into matured pilin requires both the cleavage by the host leader peptidase B (LepB) and the N-terminal acetylation by TraX [49,50]. This maturation process ensures the availability of pilin subunits into the inner membrane before the assembly of the pilus by TraE, TraK, TraB, TraV, TraC, TraW, TraG, TraF, TraH, TraL, TrbC encoded by the transfer region [51–53]. Mutational experiments showed that this set of proteins can be separate in different functions. Briefly, TraE, K, C, G, L are responsible for the assembly of the tip, while TraB, V, W, F, and H are important for the pilus extension and TrbI for pilus retraction [52,53]. The pathway of F-pilus biosynthesis has been extensively reviewed [52,54] and will not be detailed here (Table 1).

Protein	Proposed function	Description	Localization	Homologue	Reference
TraM	Relaxosome	<i>oriT</i> binding, TraI stimulation, Interaction with TraD	C		[17,52,55,56]
TraJ	Regulation	Transcription factor (anti-silencer / activator of P _γ)	C		[17,55,57]
TraY	Relaxosome Regulation	<i>oriT</i> binding, Transcription factor (activator of P _M)	C		[17,52,55,57]
TraA	Pilin	Major subunit of the pilus	IM	VirB2 (pTi) TrbC (RP4)	[52,55,56]
TraL	Pilus assembly	Pilus assembly	OM	VirB3 (pTi) TrbD (RP4)	[52,53,55,57,58]
TraE	Pilus assembly	Pilus assembly	IM / P	VirB5 (pTi)	[52,53,55]
TraK	Pilus assembly	Cell envelope-spanning channel	IM / P	VirB9 (pTi)	[52,53,55,56]
TraB	Pilus extension	Cell envelope-spanning channel	IM	VirB10 (pTi) TrbI (RP4)	[52,53,55,56,58]
TraP	Pilus extension	Extended pilus stabilization	IM		[55,57]
TraG	Pilus assembly	Pilus tip assembly	IM	VirB6/VirB8 (pTi)	[52,53,55,57,59]
	Mating-pair stabilization	Stabilization via C-terminal Interaction with TraN,			
	Exclusion	Interaction with TraS			
TraV	Pilus extension	Lipoprotein	OM / P	VirB7 (pTi)	[52,53,55,56]
TraR	Regulation	Transcription regulator by binding to RNA polymerase	C		[58,60]
TraC	Pilus assembly	NTPase	IM	VirB4 (pTi) TrbE (RP4)	[52,53,55,56]
TraW	Pilus extension	Pilus synthesis	P		[52,53,56,57,61]
TraU	DNA transfer	DNA transfer	P		[52,55,56]
TraN	Mating-pair stabilization	Stabilization of OmpA and Lps binding	OM		[52,55,56]
	Exclusion system	Interaction with TraG			
TraF	Pilus extension	Disulfide bonds for T4SS assembly	P		[52,53,55,56]
TraQ	Pilin maturation	Chaperone-like	IM		[55–57]
TraH	Pilus extension	Interaction with TraF and TraU	P		[52,55]
TraG	Pilus assembly	Pilus tip assembly	IM	VirB6/VirB8 (pTi)	[52,55,57,59]
	Mating-pair stabilization	Stabilization via C-terminal Interaction with TraN,			
	Exclusion	Interaction with TraS			
TraS	Entry Exclusion (Eex)	Interaction with TraG	IM		[55,56]

TraT	Surface exclusion (Sfx)	Disaggregation of mating-pair after DNA transfer, Interferes with TraN-OmpA interaction	OM		[55,56,62,63]
TraD	T4CP	Coupling protein / DNA dependent ATPase Interaction with TraM	IM	VirD4 (pTi) TraG (RP4) TrwB (R388)	[55-57,64]
TraI	Relaxosome	Relaxase, transesterase and helicase	C	VirD2 (pTi) TrwC (R388)	[55,57,65,66]
TraX	Pilin maturation	N-terminal acetylase	IM	TrbP (RP4)	[55-57](1)(3)(8)

Table 1 : Tra proteins description. Proteins are presented following the order of the corresponding *tra* genes in the *tra* region of the F plasmid. The proposed function, the description of their biological activity, the subcellular localization (IM : Inner membrane; OM : Outer membrane; C : Cytoplasm; P: Periplasm), and the homologues in RP4, pTI or R388 plasmids are shown.

2.2.2. Pilus biological function

The role of the F-pilus in conjugation has been actively debated. It was early proposed that the F-pilus extends to contact the recipient cell, then retracts to bring together the donor and recipient cells and form the mating pair [41,67,68]. This idea was convincingly supported by the direct visualization of the F pilus dynamics in live-cell, using a fluorescently labeled R17 bacteriophage that specifically binds along the pilus sides [69]. This work showed that donor cells produce flexible pili that continuously undergo cycles of extension and retracting, thereby probing the surroundings, regardless of the absence or presence of recipient cells. Yet, when contact is established with a recipient, the pilus retraction draws the cells together, resulting in the formation of mating pair [69]. In liquid culture, the pili mediate the formation of larger mating aggregates containing donor and recipient cells in tight wall-to-wall contact [70,71].

Allowing the formation of the wall-to-wall contact between the mating partners might not be the only role of the F-pilus, which was also proposed to serve as a channel conducting the single-stranded DNA during conjugation between distant donor and recipient cells [72]. Undeniably, the pilus axial hole has a diameter (30 Å) large enough to accommodate the DNA molecule [41,44]. Yet, only a few reports provide evidence for conjugational DNA transfer between mating partners spatially separated from each other. It was shown that DNA transfer could occur between donor and recipient separated by 6 microns membrane with pores 0.01 to 0.1 micron in diameter [73]. Also, microscopy imaging provided some evidence for DNA acquisition by recipient cells that are not in direct contact with a donor cell [74]. Yet, F-pilus ability to conduct DNA is still questioned and awaits for the clear visualization of DNA transfer between distant donor and recipient cells only connected by a pilus.

2.2.3. Factors involved in the specificity of donor-recipient interactions

The pilus ability to establish contact between donor and recipient cells can be considered as the first rate-limiting step in the conjugation process, and a key determinant of plasmid host-range specificity. In the seventies, numerous studies attempted to identify specific recipient receptor required for F plasmid transfer [70,75-80]. The results revealed that mutations localized in the fourth external loop of the major porin OmpA or that alter the inner core composition of the lipopolysaccharide (LPS) affect the transfer of the plasmid F and other IncF-like plasmids such as R386, R538-1drd, and R1-19 but not the IncFII-type plasmid R100-1 [81-84]. Analysis of several *ompA* and LPS mutants revealed that they do not affect pili binding but result in defective mating-pair

stabilization [85]. Further investigation excluded that TraA is the donor component responsible for specific recognition of the recipient receptors [86] and uncovered the mating-pair stabilization function of the outer membrane protein TraN, whose three external loops have been reported to interact with OmpA and LPS [87–89]. These findings explain that *ompA* and LPS mutations do not alter the conjugation efficiency of the closely related IncFII-type plasmid R100, which TraN central region amino acid sequence is highly divergent from F-encoded TraN [81,84,90]. Mating pair stabilization additionally involves the multifunctional inner membrane protein TraG, which N-terminal part is also a role in piliation and surface exclusion [90–93].

OmpA or LPS receptor factors cannot be considered as strictly essential to F conjugation, since their mutation only decrease transfer efficiency by a 2-3 log. Moreover, mating-pair stabilization defect can be overridden by performing mating on solid media, suggesting that interactions with OmpA or LPS are needed to stabilize the mating pair formation only in liquid mating [77,85,94]. One can then wonder if the F plasmid is an exception concerning the need for recipient factors in some conditions? A recent study identified in *Klebsiella pneumoniae* the outer membrane OmpK36 homolog of the *E. coli* outer membrane OmpC as a receptor mediating conjugation of the IncFII plasmid pKpQIL [95]. As observed in the case of the plasmid F, TraN_{pKpQIL} would eventually mediate the OmpK36-receptor specificity, while complementing a *traN*_{pKpQIL} mutant with TraN_{R100} abolishes this dependency, demonstrating that recipient receptors might be highly specific of the transferred plasmid. Beside the IncF-type, the conjugation process of IncI-type plasmids was also shown to be sensitive to LPS mutations, and interestingly some LPS mutations affecting IncI plasmid transfer do not impact the entry of the F plasmid while some others affect both type of plasmid [96,97]. Recently, the PilV adhesin encoded by the IncI1-type plasmid R64 has been identified as the donor factor binding to recipient cell LPS [98]. This adhesin is thought to locate at the tip of the thin Type IV pilus required only in liquid mating conditions rendering the interaction between PilV and LPS important only under these conditions as observed for the TraN and OmpA or LPS interactions in the case of the plasmid F. In contrast, no such recipient receptors could be identified for the broad-host-range plasmids RP4 and R388. Indeed, the *ompA* and LPS recipient mutants decreasing the efficiency of F conjugation drastically do not affect RP4 conjugation efficiency [99]. Large-scale mutants screening using the *E. coli* Keio collection or random *E. coli* insertion mutant library failed to identify recipient mutants affecting the transfer of the plasmid RP4 [100] or plasmid R388 [101].

Remarkably, some broad-host range IncP-like plasmids can also be transferred in archaea [102] and eukaryotes such as yeast [103] and mammalian cells [104]. Although the efficiency of conjugation varies among recipient type cells, these findings strongly suggest that plasmid transfer does not require any specific factors or active mechanism on the recipients side. Alternatively, a “shoot and pump” conjugation model envisages that the type IV secretion apparatus could act like a syringe able to inject DNA in any kind of membranes, using the pilus as a needle [105]. Perforation of the recipient bilayer cell-wall could be achieved by force or by dedicated enzymatic activity exposed at the pilus tip. The lack of requirement for specific receptors on the surface of the target cell is not an exception of the conjugative T4SS. Indeed, although the structural components of the Type VI secretion system machinery have been widely documented, no studies have yet characterized genetic factors that could act as target receptors on the prey cell surface.

2.3. Plasmid processing by the relaxosome

The initiation of conjugation requires the assembly and activity of a protein complex, the relaxosome, that processes the plasmid before DNA transfer (Figure 1, step ii). The relaxosome introduces a site- and strand-specific DNA cut (nick) at the *nic* site located in the origin of transfer *oriT* and is required for the extrusion of the single-stranded DNA to be transferred (T-strand) [106–108]. In the F system, these two reactions are performed by the multifunctional TraI relaxase protein, which has both a transesterase domain that catalyzes the *nic* reaction and a DNA helicase domain

that unwinds the plasmid DNA [64,109–114]. Crucially, TraI recruitment and activity are governed by auxiliary proteins, including the integration host factor IHF and the plasmid-encoded TraY and TraM proteins, which have distinct roles in the relaxosome formation and activity at *oriT* [115–117]. Binding of IHF and TraY to their respective cognate binding sites *ihfA/ihfB* and *sbyA* located at *oriT* modulate the architecture of the DNA, thereby stimulating the loading of TraI [118–122]. TraM is a small protein that regulates its own expression by binding to *sbmA* and *sbmB* sites located in PM promoter, and that stimulates the DNA relaxation reaction through direct interaction with TraI after binding to *sbmC* site located in *oriT* region [123–127]. TraI nicking reaction involves a catalytic active Tyr residue [65,128] and results in the relaxation of the plasmid dsDNA, where the 5'-phosphate (P) end of the nicked strand (or T-strand) remains covalently bound to TraI (Figure 1, step iii) [111,113,129–132]. After the nicking reaction, the circular ssDNA conjugative plasmid is converted into dsDNA by Rolling Circle Replication (RCR) in the donor, while the linearized T-strand DNA bound to TraI at the 5' end is transferred into the recipient cell through the conjugative pore (Figure 1, step iv).

2.4. Initiation of rolling circle replication in the donor cell

Rolling circle replication RCR mechanism is employed for the vegetative replication of some bacterial plasmids and has been very well-reviewed [133–135]. RCR is key to the transfer process of many Gram-negative and Gram-positive conjugative plasmids, but also to the infection cycle of other mobile genetic elements like DNA/RNA viruses and bacteriophages [128,136]. RCR reactions involved in vegetative replication or in plasmid transfer are very similar. The initiation and termination of RCR reactions performed by the Rep protein during vegetative plasmid replication are achieved by the relaxase protein during conjugation. Indeed, Rep and relaxases exhibit closely related functions, primarily allowing RCR initiation by nicking the double-strand DNA at the *dso* site (double-stranded-origin) or the *oriT* site, respectively [137]. The nicking reaction generates a 5'-P end that remains covalently bound to Rep or TraI and a 3'OH end used as a primer for the host DNA polymerase III. While the DNA polymerase III performs leading strand elongation, the parental double helix is unwound, and RCR ends by a second nicking reaction that releases the newly synthesized strand DNA (Figure 1, step v). In the case of vegetative RCR, DNA unwinding is performed by a host DNA helicase recruited by the Rep protein, while Rep itself ensures termination and second nicking reaction. One major specificity of conjugation-associated RCR is that replication of the two ssDNA strands occurs in different cells, i.e., the leading strand is replicated in the donor, while the T-strand (lagging strand) is transferred and replicated in the recipient cell (Figure 1). Because the relaxase that initiates the nicking reaction is transferred together with the T-strand [65,138,139], a second relaxase protein is required in the donor to perform the DNA unwinding as well as the second nicking reaction [3,65,128]. Consistently, structure analysis by Cryo-EM shows that two relaxase molecules bind the *oriT*, one associated with the 5' end that is in open transesterase conformation and one associated with the 3' end that is in closed helicase conformation [140].

2.5. T4CP connects the relaxosome to the T4SS

After processing by the relaxosome complex, the nucleoprotein complex composed of the T-strand and the covalently bound TraI needs to be recruited to the conjugative pore for transfer (Figure 1, step iii). This connection is mediated by the interaction between the relaxosome and the Type IV Coupling Protein (T4CP) located at the cell membrane [3,141]. All conjugative systems have their own T4CP, such as TraD, TraG, and TrwB, respectively, for the F, RP4, and R388 plasmids. T4CP are not required for pilus production or DNA processing, yet they are key to substrate specificity [142]. Our understanding of the molecular interactions required for specific substrate recognition and translocation is still incomplete. A great deal of information has been provided by comparing various T4SS involved in protein or nucleoprotein transport [143–146]. It appears likely that conjugation systems derive from ancestral protein translocation machinery that evolved to

translocate DNA coincidentally. In this view, the T4CP would serve as the substrate receptor that interacts with one or several relaxosome components to recruit the T-strand to the T4SS. In the case of the F plasmid and some other plasmid systems, it is well-established that TraD interacts with TraM relaxosome protein [147–151]. Interaction between the T4CP and the relaxase has been demonstrated in the case of RP4 [152] and R388 plasmids [153]. Such interaction has been speculated in the F system but remains elusive [112,154].

T4CP are DNA-dependent ATPases anchored to the cell membrane via their N-terminal domain and have been shown to interact with the T4SS components in R27 [155] and R388 plasmid systems [153]. T4CPs show similarities with membrane-anchored ring DNA translocases, such as SpoIIIE and FtsK involved in chromosome DNA translocation during sporulation and cell division, respectively [141,156,157]. T4CP binds non-specifically to DNA, with a higher affinity for ssDNA [152,158,159], on which it forms oligomers with enhanced ATPase activity [160,161].

Altogether, these findings led to the model that membrane-anchored T4CPs interact directly with the relaxosome and form hexameric structures on the T-strand, which they actively translocate through the conjugation pore during transfer. Yet, it remains unclear whether a signal is required to activate the coupling function of the T4CP in a donor cell where the T4SS and the relaxosome are already assembled and functional [3]. It has been suggested that the stability of the TraD oligomeric complex depends on a yet unidentified F-encoded protein, which could then be a key regulator of plasmid transfer activation [160]. It has also been suggested that the formation of the mating pair could transduce a signal to activate the T4CP and trigger the transfer of the processed T-strand [3]. Of note, it was shown the relaxase has to be unfolded to be translocated into recipient cells [139]. In *A. tumefaciens*, as in many other Gram-negative and Gram-positive systems, unfolding of translocated proteins is proposed to be performed by the VirB11-like ATPase, absent from the F plasmid system [52,145,162]. However, one can reasonably presume that TraI unfolding also required ATPase activity involving one of the ATPases of the F T4SS.

3. WITHIN THE RECIPIENT CELL

3.1. Plasmid circularization by TraI

The relaxase is transferred into the recipient, where it is refolded and able to perform several activities required for completion of the conjugation process (Figure 1, step *a*) [65,138,163]. The helicase activity of the internalized relaxase is thought to perform 5' to 3' tracking of the T-strand. Pulling by the relaxase from the recipient together with pushing by the T4CP from the donor presumably facilitates the transport of the T-strand through the conjugation pore [3]. Once both extremities of the *oriT* are brought together in the recipient, the relaxase performs the joining reaction, resulting in the recircularization of the ssDNA plasmid (Figure 1, step *b*) [65,138,163,164]. An alternative model proposes that the nicking of the newly synthesized *oriT* occurs in the donor cell before transfer [165]. There is no evidence for the requirement of additional host or plasmid factors in the circularization of the internalized T-strand. After completion of the recircularization reaction, the recipient cell possesses a single-stranded circular copy of the conjugative plasmid.

3.2. Avoiding host defense systems against foreign DNA

The newly acquired ssDNA conjugative plasmid might be considered as a foreigner DNA, against which host bacteria have developed defense mechanisms, such as restriction-modification, exonucleases, and recombination system or adaptive immunity as the CRISPR-Cas [166]. Despite these defense mechanisms, horizontal gene transfer plays an important role in genomic evolution (5-6% of bacterial genomes and up to 20% in some organisms) [6,167,168], implying that transferrable plasmids have evolved adaptive mechanisms to counteract these host defenses.

Restriction-modification system (RM) is an ubiquitous defense mechanism found in 90% of sequenced bacterial genomes and other prokaryotes based on restriction enzymes and methylation [169]. Restriction-modification mechanisms are described as a « primitive immunity system » against exogenous DNA [170]. These systems are based on restriction enzymes that target specifically unmethylated DNA sequences located on the newly acquired mobile genetic elements, while the host DNA is protected by methyl groups added to specific adenine or cytosine [170]. Yet, plasmids have evolved various strategies to counteract enzymatic DNA degradation upon entry in the new host cell. IncN, IncI, and IncF plasmids encode ArdA and ArdB proteins (Alleviation of restriction of DNA) that directly inhibit the REase (Restriction Endonuclease) by mimic DNA sequences, thus competing for enzymes target [171–173]. IncW plasmids encode the ArdC protein that protects the transferred T-strand by transiently blocking the restriction sites [174]. More recently, it was shown that a *hde* operon (host defense evasion) of IncI plasmids encode 2 genes involved in anti-RM (*vcrx089* and *vcrx090*) [175]. Besides the production of inhibitory proteins, some plasmids completely lost the restriction sites, as it is the case for the RP4 plasmid [176].

CRISPR-Cas immune systems (Clustered Regularly Interspaced Short Palindromic Repeats and the CRISPR-associated protein) represent another defense mechanism against foreign DNA. CRISPR-Cas systems, found in ~45% of bacterial and up to 84% of Archaea genomes [177,178], have been described to protect against infection by bacteriophages, and more recently against plasmids acquisition [179]. It was then discovered that some phages encode an anti-CRISPR Acr protein that inhibits the activity of the CRISPR-Cas system [180,181]. Importantly, anti-CRISPR Acr *loci* have been identified in conjugative elements and plasmids of *Listeria*, *Enterococcus*, *Streptococcus*, and *Staphylococcus* [182]. These *loci* encode for the CRISPR-Cas inhibitors AcrIIA16-19 that prevent exogenous DNA nicking mediated by the Cas9 enzyme *in vivo*. Mahendra and co-workers have also shown that conjugation of a Cas9-targeted plasmid of *E. faecalis* was possible in the presence of these CRISPR-Cas inhibitors. Encoding Acr-like proteins is therefore an efficient strategy for conjugative plasmids to facilitate their dissemination by avoiding degradation by the host CRISPR-Cas immune system. Another strategy is for conjugative plasmids to encode a Bet/Exo system able to repair double-strand breaks caused by CRISPR-Cas during conjugation, as recently reported for IncC plasmid pVCR94 [175]. Expression of these genes inside the recipient bacteria after the acquisition of the *V. cholerae* pCVR94 plasmid allows survival against exogenous DNA defense mechanisms without the involvement of anti-CRISPR-Cas proteins.

3.3. Role of the leading region genes and conversion of the ssDNA plasmid into dsDNA

3.3.1. Early expression of the leading region genes

The leading region of conjugative plasmids is the first to be transferred into the recipient cell during conjugation (Figure 1, step *a*) [57]. The F plasmid leading region is well conserved, of a size of 13kb and directly adjacent to the *oriT* and encodes for at least eight proteins [183], including a homolog to the chromosomal SSB (SSB_c), PsiB and other proteins of unknown function. Importantly, plasmid *ssb* (*ssb_p*) and *psiB* genes are expressed early upon entry of the plasmid in the transconjugant bacteria, but not in the donor cells [184]. This was demonstrated using *lacZ* fusions of the two genes inside the Collb-P9 plasmid, and also by immunoassays concerning the F plasmid PsiB protein [184]. Similarly, RT-PCR studies showed that *psiB* and *ardA* genes of the IncI1 plasmid begin to be expressed 5 minutes after transfer initiation [185]. These observations suggested that leading region genes could be expressed rapidly from the newly acquired plasmid in single-stranded form, before its conversion into dsDNA. It was later shown that the leading region contains a specific 328 bp *Frho* region (for F plasmid RNA polymerase), which, when single-stranded, can form a stem-loop structure presenting -10 and -35 double-stranded boxes recognized by the host RNA polymerase that initiates the synthesis of RNA primers *in vitro* [186]. It was therefore proposed that *Frho* may serve as a single-stranded promoter that allows the early expression of the leading region genes (Figure 1, step *a*).

Frpo was also proposed to direct the single-strand to double-strand conversion of the F plasmid. *In vitro* assays showed that the RNA primers synthesized by the RNA polymerase from *Frpo* persists as a RNA-DNA duplex that is recognized by host DNA polymerase III to initiate complementary strand synthesis [186]. *Frpo*-type sequences also termed *ssi* for single-strand initiation sequence are found on various conjugative plasmids including R6K, R100, ColE1, ColE2, Col1B and RSF1010 [187–190] and are functionally comparable to *sso* sequences (single-stranded origin) involved in the rolling circle replication mechanism [128,134,191]. These findings are consistent with the previous observation that complementary strand synthesis of the ssDNA F plasmid inside the recipient bacteria involves a cooperative mechanism between host RNA and DNA polymerases III [192].

Altogether, these findings led to the model that the *Frpo* could help initiating early gene expression and the DNA synthesis reaction that converts the ssDNA plasmid into dsDNA duplex, immediately upon entry of the T-strand into the recipient cell (Figure 1, step *a-c*). Whether *Frpo* performs these functions during conjugation *in vivo* remains to be demonstrated.

3.3.2. PsiB inhibits the SOS response

In the recipient cell, the presence of abnormal amounts of ssDNA, usually associated with DNA damages, results in the induction of the SOS response [193,194]. More precisely, the loading of RecA recombination protein on ssDNA results in the formation of the pre-synaptic filament, which stimulates the autocatalytic cleavage of LexA, the repressor of the SOS regulon. The SOS response triggers the induction of the division inhibitor Sula, resulting in cell filamentation and potentially the death of the transconjugant cell. SOS also induces the production of nucleases and other DNA processing factors that could provoke the degradation or mutation of the transferred ssDNA, or its processing as a recombination intermediate [195]. To counteract these effects, several conjugative plasmids, including F, encode the PsiB protein (Plasmid SOS inhibition), which inhibits SOS induction [196]. The depletion of *psiB* has mild effects on the efficiency of conjugation but increases the host SOS response up to six-fold [184]. PsiB interacts directly with RecA, thereby inhibiting several activities such as DNA binding, LexA cleavage, and strand exchange reaction [197,198]. SOS response inhibition by PsiB is even more potent in presence of SSB_c protein coating the ssDNA. PsiB is well conserved among conjugative plasmids and is considered important for the early steps of plasmid establishment in the recipient, consistent with its early production in the transconjugant cells [57,196].

3.3.3. Roles of the host and plasmidic SSB proteins in plasmid establishment

Upon entry in the recipient cell, the transferred ssDNA plasmid is coated by the host SSB_c protein. SSB_c is an essential conserved protein that binds non-specifically to ssDNA. It is involved in various mechanisms, including DNA replication, repair and recombination, SOS-induction, and other DNA metabolic processes (for review see [199,200]). SSB binding protects the ssDNA against enzymatic degradation and increases the processivity of DNA polymerase II and III [201] during the replication reaction converting the ssDNA strand into dsDNA helix. The rapid recruitment of the host SSB_c protein onto the transferred ssDNA has recently been visualized by fluorescence microscopy [202]. This work revealed that SSB_c proteins are rapidly recruited onto the ssDNA that penetrates the recipient cell presumably protecting it and facilitating its processing (Figure 1, step *a*). Interestingly, the F plasmid, as many other conjugative plasmids encode their own SSB_P protein homologous to the *E. coli* SSB_c [203]. One can then wonder what is the benefit for conjugative plasmids to encode their own SSB_P, and what is the specific function of SSB_P compared to the chromosomal SSB_c?

SSB_P can bind ssDNA without specificity and SSB_P of different incompatibility groups (IncF, IncI, IncY, Inc9, IncT, and IncB/O) can partially complement conditional mutations of *E. coli* *ssbc* gene [204–207]. However, although producing plasmid F SSB_P *in trans* enables the growth of *ssbc* deletion mutant, complemented mutant exhibit some filamentation, growth rate reduction [208]. Moreover,

a reduced affinity to ssDNA is observed for plasmid F SSB_P in comparison to the *E. coli* SSB_C, and plasmid F SSB_P cannot stimulate the reaction of DNA synthesis by DNA polymerase III *in vitro* [209]. Sequence alignment revealed that SSB_P proteins complementing *E. coli* *ssbc* mutant share high homology only with *E. coli* SSB_C N-terminal part [209]. The SSB_C N-terminal region contains the domains for ssDNA binding and monomer-monomer interactions to cooperatively maintains the binding of the tetrameric structure of SSB_C to ssDNA. This structural conservation would explain the ability of SSB_P to bind ssDNA. However, the C-terminal domains of SSB_P are quite more homologous to each other than to SSB_C [209]. Yet, as this domain is interacting with partner proteins constituting the SSB_C interactome, one possibility would be that the interactome of SSB_P and the reaction it is involved in might be different than SSB_C.

Not all SSB_P has been shown to complement *E. coli* *ssbc* mutants [205]. The latter study found that the inability of SSB_P from IncP-like RK2 to complement *E. coli* *ssb1* mutation could be attributed to *ssbP* gene repression by the RK2 *kor* genes and that a derepressed plasmid indeed complement the thermosensitive growth of *E. coli* *ssb1* mutations [210]. It is thus to consider the possibility that the numerous SSB_P could also complement *ssbc* mutant.

To date, the function of plasmid F SSB_P in the context of conjugation is still unclear. SSB_P protein could contribute to the protection of the transferred ssDNA by inhibiting enzymatic degradation or the recruitment of host inhibiting proteins. However, its expression timing instead supports the idea that SSB_P could be involved in assisting the complementary strand synthesis of the transferred DNA, or simply increase the pool of available single-strand binding protein required for the first cycle of vegetative replication of the plasmid.

3.4. Plasmid maintenance: replication and segregation

Maintenance of the newly acquired dsDNA conjugative plasmid in the recipient cell lineage depends on two main active mechanisms, plasmid replication and segregation of the plasmid copies to daughter cells over generations.

The mechanisms of plasmid DNA replication in bacteria have been extensively studied and is the focus of well-referenced reviews [128,133,211–214]. Here we would like to stress the role of replication in conjugation host-range specificity. Indeed, the ability of the transferred plasmid to replicate in the recipient cells is considered as the primary determinant of plasmid host-range specificity and efficiency of conjugation. The plasmid F transfers efficiently to *E. coli* and relatively closed enterobacteria while no transconjugant can be recovered after mating with more distant bacteria such as *Vibrio* or *Pseudomonas*. As early 1982, host-range restriction was attributed to the plasmid's inability to replicate in the recipient bacteria rather than to the inefficiency of plasmid transfer *per se* [215]. This was demonstrated by showing that mobilizable plasmids containing a cis-acting origin of transfer of the plasmid F and an origin of replication active in the tested recipient can then be transferred by the F conjugation machinery in *Pseudomonas* [215], *V. cholerae* [216] and yeast [103]. The same approach was employed to show the ability of pCTX-M3, an IncI-like plasmid, to use its conjugation machinery to transfer a mobilizable plasmid in the host recipients where pCTX-M3 does not replicate [32]. The failure of F plasmid replication in *P. aeruginosa* comes in part from an inability of the plasmid replication protein RepE of the RepFIA replicon in complex with host DnaA-*oriS* to form a stable interaction with the host helicase DnaB [217]. In contrast, broad-host-range plasmids such as RK2 regulate their maintenance by modulating alternative strategies of replication depending on the host [218]. These findings indicate that the host-range of the conjugation machinery and the replication origins belonging to the same plasmid differ and can be mechanistically uncoupled. The specificity of narrow-host range plasmids appears limited by the specificity of their replicon rather than by their transfer range.

Maintenance of the newly acquired dsDNA plasmid also requires the timely segregation of the plasmid copies to the daughter cell during transconjugant cell division. To do so, low copy-number plasmids encode active partition systems, which mechanism, biological functions and conservation have been extensively reviewed [219–223]. In the context of conjugation, it is interesting to mention another maintenance strategy that involves the integration of the newly acquired conjugative DNA to the chromosome. Chromosome integration ensures the stable inheritance of the conjugative elements by vertical gene transfer over generations. The first characterized example was again the F plasmid, which uses insertion sequences (IS) and RecA-dependent homologous recombination to integrate into the *E. coli* genome [224]. The integrated plasmid can still initiate the conjugative transfer of the whole chromosome of the resulting “Hfr” strain (High frequency of recombination). This process of transfer is both progressive and oriented, and the order of chromosomal gene transferred depends on the position where the F plasmid was integrated. Alternatively, the integrated F plasmid can be excised out of the chromosome and recover its original autonomous form [225]. Chromosome integration is widely used by mobile DNA such as ICE (Integrative Conjugative Elements), transposons, phages, in both Gram-negative and Gram-positive bacteria, with all these systems having their particularities, especially regarding the recombination systems used for their integration/excision [226,227].

3.5. Phenotypic conversion of the transconjugant

Expression of the genes carried by the newly acquired genetic element results in the phenotypic conversion of the transconjugants into a novel type of cell that exhibits additional metabolic properties (Figure 1, step *d*). The expression of plasmid genes involved in DNA transfer converts the transconjugant into a new donor able to further transfer the plasmid in the population, thus accounting for the exponential rate of conjugative plasmid dissemination (see *tra* gene expression section 2.1). Other plasmid genes encoded by the *tra* region of the plasmid backbone are not directly involved in the transfer process per se. Indeed, conjugative plasmids often carry immunity (or exclusion) gene systems that are widespread in Gram-negative and Gram-positive organisms [93,228–231,231–235]. These immunity systems limit the plasmid-carrying cell ability to serve as a recipient for the same plasmid [42,54,62,236]. Preventing self-mating by surface exclusion is thought to avoid the metabolic cost and potential cell death associated with repeated plasmid transfer, but also to be important for plasmid stability and evolution [62,236].

The F plasmid immunity system relies on two factors, TraT and TraS proteins, none of which is required for F pilus synthesis or DNA transfer [56,237–239]. These two exclusion factors work at different levels. TraT is an abundant outer membrane protein that is thought to span the cell surface [237,238]. TraT production reduces the recipient to form stable mating aggregates, presumably by interfering with the interaction between the pilus and recipient surface receptors. Consistent with this idea, it was reported that TraT interacts with OmpA, further suggesting that it could compete with TraN, which is key to mating pair stabilization [87,88,240]. TraS is an inner membrane protein, which production only slightly reduced aggregation of mating populations, but reduced DNA transfer frequencies by 100-200-fold. It is then proposed that TraS acts by preventing DNA transfer when stable mating aggregates have already formed [237,238,241]. In F-like plasmid, it is proposed that TraS interacts with TraG to achieve the entry exclusion process [92,93]. For these reasons, TraS is referred to as an entry exclusion protein (Eex) and TraT as a surface exclusion protein (Sfx). Like other *tra* genes, *traS* and *traT* expression are controlled by TraJ, implying that transconjugant cells acquire plasmid transfer ability immunity and concomitantly during the phenotypic conversion.

In addition to genes located on the plasmid backbone, conjugative plasmids may carry additional genes that are not directly involved in conjugation, but in a variety of biological functions such as virulence, biofilm formation, symbiotic life-style, membrane trafficking, resistance to heavy metal and, most importantly, resistance to antibiotics. Acquisition of these metabolic functions potentially facilitate the bacterial adaptation and survival in changing environments, and make of conjugation

a major driver of the evolution of bacterial genomes. The successful maintenance of conjugative elements in bacterial populations shows that this selective advantage compensates for the metabolic burden associated with the metabolism of the newly acquired genetic information (highjacking of the host replication, transcription and translation machineries) [242]. The most prominent example being the acquisition of conjugative drug-resistance plasmids, which enables bacterial proliferation in microbial communities containing antibiotic-producing organisms, or in antibiotic-polluted and clinical environments. Indeed, analysis of commensal, environmental and clinical antibiotic-resistant pathogenic strains revealed a multitude of conjugative plasmids carrying one or more genes for resistance to most, if not all, classes of antibiotics currently used in clinical treatments. Conjugation is considered to be the most widespread intra- and inter-species resistance transfer mechanism, accounting for 80% of acquired resistance [243].

4. CONJUGATION IN NATURAL HABITATS, THE EXAMPLE OF BACTERIAL BIOFILMS

Gene transfer by conjugation is known to contribute to the genetic dynamics of bacterial populations living in a variety of environments, including the soil, on plant surfaces, in water and sewage, as well as in bacterial communities associated with plant or animal hosts [4]. Bacteria are generally considered as planktonic unicellular organisms, yet in natural and clinical environments, they often live in complex structures called biofilms. Biofilms shelter bacteria against external hazards, but have also been proposed to offer a niche that facilitates the dissemination of drug-resistance determinants by conjugation. Below, we review our current understanding of the interplay between biofilms and bacterial conjugation.

4.1. The biofilm as a niche promoting bacterial conjugation

In natural environments, bacteria predominantly live in spatially structured communities termed biofilms, in which a self-produced extracellular matrix holds the cells together [244]. Planktonic cells that initiate biofilm formation can adhere to a living or inert surface (surface-attached biofilms) or can be present at the air-liquid interface as a free-floating community (pellicle). Bacterial biofilms are found in virtually every ecosystem on Earth, from aquatic systems (sludge, rocks, and wastewater) to terrestrial environments (rhizosphere) and human organisms (skin, intestinal, urogenital and respiratory tracts). Furthermore, biofilms are associated with persistent and severe infections because of their ability to colonize medical devices and implants. Indeed, biofilm structures offer protection against hostile environments and, more worryingly, against antibiotic treatments. The biofilm architecture depends on the bacterial species, the surface colonized and environmental conditions, but this lifestyle is characterized by the production of extracellular polymeric substances (EPS) mainly composed of polysaccharides, proteins, lipids, and extracellular DNA. EPS matrix production is dynamic and continuous and mediates the formation of the biofilm architecture in which aggregates of microorganisms are trapped.

Many studies on bacterial conjugation show that plasmids transfer can occur in both natural and artificial biofilms, from the aquatic environment [245], phytosphere [246], animal and human host [247,248] or reactor-associated biofilms [249,250]. These researches studied plasmid transfer at the population level and mainly relied on limited cultivation-based assays that probably underestimate the extent of conjugation in natural biofilm. Further works elaborated conjugative plasmid expressing a fluorescent marker allowing direct *in situ* visualization of donors, recipients and transconjugants at the single-cell level within several types of biofilm formed at the liquid-air surface [251], on semi-solid agar surface [252–254], on filter [255], in flow-chambers [253,256,257] or in a reactor [258].

The biofilm environment provides a high cell density and close cell-to-cell proximity that may facilitate HGT through bacterial conjugation. In line with a view of biofilm as a hot spot niche of conjugation, several studies showed a higher frequency of plasmid transfer in biofilm than in the

planktonic mode of growth (reviewed in [259–262]. Although the biofilm appears as a favorable environment to HGT by conjugation, many studies at single cells level have reported limited plasmid propagation inside a preformed established biofilm beyond the contact zone between the donors and recipient (reviewed in [263]. These observations are directly linked to the complex spatial structure of the biofilm, which might have a critical impact on horizontal plasmid spread within a biofilm.

4.2. Impact of the biofilm structure on conjugation

The biofilm is a complex structure sometimes composed of mixed bacterial species. The matrix shapes the spatial organization by clustering cells in microcolonies in an architecture characterized by non-uniform cell arrangements, open channels, pores, cavities, and different layers of living cells [264,265]. Such an organization determines the formation of cell clusters/aggregates and could therefore influence the efficiency of conjugative transfer [251]. This possibility can be addressed using microscopy to analyze the distribution of cells active in conjugation within biofilms. To date, most studies investigated the spread of the GFP-tagged *Pseudomonas putida* TOL plasmid. Analysis of *P. putida* recipient biofilm established in flow chambers revealed that transconjugants appear at the top surface of the biofilm, but not in the deep lower layers, reflecting a limited invasion of the transferred plasmid [253,257]. On agar plate, transconjugants only appear at the contact zone between TOL plasmid-donors and the recipient colonies [252,253,266]. Limitation of transfer to the outer layers of biofilm was also observed for IncF, IncI and IncW plasmids in *E. coli* [254]. It has been proposed that transfer can only be efficient over a short period between metabolically active cells growing at the donor-recipient interface [253,267]. Consistently, it was shown that plasmid invasion stop in non-dividing cells [266].

Within the biofilm structures, chemical gradients of oxygen, nutrients, temperature and pH, create microenvironments that influence the metabolic activity of bacterial cells [268]. This results in physiological heterogeneity between the cells surrounding the border of the biofilm and those embedded deep inside. Also, variations of the spontaneous mutation frequency within the biofilm parts result in the emergence of variant subpopulations with genetic heterogeneity [269]. Whether and how these factors impact the spatial pattern of conjugation within biofilms remain unknown.

4.3. Impact of conjugative plasmids on biofilm formation

Several studies investigated an implication of the presence of conjugative plasmid from diverse incompatibility group on biofilm formation ability [270–274]. To initiate biofilm formation, planktonic cells produce cell appendages like flagella and adhesion factors such as type IV pili, type 1 and curli fimbriae [275,276]. Genes coding for these types of accessory factors that promote attachment to biotic or abiotic surfaces are often found on conjugative plasmids, resulting in host biofilm formation increase (reviewed in [260]). Examples include type 3 fimbriae of the IncX1 plasmid pOLA52 [277]), non-conjugative type IV pili of the IncI1 plasmid pSERB1 [278] or pilus-like structure and surface adhesins of the *Enterococcus faecalis* plasmids pBEE99 and pCF10 respectively [279–281]. In 2001, Ghigo made the unexpected observation that conjugative pili itself of derepressed plasmid F can promote biofilm of *E. coli* cells initially unable to form such structure and revealed the pilin TraA as the main adhesion factor inducing biofilm formation [270]. Microscopic structure analysis of derepressed IncF plasmid R1drd19- and F-carrying *E. coli* biofilm showed the rapid formation of a dense and mature 3D mushroom-type biofilm similar to the *P. aeruginosa* biofilm architecture [282]. This peculiar architecture formation and biofilm maturation generated by derepressed plasmids overrides the need for cell surface appendages essential to *E. coli* biofilm such as flagella, type 1 fimbriae, Ag43 or curli [282]. In contrast, maturation of 3D mushroom-type biofilm structure depends of curli production induced in *E. coli* by natural plasmid F, which does not constitutively express F pili [283]. The presence of the plasmid R1drd19 also increase *E. coli* biofilm formation by decreasing the motility and increasing the level of quorum sensing inducer AI-

2 [284,285] and IncP-9 TOL plasmid increases in *P. putida* the production of extracellular DNA known to structure the biofilm [286] and thus the biofilm formation capacity [287]. However, the genetic mechanism by which conjugative plasmids increase the biofilm formation has not been elucidated, but the presence of repressed or natural conjugative plasmids affects the global host chromosomal gene expression [283,285].

While the role of the conjugative pili has been mainly studied in *E. coli* biofilm, one may consider their impact differ depending on the host. Røder *et al.* indeed observed that the conjugative pili of the IncP-1 plasmids pKJK5 reduce the surface attachment of *P. putida* by increasing cell-cell adhesion resulting in reduced biofilm formation [288]. Further investigation will be necessary to decipher the complex interconnections between the conjugative plasmid and biofilm formation.

4.4. Influence of antibiotic treatment on conjugation within biofilms

Dynamics of conjugation within biofilms need to be more investigated, regarding the specific interconnection where biofilm and conjugative plasmids act in synergy to promote both community-building and gene transfer. This cooperation raises serious questions about the contribution of HGT in the evolution and adaptation of biofilm-forming pathogens. Because of the increase of antibiotic-resistant infections, recent advances tried to bring a new understanding of biofilm responses to antimicrobial treatments.

Sub-minimal inhibitory concentrations (sub-MIC) of aminoglycosides enhance biofilm biomass of *P. aeruginosa* strain PAO1 and clinical isolates of *E. coli* through the response regulator Arr, a predicted phosphodiesterase altering the cyclic di-guanosine monophosphate (c-di-GMP) levels [289]. Linares *et al.* further demonstrated that besides aminoglycosides, sub-MIC of tetracycline and norfloxacin also increase the formation of *P. aeruginosa* biofilm. Still, no clear causal factors were identified [290]. Interestingly, a combination of tetracycline with cephradine has a synergistic effect on biofilm formation of a mixed culture of *E. coli* and *P. aeruginosa* [291]. Tetracycline also promotes biofilm of the pathogen *Acinetobacter baumannii* and whole genome sequence analysis revealed an increase of rate mutation under sub-inhibitory drug exposition such as SNPs but also insertions and deletions [292]. Besides the accumulation of genotypic variation, biofilm treatment with a low level of antibiotics produces changes in the gene expression profile, some of which may be linked to the increase biofilm formation [290,292,293].

Recently, Diaz-Pascual *et al.*, investigated *Vibrio cholerae* biofilm at the community-scale level using a single-cell imaging system revealing changes in biofilms dynamics and architecture in response to antibiotic treatment [294]. After tetracycline exposure, they observed modifications in biofilm architecture and cell morphology, including a 2.5-fold increase in cell volume and a 29% decrease in cell density. This cell density decrease reflected a considerable alteration of the multicellular arrangement and the breakdown of the matrix within the biofilm. Furthermore, biofilms became susceptible to the colonization of their interior by new cells and the colonizer population increased until they invade the resident biofilm. Clearly, a sublethal dose of antibiotics influences the biofilm lifestyle, inducing significant modifications to the entire population. The biofilm matrix forms a shield to prevent penetration and diffusion of antimicrobials and the increase of biofilm formation in response to antibiotics, illustrated by an enhancing of biofilm matrix, seems to be a defense mechanism of the bacterial community.

In parallel, antibiotics have also been recognized as signaling molecules increasing conjugative transfer although not all antibiotics can potentiate the frequency of conjugation [295–303]. Interestingly, when conjugation occurs in a free-antibiotics environment with a donor strain pre-treated with sub-concentrations of antibiotics, conjugation frequency increased significantly [304,305]. The mechanisms by which antibiotics affect plasmid transfer remains unclear. In the literature, it is proposed that sub-MIC antibiotics treatment enhance the frequency of conjugation

through an up-regulation of the *tra* gene expression in donors [298,304–307]. However, in many studies, the increase of conjugation frequency was evaluated using the antibiotic which resistance gene is carried by the tested conjugative plasmid itself [295,297,298,300,301,307,308]. This approach makes it difficult to distinguish between the selection bias induced by the antibiotic on the mating population, and the actual effect on conjugation frequencies. Two studies rather support that antibiotics primarily acts through differential selection of the donor, recipient and transconjugant once gene transfer has occurred, rather than stimulating conjugation *per se* [202,309]. By using living cell microscopy, they were able to visualize in real time conjugation dynamics. Nolivos *et al.* showed that transfer frequency of the F plasmid, harboring the tetracycline gene resistance, is not increased by the presence of tetracycline. Lopatkin *et al.* also demonstrated that antibiotics, from six major classes, not significantly increase the conjugation efficiency of plasmids from five different incompatibility groups. The reports suggested that the direct contribution of antibiotics on gene transfer have been overestimated and proposed that antibiotics may act only as post-transfer selection drivers, favoring the growth of transconjugants over recipients. Even if a number of studies have advanced the potential stimulating effect of antibiotics on conjugation, their real impact needs to be further explored and antibiotics with different modes of action tested.

Undoubtedly, antibiotics play a role that must not be overlooked in the emergence of new multi-resistant pathogenic strains. It is troubling to admit that antibiotic treatments amplify biofilm formation, then strengthened the difficulty to heal infections biofilm-associated. Not only, antibiotics induce biofilm formation, but they improve gene transfer within the community. Furthermore, profound changes induced by antibiotics allow the invasion of the biofilm by external microorganisms [294]. As biofilms are suitable environments for conjugative transfer, we can easily imagine that antibiotics could potentiate the invasion by a potential donor harboring gene resistance and its dissemination within the biofilm acting as a synergistic factor instead of an antagonist one. Microfluidic technology represents a promising method to investigate, in real-time and without disrupting biofilm structure, the dynamic of conjugation within communities. Recent studies used microfluidic devices combined with confocal microscopy to monitor in real-time plasmid RP4 transfer in mixed *P. putida* and *E. coli* biofilms and in activated sludge [310,311]. They were able to show that the structure and composition of the biofilm could modulate gene transfer routes. Indeed, in *E. coli* biofilms, the explosive spread of transconjugants illustrated the significant role of plasmid transfer, while in the sludge community, vertical gene transfer is more predominant. Using these advanced techniques, it is now more needed to understand how antibiotics can influence gene dissemination within these complex structures.

4. Conclusion

Our current knowledge of the sequence of reactions required for plasmid conjugation is well documented, especially for model plasmids such as the F factor, but also for other plasmids like RP4, R388 or pTi. The combination of genetic and biochemistry approaches has allowed describing the function of key Tra proteins in these reactions. However, even for these systems, the mechanistic functions of the most Tra proteins remain elusive. They have mainly been described in terms of essentiality for the mating pair formation or stabilization, DNA transfer and immunity, without further understanding of their activity at the molecular scale. Furthermore, as emphasized in the review, a number of major fundamental questions remains, such as the pilus ability to conduct the DNA during distant transfer, the existence and nature of a potential signal triggered by the mating pair formation that would activate conjugation, or the role of the leading genes in the early steps of plasmid establishment, for instance. Due to its intimate connection with the dissemination of drug-resistance, conjugation is back in the focus of a global research effort worldwide. Modern experimental approaches should help gaining new insights into the mechanism of conjugation at the molecular and cellular scales, but also regarding the extend of conjugation into natural bacterial communities and its impact on the dissemination of bacterial metabolic traits.

Figure legends

Figure 1. Schematic diagram of the life-cycle of the F plasmid during conjugational transfer from the donor to the recipient cell. This F plasmid backbone is composed of the *tra* regions encoding all genes involved in conjugational transfer (light blue), the origin of transfer *oriT* (red), the leading region (green) which is the first to be transferred into the recipient cell, and the maintenance region (dark blue) involved in plasmid replication and partition. (i) The initiation of conjugation requires expression of the *tra* genes. Part of the produced Tra proteins form the T4SS and the conjugative pilus that will recruit the recipient cell and mediate mating pair stabilization. (ii) Other Tra proteins constitute the relaxosome (TraI, TraM and TraY), which, in combination with the host factor IHF, bind to the *oriT* and prepare the plasmid for transfer by inducing the nicking reaction by the TraI relaxase. (iii) Interaction between the relaxosome and the Type IV Coupling Protein (T4CP) initiate the transfer of the T-strand by the T4SS. (iv and v) Transfer of the TraI-bound T-strand in the recipient is concomitant with the conversion of the ssDNA into dsDNA by Rolling-Circle-Replication (RCR) in the donor. (a) Upon the entry into recipient, the ssDNA T-strand is coated by the host chromosomal SSB and the single-stranded promoter FrpO adopts a stem-loop structure recognized by the host RNA polymerase to initiate the synthesis of RNA primers. (b) TraI performs the circularization of the fully internalized T-strand. (c) The RNA-DNA duplex is recognized by the host DNA polymerase to initiate the complementary strand synthesis reaction. (d) Once the conversion of the ssDNA plasmid into dsDNA is completed, plasmid genes expression results in the phenotypic conversion of the recipient cell into a transconjugant cell.

Figure 2 Activation cascade of *tra* genes expression

Promoter P_J first drives the *traJ* expression (1). Translated TraJ protein binds the P_Y promoter to notably produce TraY that activate the P_M promoter (2), other Tra proteins constituting the T4SS and the relaxase TraI. Once produced, TraM autoregulates its own expression through the P_M promoter and in combination with TraY and TraI forms the relaxosome bound to *oriT*. The activation of this regulatory cascade is modulated by the FinP/FinO complex that repressed at the post-transcriptional level the translation of TraY. Dotted red arrows illustrate the transcription-translation process.

Acknowledgments: The authors thank Yoshiharu Yamaichi, Nelly Dubarry and Erwan Gueguen for helpful discussions. We are also grateful to all research groups contributing to improve our understanding of bacterial DNA conjugation and we apologize for the potentially incomplete citation list.

Author Contributions: Co-First authors C.V, K.G, S.D contributed equally to the work. C.V, K.G, S.D, S.B. and C.L. wrote the manuscript. C.V and K.G prepared the table. S.B. and C.L. prepared the figures.

Funding: This research was funded by French National Research Agency, grant number ANR-18-CE35-0008 to C.L and K. G and ANR-19-ARMB-0006-01 to C.L, S.B and S.D; the University of Lyon for funding to C.V. C. L. also acknowledges the Schlumberger Foundation for Education and Research (FSER 2019) and the Foundation for Innovation in Infectiology FINOVI (AO-2014) for financial support.

Conflicts of Interest: The authors declared no competing interests.

References

1. Lederberg, J.; Tatum, E.L. Gene recombination in *Escherichia coli*. *Nature* **1946**, *158*, 558.
2. Grohmann, E.; Muth, G.; Espinosa, M. Conjugative plasmid transfer in gram-positive bacteria. *Microbiol. Mol. Biol. Rev.* **2003**, *67*, 277–301, table of contents, doi:10.1128/mmbr.67.2.277-301.2003.
3. Cruz, F.D.L.; Frost, L.S.; Meyer, R.J.; Zechner, E.L. Conjugative DNA metabolism in Gram-negative bacteria. *FEMS Microbiology Reviews* **2010**, *34*, 18–40, doi:10.1111/j.1574-6976.2009.00195.x.
4. Davison, J. Genetic exchange between bacteria in the environment. *Plasmid* **1999**, *42*, 73–91, doi:10.1006/plas.1999.1421.

5. Lawrence, J.G. Gene transfer, speciation, and the evolution of bacterial genomes. *Curr. Opin. Microbiol.* **1999**, *2*, 519–523, doi:10.1016/s1369-5274(99)00010-7.
6. Ochman, H.; Lawrence, J.G.; Groisman, E.A. Lateral gene transfer and the nature of bacterial innovation. *Nature* **2000**, *405*, 299–304, doi:10.1038/35012500.
7. Ippen-Ihler, K.; Achtman, M.; Willetts, N. Deletion map of the Escherichia coli K-12 sex factor F: the order of eleven transfer cistrons. *J. Bacteriol.* **1972**, *110*, 857–863, doi:10.1128/JB.110.3.857-863.1972.
8. Penfold, S.S.; Simon, J.; Frost, L.S. Regulation of the expression of the traM gene of the F sex factor of Escherichia coli. *Mol. Microbiol.* **1996**, *20*, 549–558, doi:10.1046/j.1365-2958.1996.5361059.x.
9. Koraimann, G.; Wagner, M.A. Social behavior and decision making in bacterial conjugation. *Front Cell Infect Microbiol* **2014**, *4*, doi:10.3389/fcimb.2014.00054.
10. Timmis, K.N.; Andrés, I.; Achtman, M. Fertility repression of F-like conjugative plasmids: physical mapping of the R6--5 finO and finP cistrons and identification of the finO protein. *Proc. Natl. Acad. Sci. U.S.A.* **1978**, *75*, 5836–5840, doi:10.1073/pnas.75.12.5836.
11. Willetts, N. The transcriptional control of fertility in F-like plasmids. *J. Mol. Biol.* **1977**, *112*, 141–148, doi:10.1016/s0022-2836(77)80161-7.
12. Frost, L.; Lee, S.; Yanchar, N.; Paranchych, W. finP and fisO mutations in FinP anti-sense RNA suggest a model for FinOP action in the repression of bacterial conjugation by the Flac plasmid JCFL0. *Mol. Gen. Genet.* **1989**, *218*, 152–160, doi:10.1007/BF00330578.
13. Koraimann, G.; Teferle, K.; Markolin, G.; Woger, W.; Högenauer, G. The FinOP repressor system of plasmid R1: analysis of the antisense RNA control of traJ expression and conjugative DNA transfer. *Mol. Microbiol.* **1996**, *21*, 811–821, doi:10.1046/j.1365-2958.1996.361401.x.
14. van Biesen, T.; Frost, L.S. Differential levels of fertility inhibition among F-like plasmids are related to the cellular concentration of finO mRNA. *Mol. Microbiol.* **1992**, *6*, 771–780, doi:10.1111/j.1365-2958.1992.tb01527.x.
15. van Biesen, T.; Frost, L.S. The FinO protein of IncF plasmids binds FinP antisense RNA and its target, traJ mRNA, and promotes duplex formation. *Mol. Microbiol.* **1994**, *14*, 427–436.
16. Jerome, L.J.; van Biesen, T.; Frost, L.S. Degradation of FinP antisense RNA from F-like plasmids: the RNA-binding protein, FinO, protects FinP from ribonuclease E. *J. Mol. Biol.* **1999**, *285*, 1457–1473, doi:10.1006/jmbi.1998.2404.
17. Wong, J.J.W.; Lu, J.; Glover, J.N.M. Relaxosome function and conjugation regulation in F-like plasmids – a structural biology perspective. *Molecular Microbiology* **2012**, *85*, 602–617, doi:10.1111/j.1365-2958.2012.08131.x.
18. Will, W.R.; Frost, L.S. Characterization of the Opposing Roles of H-NS and TraJ in Transcriptional Regulation of the F-Plasmid tra Operon. *J Bacteriol* **2006**, *188*, 507–514, doi:10.1128/JB.188.2.507-514.2006.
19. Will, W.R.; Lu, J.; Frost, L.S. The role of H-NS in silencing F transfer gene expression during entry into stationary phase. *Mol. Microbiol.* **2004**, *54*, 769–782, doi:10.1111/j.1365-2958.2004.04303.x.

20. Ali Azam, T.; Iwata, A.; Nishimura, A.; Ueda, S.; Ishihama, A. Growth phase-dependent variation in protein composition of the Escherichia coli nucleoid. *J. Bacteriol.* **1999**, *181*, 6361–6370, doi:10.1128/JB.181.20.6361-6370.1999.
21. Frost, L.S.; Manchak, J. F-phenocopies: characterization of expression of the F transfer region in stationary phase. *Microbiology (Reading, Engl.)* **1998**, *144* (Pt 9), 2579–2587, doi:10.1099/00221287-144-9-2579.
22. Headd, B.; Bradford, S.A. The Conjugation Window in an Escherichia coli K-12 Strain with an IncFII Plasmid. *Appl. Environ. Microbiol.* **2020**, *86*, doi:10.1128/AEM.00948-20.
23. Lu, J.; Peng, Y.; Wan, S.; Frost, L.S.; Raivio, T.; Glover, J.N.M. Cooperative Function of TraJ and ArcA in Regulating the F Plasmid tra Operon. *Journal of Bacteriology* **2019**, *201*, doi:10.1128/JB.00448-18.
24. Camacho, E.M.; Serna, A.; Madrid, C.; Marqués, S.; Fernández, R.; de la Cruz, F.; Juárez, A.; Casadesús, J. Regulation of finP Transcription by DNA Adenine Methylation in the Virulence Plasmid of Salmonella enterica. *J Bacteriol* **2005**, *187*, 5691–5699, doi:10.1128/JB.187.16.5691-5699.2005.
25. Zahrl, D.; Wagner, A.; Tscherner, M.; Koraimann, G. GroEL plays a central role in stress-induced negative regulation of bacterial conjugation by promoting proteolytic degradation of the activator protein TraJ. *J. Bacteriol.* **2007**, *189*, 5885–5894, doi:10.1128/JB.00005-07.
26. Dessaux, Y.; Faure, D. Quorum Sensing and Quorum Quenching in Agrobacterium: A “Go/No Go System”? *Genes (Basel)* **2018**, *9*, doi:10.3390/genes9040210.
27. Lu, Y.; Zeng, J.; Wu, B.; E, S.; Wang, L.; Cai, R.; Zhang, N.; Li, Y.; Huang, X.; Huang, B.; et al. Quorum Sensing N-acyl Homoserine Lactones-SdiA Suppresses Escherichia coli-Pseudomonas aeruginosa Conjugation through Inhibiting traI Expression. *Front Cell Infect Microbiol* **2017**, *7*, doi:10.3389/fcimb.2017.00007.
28. Cheah, K.C.; Skurray, R. The F plasmid carries an IS3 insertion within finO. *J. Gen. Microbiol.* **1986**, *132*, 3269–3275, doi:10.1099/00221287-132-12-3269.
29. Yoshioka, Y.; Ohtsubo, H.; Ohtsubo, E. Repressor gene finO in plasmids R100 and F: constitutive transfer of plasmid F is caused by insertion of IS3 into F finO. *J Bacteriol* **1987**, *169*, 619–623.
30. Poidevin, M.; Sato, M.; Altinoglu, I.; Delaplace, M.; Sato, C.; Yamaichi, Y. Mutation in ESBL Plasmid from Escherichia coli O104:H4 Leads Autoagglutination and Enhanced Plasmid Dissemination. *Front Microbiol* **2018**, *9*, 130, doi:10.3389/fmicb.2018.00130.
31. Yamaichi, Y.; Chao, M.C.; Sasabe, J.; Clark, L.; Davis, B.M.; Yamamoto, N.; Mori, H.; Kurokawa, K.; Waldor, M.K. High-resolution genetic analysis of the requirements for horizontal transmission of the ESBL plasmid from Escherichia coli O104:H4. *Nucleic Acids Res.* **2015**, *43*, 348–360, doi:10.1093/nar/gku1262.
32. Dmowski, M.; Gołębiewski, M.; Kern-Zdanowicz, I. Characteristics of the Conjugative Transfer System of the IncM Plasmid pCTX-M3 and Identification of Its Putative Regulators. *J. Bacteriol.* **2018**, *200*, doi:10.1128/JB.00234-18.
33. Kohler, V.; Goessweiner-Mohr, N.; Aufschnaiter, A.; Fercher, C.; Probst, I.; Pavkov-Keller, T.; Hunger, K.; Wolinski, H.; Büttner, S.; Grohmann, E.; et al. TraN: A novel repressor of an Enterococcus conjugative type IV secretion system. *Nucleic Acids Res.* **2018**, *46*, 9201–9219, doi:10.1093/nar/gky671.

34. Potron, A.; Poirel, L.; Nordmann, P. Derepressed transfer properties leading to the efficient spread of the plasmid encoding carbapenemase OXA-48. *Antimicrob. Agents Chemother.* **2014**, *58*, 467–471, doi:10.1128/AAC.01344-13.
35. Bradley, D.E. Derepressed plasmids of incompatibility group I1 determine two different morphological forms of pilus. *Plasmid* **1983**, *9*, 331–334, doi:10.1016/0147-619x(83)90011-2.
36. Bradley, D.E. Specification of the conjugative pili and surface mating systems of *Pseudomonas* plasmids. *J. Gen. Microbiol.* **1983**, *129*, 2545–2556, doi:10.1099/00221287-129-8-2545.
37. Bradley, D.E. Characteristics and function of thick and thin conjugative pili determined by transfer-derepressed plasmids of incompatibility groups I1, I2, I5, B, K and Z. *J. Gen. Microbiol.* **1984**, *130*, 1489–1502, doi:10.1099/00221287-130-6-1489.
38. Bradley, D.E.; Taylor, D.E.; Cohen, D.R. Specification of surface mating systems among conjugative drug resistance plasmids in *Escherichia coli* K-12. *J. Bacteriol.* **1980**, *143*, 1466–1470, doi:10.1128/JB.143.3.1466-1470.1980.
39. Costa, T.R.D.; Ilangovan, A.; Ukleja, M.; Redzej, A.; Santini, J.M.; Smith, T.K.; Egelman, E.H.; Waksman, G. Structure of the Bacterial Sex F Pilus Reveals an Assembly of a Stoichiometric Protein-Phospholipid Complex. *Cell* **2016**, *166*, 1436–1444.e10, doi:10.1016/j.cell.2016.08.025.
40. Date, T.; Inuzuka, M.; Tomoeda, M. Purification and characterization of F pili from *Escherichia coli*. *Biochemistry* **1977**, *16*, 5579–5585, doi:10.1021/bi00644a030.
41. Folkhard, W.; Leonard, K.R.; Malsey, S.; Marvin, D.A.; Dubochet, J.; Engel, A.; Achtman, M.; Helmuth, R. X-ray diffraction and electron microscope studies on the structure of bacterial F pili. *J. Mol. Biol.* **1979**, *130*, 145–160, doi:10.1016/0022-2836(79)90423-6.
42. Ippen-Ihler, K.A.; Minkley, E.G. The conjugation system of F, the fertility factor of *Escherichia coli*. *Annu. Rev. Genet.* **1986**, *20*, 593–624, doi:10.1146/annurev.ge.20.120186.003113.
43. Silverman, P.M.; Clarke, M.B. New insights into F-pilus structure, dynamics, and function. *Integr Biol (Camb)* **2010**, *2*, 25–31, doi:10.1039/b917761b.
44. Wang, Y.A.; Yu, X.; Silverman, P.M.; Harris, R.L.; Egelman, E.H. The structure of F-pili. *J. Mol. Biol.* **2009**, *385*, 22–29, doi:10.1016/j.jmb.2008.10.054.
45. Frost, L.S.; Paranchych, W.; Willetts, N.S. DNA sequence of the F traALE region that includes the gene for F pilin. *J. Bacteriol.* **1984**, *160*, 395–401, doi:10.1128/JB.160.1.395-401.1984.
46. Minkley, E.G.; Polen, S.; Brinton, C.C.; Ippen-Ihler, K. Identification of the structural gene for F-pilin. *J. Mol. Biol.* **1976**, *108*, 111–121, doi:10.1016/s0022-2836(76)80098-8.
47. Maneewannakul, K.; Maneewannakul, S.; Ippen-Ihler, K. Synthesis of F pilin. *J. Bacteriol.* **1993**, *175*, 1384–1391, doi:10.1128/jb.175.5.1384-1391.1993.
48. Harris, R.L.; Sholl, K.A.; Conrad, M.N.; Dresser, M.E.; Silverman, P.M. Interaction between the F plasmid TraA (F-pilin) and TraQ proteins. *Mol. Microbiol.* **1999**, *34*, 780–791, doi:10.1046/j.1365-2958.1999.01640.x.
49. Majdalani, N.; Ippen-Ihler, K. Membrane insertion of the F-pilin subunit is Sec independent but requires leader peptidase B and the proton motive force. *J Bacteriol* **1996**, *178*, 3742–3747.
50. Moore, D.; Hamilton, C.M.; Maneewannakul, K.; Mintz, Y.; Frost, L.S.; Ippen-Ihler, K. The *Escherichia coli* K-12 F plasmid gene traX is required for acetylation of F pilin. *J. Bacteriol.* **1993**, *175*, 1375–1383, doi:10.1128/jb.175.5.1375-1383.1993.

51. Firth, N.; Berg, T.; Skurray, R.A. Evolution of conjugative plasmids from gram-positive bacteria. *Mol. Microbiol.* **1999**, *31*, 1598–1600.
52. Lawley, T.D.; Klimke, W.A.; Gubbins, M.J.; Frost, L.S. F factor conjugation is a true type IV secretion system. *FEMS Microbiol. Lett.* **2003**, *224*, 1–15, doi:10.1016/S0378-1097(03)00430-0.
53. Manchak, J.; Anthony, K.G.; Frost, L.S. Mutational analysis of F-pilin reveals domains for pilus assembly, phage infection and DNA transfer. *Mol. Microbiol.* **2002**, *43*, 195–205, doi:10.1046/j.1365-2958.2002.02731.x.
54. Firth, N.; Ippen-Ihler, K.; Skurray, R.A. Gene Transfer : Conjugation Structure and Function of the F Factor and Mechanism of Conjugation.; 1999.
55. Koraimann, G. Spread and Persistence of Virulence and Antibiotic Resistance Genes: A Ride on the F Plasmid Conjugation Module. *EcoSal Plus* **2018**, *8*, doi:10.1128/ecosalplus.ESP-0003-2018.
56. Arutyunov, D.; Frost, L.S. F conjugation: back to the beginning. *Plasmid* **2013**, *70*, 18–32, doi:10.1016/j.plasmid.2013.03.010.
57. Frost, L.S.; Ippen-Ihler, K.; Skurray, R.A. Analysis of the sequence and gene products of the transfer region of the F sex factor. *Microbiol. Rev.* **1994**, *58*, 162–210.
58. Anthony, K.G.; Kathir, P.; Moore, D.; Ippen-Ihler, K.; Frost, L.S. Analysis of the traLEKBP sequence and the TraP protein from three F-like plasmids: F, R100-1 and ColB2. *J. Bacteriol.* **1996**, *178*, 3194–3200, doi:10.1128/jb.178.11.3194-3200.1996.
59. Anthony, K.G.; Klimke, W.A.; Manchak, J.; Frost, L.S. Comparison of Proteins Involved in Pilus Synthesis and Mating Pair Stabilization from the Related Plasmids F and R100-1: Insights into the Mechanism of Conjugation. *J Bacteriol* **1999**, *181*, 5149–5159.
60. Gopalkrishnan, S.; Ross, W.; Chen, A.Y.; Gourse, R.L. TraR directly regulates transcription initiation by mimicking the combined effects of the global regulators DksA and ppGpp. *Proc. Natl. Acad. Sci. U.S.A.* **2017**, *114*, E5539–E5548, doi:10.1073/pnas.1704105114.
61. Shala-Lawrence, A.; Bragagnolo, N.; Nowroozi-Dayeni, R.; Kheyson, S.; Audette, G.F. The interaction of TraW and TrbC is required to facilitate conjugation in F-like plasmids. *Biochem. Biophys. Res. Commun.* **2018**, *503*, 2386–2392, doi:10.1016/j.bbrc.2018.06.166.
62. Garcillán-Barcia, M.P.; de la Cruz, F. Why is entry exclusion an essential feature of conjugative plasmids? *Plasmid* **2008**, *60*, 1–18, doi:10.1016/j.plasmid.2008.03.002.
63. Achtman, M. Mating aggregates in Escherichia coli conjugation. *J. Bacteriol.* **1975**, *123*, 505–515, doi:10.1128/JB.123.2.505-515.1975.
64. Lanka, E.; Wilkins, B.M. DNA PROCESSING REACTIONS IN BACTERIAL CONJUGATION. 31.
65. Dostál, L.; Shao, S.; Schildbach, J.F. Tracking F plasmid TraI relaxase processing reactions provides insight into F plasmid transfer. *Nucleic Acids Res.* **2011**, *39*, 2658–2670, doi:10.1093/nar/gkq1137.
66. Matson, S.W.; Sampson, J.K.; Byrd, D.R.N. F Plasmid Conjugative DNA Transfer THE TraI HELICASE ACTIVITY IS ESSENTIAL FOR DNA STRAND TRANSFER. *J. Biol. Chem.* **2001**, *276*, 2372–2379, doi:10.1074/jbc.M008728200.
67. Curtiss, R. Bacterial conjugation. *Annu. Rev. Microbiol.* **1969**, *23*, 69–136, doi:10.1146/annurev.mi.23.100169.000441.
68. Marvin, D.A.; Hohn, B. Filamentous bacterial viruses. *Bacteriol Rev* **1969**, *33*, 172–209.

69. Clarke, M.; Maddera, L.; Harris, R.L.; Silverman, P.M. F-pili dynamics by live-cell imaging. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 17978–17981, doi:10.1073/pnas.0806786105.
70. Achtman, M.; Morelli, G.; Schwuchow, S. Cell-cell interactions in conjugating *Escherichia coli*: role of F pili and fate of mating aggregates. *J. Bacteriol.* **1978**, *135*, 1053–1061, doi:10.1128/JB.135.3.1053-1061.1978.
71. Dürrenberger, M.B.; Villiger, W.; Bächli, T. Conjugational junctions: morphology of specific contacts in conjugating *Escherichia coli* bacteria. *J. Struct. Biol.* **1991**, *107*, 146–156, doi:10.1016/1047-8477(91)90018-r.
72. Brinton, C.C. The structure, function, synthesis and genetic control of bacterial pili and a molecular model for DNA and RNA transport in gram negative bacteria. *Trans N Y Acad Sci* **1965**, *27*, 1003–1054, doi:10.1111/j.2164-0947.1965.tb02342.x.
73. Harrington, L.C.; Rogerson, A.C. The F pilus of *Escherichia coli* appears to support stable DNA transfer in the absence of wall-to-wall contact between cells. *J. Bacteriol.* **1990**, *172*, 7263–7264, doi:10.1128/jb.172.12.7263-7264.1990.
74. Babić, A.; Lindner, A.B.; Vulić, M.; Stewart, E.J.; Radman, M. Direct Visualization of Horizontal Gene Transfer. *Science* **2008**, *319*, 1533–1536, doi:10.1126/science.1153498.
75. Davies, J.K.; Reeves, P. Colicin tolerance and map location of conjugation-deficient mutants. *J. Bacteriol.* **1975**, *123*, 372–373, doi:10.1128/JB.123.1.372-373.1975.
76. Falkinham, J.O.; Curtiss, R. Isolation and characterization of conjugation-deficient mutants of *Escherichia coli* K-12. *J. Bacteriol.* **1976**, *126*, 1194–1206, doi:10.1128/JB.126.3.1194-1206.1976.
77. Havekes, L.M.; Hoekstra, W.P. Characterization of an *Escherichia coli* K-12 F-Con-mutant. *J. Bacteriol.* **1976**, *126*, 593–600.
78. Manning, P.A.; Puspurs, A.; Reeves, P. Outer membrane of *Escherichia coli* K-12: isolation of mutants with altered protein 3A by using host range mutants of bacteriophage K3. *J. Bacteriol.* **1976**, *127*, 1080–1084.
79. Monner, D.A.; Jonsson, S.; Boman, H.G. Ampicillin-resistant mutants of *Escherichia coli* K-12 with lipopolysaccharide alterations affecting mating ability and susceptibility to sex-specific bacteriophages. *J. Bacteriol.* **1971**, *107*, 420–432, doi:10.1128/JB.107.2.420-432.1971.
80. Skurray, R.A.; Hancock, R.E.; Reeves, P. Con--mutants: class of mutants in *Escherichia coli* K-12 lacking a major cell wall protein and defective in conjugation and adsorption of a bacteriophage. *J. Bacteriol.* **1974**, *119*, 726–735.
81. Havekes, L.M.; Hoekstra, W.P. Characterization of an *Escherichia coli* K-12 F-Con-mutant. *J. Bacteriol.* **1976**, *126*, 593–600, doi:10.1128/JB.126.2.593-600.1976.
82. Manoil, C.; Rosenbusch, J.P. Conjugation-deficient mutants of *Escherichia coli* distinguish classes of functions of the outer membrane OmpA protein. *Mol. Gen. Genet.* **1982**, *187*, 148–156, doi:10.1007/BF00384398.
83. Morona, R.; Klose, M.; Henning, U. *Escherichia coli* K-12 outer membrane protein (OmpA) as a bacteriophage receptor: analysis of mutant genes expressing altered proteins. *J. Bacteriol.* **1984**, *159*, 570–578, doi:10.1128/JB.159.2.570-578.1984.
84. Skurray, R.A.; Hancock, R.E.; Reeves, P. Con--mutants: class of mutants in *Escherichia coli* K-12 lacking a major cell wall protein and defective in conjugation and adsorption of a bacteriophage. *J. Bacteriol.* **1974**, *119*, 726–735, doi:10.1128/JB.119.3.726-735.1974.

85. Achtman, M.; Schwuchow, S.; Helmuth, R.; Morelli, G.; Manning, P.A. Cell-cell interactions in conjugating *Escherichia coli*: Con- mutants and stabilization of mating aggregates. *Molecular and General Genetics MGG* **1978**, *164*, 171–183, doi:10.1007/BF00267382.
86. Anthony, K.G.; Sherburne, C.; Sherburne, R.; Frost, L.S. The role of the pilus in recipient cell recognition during bacterial conjugation mediated by F-like plasmids. *Mol. Microbiol.* **1994**, *13*, 939–953, doi:10.1111/j.1365-2958.1994.tb00486.x.
87. Klimke, W.A.; Frost, L.S. Genetic analysis of the role of the transfer gene, traN, of the F and R100-1 plasmids in mating pair stabilization during conjugation. *J. Bacteriol.* **1998**, *180*, 4036–4043.
88. Klimke, W.A.; Rypien, C.D.; Klinger, B.; Kennedy, R.A.; Rodriguez-Maillard, J.M.; Frost, L.S. The mating pair stabilization protein, TraN, of the F plasmid is an outer-membrane protein with two regions that are important for its function in conjugation. *Microbiology* **2005**, *151*, 3527–3540, doi:10.1099/mic.0.28025-0.
89. Maneewannakul, S.; Kathir, P.; Ippen-Ihler, K. Characterization of the F plasmid mating aggregation gene traN and of a new F transfer region locus trbE. *J. Mol. Biol.* **1992**, *225*, 299–311, doi:10.1016/0022-2836(92)90923-8.
90. Anthony, K.G.; Klimke, W.A.; Manchak, J.; Frost, L.S. Comparison of proteins involved in pilus synthesis and mating pair stabilization from the related plasmids F and R100-1: insights into the mechanism of conjugation. *J. Bacteriol.* **1999**, *181*, 5149–5159.
91. Willetts, N.; Achtman, M. Genetic analysis of transfer by the *Escherichia coli* sex factor F, using P1 transductional complementation. *J. Bacteriol.* **1972**, *110*, 843–851, doi:10.1128/JB.110.3.843-851.1972.
92. Audette, G.F.; Manchak, J.; Beatty, P.; Klimke, W.A.; Frost, L.S. Entry exclusion in F-like plasmids requires intact TraG in the donor that recognizes its cognate TraS in the recipient. *Microbiology (Reading, Engl.)* **2007**, *153*, 442–451, doi:10.1099/mic.0.2006/001917-0.
93. Marrero, J.; Waldor, M.K. Determinants of entry exclusion within Eex and TraG are cytoplasmic. *J. Bacteriol.* **2007**, *189*, 6469–6473, doi:10.1128/JB.00522-07.
94. Manoil, C.; Rosenbusch, J.P. Conjugation-deficient mutants of *Escherichia coli* distinguish classes of functions of the outer membrane OmpA protein. *Mol. Gen. Genet.* **1982**, *187*, 148–156, doi:10.1007/bf00384398.
95. Low, W.W.; Wong, J.; Pena, A.; Seddon, C.; Costa, T.; Beis, K.; Frankel, G. *OmpK36 and TraN facilitate conjugal transfer of the Klebsiella pneumoniae carbapenem resistance plasmid pKpQIL*; *Microbiology*, 2020;
96. Falkinham, J.O.; Curtiss, R. Isolation and characterization of conjugation-deficient mutants of *Escherichia coli* K-12. *J. Bacteriol.* **1976**, *126*, 1194–1206.
97. Havekes, L.; Tommassen, J.; Hoekstra, W.; Lugtenberg, B. Isolation and characterization of *Escherichia coli* K-12 F- mutants defective in conjugation with an I-type donor. *Journal of Bacteriology* **1977**, *129*, 1–8, doi:10.1128/JB.129.1.1-8.1977.
98. Ishiwa, A.; Komano, T. PilV Adhesins of Plasmid R64 Thin Pili Specifically Bind to the Lipopolysaccharides of Recipient Cells. *Journal of Molecular Biology* **2004**, *343*, 615–625, doi:10.1016/j.jmb.2004.08.059.

99. Frost, L.S.; Simon, J. Studies on the pili of the promiscuous plasmid RP4. In *Molecular Mechanisms of Bacterial Virulence*; Kado, C.I., Crosa, J.H., Eds.; Developments in Plant Pathology; Springer Netherlands: Dordrecht, 1994; Vol. 3, pp. 47–65 ISBN 978-94-010-4322-9.
100. Moriguchi, K.; Zoolkefli, F.I.R.M.; Abe, M.; Kiyokawa, K.; Yamamoto, S.; Suzuki, K. Targeting Antibiotic Resistance Genes Is a Better Approach to Block Acquisition of Antibiotic Resistance Than Blocking Conjugal Transfer by Recipient Cells: A Genome-Wide Screening in *Escherichia coli*. *Frontiers in Microbiology* **2020**, *10*, doi:10.3389/fmicb.2019.02939.
101. Pérez-Mendoza, D.; de la Cruz, F. *Escherichia coli* genes affecting recipient ability in plasmid conjugation: Are there any? *BMC Genomics* **2009**, *10*, 71, doi:10.1186/1471-2164-10-71.
102. Dodsworth, J.A.; Li, L.; Wei, S.; Hedlund, B.P.; Leigh, J.A.; de Figueiredo, P. Interdomain Conjugal Transfer of DNA from Bacteria to Archaea. *Applied and Environmental Microbiology* **2010**, *76*, 5644–5647, doi:10.1128/AEM.00967-10.
103. Heinemann, J.A.; Sprague, G.F. Bacterial conjugative plasmids mobilize DNA transfer between bacteria and yeast. *Nature* **1989**, *340*, 205–209, doi:10.1038/340205a0.
104. Waters, V.L. Conjugation between bacterial and mammalian cells. *Nature Genetics* **2001**, *29*, 375–376, doi:10.1038/ng779.
105. Llosa, M.; Gomis-Rüth, F.X.; Coll, M.; de la Cruz Fd, F. Bacterial conjugation: a two-step mechanism for DNA transport. *Mol. Microbiol.* **2002**, *45*, 1–8, doi:10.1046/j.1365-2958.2002.03014.x.
106. Clewell, D.B.; Helinski, D.E. Existence of the colicinogenic factor-sex factor Coll-b-P9 as a supercoiled circular DNA-protein relaxation complex. *Biochem. Biophys. Res. Commun.* **1970**, *41*, 150–156, doi:10.1016/0006-291x(70)90481-x.
107. Cohen, A.; Fisher, W.D.; Curtiss, R.; Adler, H.I. DNA isolated from *Escherichia coli* minicells mated with F⁺ cells. *Proc. Natl. Acad. Sci. U.S.A.* **1968**, *61*, 61–68, doi:10.1073/pnas.61.1.61.
108. Willetts, N.; Skurray, R. The conjugation system of F-like plasmids. *Annu. Rev. Genet.* **1980**, *14*, 41–76, doi:10.1146/annurev.ge.14.120180.000353.
109. Dostál, L.; Schildbach, J.F. Single-Stranded DNA Binding by F TraI Relaxase and Helicase Domains Is Coordinately Regulated. *J Bacteriol* **2010**, *192*, 3620–3628, doi:10.1128/JB.00154-10.
110. Everett, R.; Willetts, N. Characterisation of an in vivo system for nicking at the origin of conjugal DNA transfer of the sex factor F. *J. Mol. Biol.* **1980**, *136*, 129–150, doi:10.1016/0022-2836(80)90309-5.
111. Matson, S.W.; Morton, B.S. *Escherichia coli* DNA helicase I catalyzes a site- and strand-specific nicking reaction at the F plasmid oriT. *J. Biol. Chem.* **1991**, *266*, 16232–16237.
112. Matson, S.W.; Ragonese, H. The F-plasmid TraI protein contains three functional domains required for conjugative DNA strand transfer. *J. Bacteriol.* **2005**, *187*, 697–706, doi:10.1128/JB.187.2.697-706.2005.
113. Reygers, U.; Wessel, R.; Müller, H.; Hoffmann-Berling, H. Endonuclease activity of *Escherichia coli* DNA helicase I directed against the transfer origin of the F factor. *EMBO J.* **1991**, *10*, 2689–2694.
114. Traxler, B.A.; Minkley, E.G. Evidence that DNA helicase I and oriT site-specific nicking are both functions of the F TraI protein. *J. Mol. Biol.* **1988**, *204*, 205–209, doi:10.1016/0022-2836(88)90609-2.

115. Howard, M.T.; Nelson, W.C.; Matson, S.W. Stepwise assembly of a relaxosome at the F plasmid origin of transfer. *J. Biol. Chem.* **1995**, *270*, 28381–28386.
116. Nelson, W.C.; Morton, B.S.; Lahue, E.E.; Matson, S.W. Characterization of the Escherichia coli F factor traY gene product and its binding sites. *J. Bacteriol.* **1993**, *175*, 2221–2228, doi:10.1128/jb.175.8.2221-2228.1993.
117. Schildbach, J.F.; Robinson, C.R.; Sauer, R.T. Biophysical characterization of the TraY protein of Escherichia coli F factor. *J. Biol. Chem.* **1998**, *273*, 1329–1333, doi:10.1074/jbc.273.3.1329.
118. Luo, Y.; Gao, Q.; Deonier, R.C. Mutational and physical analysis of F plasmid traY protein binding to oriT. *Mol. Microbiol.* **1994**, *11*, 459–469, doi:10.1111/j.1365-2958.1994.tb00327.x.
119. Nelson, W.C.; Howard, M.T.; Sherman, J.A.; Matson, S.W. The traY gene product and integration host factor stimulate Escherichia coli DNA helicase I-catalyzed nicking at the F plasmid oriT. *J. Biol. Chem.* **1995**, *270*, 28374–28380.
120. Rice, P.A.; Yang, S.; Mizuuchi, K.; Nash, H.A. Crystal structure of an IHF-DNA complex: a protein-induced DNA U-turn. *Cell* **1996**, *87*, 1295–1306, doi:10.1016/s0092-8674(00)81824-3.
121. Tsai, M.M.; Fu, Y.H.; Deonier, R.C. Intrinsic bends and integration host factor binding at F plasmid oriT. *J. Bacteriol.* **1990**, *172*, 4603–4609, doi:10.1128/jb.172.8.4603-4609.1990.
122. Williams, S.L.; Schildbach, J.F. TraY and integration host factor oriT binding sites and F conjugal transfer: sequence variations, but not altered spacing, are tolerated. *J. Bacteriol.* **2007**, *189*, 3813–3823, doi:10.1128/JB.01783-06.
123. Di Laurenzio, L.; Frost, L.S.; Paranchych, W. The TraM protein of the conjugative plasmid F binds to the origin of transfer of the F and ColE1 plasmids. *Mol. Microbiol.* **1992**, *6*, 2951–2959, doi:10.1111/j.1365-2958.1992.tb01754.x.
124. Fu, Y.H.; Tsai, M.M.; Luo, Y.N.; Deonier, R.C. Deletion analysis of the F plasmid oriT locus. *J. Bacteriol.* **1991**, *173*, 1012–1020, doi:10.1128/jb.173.3.1012-1020.1991.
125. Kupelwieser, G.; Schwab, M.; Högenauer, G.; Koraimann, G.; Zechner, E.L. Transfer protein TraM stimulates TraI-catalyzed cleavage of the transfer origin of plasmid R1 in vivo. *J. Mol. Biol.* **1998**, *275*, 81–94, doi:10.1006/jmbi.1997.1436.
126. Mihajlovic, S.; Lang, S.; Sut, M.V.; Strohmaier, H.; Gruber, C.J.; Koraimann, G.; Cabezón, E.; Moncalián, G.; de la Cruz, F.; Zechner, E.L. Plasmid r1 conjugative DNA processing is regulated at the coupling protein interface. *J. Bacteriol.* **2009**, *191*, 6877–6887, doi:10.1128/JB.00918-09.
127. Ragonese, H.; Haisch, D.; Villareal, E.; Choi, J.-H.; Matson, S.W. The F plasmid-encoded TraM protein stimulates relaxosome-mediated cleavage at oriT through an interaction with TraI. *Molecular Microbiology* **2007**, *63*, 1173–1184, doi:10.1111/j.1365-2958.2006.05576.x.
128. Wawrzyniak, P.; Plucienniczak, G.; Bartosik, D. The Different Faces of Rolling-Circle Replication and Its Multifunctional Initiator Proteins. *Front Microbiol* **2017**, *8*, 2353, doi:10.3389/fmicb.2017.02353.
129. Byrd, D.R.; Matson, S.W. Nicking by transesterification: the reaction catalysed by a relaxase. *Mol. Microbiol.* **1997**, *25*, 1011–1022, doi:10.1046/j.1365-2958.1997.5241885.x.
130. Guiney, D.G.; Helinski, D.R. Relaxation complexes of poasmid DNA and protein. III. Association of protein with the 5' terminus of the broken DNA strand in the relaxed complex of plasmid ColE1. *J. Biol. Chem.* **1975**, *250*, 8796–8803.

131. Matson, S.W.; Nelson, W.C.; Morton, B.S. Characterization of the reaction product of the oriT nicking reaction catalyzed by Escherichia coli DNA helicase I. *J. Bacteriol.* **1993**, *175*, 2599–2606, doi:10.1128/jb.175.9.2599-2606.1993.
132. Pansegrau, W.; Ziegelin, G.; Lanka, E. Covalent association of the traI gene product of plasmid RP4 with the 5'-terminal nucleotide at the relaxation nick site. *J. Biol. Chem.* **1990**, *265*, 10637–10644.
133. Khan, S.A. Rolling-circle replication of bacterial plasmids. *Microbiol. Mol. Biol. Rev.* **1997**, *61*, 442–455.
134. Khan, S.A. Plasmid rolling-circle replication: highlights of two decades of research. *Plasmid* **2005**, *53*, 126–136, doi:10.1016/j.plasmid.2004.12.008.
135. Ruiz-Masó, J.A.; Machón, C.; Bordanaba-Ruiseco, L.; Espinosa, M.; Coll, M.; Del Solar, G. Plasmid Rolling-Circle Replication. *Microbiol Spectr* **2015**, *3*, PLAS-0035-2014, doi:10.1128/microbiolspec.PLAS-0035-2014.
136. Waters, V.L.; Guiney, D.G. Processes at the nick region link conjugation, T-DNA transfer and rolling circle replication. *Mol. Microbiol.* **1993**, *9*, 1123–1130, doi:10.1111/j.1365-2958.1993.tb01242.x.
137. Lorenzo-Díaz, F.; Fernández-López, C.; Garcillán-Barcia, M.P.; Espinosa, M. Bringing them together: Plasmid pMV158 rolling circle replication and conjugation under an evolutionary perspective. *Plasmid* **2014**, *74*, 15–31, doi:10.1016/j.plasmid.2014.05.004.
138. Draper, O.; César, C.E.; Machón, C.; de la Cruz, F.; Llosa, M. Site-specific recombinase and integrase activities of a conjugative relaxase in recipient cells. *Proc Natl Acad Sci U S A* **2005**, *102*, 16385–16390, doi:10.1073/pnas.0506081102.
139. Trokter, M.; Waksman, G. Translocation through the Conjugative Type IV Secretion System Requires Unfolding of Its Protein Substrate. *J Bacteriol* **2018**, *200*, doi:10.1128/JB.00615-17.
140. Ilangovan, A.; Kay, C.W.M.; Roier, S.; El Mkami, H.; Salvadori, E.; Zechner, E.L.; Zanetti, G.; Waksman, G. Cryo-EM Structure of a Relaxase Reveals the Molecular Basis of DNA Unwinding during Bacterial Conjugation. *Cell* **2017**, *169*, 708-721.e12, doi:10.1016/j.cell.2017.04.010.
141. Llosa, M.; Gomis-Rüth, F.X.; Coll, M.; de la Cruz Fd, F. Bacterial conjugation: a two-step mechanism for DNA transport. *Mol. Microbiol.* **2002**, *45*, 1–8, doi:10.1046/j.1365-2958.2002.03014.x.
142. Cabezón, E.; Sastre, J.I.; de la Cruz, F. Genetic evidence of a coupling role for the TraG protein family in bacterial conjugation. *Mol. Gen. Genet.* **1997**, *254*, 400–406, doi:10.1007/s004380050432.
143. Cascales, E.; Christie, P.J. Definition of a bacterial type IV secretion pathway for a DNA substrate. *Science* **2004**, *304*, 1170–1173, doi:10.1126/science.1095211.
144. Christie, P.J. Type IV secretion: the Agrobacterium VirB/D4 and related conjugation systems. *Biochim. Biophys. Acta* **2004**, *1694*, 219–234, doi:10.1016/j.bbamcr.2004.02.013.
145. Christie, P.J. The Mosaic Type IV Secretion Systems. *EcoSal Plus* **2016**, *7*, doi:10.1128/ecosalplus.ESP-0020-2015.
146. Christie, P.J.; Whitaker, N.; González-Rivera, C. BBA Review Revised Mechanism and Structure of the Bacterial Type IV Secretion Systems. *Biochimica et biophysica acta* **2014**, *1843*, 1578, doi:10.1016/j.bbamcr.2013.12.019.

147. Beranek, A.; Zettl, M.; Lorenzoni, K.; Schauer, A.; Manhart, M.; Koraimann, G. Thirty-eight C-terminal amino acids of the coupling protein TraD of the F-like conjugative resistance plasmid R1 are required and sufficient to confer binding to the substrate selector protein TraM. *J. Bacteriol.* **2004**, *186*, 6999–7006, doi:10.1128/JB.186.20.6999-7006.2004.
148. Disqué-Kochem, C.; Dreiseiklmann, B. The cytoplasmic DNA-binding protein TraM binds to the inner membrane protein TraD in vitro. *J. Bacteriol.* **1997**, *179*, 6133–6137, doi:10.1128/jb.179.19.6133-6137.1997.
149. Lu, J.; Frost, L.S. Mutations in the C-terminal region of TraM provide evidence for in vivo TraM-TraD interactions during F-plasmid conjugation. *J. Bacteriol.* **2005**, *187*, 4767–4773, doi:10.1128/JB.187.14.4767-4773.2005.
150. Lu, J.; Zhao, W.; Frost, L.S. Mutational analysis of TraM correlates oligomerization and DNA binding with autoregulation and conjugative DNA transfer. *J. Biol. Chem.* **2004**, *279*, 55324–55333, doi:10.1074/jbc.M409352200.
151. Lu, J.; Wong, J.J.W.; Edwards, R.A.; Manchak, J.; Frost, L.S.; Glover, J.N.M. Structural basis of specific TraD-TraM recognition during F plasmid-mediated bacterial conjugation. *Mol. Microbiol.* **2008**, *70*, 89–99, doi:10.1111/j.1365-2958.2008.06391.x.
152. Schröder, G.; Krause, S.; Zechner, E.L.; Traxler, B.; Yeo, H.-J.; Lurz, R.; Waksman, G.; Lanka, E. TraG-like proteins of DNA transfer systems and of the *Helicobacter pylori* type IV secretion system: inner membrane gate for exported substrates? *J. Bacteriol.* **2002**, *184*, 2767–2779, doi:10.1128/jb.184.10.2767-2779.2002.
153. Llosa, M.; Zunzunegui, S.; de la Cruz, F. Conjugative coupling proteins interact with cognate and heterologous VirB10-like proteins while exhibiting specificity for cognate relaxosomes. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 10465–10470, doi:10.1073/pnas.1830264100.
154. Haft, R.J.F.; Palacios, G.; Nguyen, T.; Mally, M.; Gachelet, E.G.; Zechner, E.L.; Traxler, B. General mutagenesis of F plasmid TraI reveals its role in conjugative regulation. *J. Bacteriol.* **2006**, *188*, 6346–6353, doi:10.1128/JB.00462-06.
155. Gilmour, M.W.; Gunton, J.E.; Lawley, T.D.; Taylor, D.E. Interaction between the IncHI1 plasmid R27 coupling protein and type IV secretion system: TraG associates with the coiled-coil mating pair formation protein TrhB. *Mol. Microbiol.* **2003**, *49*, 105–116, doi:10.1046/j.1365-2958.2003.03551.x.
156. Egelman, E.H. Structural biology. Pumping DNA. *Nature* **2001**, *409*, 573, 575, doi:10.1038/35054652.
157. Gomis-Rüth, F.X.; Moncalián, G.; Pérez-Luque, R.; González, A.; Cabezón, E.; de la Cruz, F.; Coll, M. The bacterial conjugation protein TrwB resembles ring helicases and F1-ATPase. *Nature* **2001**, *409*, 637–641, doi:10.1038/35054586.
158. Moncalián, G.; Cabezón, E.; Alkorta, I.; Valle, M.; Moro, F.; Valpuesta, J.M.; Goñi, F.M.; de La Cruz, F. Characterization of ATP and DNA binding activities of TrwB, the coupling protein essential in plasmid R388 conjugation. *J. Biol. Chem.* **1999**, *274*, 36117–36124, doi:10.1074/jbc.274.51.36117.
159. Schröder, G.; Lanka, E. TraG-like proteins of type IV secretion systems: functional dissection of the multiple activities of TraG (RP4) and TrwB (R388). *J. Bacteriol.* **2003**, *185*, 4371–4381, doi:10.1128/jb.185.15.4371-4381.2003.

160. Haft, R.J.F.; Gachelet, E.G.; Nguyen, T.; Toussaint, L.; Chivian, D.; Traxler, B. In vivo oligomerization of the F conjugative coupling protein TraD. *J. Bacteriol.* **2007**, *189*, 6626–6634, doi:10.1128/JB.00513-07.
161. Tato, I.; Zunzunegui, S.; de la Cruz, F.; Cabezon, E. TrwB, the coupling protein involved in DNA transport during bacterial conjugation, is a DNA-dependent ATPase. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 8156–8161, doi:10.1073/pnas.0503402102.
162. Christie, P.J.; Atmakuri, K.; Krishnamoorthy, V.; Jakubowski, S.; Cascales, E. Biogenesis, architecture, and function of bacterial type IV secretion systems. *Annu. Rev. Microbiol.* **2005**, *59*, 451–485, doi:10.1146/annurev.micro.58.030603.123630.
163. Garcillán-Barcia, M.P.; Jurado, P.; González-Pérez, B.; Moncalián, G.; Fernández, L.A.; de la Cruz, F. Conjugative transfer can be inhibited by blocking relaxase activity within recipient cells with intrabodies. *Mol. Microbiol.* **2007**, *63*, 404–416, doi:10.1111/j.1365-2958.2006.05523.x.
164. Chandler, M.; de la Cruz, F.; Dyda, F.; Hickman, A.B.; Moncalian, G.; Ton-Hoang, B. Breaking and joining single-stranded DNA: the HUH endonuclease superfamily. *Nat. Rev. Microbiol.* **2013**, *11*, 525–538, doi:10.1038/nrmicro3067.
165. Becker, E.C.; Meyer, R. Origin and fate of the 3' ends of single-stranded DNA generated by conjugal transfer of plasmid R1162. *J. Bacteriol.* **2012**, *194*, 5368–5376, doi:10.1128/JB.00818-12.
166. Forsberg, K.J.; Malik, H.S. Microbial Genomics: The Expanding Universe of Bacterial Defense Systems. *Curr. Biol.* **2018**, *28*, R361–R364, doi:10.1016/j.cub.2018.02.053.
167. Casjens, S. Prophages and bacterial genomics: what have we learned so far? *Mol. Microbiol.* **2003**, *49*, 277–300, doi:10.1046/j.1365-2958.2003.03580.x.
168. Narra, H.P.; Ochman, H. Of what use is sex to bacteria? *Curr. Biol.* **2006**, *16*, R705-710, doi:10.1016/j.cub.2006.08.024.
169. Johnston, C.D.; Cotton, S.L.; Rittling, S.R.; Starr, J.R.; Borisy, G.G.; Dewhirst, F.E.; Lemon, K.P. Systematic evasion of the restriction-modification barrier in bacteria. *Proc. Natl. Acad. Sci. U.S.A.* **2019**, *116*, 11454–11459, doi:10.1073/pnas.1820256116.
170. Bickle, T.A. Restricting restriction. *Mol. Microbiol.* **2004**, *51*, 3–5, doi:10.1046/j.1365-2958.2003.03846.x.
171. Chen, K.; Reuter, M.; Sanghvi, B.; Roberts, G.A.; Cooper, L.P.; Tilling, M.; Blakely, G.W.; Dryden, D.T.F. ArdA proteins from different mobile genetic elements can bind to the EcoKI Type I DNA methyltransferase of *E. coli* K12. *Biochim. Biophys. Acta* **2014**, *1844*, 505–511, doi:10.1016/j.bbapap.2013.12.008.
172. McMahon, S.A.; Roberts, G.A.; Johnson, K.A.; Cooper, L.P.; Liu, H.; White, J.H.; Carter, L.G.; Sanghvi, B.; Oke, M.; Walkinshaw, M.D.; et al. Extensive DNA mimicry by the ArdA anti-restriction protein and its role in the spread of antibiotic resistance. *Nucleic Acids Res.* **2009**, *37*, 4887–4897, doi:10.1093/nar/gkp478.
173. Wilkins, B.M. Plasmid promiscuity: meeting the challenge of DNA immigration control. *Environ. Microbiol.* **2002**, *4*, 495–500, doi:10.1046/j.1462-2920.2002.00332.x.
174. Belogurov, A.A.; Delver, E.P.; Agafonova, O.V.; Belogurova, N.G.; Lee, L.Y.; Kado, C.I. Antirestriction protein Ard (Type C) encoded by IncW plasmid pSa has a high similarity to the “protein transport” domain of TraC1 primase of promiscuous plasmid RP4. *J. Mol. Biol.* **2000**, *296*, 969–977, doi:10.1006/jmbi.1999.3493.

175. Roy, D.; Huguet, K.T.; Grenier, F.; Burrus, V. IncC conjugative plasmids and SXT/R391 elements repair double-strand breaks caused by CRISPR-Cas during conjugation. *Nucleic Acids Res.* **2020**, doi:10.1093/nar/gkaa518.
176. Wilkins, B.M.; Chilly, P.M.; Thomas, A.T.; Pocklington, M.J. Distribution of restriction enzyme recognition sequences on broad host range plasmid RP4: molecular and evolutionary implications. *J. Mol. Biol.* **1996**, *258*, 447–456, doi:10.1006/jmbi.1996.0261.
177. Grissa, I.; Vergnaud, G.; Pourcel, C. The CRISPRdb database and tools to display CRISPRs and to generate dictionaries of spacers and repeats. *BMC Bioinformatics* **2007**, *8*, 172, doi:10.1186/1471-2105-8-172.
178. Makarova, K.S.; Wolf, Y.I.; Alkhnbashi, O.S.; Costa, F.; Shah, S.A.; Saunders, S.J.; Barrangou, R.; Brouns, S.J.J.; Charpentier, E.; Haft, D.H.; et al. An updated evolutionary classification of CRISPR-Cas systems. *Nat. Rev. Microbiol.* **2015**, *13*, 722–736, doi:10.1038/nrmicro3569.
179. Garneau, J.E.; Dupuis, M.-È.; Villion, M.; Romero, D.A.; Barrangou, R.; Boyaval, P.; Fremaux, C.; Horvath, P.; Magadán, A.H.; Moineau, S. The CRISPR/Cas bacterial immune system cleaves bacteriophage and plasmid DNA. *Nature* **2010**, *468*, 67–71, doi:10.1038/nature09523.
180. Bondy-Denomy, J.; Pawluk, A.; Maxwell, K.L.; Davidson, A.R. Bacteriophage genes that inactivate the CRISPR/Cas bacterial immune system. *Nature* **2013**, *493*, 429–432, doi:10.1038/nature11723.
181. Bondy-Denomy, J.; Garcia, B.; Strum, S.; Du, M.; Rollins, M.F.; Hidalgo-Reyes, Y.; Wiedenheft, B.; Maxwell, K.L.; Davidson, A.R. Multiple mechanisms for CRISPR-Cas inhibition by anti-CRISPR proteins. *Nature* **2015**, *526*, 136–139, doi:10.1038/nature15254.
182. Mahendra, C.; Christie, K.A.; Osuna, B.A.; Pinilla-Redondo, R.; Kleinstiver, B.P.; Bondy-Denomy, J. Broad-spectrum anti-CRISPR proteins facilitate horizontal gene transfer. *Nat Microbiol* **2020**, *5*, 620–629, doi:10.1038/s41564-020-0692-2.
183. Cram, D.; Ray, A.; O’Gorman, L.; Skurray, R. Transcriptional analysis of the leading region in F plasmid DNA transfer. *Plasmid* **1984**, *11*, 221–233, doi:10.1016/0147-619x(84)90028-3.
184. Jones, A.L.; Barth, P.T.; Wilkins, B.M. Zygotic induction of plasmid *ssb* and *psiB* genes following conjugative transfer of IncI1 plasmid Collb-P9. *Mol. Microbiol.* **1992**, *6*, 605–613.
185. Althorpe, N.J.; Chilly, P.M.; Thomas, A.T.; Brammar, W.J.; Wilkins, B.M. Transient transcriptional activation of the IncI1 plasmid anti-restriction gene (*ardA*) and SOS inhibition gene (*psiB*) early in conjugating recipient bacteria. *Mol. Microbiol.* **1999**, *31*, 133–142, doi:10.1046/j.1365-2958.1999.01153.x.
186. Masai, H.; Arai, K. Frpo: A Novel Single-Stranded DNA Promoter for Transcription and for Primer RNA Synthesis of DNA Replication. *Cell* **1997**, *89*, 897–907, doi:10.1016/S0092-8674(00)80275-5.
187. Honda, Y.; Sakai, H.; Komano, T. Two single-strand DNA initiation signals located in the *oriV* region of plasmid RSF1010. *Gene* **1988**, *68*, 221–228, doi:10.1016/0378-1119(88)90024-8.
188. Masai, H.; Arai, K. Mechanisms of primer RNA synthesis and D-loop/R-loop-dependent DNA replication in *Escherichia coli*. *Biochimie* **1996**, *78*, 1109–1117, doi:10.1016/s0300-9084(97)86737-5.
189. Nomura, N.; Low, R.L.; Ray, D.S. Selective cloning of CoIE1 DNA initiation sequences using the cloning vector M13 delta E101. *Gene* **1982**, *18*, 239–246, doi:10.1016/0378-1119(82)90161-5.

190. Nomura, N.; Masai, H.; Inuzuka, M.; Miyazaki, C.; Ohtsubo, E.; Itoh, T.; Sasamoto, S.; Matsui, M.; Ishizaki, R.; Arai, K. Identification of eleven single-strand initiation sequences (ssi) for priming of DNA replication in the F, R6K, R100 and ColE2 plasmids. *Gene* **1991**, *108*, 15–22, doi:10.1016/0378-1119(91)90482-q.
191. Díaz, A.; Lacks, S.A.; López, P. Multiple roles for DNA polymerase I in establishment and replication of the promiscuous plasmid pLS1. *Mol. Microbiol.* **1994**, *14*, 773–783, doi:10.1111/j.1365-2958.1994.tb01314.x.
192. Wilkins, B.M.; Hollom, S.E. Conjugational synthesis of F lac⁺ and Col I DNA in the presence of rifampicin and in Escherichia coli K12 mutants defective in DNA synthesis. *Mol. Gen. Genet.* **1974**, *134*, 143–156, doi:10.1007/BF00268416.
193. Baharoglu, Z.; Mazel, D. SOS, the formidable strategy of bacteria against aggressions. *FEMS Microbiol. Rev.* **2014**, *38*, 1126–1145, doi:10.1111/1574-6976.12077.
194. Baharoglu, Z.; Bikard, D.; Mazel, D. Conjugative DNA Transfer Induces the Bacterial SOS Response and Promotes Antibiotic Resistance Development through Integron Activation. *PLoS Genet* **2010**, *6*, doi:10.1371/journal.pgen.1001165.
195. Maslowska, K.H.; Makiela-Dzvenska, K.; Fijalkowska, I.J. The SOS system: A complex and tightly regulated response to DNA damage. *Environ. Mol. Mutagen.* **2019**, *60*, 368–384, doi:10.1002/em.22267.
196. Golub, E.; Bailone, A.; Devoret, R. A gene encoding an SOS inhibitor is present in different conjugative plasmids. *J Bacteriol* **1988**, *170*, 4392–4394.
197. Bailone, A.; Bäckman, A.; Sommer, S.; Célérier, J.; Bagdasarian, M.M.; Bagdasarian, M.; Devoret, R. PsiB polypeptide prevents activation of RecA protein in Escherichia coli. *Mol. Gen. Genet.* **1988**, *214*, 389–395, doi:10.1007/bf00330471.
198. Petrova, V.; Chitteni-Pattu, S.; Drees, J.C.; Inman, R.B.; Cox, M.M. An SOS inhibitor that binds to free RecA protein: the PsiB protein. *Mol. Cell* **2009**, *36*, 121–130, doi:10.1016/j.molcel.2009.07.026.
199. Meyer, R.R.; Laine, P.S. The single-stranded DNA-binding protein of Escherichia coli. *Microbiol. Rev.* **1990**, *54*, 342–380.
200. Shereda, R.D.; Kozlov, A.G.; Lohman, T.M.; Cox, M.M.; Keck, J.L. SSB as an organizer/mobilizer of genome maintenance complexes. *Crit. Rev. Biochem. Mol. Biol.* **2008**, *43*, 289–318, doi:10.1080/10409230802341296.
201. Sigal, N.; Delius, H.; Kornberg, T.; Gefter, M.L.; Alberts, B. A DNA-unwinding protein isolated from Escherichia coli: its interaction with DNA and with DNA polymerases. *Proc. Natl. Acad. Sci. U.S.A.* **1972**, *69*, 3537–3541, doi:10.1073/pnas.69.12.3537.
202. Nolivos, S.; Cayron, J.; Dedieu, A.; Page, A.; Delolme, F.; Lesterlin, C. Role of AcrAB-TolC multidrug efflux pump in drug-resistance acquisition by plasmid transfer. *Science* **2019**, *364*, 778–782, doi:10.1126/science.aav6390.
203. Golub, E.I.; Low, K.B. Conjugative plasmids of enteric bacteria from many different incompatibility groups have similar genes for single-stranded DNA-binding proteins. *J Bacteriol* **1985**, *162*, 235–241.
204. Kolodkin, A.L.; Capage, M.A.; Golub, E.I.; Low, K.B. F sex factor of Escherichia coli K-12 codes for a single-stranded DNA binding protein. *Proc. Natl. Acad. Sci. U.S.A.* **1983**, *80*, 4422–4426, doi:10.1073/pnas.80.14.4422.

205. Golub, E.I.; Low, K.B. Conjugative plasmids of enteric bacteria from many different incompatibility groups have similar genes for single-stranded DNA-binding proteins. *J. Bacteriol.* **1985**, *162*, 235–241.
206. Golub, E.I.; Low, K.B. Derepression of single-stranded DNA-binding protein genes on plasmids derepressed for conjugation, and complementation of an *E. coli* *ssb*-mutation by these genes. *Mol. Gen. Genet.* **1986**, *204*, 410–416, doi:10.1007/bf00331017.
207. Howland, C.J.; Rees, C.E.; Barth, P.T.; Wilkins, B.M. The *ssb* gene of plasmid ColIb-P9. *J. Bacteriol.* **1989**, *171*, 2466–2473, doi:10.1128/jb.171.5.2466-2473.1989.
208. Porter, R.D.; Black, S. The single-stranded-DNA-binding protein encoded by the *Escherichia coli* F factor can complement a deletion of the chromosomal *ssb* gene. *J. Bacteriol.* **1991**, *173*, 2720–2723, doi:10.1128/jb.173.8.2720-2723.1991.
209. Ruvolo, P.P.; Keating, K.M.; Williams, K.R.; Chase, J.W. Single-stranded DNA binding proteins (SSBs) from prokaryotic transmissible plasmids. *Proteins* **1991**, *9*, 120–134, doi:10.1002/prot.340090206.
210. Jovanovic, O.S.; Ayres, E.K.; Figurski, D.H. The replication initiator operon of promiscuous plasmid RK2 encodes a gene that complements an *Escherichia coli* mutant defective in single-stranded DNA-binding protein. *J. Bacteriol.* **1992**, *174*, 4842–4846, doi:10.1128/jb.174.14.4842-4846.1992.
211. del Solar, G.; Giraldo, R.; Ruiz-Echevarría, M.J.; Espinosa, M.; Díaz-Orejas, R. Replication and control of circular bacterial plasmids. *Microbiol. Mol. Biol. Rev.* **1998**, *62*, 434–464.
212. Pinto, U.M.; Pappas, K.M.; Winans, S.C. The ABCs of plasmid replication and segregation. *Nat. Rev. Microbiol.* **2012**, *10*, 755–765, doi:10.1038/nrmicro2882.
213. Rakowski, S.A.; Filutowicz, M. Plasmid R6K replication control. *Plasmid* **2013**, *69*, 231–242, doi:10.1016/j.plasmid.2013.02.003.
214. Nordström, K. Plasmid R1--replication and its control. *Plasmid* **2006**, *55*, 1–26, doi:10.1016/j.plasmid.2005.07.002.
215. Guiney, D.G. Host range of conjugation and replication functions of the *Escherichia coli* sex plasmid Flac. *Journal of Molecular Biology* **1982**, *162*, 699–703, doi:10.1016/0022-2836(82)90397-7.
216. Baharoglu, Z.; Bikard, D.; Mazel, D. Conjugative DNA Transfer Induces the Bacterial SOS Response and Promotes Antibiotic Resistance Development through Integron Activation. *PLoS Genetics* **2010**, *6*, e1001165, doi:10.1371/journal.pgen.1001165.
217. Zhong, Z.; Helinski, D.; Toukdarian, A. Plasmid host-range: restrictions to F replication in *Pseudomonas*. *Plasmid* **2005**, *54*, 48–56, doi:10.1016/j.plasmid.2004.11.001.
218. Adamczyk, M.; Jagura-Burdzy, G. Spread and survival of promiscuous IncP-1 plasmids. *Acta Biochim. Pol.* **2003**, *50*, 425–453, doi:035002425.
219. Baxter, J.C.; Funnell, B.E. Plasmid Partition Mechanisms. *Microbiology Spectrum* **2014**, *2*, doi:10.1128/microbiolspec.PLAS-0023-2014.
220. Ebersbach, G.; Gerdes, K. Plasmid segregation mechanisms. *Annu. Rev. Genet.* **2005**, *39*, 453–479.
221. Gerdes, K.; Howard, M.; Szardenings, F. Pushing and pulling in prokaryotic DNA segregation. *Cell* **2010**, *141*, 927–942, doi:10.1016/j.cell.2010.05.033.

222. Møller-Jensen, J.; Jensen, R.B.; Gerdes, K. Plasmid and chromosome segregation in prokaryotes. *Trends Microbiol.* **2000**, *8*, 313–320, doi:10.1016/s0966-842x(00)01787-x.
223. Schumacher, M.A. Bacterial plasmid partition machinery: a minimalist approach to survival. *Curr. Opin. Struct. Biol.* **2012**, *22*, 72–79, doi:10.1016/j.sbi.2011.11.001.
224. Deonier, R.C.; Davidson, N. The sequence organization of the integrated F plasmid in two Hfr strains of Escherichia coli. *J. Mol. Biol.* **1976**, *107*, 207–222, doi:10.1016/s0022-2836(76)80002-2.
225. Deonier, R.C.; Mirels, L. Excision of F plasmid sequences by recombination at directly repeated insertion sequence 2 elements: involvement of recA. *Proc Natl Acad Sci U S A* **1977**, *74*, 3965–3969.
226. Johnson, C.M.; Grossman, A.D. Integrative and Conjugative Elements (ICEs): What They Do and How They Work. *Annual Review of Genetics* **2015**, *49*, 577–601, doi:10.1146/annurev-genet-112414-055018.
227. Shao, Q.; Hawkins, A.; Zeng, L. Phage DNA Dynamics in Cells with Different Fates. *Biophys J* **2015**, *108*, 2048–2060, doi:10.1016/j.bpj.2015.03.027.
228. Gago-Córdoba, C.; Val-Calvo, J.; Miguel-Arribas, A.; Serrano, E.; Singh, P.K.; Abia, D.; Wu, L.J.; Meijer, W.J.J. Surface Exclusion Revisited: Function Related to Differential Expression of the Surface Exclusion System of Bacillus subtilis Plasmid pLS20. *Front Microbiol* **2019**, *10*, 1502, doi:10.3389/fmicb.2019.01502.
229. Galli, D.M.; Chen, J. Entry exclusion activity on conjugative plasmid pVT745. *Plasmid* **2006**, *55*, 158–163, doi:10.1016/j.plasmid.2005.07.003.
230. Gunton, J.E.; Ussher, J.E.R.; Rooker, M.M.; Wetsch, N.M.; Alonso, G.; Taylor, D.E. Entry exclusion in the IncHI1 plasmid R27 is mediated by EexA and EexB. *Plasmid* **2008**, *59*, 86–101, doi:10.1016/j.plasmid.2007.11.004.
231. Haase, J.; Kalkum, M.; Lanka, E. TrbK, a small cytoplasmic membrane lipoprotein, functions in entry exclusion of the IncP alpha plasmid RP4. *Journal of Bacteriology* **1996**, *178*, 6720–6729, doi:10.1128/jb.178.23.6720-6729.1996.
232. Humbert, M.; Huguët, K.T.; Coulombe, F.; Burrus, V. Entry Exclusion of Conjugative Plasmids of the IncA, IncC, and Related Untyped Incompatibility Groups. *J. Bacteriol.* **2019**, *201*, doi:10.1128/JB.00731-18.
233. Kraushaar, B.; Appel, B.; Lanka, E.; Strauch, E. Entry exclusion and oriT of a conjugative system encoded by the cryptic plasmid p29930 of Yersinia enterocolitica. *Plasmid* **2010**, *64*, 79–84, doi:10.1016/j.plasmid.2010.05.001.
234. Pohlman, R.F.; Genetti, H.D.; Winans, S.C. Entry exclusion of the IncN plasmid pKM101 is mediated by a single hydrophilic protein containing a lipid attachment motif. *Plasmid* **1994**, *31*, 158–165, doi:10.1006/plas.1994.1017.
235. Possoz, C.; Gagnat, J.; Sezonov, G.; Guérineau, M.; Pernodet, J.-L. Conjugal immunity of Streptomyces strains carrying the integrative element pSAM2 is due to the pif gene (pSAM2 immunity factor). *Mol. Microbiol.* **2003**, *47*, 1385–1393, doi:10.1046/j.1365-2958.2003.03380.x.
236. Thomas, C.M.; Nielsen, K.M. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nat. Rev. Microbiol.* **2005**, *3*, 711–721, doi:10.1038/nrmicro1234.
237. Achtman, M.; Kennedy, N.; Skurray, R. Cell-cell interactions in conjugating Escherichia coli: role of traT protein in surface exclusion. *Proc Natl Acad Sci U S A* **1977**, *74*, 5104–5108.

238. Manning, P.A.; Beutin, L.; Achtman, M. Outer membrane of *Escherichia coli*: properties of the F sex factor traT protein which is involved in surface exclusion. *J. Bacteriol.* **1980**, *142*, 285–294, doi:10.1128/JB.142.1.285-294.1980.
239. Minkley, E.G.; Ippen-Ihler, K. Identification of a membrane protein associated with expression of the surface exclusion region of the F transfer operon. *J. Bacteriol.* **1977**, *129*, 1613–1622, doi:10.1128/JB.129.3.1613-1622.1977.
240. Riede, I.; Eschbach, M.L. Evidence that TraT interacts with OmpA of *Escherichia coli*. *FEBS Lett.* **1986**, *205*, 241–245, doi:10.1016/0014-5793(86)80905-x.
241. Jalajakumari, M.B.; Guidolin, A.; Buhk, H.J.; Manning, P.A.; Ham, L.M.; Hodgson, A.L.; Cheah, K.C.; Skurray, R.A. Surface exclusion genes traS and traT of the F sex factor of *Escherichia coli* K-12. Determination of the nucleotide sequence and promoter and terminator activities. *J. Mol. Biol.* **1987**, *198*, 1–11, doi:10.1016/0022-2836(87)90452-9.
242. San Millan, A.; MacLean, R.C. Fitness Costs of Plasmids: a Limit to Plasmid Transmission. *Microbiol Spectr* **2017**, *5*, doi:10.1128/microbiolspec.MTBP-0016-2017.
243. Barlow, M. What antimicrobial resistance has taught us about horizontal gene transfer. *Methods Mol. Biol.* **2009**, *532*, 397–411, doi:10.1007/978-1-60327-853-9_23.
244. Hall-Stoodley, L.; Costerton, J.W.; Stoodley, P. Bacterial biofilms: from the natural environment to infectious diseases. *Nat. Rev. Microbiol.* **2004**, *2*, 95–108, doi:10.1038/nrmicro821.
245. Bale, M.J.; Fry, J.C.; Day, M.J. Plasmid transfer between strains of *Pseudomonas aeruginosa* on membrane filters attached to river stones. *J. Gen. Microbiol.* **1987**, *133*, 3099–3107, doi:10.1099/00221287-133-11-3099.
246. Lilley, A.K.; Bailey, M.J. Impact of Plasmid pQBR103 Acquisition and Carriage on the Phytosphere Fitness of *Pseudomonas fluorescens* SBW25: Burden and Benefit. *Appl. Environ. Microbiol.* **1997**, *63*, 1584–1587.
247. Munck, C.; Sheth, R.U.; Freedberg, D.E.; Wang, H.H. Recording mobile DNA in the gut microbiota using an *Escherichia coli* CRISPR-Cas spacer acquisition platform. *Nat Commun* **2020**, *11*, 95, doi:10.1038/s41467-019-14012-5.
248. Ronda, C.; Chen, S.P.; Cabral, V.; Yaung, S.J.; Wang, H.H. Metagenomic engineering of the mammalian gut microbiome in situ. *Nat. Methods* **2019**, *16*, 167–170, doi:10.1038/s41592-018-0301-y.
249. Ehlers, L.J.; Bouwer, E.J. RP4 plasmid transfer among species of *pseudomonas* in a biofilm reactor. *Water Science and Technology* **1999**, *39*, 163–171, doi:10.2166/wst.1999.0353.
250. Yang, D.; Wang, J.; Qiu, Z.; Jin, M.; Shen, Z.; Chen, Z.; Wang, X.; Zhang, B.; Li, J.-W. Horizontal transfer of antibiotic resistance genes in a membrane bioreactor. *J. Biotechnol.* **2013**, *167*, 441–447, doi:10.1016/j.jbiotec.2013.08.004.
251. Hausner, M.; Wuertz, S. High rates of conjugation in bacterial biofilms as determined by quantitative in situ analysis. *Appl. Environ. Microbiol.* **1999**, *65*, 3710–3713.
252. Christensen, B.B.; Sternberg, C.; Molin, S. Bacterial plasmid conjugation on semi-solid surfaces monitored with the green fluorescent protein (GFP) from *Aequorea victoria* as a marker. *Gene* **1996**, *173*, 59–65, doi:10.1016/0378-1119(95)00707-5.

253. Haagenen, J.A.J.; Hansen, S.K.; Johansen, T.; Molin, S. In situ detection of horizontal transfer of mobile genetic elements. *FEMS Microbiol. Ecol.* **2002**, *42*, 261–268, doi:10.1111/j.1574-6941.2002.tb01016.x.
254. Reisner, A.; Wolinski, H.; Zechner, E.L. In situ monitoring of IncF plasmid transfer on semi-solid agar surfaces reveals a limited invasion of plasmids in recipient colonies. *Plasmid* **2012**, *67*, 155–161, doi:10.1016/j.plasmid.2012.01.001.
255. Lilley, A.K.; Bailey, M.J. The transfer dynamics of *Pseudomonas* sp. plasmid pQBR11 in biofilms. *FEMS Microbiol. Ecol.* **2002**, *42*, 243–250, doi:10.1111/j.1574-6941.2002.tb01014.x.
256. Aspray, T.J.; Hansen, S.K.; Burns, R.G. A soil-based microbial biofilm exposed to 2,4-D: bacterial community development and establishment of conjugative plasmid pJP4. *FEMS Microbiol. Ecol.* **2005**, *54*, 317–327, doi:10.1016/j.femsec.2005.04.007.
257. Christensen, B.B.; Sternberg, C.; Andersen, J.B.; Eberl, L.; Moller, S.; Givskov, M.; Molin, S. Establishment of new genetic traits in a microbial biofilm community. *Appl. Environ. Microbiol.* **1998**, *64*, 2247–2255.
258. Nancharaiah, Y.V.; Wattiau, P.; Wuertz, S.; Bathe, S.; Mohan, S.V.; Wilderer, P.A.; Hausner, M. Dual labeling of *Pseudomonas putida* with fluorescent proteins for in situ monitoring of conjugal transfer of the TOL plasmid. *Appl. Environ. Microbiol.* **2003**, *69*, 4846–4852, doi:10.1128/aem.69.8.4846-4852.2003.
259. Abe, K.; Nomura, N.; Suzuki, S. Biofilms: hot spots of horizontal gene transfer (HGT) in aquatic environments, with a focus on a new HGT mechanism. *FEMS Microbiol. Ecol.* **2020**, *96*, doi:10.1093/femsec/fiaa031.
260. Madsen, J.S.; Burmølle, M.; Hansen, L.H.; Sørensen, S.J. The interconnection between biofilm formation and horizontal gene transfer. *FEMS Immunol. Med. Microbiol.* **2012**, *65*, 183–195, doi:10.1111/j.1574-695X.2012.00960.x.
261. Molin, S.; Tolker-Nielsen, T. Gene transfer occurs with enhanced efficiency in biofilms and induces enhanced stabilisation of the biofilm structure. *Curr. Opin. Biotechnol.* **2003**, *14*, 255–261, doi:10.1016/s0958-1669(03)00036-3.
262. Nesse, L.L.; Simm, R. Biofilm: A Hotspot for Emerging Bacterial Genotypes. *Adv. Appl. Microbiol.* **2018**, *103*, 223–246, doi:10.1016/bs.aams.2018.01.003.
263. Stalder, T.; Top, E. Plasmid transfer in biofilms: a perspective on limitations and opportunities. *NPJ Biofilms Microbiomes* **2016**, *2*, doi:10.1038/npjbiofilms.2016.22.
264. Fuchs, F.M.; Holland, G.; Moeller, R.; Laue, M. Directed freeze-fracturing of *Bacillus subtilis* biofilms for conventional scanning electron microscopy. *J. Microbiol. Methods* **2018**, *152*, 165–172, doi:10.1016/j.mimet.2018.08.005.
265. Serra, D.O.; Richter, A.M.; Klauck, G.; Mika, F.; Hengge, R. Microanatomy at cellular resolution and spatial order of physiological differentiation in a bacterial biofilm. *mBio* **2013**, *4*, e00103-00113, doi:10.1128/mBio.00103-13.
266. Seoane, J.; Yankelevich, T.; Dechesne, A.; Merkey, B.; Sternberg, C.; Smets, B.F. An individual-based approach to explain plasmid invasion in bacterial populations. *FEMS Microbiol. Ecol.* **2011**, *75*, 17–27, doi:10.1111/j.1574-6941.2010.00994.x.
267. Licht, T.R.; Christensen, B.B.; Krogfelt, K.A.; Molin, S. Plasmid transfer in the animal intestine and other dynamic bacterial populations: the role of community structure and

- environment. *Microbiology (Reading, Engl.)* **1999**, *145* (Pt 9), 2615–2622, doi:10.1099/00221287-145-9-2615.
268. Serra, D.O.; Hengge, R. Stress responses go three dimensional - the spatial order of physiological differentiation in bacterial macrocolony biofilms. *Environ. Microbiol.* **2014**, *16*, 1455–1471, doi:10.1111/1462-2920.12483.
269. Stewart, P.S.; Franklin, M.J. Physiological heterogeneity in biofilms. *Nat. Rev. Microbiol.* **2008**, *6*, 199–210, doi:10.1038/nrmicro1838.
270. Ghigo, J.M. Natural conjugative plasmids induce bacterial biofilm development. *Nature* **2001**, *412*, 442–445, doi:10.1038/35086581.
271. Lim, J.Y.; La, H.J.; Sheng, H.; Forney, L.J.; Hovde, C.J. Influence of plasmid pO157 on *Escherichia coli* O157:H7 Sakai biofilm formation. *Appl. Environ. Microbiol.* **2010**, *76*, 963–966, doi:10.1128/AEM.01068-09.
272. Liu, Z.; Que, F.; Liao, L.; Zhou, M.; You, L.; Zhao, Q.; Li, Y.; Niu, H.; Wu, S.; Huang, R. Study on the promotion of bacterial biofilm formation by a *Salmonella* conjugative plasmid and the underlying mechanism. *PLoS ONE* **2014**, *9*, e109808, doi:10.1371/journal.pone.0109808.
273. Reisner, A.; Höller, B.M.; Molin, S.; Zechner, E.L. Synergistic effects in mixed *Escherichia coli* biofilms: conjugative plasmid transfer drives biofilm expansion. *J. Bacteriol.* **2006**, *188*, 3582–3588, doi:10.1128/JB.188.10.3582-3588.2006.
274. Shi, H.; Zhou, X.; Zou, W.; Wang, Y.; Lei, C.; Xiang, R.; Zhou, L.; Liu, B.; Zhang, A.; Wang, H. Co-occurrence of biofilm formation and quinolone resistance in *Salmonella enterica* serotype typhimurium carrying an IncHI2-type oqxAB-positive plasmid. *Microb. Pathog.* **2018**, *123*, 68–73, doi:10.1016/j.micpath.2018.06.006.
275. Beloin, C.; Roux, A.; Ghigo, J.M. *Escherichia coli* biofilms. *Curr. Top. Microbiol. Immunol.* **2008**, *322*, 249–289, doi:10.1007/978-3-540-75418-3_12.
276. Craig, L.; Forest, K.T.; Maier, B. Type IV pili: dynamics, biophysics and functional consequences. *Nat. Rev. Microbiol.* **2019**, *17*, 429–440, doi:10.1038/s41579-019-0195-4.
277. Burmølle, M.; Bahl, M.I.; Jensen, L.B.; Sørensen, S.J.; Hansen, L.H. Type 3 fimbriae, encoded by the conjugative plasmid pOLA52, enhance biofilm formation and transfer frequencies in Enterobacteriaceae strains. *Microbiology (Reading, Engl.)* **2008**, *154*, 187–195, doi:10.1099/mic.0.2007/010454-0.
278. Dudley, E.G.; Abe, C.; Ghigo, J.-M.; Latour-Lambert, P.; Hormazabal, J.C.; Nataro, J.P. An IncII plasmid contributes to the adherence of the atypical enteroaggregative *Escherichia coli* strain C1096 to cultured cells and abiotic surfaces. *Infect. Immun.* **2006**, *74*, 2102–2114, doi:10.1128/IAI.74.4.2102-2114.2006.
279. Bhatti, M.; Cruz, M.R.; Frank, K.L.; Gomez, J.A.L.; Andrade, F.; Garsin, D.A.; Dunny, G.M.; Kaplan, H.B.; Christie, P.J. Enterococcus faecalis pCF10-encoded surface proteins PrgA, PrgB (aggregation substance) and PrgC contribute to plasmid transfer, biofilm formation and virulence. *Mol. Microbiol.* **2015**, *95*, 660–677, doi:10.1111/mmi.12893.
280. Coburn, P.S.; Baghdayan, A.S.; Craig, N.; Burroughs, A.; Tendolkar, P.; Miller, K.; Najjar, F.Z.; Roe, B.A.; Shankar, N. A novel conjugative plasmid from *Enterococcus faecalis* E99 enhances resistance to ultraviolet radiation. *Plasmid* **2010**, *64*, 18–25, doi:10.1016/j.plasmid.2010.03.001.

281. Tendolkar, P.M.; Baghdayan, A.S.; Shankar, N. Putative surface proteins encoded within a novel transferable locus confer a high-biofilm phenotype to *Enterococcus faecalis*. *J. Bacteriol.* **2006**, *188*, 2063–2072, doi:10.1128/JB.188.6.2063-2072.2006.
282. Reisner, A.; Haagensen, J.A.J.; Schembri, M.A.; Zechner, E.L.; Molin, S. Development and maturation of *Escherichia coli* K-12 biofilms. *Mol. Microbiol.* **2003**, *48*, 933–946, doi:10.1046/j.1365-2958.2003.03490.x.
283. May, T.; Okabe, S. *Escherichia coli* harboring a natural IncF conjugative F plasmid develops complex mature biofilms by stimulating synthesis of colanic acid and Curli. *J. Bacteriol.* **2008**, *190*, 7479–7490, doi:10.1128/JB.00823-08.
284. Barrios, A.F.G.; Zuo, R.; Ren, D.; Wood, T.K. Hha, YbaJ, and OmpA regulate *Escherichia coli* K12 biofilm formation and conjugation plasmids abolish motility. *Biotechnol. Bioeng.* **2006**, *93*, 188–200, doi:10.1002/bit.20681.
285. Yang, X.; Ma, Q.; Wood, T.K. The R1 conjugative plasmid increases *Escherichia coli* biofilm formation through an envelope stress response. *Appl. Environ. Microbiol.* **2008**, *74*, 2690–2699, doi:10.1128/AEM.02809-07.
286. Okshevsy, M.; Meyer, R.L. The role of extracellular DNA in the establishment, maintenance and perpetuation of bacterial biofilms. *Crit. Rev. Microbiol.* **2015**, *41*, 341–352, doi:10.3109/1040841X.2013.841639.
287. D’Alvise, P.W.; Sjøholm, O.R.; Yankelevich, T.; Jin, Y.; Wuertz, S.; Smets, B.F. TOL plasmid carriage enhances biofilm formation and increases extracellular DNA content in *Pseudomonas putida* KT2440. *FEMS Microbiol. Lett.* **2010**, *312*, 84–92, doi:10.1111/j.1574-6968.2010.02105.x.
288. Røder, H.L.; Hansen, L.H.; Sørensen, S.J.; Burmølle, M. The impact of the conjugative IncP-1 plasmid pKJK5 on multispecies biofilm formation is dependent on the plasmid host. *FEMS Microbiol. Lett.* **2013**, *344*, 186–192, doi:10.1111/1574-6968.12175.
289. Hoffman, L.R.; D’Argenio, D.A.; MacCoss, M.J.; Zhang, Z.; Jones, R.A.; Miller, S.I. Aminoglycoside antibiotics induce bacterial biofilm formation. *Nature* **2005**, *436*, 1171–1175, doi:10.1038/nature03912.
290. Linares, J.F.; Gustafsson, I.; Baquero, F.; Martinez, J.L. Antibiotics as intermicrobial signaling agents instead of weapons. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 19484–19489, doi:10.1073/pnas.0608949103.
291. Salcedo, D.E.; Lee, J.H.; Ha, U.H.; Kim, S.P. The effects of antibiotics on the biofilm formation and antibiotic resistance gene transfer. *Desalination and Water Treatment* **2015**, *54*, 3582–3588, doi:10.1080/19443994.2014.923206.
292. Penesyan, A.; Nagy, S.S.; Kjelleberg, S.; Gillings, M.R.; Paulsen, I.T. Rapid microevolution of biofilm cells in response to antibiotics. *NPJ Biofilms Microbiomes* **2019**, *5*, 34, doi:10.1038/s41522-019-0108-3.
293. Bagge, N.; Schuster, M.; Hentzer, M.; Ciofu, O.; Givskov, M.; Greenberg, E.P.; Høiby, N. *Pseudomonas aeruginosa* biofilms exposed to imipenem exhibit changes in global gene expression and beta-lactamase and alginate production. *Antimicrob. Agents Chemother.* **2004**, *48*, 1175–1187, doi:10.1128/aac.48.4.1175-1187.2004.
294. Díaz-Pascual, F.; Hartmann, R.; Lempp, M.; Vidakovic, L.; Song, B.; Jeckel, H.; Thormann, K.M.; Yildiz, F.H.; Dunkel, J.; Link, H.; et al. Breakdown of *Vibrio cholerae* biofilm

- architecture induced by antibiotics disrupts community barrier function. *Nat Microbiol* **2019**, 4, 2136–2145, doi:10.1038/s41564-019-0579-2.
295. al-Masaudi, S.B.; Day, M.J.; Russell, A.D. Effect of some antibiotics and biocides on plasmid transfer in *Staphylococcus aureus*. *J. Appl. Bacteriol.* **1991**, 71, 239–243, doi:10.1111/j.1365-2672.1991.tb04454.x.
296. Barr, V.; Barr, K.; Millar, M.R.; Lacey, R.W. Beta-lactam antibiotics increase the frequency of plasmid transfer in *Staphylococcus aureus*. *J. Antimicrob. Chemother.* **1986**, 17, 409–413, doi:10.1093/jac/17.4.409.
297. Feld, L.; Schjørring, S.; Hammer, K.; Licht, T.R.; Danielsen, M.; Krogfelt, K.; Wilcks, A. Selective pressure affects transfer and establishment of a *Lactobacillus plantarum* resistance plasmid in the gastrointestinal environment. *J. Antimicrob. Chemother.* **2008**, 61, 845–852, doi:10.1093/jac/dkn033.
298. Liu, G.; Bogaj, K.; Bortolaia, V.; Olsen, J.E.; Thomsen, L.E. Antibiotic-Induced, Increased Conjugative Transfer Is Common to Diverse Naturally Occurring ESBL Plasmids in *Escherichia coli*. *Front Microbiol* **2019**, 10, 2119, doi:10.3389/fmicb.2019.02119.
299. Lu, Y.; Zeng, J.; Wang, L.; Lan, K.; E, S.; Wang, L.; Xiao, Q.; Luo, Q.; Huang, X.; Huang, B.; et al. Antibiotics Promote *Escherichia coli*-*Pseudomonas aeruginosa* Conjugation through Inhibiting Quorum Sensing. *Antimicrob. Agents Chemother.* **2017**, 61, doi:10.1128/AAC.01284-17.
300. Ma, H.; Bryers, J.D. Non-invasive determination of conjugative transfer of plasmids bearing antibiotic-resistance genes in biofilm-bound bacteria: effects of substrate loading and antibiotic selection. *Appl. Microbiol. Biotechnol.* **2013**, 97, 317–328, doi:10.1007/s00253-012-4179-9.
301. Ohlsen, K.; Ternes, T.; Werner, G.; Wallner, U.; Löffler, D.; Ziebuhr, W.; Witte, W.; Hacker, J. Impact of antibiotics on conjugational resistance gene transfer in *Staphylococcus aureus* in sewage. *Environ. Microbiol.* **2003**, 5, 711–716, doi:10.1046/j.1462-2920.2003.00459.x.
302. Xia, Z.-J.; Wang, J.; Hu, W.; Liu, H.; Gao, X.-Z.; Wu, Z.-H.; Zhang, P.-Y.; Li, Y.-Z. Improving conjugation efficacy of *Sorangium cellulosum* by the addition of dual selection antibiotics. *J. Ind. Microbiol. Biotechnol.* **2008**, 35, 1157–1163, doi:10.1007/s10295-008-0395-9.
303. Zhang, P.-Y.; Xu, P.-P.; Xia, Z.-J.; Wang, J.; Xiong, J.; Li, Y.-Z. Combined treatment with the antibiotics kanamycin and streptomycin promotes the conjugation of *Escherichia coli*. *FEMS Microbiol. Lett.* **2013**, 348, 149–156, doi:10.1111/1574-6968.12282.
304. Møller, T.S.B.; Liu, G.; Boysen, A.; Thomsen, L.E.; Lüthje, F.L.; Mortensen, S.; Møller-Jensen, J.; Olsen, J.E. Treatment with Cefotaxime Affects Expression of Conjugation Associated Proteins and Conjugation Transfer Frequency of an IncI1 Plasmid in *Escherichia coli*. *Front Microbiol* **2017**, 8, 2365, doi:10.3389/fmicb.2017.02365.
305. Shun-Mei, E.; Zeng, J.-M.; Yuan, H.; Lu, Y.; Cai, R.-X.; Chen, C. Sub-inhibitory concentrations of fluoroquinolones increase conjugation frequency. *Microb. Pathog.* **2018**, 114, 57–62, doi:10.1016/j.micpath.2017.11.036.
306. Cantas, L.; Midtlyng, P.J.; Sørum, H. Impact of antibiotic treatments on the expression of the R plasmid *tra* genes and on the host innate immune activity during pRAS1 bearing *Aeromonas hydrophila* infection in zebrafish (*Danio rerio*). *BMC Microbiol.* **2012**, 12, 37, doi:10.1186/1471-2180-12-37.

307. Lu, Y.; Zeng, J.; Wang, L.; Lan, K.; E, S.; Wang, L.; Xiao, Q.; Luo, Q.; Huang, X.; Huang, B.; et al. Antibiotics Promote Escherichia coli-Pseudomonas aeruginosa Conjugation through Inhibiting Quorum Sensing. *Antimicrob. Agents Chemother.* **2017**, *61*, doi:10.1128/AAC.01284-17.
308. Kim, S.; Yun, Z.; Ha, U.-H.; Lee, S.; Park, H.; Kwon, E.E.; Cho, Y.; Choung, S.; Oh, J.; Medriano, C.A.; et al. Transfer of antibiotic resistance plasmids in pure and activated sludge cultures in the presence of environmentally representative micro-contaminant concentrations. *Sci. Total Environ.* **2014**, *468–469*, 813–820, doi:10.1016/j.scitotenv.2013.08.100.
309. Lopatkin, A.J.; Huang, S.; Smith, R.P.; Srimani, J.K.; Sysoeva, T.A.; Bewick, S.; Karig, D.K.; You, L. Antibiotics as a selective driver for conjugation dynamics. *Nat Microbiol* **2016**, *1*, 16044, doi:10.1038/nmicrobiol.2016.44.
310. Li, B.; Qiu, Y.; Zhang, J.; Huang, X.; Shi, H.; Yin, H. Real-Time Study of Rapid Spread of Antibiotic Resistance Plasmid in Biofilm Using Microfluidics. *Environ. Sci. Technol.* **2018**, *52*, 11132–11141, doi:10.1021/acs.est.8b03281.
311. Qiu, Y.; Zhang, J.; Li, B.; Wen, X.; Liang, P.; Huang, X. A novel microfluidic system enables visualization and analysis of antibiotic resistance gene transfer to activated sludge bacteria in biofilm. *Sci. Total Environ.* **2018**, *642*, 582–590, doi:10.1016/j.scitotenv.2018.06.012.