

Article

# A cross between bread wheat and a 2D(2R) disomic substitution triticales line leads to the formation of a novel disomic addition line and provides information of the role of rye secalins on breadmaking characteristics

Francesco Sestili <sup>1</sup>, Benedetta Margiotta <sup>2</sup>, Patrizia Vaccino <sup>3</sup>, Salvatore Moscaritolo <sup>4</sup>, Debora Giorgi <sup>5</sup>, Sergio Lucretti <sup>5</sup>, Samuela Palombieri <sup>1</sup>, Stefania Masci <sup>1</sup> and Domenico Lafiandra <sup>1,\*</sup>

<sup>1</sup> Department of Agriculture and Forest Sciences, University of Tuscia, Viterbo, Italy;

[francescosestili@unitus.it](mailto:francescosestili@unitus.it); [palombieri@unitus.it](mailto:palombieri@unitus.it); [masci@unitus.it](mailto:masci@unitus.it); [lafiandr@unitus.it](mailto:lafiandr@unitus.it)

<sup>2</sup> Institute of Biosciences and Bioresources of the National Research Council, Bari, Italy;

[benedetta.margiotta@ibbr.cnr.it](mailto:benedetta.margiotta@ibbr.cnr.it)

<sup>3</sup> CREA Research Centre for Cereal and Industrial Crops, Vercelli, Italy; [patrizia.vaccino@crea.gov.it](mailto:patrizia.vaccino@crea.gov.it)

<sup>4</sup> CREA Research Centre for Engineering and Agro-Food Processing, Rome, Italy;

[salvatore.moscaritolo@crea.gov.it](mailto:salvatore.moscaritolo@crea.gov.it)

<sup>5</sup> ENEA, CASACCIA Research Center, Laboratory Biotechnologies, Rome, Italy; [debora.giorgi@enea.it](mailto:debora.giorgi@enea.it);

[sergio.lucretti@enea.it](mailto:sergio.lucretti@enea.it)

\* Correspondence: [lafiandr@unitus.it](mailto:lafiandr@unitus.it); Tel.: +39-0761-357-243

**Abstract:** A bread wheat line (N11) and a disomic 2D(2R) substitution triticales line have been crossed and backcrossed four times. At each step electrophoretic selection for the seeds that possessed, simultaneously, the complete set of high molecular weight glutenin subunits of N11 and the two high molecular weight secalins of rye, present in the 2D(2R) line, was carried out. Molecular cytogenetic analyses of the BC<sub>4</sub>F<sub>5</sub> generation have revealed that the selection carried out had produced a disomic addition line (2n=44). The pair of additional chromosomes consisted of the long arm of chromosome 1R (1RL) from rye fused with the satellite body of the wheat chromosome 6B. Rheological analyses revealed that the dough obtained by the new addition line had higher quality characteristics when compared with the two parents. The role of the two additional high molecular weight secalins, present in the disomic addition line, in influencing improved dough characteristics is discussed.

**Keywords:** rye, secalins, glutenins, chromosome rearrangements, dough quality

## 1. Introduction

Wheat represents one of the most important crops for global food security, providing 20% of the total calories and 21% of the daily dietary protein consumed by the human population [1]. The global production has reached about 735 million tonnes in 2018, making the wheat the third most important crop after maize and rice (<http://www.fao.org/faostat/en/#data/QC/visualize>).

The most cultivated wheat species is *Triticum aestivum*, a hexaploid species with genome AABBDD (2n=6X=42) usually known as bread wheat. The success of wheat depends on its adaptability to different climatic conditions (from cold countries as Scandinavia and Russia to Argentina, including elevated regions in the tropics and subtropics) [2] and the unique visco-elastic properties of wheat doughs that permit to produce a vast range of end products, unmatched by the other two major cereals maize and rice, such as different kinds of bread, pasta, noodles, pizza, cakes, snacks and other bakery foods [3].

Current global genetic gains in wheat have been reported to be around 1% per annum which is insufficient to meet the future demand in the face of a constantly increasing world population [4]. Modern wheat varieties are affected by a reduced genetic diversity, as consequence of different processes such as polyploidization, domestication and modern plant breeding, the need to introduce new genetic diversity into breeding programs, to satisfy the demand of varieties with good yield and superior processing and nutritional characteristics, represents a primary objective, also in light of the new challenges posed by the global climate change [5].

Wheat, along with rye and barley, is part of the *Triticeae* tribe which includes approximately 325 annual and perennial species [6], forming a large germplasm reservoir of genes that can be used for cereal crop improvement [7, 8].

Wide hybridization represent an important approach to introduce desirable alien genes from distant relatives in wheat and the first step in this endeavor is represented by the production of wheat–alien chromosome addition or substitution lines. These lines are used as a bridge to generate useful translocations lines and many and numerous such lines have been produced using barley, rye, *Dasypyrum villosum*, *Aegilops* and several species of the genera *Agropyron* and *Thinopyrum* [9-12].

In order to produce useful translocations several methodologies are available and have been described in many papers, such as: use of *Ph* gene mutants to induce homoeologous recombination, use of radiation or gametocidal gene, transgenic approach [11, 13-15].

Rye represents an important source of useful alien genes for wheat improvement, and at this regard the intergeneric introgressions of the Robertsonian 1BL/1RS wheat-rye translocation represents one of the most successful achievement, as also witnessed by the high number of wheat variety released worldwide [16]. In this material the short arm of wheat chromosome 1B is replaced by its counterpart from the rye genome which arose spontaneously from centromeric breakage reunion and fusion of two chromosome arms broken at the centromere conferring to wheat resistance to several diseases, high yield performance, wide adaptation, resistance to drought, increasing amount of micronutrient, protein content and arabinoxylan content [17-19].

Results of several studies carried out on wheat-rye substitution, addition lines, triticale and their progeny from crosses between triticale and wheat have demonstrated that deletions and translocations of chromosomal regions and chromosome arms are among the most common spontaneous chromosomal changes generated. In addition to these rearrangements formation of minichromosomes and chromosomes with multiple centromeres have also been observed, as well as elimination of most or all of the D genome chromosomes [20].

In some cases the composition of the gluten proteins, and particularly of the high molecular weight glutenin subunits (HMW-GS), was modified as consequences of gene deletions, silencing or coding gene sequence rearrangements [21-23], resulting in the generation of new genetic diversity in this group of proteins.

Gluten proteins include the monomeric gliadins and the polymeric glutenins, important determinants of dough viscoelastic properties and breadmaking characteristics of different wheat varieties [24].

Glutenin polymers are the major responsible of dough elasticity and strength and are formed by HMW-GS and low molecular weight glutenin subunits (LMW-GS), held together by intermolecular disulfide bonds. The existence of highly positive correlation between large sized polymers as measured by the relative amount of unextractable proteins (%UPP) and dough processing properties has been investigated by many authors and it has been demonstrated that the HMW-GS play a major role compared to LMW-GS [25-28].

As far as HMW-GS concern intensive genetic researches have revealed that, in hexaploid wheat, HMW-GS are encoded by orthologous gene sets, present at the *Glu-A1*, *Glu-B1* and *Glu-D1* loci located on the long arms of the homeologous group 1 chromosomes [29]. Each locus contains two tightly linked paralogous genes encoding two types of HMW glutenin subunits, designated as x- and y- type subunits, with higher and lower molecular weight, respectively [24]. The complete sequences of numerous HMW-GS genes, have revealed that these subunits have a highly conserved structures, comprising short repetitive N- and C-terminal domains, where most of the cysteine residues are present, flanking a repetitive domain of variable size [30].

Generally, bread wheat varieties differ in number of HMW-GS, ranging from three to five, and type of subunits, with silencing of some *Glu-1* genes and allelic variation present at each locus, accounting for these differences [24]. Usually, the number of cysteine residues in x and y type subunits are four (three in the N-terminal domain and one in the C-terminal domain) and seven (five in the N-terminal domain, one in the repetitive domain and one in the C-terminal domain), respectively [24]. Exceptions are represented by the subunit 1Dx5 and 1Bx20, with the first possessing an extra cysteine residue in the repetitive domain, close to the N terminal region and the latter having only two cysteine residues, one in the N-terminal and one in the C-terminal region [31].

These cysteine residues form disulfide bonds within and between subunits, which are important in determining structure and properties of the gluten complex in wheat dough [24].

Allelic variation has been identified at each *Glu-1* locus with both number and type of subunits impacting breadmaking quality. In particular dough mixing and baking characteristics improved when the number of subunits increased from three to five [29,32], thanks to their contribution to the formation of large-sized glutenin polymers, as deduced by the proportion of %UPP [26,33].

Gupta and MacRitchie [34] reported that the genotypes with the pair of subunits 5+10, associated at the *Glu-D1* locus, had a significant higher size distribution of polymeric glutenins compared to those with the pair of allelic subunit 2+12, suggesting that the differences in number of cysteine residues between 1Dx5 (5 cysteines) and 1Dx2 (4 cysteines) was responsible for the superior characteristics associated to varieties possessing the pair of subunits 5+10.

Pirozi et al. [35], compared quality characteristics of two biotypes of the bread wheat variety Avocet differing only in the HMW-GS composition. Avocet A, possessed HMW-GS composition: null (*Glu-A1*), 7+8 (*Glu-B1*), 5+10 (*Glu-D1*), whereas HMW-GS composition of Avocet C was: 1, 20x+20y, 5+10; Avocet A exhibited an increase in polymeric protein, a decrease in the gliadin-to-glutenin ratio, and a marked decrease in %UPP. Avocet C doughs exhibited greater extensibility and shorter Mixograph, dough development times, and baked into smaller loaves compared to those from Avocet A, in spite of the fact that, compared to the former, the latter possesses an additional HMW-GS (subunit 1 at the *Glu-A1* locus).

SDS-PAGE separations of glutenin subunits present in a large collection of Chinese bread wheat variety carried out by Wang et al. [36] led to the identification of a novel pattern of HMW-GS in the bread wheat variety Xiaoyanmai 7 (XY7). This variety showed four HMW-GS, two of which resemble the pair of HMW-GS 7+8, encoded by genes present at the *Glu-B1* locus, whereas the other two subunits could not be classified. Cytogenetical, biochemical and molecular analysis demonstrated that XY7 was a wheat-rye secondary substitution line in which six pairs of rye chromosomes had replaced their homoeologs of the D genome, except the pair of the 2D chromosomes. In addition, it was also demonstrated that the two additional subunits, previously unidentified, corresponded to HMW secalins, normally associated with genes present on the long arm of the 1R chromosome at the *Sec-3* locus [37].


The work described in this paper reports novel findings of chromosomal rearrangements, resulting from a cross followed by backcrossing between wheat line N11 and a bread XY7.

## 2. Results

### 2.1 Electrophoretic Analysis of Gliadins and HMW-GS

The electrophoretic separation of gliadins and glutenin subunits present in the two parents XY7, N11 and in the derived N11<sup>XY7</sup> line are reported in Figure 1a and 1b. In a previous work we have shown that XY7 can be considered a substituted triticale possessing the set of chromosomes of the A and B genomes, six pair of rye chromosomes and the 2D(2R) substitution [37]. The analysis of the gliadin of XY7 shows the absence of the  $\omega$  components, with slow electrophoretic mobility, encoded by genes present at the *Gli-D1* locus on the short arm of the 1D chromosome, and visible in N11. In addition, the  $\omega$ -secalins encoded by genes present on the short arm of the 1R chromosomes, at the *Sec-1* locus are present in XY7. The line N11<sup>XY7</sup> shows an identical gliadin pattern to the line N11 as result of the repeated backcrosses and selection. Analysis of the electrophoretic separation of glutenin subunits reveal major differences in the LMW-GS between XY7


and N11. The absence of the 1D chromosome, in addition to the gliadin, is also responsible for the loss of the LMW-GS associated at the *Glu-D3* locus. On the contrary the LMW-GS of N11<sup>XY7</sup> result identical to N11. SDS-PAGE separation of the glutenin subunits shows that N11 has HMW-GS 1, 7+8 and 2+12 encoded by genes present at the *Glu-A1*, *Glu-B1* and *Glu-D1*, respectively, present on the long arm of the homoeologous group 1 chromosomes. The line XY7 has only the pair of subunits 7+8 and two HMW secalin (Ry and Rx) subunits, previously identified [37], encoded by genes present at the *Sec-3* locus. In the XY7 y-type subunit is larger than x-type subunit, as deduced by DNA sequencing of their corresponding genes, contrary to what normally observed in bread wheat. The line N11<sup>XY7</sup> shows seven glutenin subunits, the five HMW-GS of N11 and the two HMW-secalins from XY7.



**Figure 1.** Comparison of gliadin (a) and glutenin (b) composition of N11, N11<sup>XY7</sup> and XY7: (a) A-PAGE separation of gliadins, (b) SDS-PAGE separation of glutenins. Lane 1, N11; lane 2, N11<sup>XY7</sup>; lane 3, XY7.

## 2.2 Cytogenetic Analysis by GISH and FISH


The cytogenetic analysis of the N11<sup>XY7</sup> line revealed a chromosome numbers of  $2n=44$  (Figure S1) instead of the standard bread wheat complement of  $2n=42$  chromosomes. In order to better discriminate among wheat and rye chromosomes and identify the additional chromosomes pair, GISH and single and double target FISH experiments were performed on metaphase spreads of N11<sup>XY7</sup> and parental lines. Genomic hybridization analysis of the N11<sup>XY7</sup> line (Figure 2a) with total rye DNA, showed a clear hybridization signal on the whole long arm of a chromosomes pair, the candidate additional one. Moreover, a peculiar DAPI band was clearly visible at the telomeres of the same chromosomes pair, which facilitate their identification as a secale-derived translocation in the wheat background (Figure 2a). In XY7 six labeled chromosomes pairs were detected, as expected (Figure 2b).



**Figure 2.** Genomic *in situ* hybridization (GISH) with total ryeCy3 labeled DNA as probe (red) on N11<sup>XY7</sup> (a) and XY7 (b) metaphase spreads. DNA is counterstained with DAPI. Rye chromosomes or chromosome's arms are red. In the box a telomeric DAPI band present in the rye chromosome arm is highlighted by a white arrow.

FISH experiments were performed sequentially after GISH using the (GAA)<sub>7</sub> oligonucleotide, and the rDNA probe pTa71; both probes have been widely exploited to characterize and discriminate wheat and rye chromosomes [38,39]. Our results showed the parental lines exhibiting a (GAA)<sub>7</sub> distribution in agreement with previous reports [40], while the N11<sup>XY7</sup> line disclosed the lack of the satellite body on chromosome 6B and the following lack of the (GAA)<sub>7</sub> signal localized on the satellite (Figure 3).

To further investigate on satellite chromosome morphology and structure the pTa71 ribosomal probe was used on the two parents XY7, N11 and the derived N11<sup>XY7</sup> line. In bread wheat, hybridization signals with the pTa71 probe are usually localized at the secondary constriction of satellites chromosomes pairs 1B and 6B and on chromosomes pair 5D, while in rye pTa71 labeling is present at the secondary constriction of the chromosome 1R [41]. In our analysis the pTa71 probe showed a standard distribution on the parental lines N11 and XY7 (Figure 3); on the N11<sup>XY7</sup> line the signal was visible on chromosome 5D and at the secondary constriction of chromosome 1B, as expected, but on chromosome 6B, which was lacking of both the secondary constriction and the satellite body, the pTa71 was localized at the telomeric region of the short arm. Besides two centromeric labeling spots were visible on the extra "chromosomes pair" (Figure 3). By comparing GISH and FISH labelling pattern of the parental lines with those shown in N11<sup>XY7</sup>, we hypothesized the translocation of the satellite body from wheat chromosome 6B to the additional rye chromosomes pair present in this new N11<sup>XY7</sup> line. Our hypothesis explains the presence of the pTa71 hybridization signal at the centromeric region of the additional chromosomes pair and the uncommon localization of pTa71 probe at the telomeric region of the short arm of chromosome 6B which probably, in spite the lack of the large part of the satellite body, still retains some copies of the rDNA genes. In addition, two (GAA)<sub>7</sub> hybridization spots expected on 6B satellite body, were present on the "short arm" of the extra chromosomes pair (data not shown).



**Figure 3.** Double target FISH analysis with rDNA pTa71 probe (red) and  $(GAA)_7$  oligonucleotides (green) on N11 (a), XY7 (b) and N11<sup>XY7</sup> (c) metaphase spreads. Chromosome rDNA localization is indicated by arrows.


### 2.3 Qualitative Characterization

The values and means of the qualitative traits for the genotypes under study and growing season are reported in Table 1 and Table S1. As regards the test weight (TW), the parental lines had contrasting values, very high for line N11 (82.6 kg/hL), and very low for XY7 (69.2 kg/hL). N11<sup>XY7</sup>, showed very good values, in both years, with an average of 80.1 kg/hL, comparable to N11. No significant variation was found among the three lines for the hardness, all them showing hard texture (hardness values ranging from 89 to 100). A significant variation was observed between the growing seasons, with Y1 resulting in an increase of the hardness index by 28% in comparison to Y2. The Protein content (PC) of both parental lines was significantly lower than the one observed in N11<sup>XY7</sup> (14.2% d.m. on average), with no significant differences between the growing seasons. As regards the Gluten Index, N11<sup>XY7</sup> had a very high value (99.5), significantly higher than N11 (86.5) and XY7 (64). No significant variation between growing seasons was observed. The rheological characteristics, as determined by the alveographic test, showed large differences among the three lines (Figure 4). In this test, line N11<sup>XY7</sup> showed on average very high extensibility (L: 175 mm), significantly higher than that of N11 (80 mm) and XY7 (55 mm). Dough strength (W) of N11<sup>XY7</sup>, was twice that of the best parent, N11, and four times higher than that of XY7. The differences were confirmed in both growing seasons, being the differences not significant between Y1 and Y2 for all the alveographic parameters.

As for the Brabender farinographic test, the analyses showed that, in spite of its high protein content, N11<sup>XY7</sup> water absorbance was similar to that of N11 and slightly higher than that of XY7 (60.9% vs 60.8%, and 54.1%, respectively). In comparison to the parental lines, N11 and XY7, N11<sup>XY7</sup> had significantly higher development time (9.7 min. vs 2.8 min. and 1.4 min, respectively) and dough stability (11.8 min. vs 5.3 min. and 3.6 min, respectively). The degree of softening was similar for N11 and N11<sup>XY7</sup> (62 and 51 BU, respectively), while XY7 had a significant higher value (119 BU). The FQN, as expected, was highly correlated with the dough development time, and was significantly higher in N11<sup>XY7</sup> (171 mm) than in N11 (62 mm) and XY7 (50 mm). For all the farinograph parameters, no significant differences were observed between growing seasons.

**Table 1.** Mean of qualitative parameters determined in two different seasons (Y1: 2016-2017 and Y2: 2017-2018). TW: test weight; PC: protein content; GI: gluten index; P: dough tenacity; L: dough extensibility; W: dough strength; BU: Brabender Units; FQN: Farinograph quality number.

	Chopin alveograph							Brabender farinograph					
	TW (kg/hL)	Hardness	PC % d.m.	GI	P (mm)	L (mm)	P/L	W (J × 10 <sup>-4</sup> )	Water Abs (%)	Dough Development time (min)	Dough Stability (min)	Degree of softening (BU)	FQN (mm)
<b>N11</b>	82.6 <sup>a</sup>	89	10.6 <sup>b</sup>	86.5 <sup>ab</sup>	100 <sup>a</sup>	80 <sup>b</sup>	1.33	231 <sup>b</sup>	60.9 <sup>a</sup>	2.8 <sup>b</sup>	5.3 <sup>b</sup>	62 <sup>b</sup>	75.5 <sup>b</sup>
<b>XY7</b>	69.2 <sup>b</sup>	100	11.0 <sup>b</sup>	64 <sup>b</sup>	60 <sup>b</sup>	55 <sup>b</sup>	1.09	102 <sup>c</sup>	54.1 <sup>b</sup>	1.4 <sup>b</sup>	3.6 <sup>b</sup>	119 <sup>a</sup>	49.5 <sup>b</sup>
<b>N11<sup>XY7</sup></b>	80.1 <sup>a</sup>	96	14.2 <sup>a</sup>	99.5 <sup>a</sup>	83 <sup>ab</sup>	175 <sup>a</sup>	0.48	466 <sup>a</sup>	60.8 <sup>a</sup>	9.7 <sup>a</sup>	11.8 <sup>a</sup>	51 <sup>b</sup>	171.0 <sup>a</sup>
<b>P</b>	0.04	0.79	0.01	0.016	0.022	0.007	0.13	0.001	0.0029	0.009	0.011	0.015	0.0042
<b>sem</b>	0.95	11.69	0.37	3.89	4.77	10.15	0.22	7.02	0.46	0.77	0.81	7.69	8.63
<b>Y1</b>	76.5	106	12.1	86.7	83	95	1.09	265	58.4	4.4	6.3	83	92.0
<b>Y2</b>	78.0	83	11.7	80.0	79	111	0.84	268	58.7	4.8	7.4	71	103.3
<b>P</b>	0.81	0.01	0.80	0.68	0.83	0.78	0.57	0.98	0.93	0.91	0.77	0.70	0.84
<b>sem</b>	4.13	3.40	1.14	10.47	11.99	36.89	0.28	106.79	2.27	2.62	2.54	21.42	37.54



**Figure 4.** Alveograph profiles of XY7 (a), N11<sup>XY7</sup> (b) and (c) N11 relative to the 2017-2018 growing season.

### 3. Discussion

Genomic *in situ* hybridization (GISH) is a widely used approach for detecting alien chromatin in wheat genetic background [42]. The alien DNA can be further analyzed by Fluorescence *in situ* hybridization (FISH) using repeated sequences as probes [43]. Among the different classes of repeated sequences, ribosomal DNA (rDNA) and Simple Sequence Repeats (SSR) are the most valuable cytological markers in discrimination of plant chromosomes, and they have been widely used in rye and wheat cytogenetic analysis. In this work a new bread wheat line (N11<sup>XY7</sup>) derived from the cross between the bread wheat line N11 and XY7, with the latter harbouring six pair of chromosomes of rye [37], have been characterized by means of GISH analysis on N11<sup>XY7</sup> metaphase using total rye DNA as probe, thus enabling the identification of an additional chromosome pair to the bread wheat standard complement.

Further biochemical analysis and the cytogenetic chromosomes characterization by FISH labeling using rDNA and the SSR (GAA)<sub>7</sub> revealed that the additional pairs was, in fact, composed of the whole long arm of chromosome 1 of rye (1RL) fused with the satellite body derived from the chromosome 6B of wheat (6B SAT).

The presence of chromosomal rearrangements in the crosses between wheat x wild relatives has been particularly observed and described in several studies conducted on wheat-rye substitution and addition lines, triticale and their progeny from crosses triticale x wheat [44-46].

McClintock [47] pointed out that major restructuring of chromosome components may arise in wide species crosses and continue to arise in its progeny, as consequences of 'genome shock'.

These processes are associated with translocations, inversions, deficiencies, duplications with new stable or relatively stable "species" or "genera" derived from such initial hybrids. Triticale is described as an example by McClintock [47] in which the combined set of chromosomes of wheat and rye are not entirely stable and further selection in subsequent generations does not completely eliminate this instability which resulted from the loss of parental bands in triticale or, conversely, novel bands appearing in the allopolyploid that were not seen in the parental genomes.

Tang et al. [46] reported that wheat chromosomal alterations could be easily induced by rye chromosomes in wheat-rye hybrids, suggesting that the stability of the wheat genome could be effectively suppressed by the R genome.

Lukaszewky and Gustafson [48] carried out a large number of cytological analysis of four triticale X wheat populations and found that among 785 karyotyped plants 195 wheat/rye and 64 rye/rye translocated chromosomes were present, in addition modification of rye chromosomes by deletion or amplification of telomeric heterochromatin was identified. Out of 39 identified wheat/rye translocations, 10 occurred between homoeologous and 29 between non-homoeologous

chromosomes, five involved A-genome chromosomes, six B-genome chromosomes and the remaining 28 involved D-genome chromosomes. It has been suggested that very likely most of the translocations described resulted from mis-division of univalents at meiosis and subsequent random fusion of telocentric chromosomes [48].

Darlington [49] suggested that univalents in meiosis have a tendency to mis-divide at the centromeres, producing telocentric chromosomes. The 1RS/1BL translocations, a well-known example of the use of alien chromosomes in wheat breeding, are centric translocations formed by mis-division followed by the fusion of the mis-division products, in which the short arm of wheat chromosome 1B is replaced by the short arm of 1R rye chromosome [50]. Fu et al. [44,45] observed that even single alien rye chromosome added to wheat genome could induce alterations of wheat and rye chromosomes. They investigated the alterations and abnormal mitotic behaviours of wheat chromosomes, induced by wheat-rye 4R, 6R and 7R monosomic addition lines [44] and by wheat-rye 1R monosomic addition lines [45]. In the addition lines containing the whole 1R chromosome or 1R chromosome arms and in their self-progeny lagging chromosomes, micronuclei and chromosomal bridges were observed.

Beside centromere (primary constriction) the secondary constrictions may represent a possible region of breakage. It is known, in fact, that chromosome constrictions are points of weakness in the chromosome [51]. In addition, secondary constriction often coincides with the nucleolar organized region (NOR) containing rDNA gene. The 45S rDNA in plant is considered a fragile site [52], that is a large and highly sensitive region which is inclined to form abnormal poor-staining lesions with gaps, constrictions or breaks in metaphase chromosomes.

In our lines the mitotic instability induced by the presence of rye chromosome in wheat background join to the chromosome secondary constriction "weakness" might have led to the centromeric breakage of chromosome 1R followed by the fusion of its long arm with the satellite body of the wheat chromosome 6B. A similar translocation line was observed by Badaeva et al. [53] in 2 related strains of French tetraploid triticale. The translocation line indicated as T6A:1B-1, was formed by the break occurred in the short arm of chromosome 6A near the centromere and the second break within the region of secondary constriction of chromosome 1B, which resulted in the emergence of a new satellite chromosome formed by the long arm of 6A and the satellite of 1B [53].

Already in 1962, Fergusson-Smith [54] observed that in human metaphase chromosomes with secondary constrictions are close to one another more often than would be expected from mere chance distribution. This phenomenon has been termed satellite association. Rye chromosome 1R and wheat chromosome 6B are both satellite chromosome holding secondary constriction, their possible proximity in the nucleus could explain why the fusion occurred among these two non-homoeologous chromosomes. On the other hand, non-homoeologous wheat/rye translocation have been previously reported [48].

Changes in gluten protein composition, as consequences of gene deletions, silencing or coding gene sequence rearrangements have also been reported by different authors. Two hexaploid lines of triticale originated from the cross between the bread wheat cultivar M8003 and Austrian rye (*Secale cereale* L.) were used by Li et al. [55] to evaluate morphologic traits, chromosome composition and storage protein composition. The two triticale lines differed for plant height, spike length and number of spikelets. Cytological analyses showed the complete elimination of the D-genome chromosomes and the presence of major chromosomal rearrangements, involving chromosomes of the A and B genomes in both triticales. In addition, large variation in HMW-GS, LMW-GS and secalins between the two triticales was detected. At last, novel patterns that distinctly differentiated the two triticale lines from the parents also appeared, indicating that coding sequence rearrangements, activation of novel gene expression, gene silencing or deletions could be responsible for the changes observed.

Yuan et al. [21] analyzed by SDS-PAGE the HMW-GS composition of F<sub>1</sub> and F<sub>2</sub> progeny of a cross between a Japanese bread wheat landraces and rye. Results of their analyses revealed absence and/or additional novel HMW-GSs in the endosperm of F<sub>1</sub> and F<sub>2</sub> hybrid seeds. Analysis of cloned sequences of HMW-GS revealed that deletions involving the in-frame stop codon, present in the *Glu-Ax1* null allele of the bread wheat parent, had occurred. This produced a novel active *Glu-Ax1* allele in some F<sub>1</sub> and F<sub>2</sub> plants corresponding to one of the novel subunits detected by SDS-PAGE, as

demonstrated by heterologous expression in *E. coli*. The results of this work lead Yuan et al. [21] to suggest that mitotic illegitimate recombination between two copies of a short repeat sequence had resulted in the observed deletions with consequent change in HMW-GS compositions.

In a different work Yuan et al. [22], in order to investigate the possibility of using a wheat-rye hybridization method for inducing *de novo* phenotypes, crossed the bread wheat cultivar Shinchunaga, possessing the very large HMW-GS 1Dx2.2 associated with the *Glu-D1* locus, with the Chinese rye landrace Qinling. Electrophoretic analysis of HMW-GS of F<sub>2</sub> seeds, together with PCR amplification of the HMW-GS genes, their cloning and sequencing permitted to identify a subunit, designated Glu-1Dx2.2<sup>v</sup> derived from the 1Dx2.2 gene through deletion of a direct repeat of 295 bp length and an intervening sequence of 101 bp present in its central repetitive region.

Also in this case Yuan et al. [22] postulated that illegitimate recombination was, very likely, the mechanism responsible for the appearance of the new subunit Glu-1Dx2.2<sup>v</sup>.

Differently from the changes above reported, our work proves that the pressure exerted during the backcrossing process resulted in the production of a disomic addition line in which the pair of the novel chromosomes are the result of major rearrangements.

The presence of the additional chromosomes pair in N11<sup>XY7</sup> and associated HMW secalins resulted in an unexpected rheological behavior, but also in an exceptionally high technological quality of the dough, which allows N11<sup>XY7</sup> to be classified as improver wheat in Italy, excellent or class E wheat in France and Germany, hard red wheat in the USA. Moreover, in spite of its very high dough strength, N11<sup>XY7</sup> keeps a very good extensibility, which is one of the main objectives of bread wheat quality improvement. Usually, in fact, strong gluten often suffers for low extensibility, which is a serious problem, since adequate extensibility is required for proper dough handling and baking performance [56].

Extensibility is highly influenced by gliadins and protein content [57-59], but the gliadin pattern of N11<sup>XY7</sup> is not different from the pattern of the parent N11, thus excluding gliadins to be responsible for this change. In addition, in our opinion, the extensibility of N11<sup>XY7</sup> seems too high to be justified only by the result of the higher protein content found in the line.

The farinographic behavior of N11<sup>XY7</sup> was quite peculiar, as well. In spite of its higher protein content, in comparison to both parents, N11<sup>XY7</sup> did not show a proportional increase in water absorbance. On the contrary, great differences were observed in development time, with almost 10 minutes, on average, necessary to reach optimum dough consistency.

Unlike HMW-GS the role of HMW secalins in influencing breadmaking quality has been addressed in a very limited way. Kipp et al. [60] found that rheological properties of wheat gluten were negatively affected, when purified HMW secalins were incorporated in wheat dough, with a strong reduction in maximum resistance and a slight increase in extensibility. The authors suggested that the observed effects were associated with the more frequent occurrence of cysteine residues in the HMW secalins affecting the formation of large polymeric glutenins.

Amiour et al. [61] analyzed substitution lines 1R(1A), 1R(1B) and 1R(1D), in which the entire 1R chromosome replaces the homoeologous chromosome of the group 1 of wheat, in order to compare the contribution of the HMW secalin subunits, encoded by genes present at the *Sec-3* locus, to that of HMW-GS on the technological characteristics of the doughs.

Rheological studies demonstrated that the contribution of the three substitutions on breadmaking quality was different with the substitution 1R(1A) showing a significant positive effect compared to 1R(1B) and 1R(1D) substitutions. The alleles encoding HMW secalin subunits at the *Glu-R1* had an intermediate value between the null allele and subunit 2\* encoded at the *Glu-A1* locus. Negative effects were associated with the 1R(1B) and 1R(1D). It should be emphasized that these results are influenced by the presence of the monomeric ω-secalins and the 40k group of γ-secalins encoded by genes present at the *Sec-1* locus, on the short arm of the 1R chromosome [62].

A negative effect of the HMW secalins, on wheat quality characteristics, was also reported by Graybosch [63]. This result was obtained comparing 1B(1R) substitution lines with the 1BL/1RS sister lines. The presence of the 1RL resulted in further reductions in dough strength parameters compared to 1RS alone.

Agronomic and quality data were obtained by Kumlay, et al. [64] using the three translocation lines 1AS/1RL, 1BS/1RL and 1DS/1RL, in which the long arm of chromosome 1R replaced the long arms of the homeologous group 1 chromosomes of the bread wheat variety Pavon. This resulted in the introduction of the *Sec-3* locus, carrying genes encoding HMW secalins and elimination of the three different *Glu-1* loci carrying genes for the HMW-GS. Rheological data indicated that among the three translocation lines the 1AS/1RL was better than the control bread wheat variety Pavon carrying the subunit 1Ax1 associated at the *Glu-A1* locus.

Also in our case the N11<sup>XY7</sup> differs from N11 only for the two additional HMW secalins and the drastic changes in rheological characteristics observed, in addition to the high protein content, must in some way be associated to their presence.

In addition, the sequence of 1Ry gene present in XY7 included eight cysteine residues [37] rather than the seven normally present in y-type subunits in bread wheat, with the additional cysteine located within the C-terminal domain upstream of the one normally present in bread wheat; this cysteine residue has been suggested to be restricted to y-type HMW secalins by De Bustos and Jouve [65].

In bread wheat it has been widely demonstrated that number of subunits in addition to the number and distribution of cysteine residues, available for intermolecular bonds, are responsible for differences of breadmaking quality of different wheat varieties, as also mentioned in the introduction. Results have been obtained that show that additional cysteines not always influence positively the bread-making quality of flour, as in the case of the 1Dx5 subunit [66-68]. On the basis of the rheological data obtained we can hypothesize that the additional cysteine present in the 1Ry subunit contributes to modulate differently the size distribution of glutenin polymers, producing a dough with a very large extensibility.

## 4. Materials and Methods

### 4.1 Plant Materials

A wheat line was produced by crossing the bread wheat line N11 and the 2D(2R) disomic triticales XY7, followed by 4 backcrosses with N11 and designated N11<sup>XY7</sup>. The HMW-GS present in the seeds obtained after each backcross were analyzed by one dimensional polyacrylamide gel electrophoresis in SDS (SDS-PAGE), according to Feng et al. [37], and those containing seven HMW-GS subunits (5 typical of N11 plus the two HMW secalin subunits from rye) were selected for the next step of backcrossing. The gliadins were analyzed by acid polyacrylamide gel electrophoresis (A-PAGE) as described by Camerlengo et al. [69].

### 4.2 Field Trials

Bread wheat N11, XY7 and BC4F<sub>8</sub> seeds of N11<sup>XY7</sup> were grown in open field during 2016/2017 (Y1) and 2017/2018 (Y2) growing seasons at the Experimental Farm of the University of Tuscia, located in Viterbo, Italy (lat. 42°26' N, long. 12° 04' E, altitude 310 m a.s.l.), in 10 m<sup>2</sup> plot trials, with three replications. The experimental site is characterized by a Mediterranean climate with mean annual maximum and minimum temperatures of 19 and 8 °C, respectively, and annual rainfall of 743 mm. The agronomic practices were those recommended for high yields, including sowing 450 germinating seeds m<sup>-2</sup>, seed dressing, chemical control of weeds, 150 kg/ha of nitrogen applied in three top dressing, no application of growth regulators, fungicides, and pesticides.

After harvesting, test weight (TW) was determined by means of a Dickey-John GAC2000 grain analysis meter (Dickey-John Corp. Auburn, IL, USA), according to the supplied programme.

### 4.3 Qualitative Analyses

Grain samples (50 g) from each plot were ground to wholemeal using a 1-mm-sieve Cyclotec mill (Foss Tecator AB, Höganäs, Sweden). Grain hardness (Ha) (AACC 39-70) [70] was determined on the resulting wholemeal by a NIR System Model 6500 (FOSS NIR Systems, Laurel, MD).

The three replications for each line were bulked and milled with a Bona 4RB (Bona, Italy) experimental mill after tempering according to their hardness. Flour protein content (PC) (N × 5.7, dry

matter basis, AACC 39–11) was determined by NIRS. The gluten index was evaluated on flour according to ICC-158 [71] using the Glutomatic System (Perten Instruments, Hagersten, Sweden). All the analyses were repeated twice (technical replicates) for each sample.

The rheological properties of flour were evaluated using a Chopin alveograph, according to ICC-121 and a Brabender farinograph, according to ICC-115-D [71].

#### 4.4 Cell Cycle Synchronization

Wheat (*Triticum aestivum* L.): bread wheat line N11, XY7, and the derived line N11<sup>XY7</sup> and rye seeds (*Secale cereale* L.) were soaked in aerated water for 8 h and germinated on moistened filter paper for 2-3 days in the dark at 19°C. Root tips from 1-2 cm long roots were used for further analysis.

In root tips, cell cycle synchronization and accumulation of metaphases was accomplished according to Doležel et al. [72] with minor modifications. Young seedlings were transferred in an aerated Hoagland's solution [72] at 25°C and exposed to 1.25 mM hydroxyurea for 18 h, followed by a recovery period of 4 h. After blocking at metaphase by a 2 h treatment with 2.5 µM amiprophos-methyl, roots were incubated overnight in ice water and fixed in 3:1 ethanol-glacial acid. Metaphase spreads for Genomic *In Situ* Hybridization (GISH) and Fluorescence *In Situ* Hybridization (FISH) analysis were prepared from synchronized root tips as described by Schwarzacher and Heslop-Harrison [73]. Pre-treatments and stringency washes were applied only to the slides hybridized with total rye DNA (GISH) and pTa71 probe (standard FISH) while these steps were omitted in Non Denaturing FISH (ND-FISH) with oligonucleotides as probe.

#### 4.5 DNA Probes

Both the rye total genomic DNA, used as probe in GISH analysis, and the 18S-5.8S-26S rDNA clone pTa71 [74], probe in FISH experiments, were labeled with Cy3 (Cyanine 3) by nick-translation according to manufacturer's instructions (Nick Translation Mix, Roche). The oligonucleotide 5'-FITC-(GAA)<sub>7</sub>-3'-FITC was synthesized and labeled by Eurofins MWG Operon (Ebersberg, Germany).

#### 4.6 Non Denaturing FISH ND-FISH

Fast FISH labeling in non denaturing conditions was performed, using the synthetic oligonucleotides (GAA)<sub>7</sub> as probes, according to Cuadrado and Jouve [41] with the modifications as described by Giorgi et al. [75]. The hybridization mixture was composed of 1.6 ng/µl of probe DNA in a total volume of 60 µl of 2XSSC (300 mM sodium chloride, 0.3 mM trisodium citrate) per microscope slide. After incubation at room temperature for 1 h the slides were washed for 10 min in 4XSSC with 0.2% Tween20 and the chromosomal DNA were counterstained with DAPI (49,6-diamidino-2-phenylindole) and mounted in an antifade solution (Vectashield, Vector Labs, Burlingame CA).

#### 4.7 FISH

FISH with the probe pTa71 was performed on metaphase spreads according to Schwarzacher and Heslop-Harrison [73]. The hybridization mixture (50 µL) was made of 50% formamide, 2xSSC (0.30M NaCl, 0.030 M sodium citrate), 10% dextran sulfate, 30 ng mL<sup>-1</sup> denatured salmon sperm DNA, 100 ng probe and water to the final volume. The mix was spread on slides and DNA was denatured at 80 °C for 5 min in a thermalcycler. Hybridization was carried on overnight at 37°C in a moist chamber and the most stringent washes were performed twice in 20% formamide in 0.1xSSC at 42°C for 5 min. After DNA counterstaining with DAPI (0.2 µg mL<sup>-1</sup>), the slides were mounted in Vectashield antifade solution.

#### 4.8 GISH

GISH was performed according to Ceoloni et al. [42] with some modifications: wheat blocking DNA was used in a ratio of 20:1 to the probe, consisting of rye genomic DNA labeled by nick translation. Hybridization conditions and following steps were same as for FISH with the pTa71 probe.

## 5. Conclusions

Chromosome rearrangements are frequent in *Triticeae* and occur widely in natural and resynthesized polyploids [76-81].

The use of molecular markers, sophisticated cytogenetic approaches and biochemical analysis have greatly contributed to unveil molecular changes following wide crosses and demonstrated that the chromosomal rearrangements following the polyploidization processes, such as those involving triticale formation, are the result of changes of coding and noncoding sequences, regulatory elements, promoter regions and repetitive sequences [82,83]. The chromosomal rearrangements arising out of wheat x triticale or wheat x rye crosses have the potential to result in novel gene diversity affecting different agronomic characteristics, as well as processing and nutritional characteristics suggesting that many of the chromosomal changes may prove useful in the breeding improvement of the wheat crop.

According to Lukaszewsky and Gustafson [48] wheat/rye translocations can be produced in sufficient numbers to allow the use of this method for the introduction of alien variation into wheat research programs.

Recently, the detection of rye translocation lines, present in the bread wheat Pavon, with elevated Zn and Fe content has been reported by Velu et al. [18], whereas Szakács et al. [19] using a new 1RS/1BL translocation line, whose 1RS arm originated from the perennial *Secale cereanum*, found that the novel translocation had improved spike morphology traits, higher tillering capacity, fertility, winter hardiness and resistance against stripe rust and leaf rust. In addition, a large increase in arabinoxylans content (both total and water soluble) was found, compared to the parental bread wheat genotype improving the nutritional value of the material containing the novel 1RS/1BL translocation line. Johansson et al. [84] analyzing a diverse set of wheat-alien introgression lines with mainly introgressions of rye, but also of *Leymus* spp. and *Thinopyrum junceiforme*, found that resistance to diseases (rusts, powdery mildew, *S. triticae*, and *Fusarium* head blight), pests (Hessian fly and aphids), agronomic performance, yield potential, microelements (Fe, Zn, etc.), fertility, alpha amylase activity were increased. High variation was found both in grain protein concentration and gluten strength, measured as %UWP, within the lines, indicating large variation in bread-making quality and positive end-use quality.

De Bustos and Jouve [65] reported that the wild rye relatives *S. cereale* ssp. *afghanicum* and *dighoricum* have an additional cysteine residue in both Rx and Ry subunits compared to x- and y-type HMW-GS present in wheat, a situation never reported in bread wheat. Transfer of these gene from rye to wheat could provide useful material to investigate role of the increased number and position of cysteine on rheological and bread making characteristics of different wheat varieties.

In addition, to the many agronomical genes transferred in wheat, rye has also the potential to contribute to the improvement of the nutritional and processing characteristics of wheat.

**Supplementary Materials:** Supplementary materials can be found at [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1).

**Author Contributions:** Conceptualization, D.L.; Data curation, F.S., B.M., P.V., S.M., D.G., S.L., S.P., S.M., and D.L.; Funding acquisition, D.L., F.S. and S.M.; Investigation, D.L., F.S., and S.M.; Methodology, P.V., S.M., D.G., S.L., S.P., S.P.; Project administration, D.L., F.S., S.M.; Supervision D.L.; Validation, F.S., B.M., P.V., S.M., D.G., S.P., S.M.; Visualization, D.L.; F.S. and S.M. Writing—original draft, D.L., F.S. P.V., D.G.; Writing—review and editing, F.S., B.M., P.V., D.G., S.L., S.P., S.M., and D.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was partially funded by Italian Ministry of Education, University and Research (MIUR) in the frame of the MIUR initiative “Departments of excellence”, Law 232/2016.

**Acknowledgments:** The author wish to thank Giuseppe Colaprico for his technical assistance in making crosses, Alessandro Cammerata for his technical assistance in the rheological analyses and Anna Farina for her help in the cytogenetic analyses.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

A-PAGE      Acid polyacrylamide gel electrophoresis

FISH	Fluorescence <i>In Situ</i> Hybridization
FQN	Farinograph quality number
GISH	Genomic <i>In Situ</i> Hybridization
HMW-GS	High molecular weight glutenin subunit
LMW-GS	Low molecular weight glutenin subunit
SDS-PAGE	Polyacrylamide gel electrophoresis in SDS
TW	Test weight
UPP	Unextractable proteins

## References

1. Shiferaw, B.; Smale, M.; Braun, H.J.; Duveiller, E., Reynolds, M.; Muricho G. Crops that feed the world 10. Past successes and future challenges to the role played by wheat in global food security. *Food Secur.* **2013** *5*, 291–317.
2. Feldman, M.; Lupton, F.G.H.; Miller, T. E. Wheats. In *Evolution of crop plants*. Smartt J., Simmonds, N.W., Eds.; 3rd ed. Longman Scientific and Technical, London, 1995; pp. 184–192.
3. Shewry, P.R.; Hey, S.J. The contribution of wheat to human diet and health. *Food Energy Secur.* **2015** *4*, 178–202.
4. Dixon, J.; Braun, H.J.; Crouch, J. Transitioning wheat research to serve the future needs of the developing world. In *Wheat Facts and Futures*. Dixon, J., Braun, H.J., Kosina, P., Eds.; Mexico City (Mexico): CIMMYT; 2009; pp. 1-19.
5. Venske, E.; dos Santos, R.S.; Busanello, C.; Gustafson, P.; Costa de Oliveira, A. Bread wheat: a role model for plant domestication and breeding. *Hereditas* **2019** *156*, 16.
6. Dewey, D.R. The genomic system of classification as a guide to intergeneric hybridisation with the perennial *Triticeae*. In *Gene manipulation in plant improvement*. Gustafson, J.P. Ed.; New York, USA, Plenum Press 1984 pp. 209-279.
7. Feuillet, C.; Langridge, P.; Waugh, R. Cereal breeding takes a walk on the wild side. *Trends Genet.* **2008** *24*, 24-32.
8. Mujeeb-Kazi, A.; Kazi, A.; Dundas, I.; Rasheed, A.; Ogbonnaya, F.; Kishii, M.; Bonnett, D.; Wang, R.; Xu, S.; Chen, R.; Mahmood, T.; Bux, H.; Farrakh, S. Genetic diversity for wheat improvement as a conduit to food security. *Adv. Agron.* **2013** *122*, 179-257.
9. Vaccino, P.; Banfi, R.; Corbellini, M.; De Pace, C. Improving the wheat genetic diversity for end-use grain quality by chromatin introgression from the wheat wild relative *Dasypyrum villosum*. *Crop Sci.* **2010** *50*, 528-540.
10. Han, H.; Bai, L.; Su, J.; Zhang, J.; Song, L.; et al. Genetic rearrangements of six wheat–*Agropyron cristatum* 6P addition lines revealed by molecular markers. *PLoS ONE* **2014** *9*, e91066.
11. Rather, S.A., Sharma, D., Pandey, I., Joshi, N. Alien Gene Introgression in Wheat. In *Wheat a premier food crop*. Kumar A., Kumar A., Prasad B., Eds; Kalyani Publishers, New Delhi, pp. 90-109.
12. Li, Q.; Lu, Y.; Pan, C.; Wang, Z.; Liu, F.; Zhang, J.; Yang, X.; Li, X.; Liu, W.; Li, L. Characterization, identification and evaluation of a novel wheat-*Agropyron cristatum* (L.) Gaertn. disomic addition line II-30-5. *Genet. Resour. Crop Evol.* **2020** In press.
13. Sears, E.R. Genetic control of chromosome pairing in wheat. *Ann. Rev. Genet* **1976** *10*, 31-51.
14. Sears, E.R.; Gustafson, J.P. Use of radiation to transfer alien chromosome segments to wheat. *Crop Sci.* **1993** *33*, 897-901.
15. Endo, T.R. The gametocidal chromosome as a tool for chromosome manipulation in wheat. *Chromosome Res* **2007** *15*, 67-75.
16. Rabinovich, S.V. Importance of wheat-rye translocations for breeding modern cultivars of *Triticum aestivum* L. *Euphytica* **1998** *100*, 323–340.
17. Lukaszewski, A.J. Introgressions between wheat and rye. In *Alien Introgression in Wheat-Cytogenetics, Molecular Biology, and Genomics*. Molnár-Láng, M., Ceoloni, C., Doležel, J., Eds.; Springer: Berlin/Heidelberg, Germany, 2015 pp. 163-189.

18. Velu, G.; Crespo Herrera, L.; Guzman, C.; Huerta, J.; Payne, T.; Singh, R.P. Assessing genetic diversity to breed competitive biofortified wheat with enhanced grain Zn and Fe concentrations. *Front. Plant Sci.* **2019** *9*, 1971.
19. Szakács, È.; Szóke-Pázsai, K.; Kalapos, B.; Schneider, A.; Ivanizs, L.; Rakszegi, M.; Vida, G.; Molnár, I.; Molnár-Láng, M. 1RS arm of *Secale cereale* 'Kriszta' confers resistance to stripe rust, improved yield components and high arabinoxylan content in wheat. *Sci. Rep.* **2020** *10*, 1972.
20. Evtushenko, E.V.; Lipikhina, Y.A.; Stepochkin, P.I.; Vershinin, A.V. Cytogenetic and molecular characteristics of rye genome in octoploid triticale ( $\times$  Triticosecale Wittmack). *Comp. Cytogen.* **2019** *13*, 423–434.
21. Yuan, Z.; Liu, D.; Zhang, L.; Zhang, L.; Chen, W.; Yan, Z.; Zheng, Y.; Zhang, H.; Yen, Y. Mitotic illegitimate recombination is a mechanism for novel changes in high-molecular-weight glutenin subunits in wheat–rye hybrids. *PLoS One* **2011** *6*, e23511.
22. Yuan, Z.; Liu, M.; Ouyang, Y.; Zeng, X.; Hao, M.; Zhang, L.; Ning, N.; Yan, Z.; Liu, D. The detection of a de novo allele of the *Glu-1Dx* gene in wheat–rye hybrid offspring. *Theor. Appl. Genet.* **2014** *27*, 2173–2182.
23. Li, H.; Guo, X.; Wang, C.; Ji, W. Spontaneous and divergent hexaploid triticales derived from common wheat/rye by complete elimination of D-genome chromosomes. *PLoS ONE* **2015** *10*, e0120421.
24. Shewry, P.R.; Halford, N.G.; Lafiandra, D. Genetics of wheat gluten proteins. *Adv. Genet.* **2003** *49*, 111–184.
25. Field, J.M.; Shewry, P.R.; Mifflin, B.J. Solubilization and characterization of wheat gluten proteins. Correlations between the amount of aggregated proteins and baking quality. *J. Sci. Food Agric.* **1983** *34*, 370–377.
26. Gupta, R.B.; Popineau, Y.; Lefebvre, J.; Cornec, M.; Lawrence, G.J.; MacRitchie, F. Biochemical basis of flour properties in bread wheats. II. Changes in polymeric protein formation and dough/gluten properties associated with the loss of low Mr or high Mr glutenin subunits. *J. Cereal Sci.* **1995** *21*, 103–116.
27. Weegels, P.L.; Hamer, R.J.; Schofield, J.D. Functional properties of wheat glutenin. *J. Cereal Sci.* **1996** *23*, 1–17.
28. Lafiandra, D.; Masci, S.; Blumenthal, C.; Wrigley, C.W. The formation of glutenin polymer in practice. *Cereal Foods World* **1999** *44*, 572–578.
29. Payne, P.I. (1987) Genetics of wheat storage proteins and the effect of allelic variation on bread-making quality. *Annu. Rev. Plant Physiol.* **1987** *38*, 141–153.
30. Shewry, P.R.; D'Ovidio, R.; Lafiandra, D.; Jenkins, J.A.; Mills, E.C.; Békés, F. Wheat grain proteins. In *Wheat: Chemistry and Technology*, 4th ed.; Khan K., Shewry P.R., Eds.; Elsevier Inc. in cooperation with AACC International 2009 pp. 223–298.
31. Santagati, V.D.; Sestili, F.; Lafiandra, D.; D'Ovidio, R.; Rogniaux, H.; Masci, S., 2016. Characterization of durum wheat high molecular weight glutenin subunits Bx20 and By20 sequences by a molecular and proteomic approach. *J. Mass Spect.* **2016** *51*, 512–517.
32. Lawrence, G.J.; MacRitchie, F.; Wrigley, C.W. Dough and baking quality of wheat lines deficient in glutenin subunits controlled by the *Glu-A1*, *Glu-B1* and *Glu-D1* loci. *J. Cereal Sci.* **1988** *7*, 109–112.
33. Halford, N.G.; Field, J.M.; Blair, H.; Urwin, P.; Moore, K.; Robert, L.; Thompson, R.; Flavell, R.B.; Tatham, A.S.; Shewry, P. Analysis of HMW glutenin subunits encoded by chromosome 1A of bread wheat (*Triticum aestivum* L.) indicates quantitative effects on grain quality. *Theor. Appl. Genet.* **1992** *83*, 373–378.
34. Gupta, R.B.; MacRitchie, F. Allelic variation at glutenin subunit and gliadin loci, *Glu-1*, *Glu-3* and *Gli-1* of common wheats. II. Biochemical basis of the allelic effects on dough properties. *J. Cereal Sci.* **1994** *19*, 19–29.
35. Pirozi, M.R.; Margiotta, B.; Lafiandra, D.; MacRitchie, F. Composition of polymeric proteins and bread-making quality of wheat lines with allelic HMW-GS differing in number of cysteines. *J. Cereal Sci.* **2008** *48*, 117–122.
36. Wang, G.; Snape, J.W.; Hu, H.; Rogers, W.J. The high-molecular-weight glutenin subunit compositions of Chinese bread wheat varieties and their relationship with bread-making quality. *Euphytica* **1993** *68*, 205–212.
37. Feng, B.; Sestili, F.; Masci, S.; Margiotta, B.; Xu, Z.; Yang, Z.; Xiang, C.; Zhou, C.; Lafiandra, D.; Wang, T. The Chinese bread wheat cultivar Xiaoyanmai 7 harbours genes encoding a pair of novel high-molecular-weight glutenin subunits inherited from cereal rye. *Crop Pasture Sci.* **2016** *67*, 29–36.
38. Cuadrado, Á.; Schwarzacher, T. The chromosomal organization of simple sequence repeats in wheat and rye genomes. *Chromosoma* **1998** *107*, 587–594.
39. Cuadrado, Á.; Cardoso, M.; Jouve, N. Physical organisation of simple sequence repeats (SSRs) in *Triticeae*: structural, functional and evolutionary implications. *Cytogenet. Genome Res.* **2008** *120*, 210–219.

40. Pedersen, C.; Rasmussen, S.K.; Lindelaursen, I. Genome and chromosome identification in cultivated barley and related species of the *Triticeae* (*Poaceae*) by *in situ* hybridization with the GAA-satellite sequence. *Genome* **1996** *39*, 93–104.
41. Cuadrado, A.; Jouve, N. Evolutionary trends of different repetitive DNA sequences during the speciation of the genus *Secale*. *J. Hered.* **2002** *93*, 339–345.
42. Ceoloni, C.; Biagetti, M.; Ciaffi, M.; Forte, P.; Pasquini, M. Wheat chromosome engineering at the 4x level: the potential of different alien gene transfers into durum wheat. *Euphytica* **1996** *89*, 87–97.
43. Islam-Faridi, M.N.; Mujeeb-Kazi, A. Visualization of *Secale cereale* DNA in wheat germ plasm by fluorescent *in situ* hybridization. *Theor. Appl. Genet.* **1995** *90*, 595–600.
44. Fu, S.L.; Yang, M.Y.; Fei, Y.Y.; Tan, F.Q.; Ren, Z.L.; et al. Alterations and abnormal mitosis of wheat chromosomes induced by wheat-rye monosomic addition lines. *PLoS ONE* **2013** *8*, e70483.
45. Fu, S.L.; Yang, M.Y.; Ren, Z.L.; Yan, B.J.; Tang, Z.X. Abnormal mitosis induced by wheat-rye 1R monosomic addition lines. *Genome* **2014** *57*, 21–28.
46. Tang, Z.; Li, M.; Chen, L.; Wang, Y.; Ren, Z.; Fu, S. New types of wheat chromosomal structural variations in derivatives of wheat-rye hybrids. *PLoS ONE* **2014** *9*, e110282.
47. McClintock, B. The significance of responses of the genome to challenge. *Science* **1984** *226*, 792–801.
48. Lukaszewski, A.J.; Gustafson, J.P. Translocations and modifications of chromosomes in triticalexwheat hybrids. *Theor. Appl. Genet.* **1983** *64*, 239–248.
49. Darlington, C.D. Misdivision and the genetics of the centromere. *J. Genet.* **1939** *37*, 341–364.
50. Wang, J.; Liu, Y.; Su, H.; Guo, X.; Han, F. Centromere structure and function analysis in wheat-rye translocation lines. *Plant J.* **2017** *91*, 199–207.
51. Therman-Suomalainen, E. Investigation on secondary constriction in *Polygonatum*. *Hereditas* **1949** *35*, 86–108.
52. Huang, M.; Li, H.; Zhang, L.; Gao, F.; Wang, P.; et al. Plant 45S rDNA clusters are fragile sites and their instability is associated with epigenetic alterations. *PLoS ONE* **2012** *7*, e35139.
53. Badaeva, E.D.; Dedkova, O.S.; Gay, G.; Pukhalskyi, V.A.; Zelenin, A.V.; Bernard, S.; Bernard, M. Chromosomal rearrangements in wheat: their types and distribution. *Genome* **2007** *50*, 907–926.
54. Ferguson-Smith, M.A. Chromosome Abnormalities. *Proc. R. Soc. Med.* **1962** *55*, 471–477.
55. Li, H.; Guo, X.; Wang, C.; Ji, W. Spontaneous and divergent hexaploid triticales derived from common wheat × rye by complete elimination of D-genome chromosomes. *PLoS ONE* **2015** *10*, e0120421.
56. Nash, D.; Lanning, S.P.; Fox, P.; Martin, J.M.; Blake, N.K.; Souza, E.; Graybosch, R.A.; Giroux, M.J.; Talbert, L.E. Relationship of dough extensibility to dough strength in a spring wheat cross. *Cereal Chem.* **2006** *83*, 255–258.
57. Fido, R.J.; Békés, F.; Gras, P.W.; Tatham, A.S. Effects of  $\alpha$ -,  $\beta$ -,  $\gamma$ - and  $\omega$ -gliadins on the dough mixing properties of wheat flour. *J. Cereal Sci.* **1997** *26*, 271–277.
58. Uthayakumaran, S.; Tömösközi, S.; Tatham, A.S.; Savage, A.W.J.; Gianibelli, M.C.; Stoddard, F.L.; Bekes, F. Effects of gliadin fractions on functional properties of wheat dough depending on molecular size and hydrophobicity. *Cereal Chem.* **2001** *78*, 38–141.
59. Gupta, R.B.; Batey, I.L.; MacRitchie, F. Relationships between protein composition and functional properties of wheat flours. *Cereal Chem.* **1992** *69*, 125–131.
60. Kipp, B.; Belitz, H.D.; Seilmeier, W.; Wieser, H. Comparative studies of high molecular-weight, subunits of rye and wheat. I. Isolation and biochemical characterisation and effects on gluten extensibility. *J. Cereal Sci.* **1996** *23*, 227–234.
61. Amiour, N.; Jahier, J.; Tanguy, A.M.; Chiron, H.; Branlard, G. Effect of 1R(1A), 1R(1B) and 1R(1D) substitution on technological value of bread wheat. *J. Cereal Sci.* **2002** *35*, 149–160.
62. Shewry, P.R.; Bradberry, D.; Franklin, J.; White, R.P. The chromosomal locations and linkage relationships of the structural genes for the prolamins storage proteins (secalins) of rye. *Theor. Appl. Genet.* **1984** *69*, 63–69.
63. Graybosch, R.A. Uneasy unions: quality effects of rye chromatin transfers to wheat. *J. Cereal Sci.* **2001** *33*, 3–16.
64. Kumlay, A.M.; Baenziger, P. S.; Gill, K.S.; Shelton, D.R.; Graybosch, R.A.; Lukaszewski, A.J.; Wesenberg, D.W. Understanding the effect of rye chromatin in bread wheat. *Crop Sci.* **2003** *43*, 1643–1651.
65. De Bustos, A.; Jouve, N. Characterisation and analysis of new HMW-glutenin alleles encoded by the *Glu-R1* locus of *Secale cereale*. *Theor. Appl. Genet.* **2003** *107*, 74–83.
66. Lindsay, L.; Tamas, L.; Appels, R.; Skerritt, J.H. direct evidence that the number and location of cysteine residues affect glutenin polymer structure. *J. Cereal Sci.* **2000** *31*, 321–333.

67. Juhasz, A.; Larroque, O.R.; Tamas, L.; Hsam, S.L.K.; Zeller, F.J.; Bekes, F.; Bedo, Z. Bánkúti 1201 - An old Hungarian wheat variety with special storage protein composition. *Theor. Appl. Genet.* **2003** *107*, 697–704.
68. Gao, X.; Appelbee, M.J.; Mekuria, G.T.; Chalmers, K.J.; Mather, D.E. A second 'overexpression' allele at the *Glu-B1* high-molecular-weight glutenin locus of wheat: sequence characterisation and functional effects *Theor. Appl. Genet.* **2012** *124*, 333–343.
69. Camerlengo, F.; Sestili, F.; Silvestri, M.; Colaprico, G.; Margiotta, B.; Ruggeri, R. et al Production and molecular characterization of bread wheat lines with reduced amount of  $\alpha$ -type gliadins. *BMC Plant Biol.* **2017** *17*, 248.
70. AACC. 2000. Approved methods of the American Association of Cereal Chemists. 10th ed. The Association, St Paul, MN
71. ICC. 1992. Standard methods of the International Association for Cereal Chemistry. ICC, Vienna, Austria.
72. Doležel, J.; Číhalíková, J.; Weiserová, J.; Lucretti, S. Cell cycle synchronization in plant root meristems. *Meth. Cell Sci.* **1999** *21*, 95–107.
73. Schwarzbacher, T.; Heslop-Harrison, P. Practical *in situ* hybridization. Oxford, UK; 2000 BIOS Scientific Publishers Ltd.
74. Gerlach, W.L.; Bedbrook, J.R. Cloning and characterization of ribosomal RNA genes from wheat and barley. *Nucleic Acids Res.* **1979** *7*, 1869–1885.
75. Giorgi, D.; Farina, A.; Grosso, V.; Gennaro, A.; Ceoloni, C.; Lucretti, S. FISHIS: Fluorescence *in situ* hybridization in suspension and chromosome flow sorting made easy. *PLoS ONE* **2013** *8*, e57994.
76. Shaked, H.; Kashkush, K.; Ozkan, H.; Feldman, M.; Levy, A.A. Sequence elimination and cytosine methylation are rapid and reproducible responses of the genome to wide hybridization and allopolyploidy in wheat. *Plant Cell* **2001** *13*, 1749–1759.
77. Ozkan, H.; Levy, A.A.; Feldman, M. Allopolyploidy-induced rapid genome evolution in the wheat (*Aegilops-Triticum*) group. *Plant Cell* **2001** *13*, 1735–1747.
78. Kashkush, K.; Feldman, M.; Levy, A.A. Gene loss, silencing and activation in a newly synthesized wheat allotetraploid. *Genetics* **2002** *160*, 1651–1659.
79. Feldman, M.; Levy, A. Genome evolution due to allopolyploidization in wheat. *Genetics* **2012** *192*, 763–774.
80. Wang, Q.X.; Gao, A.N.; Yang, X.M.; Li, L.H. (2014) Chromosome changes after polyploidization in *Triticeae*. *J. Syst. Evol.* **2014** *52*, 790–793.
81. Bariah, I.; Keidar-Friedman, D.; Kashkush, K. Identification and characterization of large-scale genomic rearrangements during wheat evolution. *PLoS ONE* **2020** *15*, e0231323.
82. Bento, M.; Gustafson, J.P.; Viegas, W.; Silva, M. Size matters in *Triticeae* polyploids: Larger genomes have higher remodeling. *Genome* **2011** *54*, 175–183.
83. Zheng, X.L.; Zhou, J.P.; Zang, L.L.; Tang, A.T.; Liu, D.Q.; Deng, K.J.; Zhang, Y. Genetic and epigenetic alterations induced by different levels of rye genome integration in wheat recipient. *Genet. Mol. Res.* **2016** *15*, gmr.150280.
84. Johansson, E.; Henriksson, T.; Prieto-Linde, M.L.; Andersson, S.; Ashraf, R.; Rahmatov, M. Diverse wheat-alien introgression lines as a basis for durable resistance and quality characteristics in bread wheat. *Front. Plant Sci.* **2019** *11*, 1067.