

The Dirac electron consistent with proper gravitational and electromagnetic field of the Kerr-Newman solution

Alexander Burinskii

Theor. Phys. Laboratory, NSI, Russian Academy of Sciences,
B. Tulkaya 52 Moscow 115191 Russia, email: burinskii@mail.ru

January 3, 2021

Abstract

We consider the Dirac electron as a particle-like solution consistent with its own Kerr-Newman (KN) gravitational field. The regularized by Israel-López KN solution forms a bag model – the thin superconducting disk coupled with circular string placed along its perimeter. Using unique features of the Kerr-Schild coordinate system, which linearizes Dirac equation in KN space, we obtain solutions of the Dirac equation consistent with KN gravitational and electromagnetic field, and show that the corresponding solution take the form of a massless relativistic string. Obvious parallelism with quantum theory explains remarkable properties of the electron in relativistic scattering processes.

1 Introduction

One of the main points of confrontation between Gravity and Quantum theory is the structure of elementary particles, which are considered in quantum theory as structureless, like a point-like electron in Dirac theory, but must be represented as an extended field model in configuration space for compatibility with the stress-energy tensor of Einstein's equations.

A revolutionary step towards unification quantum with gravity was taken in superstring theory, which represented particles as extended strings. Gravitational black holes (BH) have been considered as candidates for elementary particles repeatedly since 1980, and since the 1990s, they have also attracted attention in the theory of superstrings.

However, as one of its founders, John Schwartz, noted, "... Since 1974, superstring theory has ceased to be regarded as particle physics... " and "... a realistic model of elementary particles still seems a distant dream ..." [1].

Meanwhile, a renewed interest to relationships between black holes and elementary particles has been obtained recently in the works [2, 3, 4, 5].

Formation of BHs is related with gravitational effect of frame-dragging. In the rotating Kerr-Newman BH solution, with parameters J, m, a corresponding to spin, mass and Kerr's rotational parameter a of elementary particle, spin creates a giant over-rotating dragging of space, which is directed along of direction of rotation, leading to a new important effect, formation of the closed Wilson loop, which never was used in particle physics before.

In contrast to considered earlier cases of the Schwarzschild or Reissner-Nordström gravity, the characteristic scale of the KN gravity is essentially increases, because it is determined by radius of the Kerr singular ring

$$a = \frac{J}{mc}, \quad (1)$$

which corresponds to the reduced Compton wave length of the particle.

This fact, established already in the first models of an electron based on the Kerr geometry [6, 7, 8, 9, 10, 11, 12] was remarkable itself, because it was known, but was not timely estimated as one of the first evidences of the correspondence between KN particle and quantum theory.




Figure 1: Kerr congruence and Kerr singular ring generated by null congruence k^μ .

The gigantic ratio between the spin and mass values for elementary particles in KN geometry violated the generally accepted concept of the weakness of gravity, based on the earlier estimations of gravitational radius of the Schwarzschild solution

$$r_g = 2Gm. \quad (2)$$

Gravitational field of an electron corresponding to the Kerr-Newman solution was singular and changed topology of space at the Compton distance.

In 1968, Carter obtained that the Kerr-Newman (KN) solution for a charged and rotating black hole (BH) has gyromagnetic ratio $g = 2$ – just the same as that of the Dirac electron [6]. It gave rise to study of the electron model based on the KN solution, see [6, 7, 8, 9, 10, 11, 12, 13, 14] and so on.

It should be noted that the KN electron model is not actually a black hole, because taking the parameters of KN solution in correspondence with parameters of an electron, mass m , charge e and angular momentum $J = ma \sim \hbar/2$, we obtain the relation $a^2 \gg (m^2 + e^2)$ which shows that the rotation parameter a is so large, that all horizons of the BH solution disappear. There appears the Kerr singular ring, which was hidden earlier behind the horizon of the KN solution. This ring forms a type of the door that opens the way to another sheet of the Kerr space. The space becomes two-sheeted, having the basic background and some kind of the mirror Alice world behind the Kerr ring.

In previous papers [15, 16, 17] we developed the line started by W. Israel [8], who suggested to truncate the second sheet of the Kerr geometry along the *disk* spanned by the Kerr singular ring. After analysis of the Israel source by Hamity [9], a modified *disk-like source*




Figure 2: Disk-like source of the regularized solution of KN as a bag model. Deformation of the disk at different ratios of parameters $R = r_e$ and a . For $r_e = e^2/2m$ disk is very thin, and $r_e/a = \alpha$ corresponds to the fine structure const.

was suggested by C. López [10] as an ellipsoidal vacuum bubble – a thin shell covering the Kerr singular ring and matching with the external KN solution.

In the papers [15, 16, 17] we considered a generalization of the López model, in which the KN bubble is formed of the Higgs field, which is in a superconducting vacuum state. The thin shell of bubble is replaced by a domain-wall solution, which is described by the Landau-Ginzburg (LG) supersymmetric model of phase transition. Domain Wall (DW) interpolates between the superconducting (and supersymmetric) internal vacuum state and the external exact gravitational KN solution.

The obtained by C.Lopez bubble source of the KN geometry [10] (see Fig.2) was presented in our works as a supersymmetric and superconducting bag model. The reason for interpretation of the KN source as a bag model, is their ability to elasticity and deformations under influence of external conditions, which is known from the behavior of the well-known MIT and SLAC bag models [18, 19]. It was assumed also that bags are similar to strings and can turn into strings under strong deformations [20, 21].

Meanwhile, one feature of the KN bag significantly distinguishes it from the MIT and SLAC bag models – the usual bag models form *a cavity in superconductor*, while the KN bag must have an internal superconducting state. This feature is the source of problems, that force us to use a supersymmetric LG field model of phase transition.

The corresponding Hamiltonian was reduced to Bogomolnyi form, and it was shown that this soliton forms a supersymmetric, PBS saturated state. In this letter we would like pay attention to the used intricate method reduction to Bogomolnyi form, which was apparently first suggested in [22] for a two-dimensional kink solution, and then successfully used for the planar DW in [23, 24, 25, 26, 27]. We generalize this method to the much more complex case of the Kerr geometry, in which the source is spinning and bounded by the DW of ellipsoidal form. Besides, it is formed by the system of chiral fields, when one of them depends on the Kerr angular coordinate and time. With respect to previous treatment [15, 16], we obtain new very important feature of the DW source – formation of the DW-antiDW (breather) structure, which is very essential for the discussed here solutions of the Dirac equations.

Our task here is to obtain a self-consistent solution of the Dirac equation embedded in the proper gravitational and electromagnetic field of the electron corresponding to Kerr-

Newman solution. By solving this problem, we obtain that Kerr-Schild coordinate system is unique, in the sense that it allows us to use γ - matrices of the auxiliary Minkowski space, where the Dirac equations in proper gravitational and electromagnetic field are linearized.

We obtain that solutions of the Dirac equations take the form of a massless relativistic string based on an orientifold structure discussed in one of our old works [28].

2 Kerr-Schild geometry and structure of KN solution

Specific feature of the Kerr-Schild approach is the use of the auxiliary Minkowski space $\mathbb{M}^4$, (signature $(-+++)$), with Cartesian coordinates $x = x^\mu = (t, x, y, z)$.

In these coordinates, metric of the KN solutions is [7]

$$g_{\mu\nu} = \eta_{\mu\nu} + 2Hk_\mu k_\nu, \quad (3)$$

where $\eta_{\mu\nu}$ is flat metric of the auxiliary Minkowski space, and H is the scalar function which for the KN solution takes the form

$$H_{KN} = \frac{mr - e^2/2}{r^2 + a^2 \cos^2 \theta}. \quad (4)$$

The KN vector potential is given as

$$A_\mu = \frac{-er}{(r^2 + a^2 \cos^2 \theta)} k_\mu. \quad (5)$$

The field $k^\mu(x)$ forms a Principal Null Congruence (PNC), $k_\mu k^\mu = 0$, shown on Fig.1. In terms of BH geometry this field shows a local direction of dragging the frame, that in the case of overrotating HB solutions produces closed Wilson lines surrounding the source of KN geometry, see Fig.3.




Figure 3: Deformations of the Kerr coordinate $\phi = \text{const.}$ caused by dragging of space in angular direction near the Kerr singular ring.

Kerr's congruence can be represented as an electromagnetic radiation which propagates (with twist) from infinity towards the Kerr ring, penetrates it, and coming out on the other sheet of the Kerr geometry goes out again to infinity. In Cartesian coordinates $x^\mu \in M^4$, the form $k_\mu dx^\mu$ shows local direction of frame-dragging.

In the Kerr angular coordinates PNC is presented in [7] by the form

$$k_\mu dx^\mu = dr - dt - a \sin^2 \theta d\phi_K. \quad (6)$$

The relation between Cartesian coordinates and Kerr's angular coordinates is the following

$$\begin{aligned} x + iy &= (r + ia) \exp\{i\phi_K\} \sin \theta, \\ z &= r \cos \theta, \quad \rho = r - t, \end{aligned} \quad (7)$$

The incoming PNC is directed to the Kerr ring. Rays lying in equatorial plane ($\cos \theta = 0$) focus on the Kerr singular ring. Other incoming rays, passing through the ring, turn into out-going rays propagating on another (say "negative") sheet of the Kerr space. Thus, the Kerr solution in the KS form describes two different sheets of space-time with two different congruences

$$k_\mu^\pm dx^\mu = \pm dr - dt - a \sin^2 \theta d\phi_K \quad (8)$$

and two different metrics

$$g_{\mu\nu}^\pm = \eta_{\mu\nu} + 2H k_\mu^\pm k_\nu^\pm \quad (9)$$

on the same Minkowski background $x^\mu \in M^4$. Working with *outgoing* Kerr field corresponding to *retarded* potentials, we choose sign plus in (8), and following [7] we take $k_\mu = k_\mu^+$.

The Kerr theorem.

Kerr theorem defines two fields of PNC, $k^+(x)$ and $k^-(x)$, in terms of Penrose's twistor theory [29, 30, 31]. Kerr theorem presents two complex analytic solutions $Y^\pm$ of the equation

$$F(T^A) = 0, \quad (10)$$

where F is quadratic holomorphic function of the projective twistor coordinates $T^A = \{Y, \zeta - Yv, u + Y\bar{\zeta}\}$, $A = 1, 2, 3$, and

$$\begin{aligned} 2^{\frac{1}{2}}\zeta &= x + iy, & 2^{\frac{1}{2}}\bar{\zeta} &= x - iy, \\ 2^{\frac{1}{2}}u &= z + t, & 2^{\frac{1}{2}}v &= z - t, \end{aligned} \quad (11)$$

are the null Cartesian coordinates of the auxiliary Minkowski space $x^\mu \in \mathbb{M}^4$.

In the class of quadratic in Y functions $F(T^A)$, the Kerr theorem gives two analytic solutions $Y^\pm(x^\mu)$, of the equation (10), which correspond to two projective spinor coordinates

$$Y^+ = \xi^{\dot{1}}/\xi^{\dot{0}}, \quad Y^- = \eta_1/\eta_0, \quad (12)$$

which are antipodically conjugate

$$Y^+ = -1/\bar{Y}^-, \quad (13)$$

and the corresponding Weyl spinors $\xi^{\dot{\alpha}}$ and η_α define two antipodal fields of the principal null directions

$$k^{\mu+} = \bar{\xi}^\alpha \sigma_{\alpha\dot{\alpha}}^\mu \xi^{\dot{\alpha}}, \quad k^{\mu-} = \bar{\eta}_{\dot{\alpha}} \bar{\sigma}^{\mu\dot{\alpha}\alpha} \eta_\alpha. \quad (14)$$

3 Shape of the KN bag model and Wilson loop

The López boundary of the bubble, where the KN space can be matched continuously with the flat internal metric $\eta_{\mu\nu}$, is unambiguously determined by the Kerr-Shild metric form (3), as the surface where $H = 0$. Setting $H_{KN} = 0$ we obtain

$$r = r_e = e^2/2m, \quad (15)$$

that gives us the “classical” electron radius.

Since r is the Kerr radial coordinate, we obtain that the bag boundary represents indeed an oblate ellipsoidal surface – a thin disk of the radius a , which is about the reduced Compton wave length, and the thickness of the disk r_e , which is equal to classical electron radius. One sees that degree of oblateness of the disk is $r_e/a = 1/137$ that corresponds to the fine structure constant α .

Therefore, the Kerr-Newman spin parameter a leads to a strong deformation of the shape of the bag model, and this deformation of the bag leads to the appearance of a relativistic string at the sharp edge of the KN disk (see Fig.4).

The existence of this string is evidenced by the Wilson loop of the vector potential placed along border of the bag, which was obtained first in [32] and then discussed in [15, 33, 34].

From (5) and (6) we obtain that vector-potential of the regularized KN solution takes its maximal value in the equatorial plane ($\cos\theta = 0$) at the bag border $r = r_e$,

$$A_\mu^{max} dx^\mu = -\frac{2m}{e}(dr - dt - a d\phi_K). \quad (16)$$

This potential is tangent to the bag border $r = r_e$, and for the fixed time $t = const.$, it forms the closed Wilson loop $C : \phi_K \in [0, 2\pi]$, so that the loop integral $W(C) = P \exp e \oint_C A_\mu^{max} dx^\mu$, gives the following incursion of the potential

$$\delta\phi = e \oint_C A_{\phi_K}^{max} d\phi_K. \quad (17)$$

Integration gives $\delta\phi = 4\pi ma$, and using relation $J = ma$ we obtain

$$\delta\phi = 4\pi J. \quad (18)$$

Definiteness of potential requires $\delta\phi = 4\pi J$, leading to quantum condition $J = \frac{1}{2}$.




Figure 4: Disk-like shape of the Kerr-Newman bag model. Border of the superconducting bag ends with Wilson loop forming a closed circular string.

4 String structure and superficial currents on the border of KN bag

4.1 Scattering process and orientation of the Kerr disk

On the initially two-sheeted KN space-time, the directed in future vortex field of the Kerr congruence k^+ forms an *in-going* on the negative sheet of the Kerr space $r^- < 0$, where it is directed towards the Kerr singular ring. Penetrating through the ring, this field continues analytically on the second sheet $r^+ > 0$, turning into an *out-going* where another coordinate system is used, (7).

Although, in the regularized KN solution, the passage to the r^- sheet is closed, consideration of the analog of this sheet is relevant in the scattering process, when we observe the in-going field incident on the source of the KN solution before the scattering, and then the signal reflected in the scattering process in the form of an out-going field.

In contrast to the case with the negative sheet of the Kerr solution, in this case we use the same Kerr's coordinate system (7) for both in-going and out-going fields, in which we do a replacement of $r \rightarrow -r$, getting an equivalent coordinate transformation for the in-going field on r^- ,

$$\begin{aligned} x + iy &= (r - ia) \exp\{-i\phi_K\} \sin \theta, \\ z &= -r \cos \theta, \quad \rho = -r - t, \end{aligned} \quad (19)$$

compatible with metric

$$g_{\mu\nu}^- = \eta_{\mu\nu} + 2H k_\mu^- k_\nu^-, \quad (20)$$

and with in-going Kerr congruence k_μ^- .

This process shows that disk-like source of KN field has two faces: one from the side of the in-going fields k_μ^- , and the other from the side of the out-going fields k_μ^+ . These two sides are related with reverse sign of the disk rotation $a \rightarrow -a$, and change the orientation angle $\phi_k \rightarrow -\phi_k$ for the incoming field.

The corresponding string-like structure, was suggested in [28] as an orientifold string. This string forms the Kerr's light-like world-sheet $X = X_L(\tau + \sigma)$, containing only the left modes on the fundamental interval $[0, \pi]$. For a static picture of the Kerr disk at $t = 0$, the orientifold string is formed as a parity operator $\Omega : [\sigma \rightarrow 2\pi - \sigma]$, which covers the string world sheet twice: first time on the interval $[0, \pi]$, and second time on the interval $[2\pi - \sigma]$ in opposite direction.

The full orientifold world-sheet is formed as a folded string on the doubled interval $\sigma \in [0, 2\pi]$, and contains the sum of the left and right modes $X = X_L(\tau + \sigma) + X_R(\tau - \sigma)$.

The orientifold string is left-right symmetric in the static representation, $t = \text{const.}$, which in quantum theory is called as Heisenberg picture, however the symmetry Ω is broken on the rotating disk.

4.2 Surface currents caused by Wilson loop, Higgs phases and stringy parametrization

The Kerr-Newman solution demonstrates an intrinsic connection to string theory. The role of the string is played by the singular ring of the Kerr solution [12, 13]. A regularized version

of this string occurs at the sharp edge of the disk-like boundary of the ellipsoidal bag forming the regular source of the Kerr-Newman solution.




Figure 5: KN disk as a carrier of the orientifold string in the static $t = \text{const.}$ picture. Kinematic relations in the equatorial plane of KN disk. The shown out-going light-like beam of the Kerr congruence k^μ is tangent to Kerr singular ring and crosses the edge of the disk at the angle $\theta_r = \arctan r_e/a \approx \alpha$.

As is known, the Higgs field model coincides with the Landau-Ginzburg (LG) model for a phase transition in a superconducting medium [35]. Similarly, the supersymmetric Higgs model is described by a supersymmetric (or generalized) Landau-Ginzburg (LG) model, [22, 26].

The corresponding supersymmetric bag model is formed by the Domain Wall (DW), which separates the external gravitational field KN from the flat inner space filled with supersymmetric vacuum of the Higgs field.

Although the consistent description of this phase transition requires a supersymmetric scheme of phase transition with several chiral fields, [15, 16, 17], the simple LG field model with one chiral field can describe each separate process of the phase transition with creation superficial currents on the boundary of the bag. Corresponding Lagrangian with one Higgs fields is, [35],

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}(\mathcal{D}_\mu\Phi)(\mathcal{D}^\mu\Phi)^* - V(|\Phi|), \quad (21)$$

where $\mathcal{D}_\mu = \nabla_\mu + ieA_\mu$ are covariant derivatives with vector-potential A_μ , $F_{\mu\nu} = A_{\mu,\nu} - A_{\nu,\mu}$, and

$$V = \lambda(\Phi\Phi^* - \eta^2)^2, \quad (22)$$

where η is the v.e.v. of the Higgs field Φ , $\eta = \langle |\Phi| \rangle$.

Superconducting vacuum state of the Higgs field inside the Bag leads to equations

$$\square A_\mu = I_\mu = e|\Phi|^2(\chi_{,\mu} + eA_\mu), \quad (23)$$

which shows that inside the superconductor current I_μ is pushed out, $I_\mu = 0$, and is concentrated in a surface layer with a depth of penetration δ , [36]). Potential of the KN field (5) increases near the bag boundary, and takes maximum in the equatorial plane, near the bag boundary $r = R = e^2/2m$, $\cos \theta = 0$.

According the Wess-Zumino model, the supersymmetric QED is described by two Higgs fields $\Phi_+ = |\Phi_+|e^{-ie\chi_+}$ and $\Phi_- = |\Phi_-|e^{ie\chi_-}$, [37], and the equation (23) allows us to connect two phases of the Higgs fields χ_+ and χ_- with two boundaries of the KN bag model A_μ^- and A_μ^+ , which were obtained in the double-face structure of the KN disk [17], related with congruences k^+ and k^- , and forming a DW-AntiDW structure, known also as "breather" [38].

Integration of the LG equations for out-going phase of the Higgs field, placed on the boundary $r = r^+$, gives $\chi_+|_{r^+} = 2m(t + a\phi_K)$,

while for the in-going Higgs phase, placed on the boundary $r = r^-$, we obtain

$$\chi_-|_{r^-} = 2m(t - a\phi_K),$$

where we took also into account the change of charge sign by the transition $a \rightarrow -a$.

Therefore, on the boundary $r = r^+$ we obtain the potential $eA_0 = 2m$, $eA_{\phi_K} = 2ma$, and on the boundary $r = r^-$ the potential $eA_0 = -2m$, $eA_{\phi_K} = -2ma$.

Applying these solutions to the out-going vector field $A_\mu^+(r_e^+)$ on the boundary $r = r_e^+$, which is dragged by gravitational field of the Kerr congruence, forming the closed Wilson loop $C^+ : t = \text{const.}$ on the border $r = r_e^+$ we obtain:

1) incursion of the potential A_μ^+ along the loop C^+ is controlled by the Higgs phase χ^+ , and integration of the equations $I_\mu^+ = 0 \Rightarrow \chi_{+, \mu} + eA_\mu^+ = 0$ gives

$$\chi_+|_{r^+} = 2m(t^+ + a\phi_K^+), \quad (24)$$

2) similarly, the out-going potential A_μ^+ , acting on the boundary r^- gives

$$\chi_-|_{r^-} = -2m(t^- - a\phi_K^-), \quad (25)$$

and therefore, the phases of the Higgs fields $(t^+ + a\phi_K^+)$ and $(t^- - a\phi_K^-)$ behave like parametrization of the left and right modes of a relativistic string, see [34, 17].

The formation of a Wilson loop around a singular ring is a characteristic feature of the string models with a tension mechanism in the form of a tube of force lines, [39].

The singular ring of the Kerr solution is a light-like line, since the light direction of the Kerr congruence k^μ touches the singular ring. Note, that the existence of "right" and "left" excitation modes is an indispensable condition for the formation of a string as a *world sheet*. A string described by only one mode, say the right one, turns into *world line* that depends on only one parameter. For this reason, a simple light-like Kerr's singular ring does not form a world sheet, and strictly speaking is not a string, but a world line. In the regularized Kerr solution, string is formed on the border of disk, where the null line of the Kerr congruence crosses the string at an angle $\theta_c = \arctan r_e/a$, and kinematic relations show (see Fig.5) that the speed at the edge of the disk is only slightly less than the speed of light, $v = c \cos \theta_c < c$. A simple null-string is replaced by a string of the 'orientifold' type considered in [28]. Two-faced structure of the KN disk is analogous to the DW-AntiDW field model of the KN source considered in [17], i.e. an oscillating solution of the type 'breather' [38].

5 The Dirac equations for an electron interacting with its own gravitational and electromagnetic field

We come to the main part of our consideration – the solutions of the Dirac equations for fermionic string which emerge on the border of the regularized KN disk interacting with the consistent gravitational and electromagnetic KN field.

We see that our analysis unavoidably leads us to the typical features of the quantum considerations: importance of the separable analysis of the Heisenberg and Schrödinger pictures, the corresponding state vectors, unitary transformation, plane waves and the scattering process.

5.1 The KN disk-like source as electron in the Heisenberg picture.

The KN gravitational field can only be consistent with one of two types of the Kerr congruence, either in-going or out-going, [46], and we choose the out-going variant, that is consistent with the retarded electromagnetic fields.

Out-going Kerr congruence

$$k_{\mu}^{+} dx^{\mu} = dr - dt - a \sin^2 \theta d\phi_K. \quad (26)$$

propagates from the both sides of the disk r^{+} and r^{-} towards direction $+\infty$. The in-going congruence

$$k_{\mu}^{-} dx^{\mu} = -dr - dt - a \sin^2 \theta d\phi_K \quad (27)$$

propagates from $-\infty$ towards the disk and focus at the both sides of the disk r^{+} and r^{-} .

The transition from out-going picture to in-going is connected with the replacement $r \rightarrow -r$, that in the coordinate transformation (7) corresponds to the replacement $\rho^{+} \rightarrow \rho^{-}$, changing in the direction of rotation $a \rightarrow -a$, and in the orientation angle ϕ_K .

The Kerr disk is located at the scattering boundary $t = 0$, which corresponds to the state vector in the Heisenberg picture. Orientation of the disk is changed under transition from $r \rightarrow -0$ to $r \rightarrow +0$, and the string on the border of KN disk acquires the properties of the orientifold string with two faces r^{+} and r^{-} .

The regularized Kerr's disk has a finite thickness $|r| = r_e$ which is determined by the physics of scattering process.

The role of the equations of motion in the Heisenberg picture, is played by the Dirac equations.

5.2 Uniqueness of the Kerr-Schild coordinates

The Dirac equation in the Kerr-Newman gravitational field was studied in many works, in particular in [40, 41, 42, 43, 44, 45]. The freedom to choice a coordinate system is an important aspect of general relativity.

It should be particularly noted the uniqueness of the description of the KN solution in the Kerr-Schild coordinate system, which allows us to linearize both electromagnetic and gravitational equations on the background of the KN solution.

The Dirac operator in the charged and curved space-time is defined by the replacement [42]

$$\gamma^{\mu}(p_{\mu} - eA_{\mu}) \rightarrow \gamma^{\mu}(p_{\mu} - eA_{\mu}) + B, \quad (28)$$

where $B = \frac{1}{2} \nabla_{\mu} \gamma^{\mu}$ can be represented in the form

$$B = \frac{1}{2\sqrt{|g|}} \partial_{\mu} (\sqrt{|g|} \gamma^{\mu}), \quad (29)$$

and is canceled because $|g| = 1$.

As a consequence, γ -matrices of the auxiliary Minkowski space can be used in the Dirac equation.

The same argument leads also to the linearization of the electromagnetic field by the use of the Kerr-Schild coordinates in the KN solution, [12].

5.3 The Dirac equations in the Weyl representation

The Dirac equations in the Weyl representation decompose into two equations

$$(p_\mu - eA_\mu^+) \sigma_{\alpha\dot{\alpha}}^\mu \xi^{\dot{\alpha}} = m\eta_\alpha, \quad (30)$$

$$(p_\mu - eA_\mu^-) \bar{\sigma}^{\mu\dot{\alpha}\alpha} \eta_\alpha = m\xi^{\dot{\alpha}}, \quad (31)$$

where $\bar{\sigma}^0 = \sigma^0$, and $\bar{\sigma}^{1,2,3} = -\sigma^{1,2,3}$, or $\bar{\sigma} = \sigma^{1,2,3}$.

Where the Wilson loop potential $eA_\mu^\pm dx^\mu = 2m(dr^\pm - dt \mp a d\phi_K)$ acts on both $r^\pm$ -boundaries (faces) of the KN disk.

In the considered earlier analogous physical model of the rotating KN disk-like source, the in-going and out-going Kerr congruences are controlled by two related phases of the Higgs field, $\chi^+ = -\chi^-$, and the momentum p_μ of the string solution must be completed by an "internal" angular momentum of two semi-strings $p^s = p^{s+} + p^{s-}$, associated with rotation of the KN disk under its evolution in time,

$$p_\mu \rightarrow p_\mu + p_\mu^s. \quad (32)$$

According (14), the spinors $\xi^{\dot{\alpha}}$ and η_α have different helicities with respect to helicity operator $\frac{1}{2}(\mathbf{k}\vec{\sigma})$, and the Weyl spinor $\xi^{\dot{\alpha}}$ is aligned with out-going direction $k^+ = (1, \mathbf{k})$, while the spinor η_α is aligned with in-going direction $k^- = (1, -\mathbf{k})$. The sign of p^s is already taken into account in the Dirac equations (30) and (31).

In the same time, the both vector fields of the Wilson lines $eA_\mu^\pm dx^\mu$ are out-going and, being emanated from the boundaries r^+ and r^- , they are related with spinors of different chirality. As a result, the electromagnetic contribution from Wilson line $eA_\mu^- dx^\mu$ should change the sign in the equation (31).

The spinor string is formed of two semi-strings of opposite helicities $\xi^{\dot{\alpha}}$ and η_α , which have the unique common point corresponding to the point where the orientation of the string changes, $a \rightarrow -a$.

Integrating the Ginzburg-Landau equations for the out-going phase of the Higgs field and $r = r^+$, we obtained $\chi^+|_{r^+} = 2m(t + a\phi)$, which for $J = ma = 1/2$ gives $p_\mu^s|_{r^+} = (2m, \partial_{\phi_K})$. The corresponding vector potential is $eA_\mu = (2m, eA_{\phi_K})$.

For the boundary $r = r^-$, we have the opposite sign of charge, which corresponds to the Wess-Zumino supersymmetric QED model, and also corresponds to integration of the BPS equations considered in [17]. The change of orientation, $a \rightarrow -a$, is accompanied by the potential of the Wilson line in the form $-eA_0 = -2m$, $-eA_{\phi_K} = 2ma$.

To simplify notations we will omit further the index K in the Kerr angular coordinate ϕ_K .

5.4 The Dirac equations in the Heisenberg picture for $t = \text{const.}$

Taking the Weyl representation for γ -matrices, we can write the Dirac equations in Heisenberg picture for $t = \text{const.}$

Setting $p_\mu = (\epsilon, \mathbf{p} + \mathbf{p}^s)$ with $\epsilon = p_0$, and $\mathbf{p} = 0$, we obtain the Dirac equations in the rest frame,

$$(p_0 - 2m)\sigma^0\xi^{\dot{\alpha}} + (p_\phi^s - 2ma\phi)\vec{\sigma}\xi^{\dot{\alpha}} = m\eta_\alpha \quad (33)$$

$$(p_0 + 2m)\sigma^0\eta_\alpha - (p_\phi^s + 2ma\phi)\vec{\sigma}\eta_\alpha = m\xi^{\dot{\alpha}}. \quad (34)$$

Where $\xi^{\dot{\alpha}}$ and η_α are normalized spinors $\bar{\xi}I\xi = \bar{\eta}I\eta = -1$, and I is unit matrix. For any $\xi^{\dot{\alpha}}, \eta_\alpha$ and $m = 0$, the first equation is identically satisfied when

$$p_0 - eA_0 = 0, \quad p_\phi^s - eA_\phi = 0, \quad (35)$$

and the second equation is identically satisfied when

$$p_0 + eA_0 = 0, \quad p_\phi^s + eA_\phi = 0. \quad (36)$$

Spinors $|u_p\rangle = \begin{pmatrix} \xi^{\dot{\alpha}} \\ \eta_\alpha \end{pmatrix}$ are normalized as $\langle \bar{u}_p | u_p \rangle = 2m$.

In the Heisenberg picture presenting the KN string at fixed time $t = \text{const.}$ we have:

1) the spinor string $\xi^{\dot{\alpha}}(\phi)$ which is a massless half-string, created by the out-going light-like directions $k^+ = (1, \mathbf{k})$ and emanated from Wilson's counter $\phi \in [0, 2\pi]$ placed at $r = r_e^+$, and also the one more massless half-string, created by the out-going light-like directions $k^+ = (1, \mathbf{k})$ emanated from Wilson's counter $\phi_K \in [-2\pi, 0]$ placed at $r = r_e^-$.

2) the spinor string $\eta_\alpha(\phi)$ representing the second massless half-string, created by the in-going light-like directions $k^- = (1, -\mathbf{k})$ towards the Wilson counter $\phi \in [-2\pi, 0]$ placed at $r = r_e^-$.

Since $m = 0$, is everywhere, for exclusion of singular point $\phi = 0$ where the oppositely directed semi-strings are joined, the both half-strings are massless and do not interact, except for the point $\phi = 0$, where $a \rightarrow -a$, and the mass term is presented as a delta-function $m = m\delta(\phi)$.

The potential energy of the semi-strings tension is determined by the Wilson loop at the boundaries $r^\pm$,

$$eA_0 = \pm 2m, \quad eA_\phi = \pm 2ma, \quad (37)$$

and the full energy of the semi-strings is cancelled, as it was shown when integrating the BPS equations for the DW-AntiDW (breather) source of the KN solution, [17].

In the Weyl representation for matrices γ^μ , the out-going and in-going fields are ordered in time, and the fields with negative frequencies do not arise.

5.5 The Schrodinger picture, plane waves and string in the Kerr-Schild coordinates

In the Schrodinger picture the plane waves and in the Kerr-Schild coordinates are described by wave function [47]

$$\psi_p = \frac{1}{\sqrt{2\epsilon}} u_p e^{-ipx}, \quad (38)$$

where $-px = -p_\mu x^\mu = p_0 x_0 - \mathbf{p}\mathbf{x}$, and $x^\mu = (t, \mathbf{x})$, $p^\mu = (p^0, \mathbf{p})$, $\epsilon = p_0 = +\sqrt{\mathbf{p}^2 + m^2}$.

The spinor ψ_p satisfies the Dirac equations

$$(\gamma^\mu \frac{\partial}{\partial x_\mu} + m)\psi_p = 0. \quad (39)$$

In the rest system, $\epsilon = m$, $\mathbf{p} = 0$, functions ψ_p and u_p are connected by unitary transformation $\mathbf{U} = e^{-iHt}$, where $H = m$ is the Hamiltonian of the system.

We consider $\mathbf{U}$ as operator acting on a state vector $|u_p\rangle = \begin{pmatrix} \xi^{\dot{\alpha}} \\ \eta_\alpha \end{pmatrix}$, in the static Heisenberg picture, while the plane wave

$$\psi_p = \mathbf{U}u_p = e^{-imt} \begin{pmatrix} \xi^{\dot{\alpha}} \\ \eta_\alpha \end{pmatrix}, \quad (40)$$

represents the state vector $|\psi_p\rangle$ in the dynamic Schrodinger picture.

In the Schrodinger picture the string turns out to be asymmetric:

the semi-string $\xi^{\dot{\alpha}}(\phi)$ covering the interval $\phi \in [0, 2\pi]$ gives $eA_0^+ = m$, $eA_\phi^+ = ma$, and semi-string $\eta_\alpha(\phi)$ covering the interval $\phi \in [-2\pi, 0]$ gives $eA_0^- = -3m$, $eA_\phi^- = -3ma$.

When $\phi = 0$, the semi-strings are joined, $\xi^{\dot{\alpha}}(0) = \pm\eta_\alpha(0)$.

Although in Heisenberg picture electron represents a string about the Compton size (or Wilson Loop) surrounding the Kerr-Newman disk, this is a static picture corresponding to shape of the string at fixed time $t = \text{const}$. Really, the electron is an ultra-relativistic object, and any external observer (including other particles) perceives the electron in Schrodinger picture, where its wave function is multiplied by unitary factor e^{iHt} .

The consistent with gravity Compton radius of an electron is not its effective scattering radius, and the people working in the high energy physics perceive it as a micro-vortex, which works like a scalpel.

6 Conclusion

This preliminary analysis shows that, following to pioneering works by Carter, Israel and López, the Dirac electron can be described as an over-rotating KN gravitating BH solution described by the Dirac equations interacting with its proper gravitational and electromagnetic field in the Kerr-Schild coordinate system.

Such description is important both from point of view of the unification gravity with quantum theory, and from point of view of the nonperturbative model of the extended electron based on the Higgs mechanism of the spontaneously broken gauge theory.

This treatment shows that the existing theories and models of elementary particles are at least incomplete, and do not take into account a number of important effects associated with the gravitational process of the frame-dragging in the spinning gravitational space-time, in particular, the strong influence of the Wilson loop in the Kerr-Newman gravitational field.

The main new lessons that this model provides are as follows:

1. The gravitational field of a particle with spin must be described by the regularized Kerr-Newman solution, which distorts space on the Compton scale, and increases the zone of influence of gravity by about 22 orders of magnitude.

2. The supersymmetric Higgs model (Landau-Ginzburg field model) allows us to resolve the known conflict between gravity and quantum theory without changes of the Einstein equations.

3. The regularized electron model takes the form of a superconducting disk-like a bag formed from the Higgs field in a supersymmetric vacuum state.
4. The Kerr-Schild coordinates are exceptional, since the Dirac and Maxwell equations are linearized in these coordinates on the gravitational KN background.
5. The Dirac equations in the KN gravitational field take a stringy form, and the corresponding electron's dynamics reproduces the dynamics of a massless relativistic string.

When this paper was finished, I found the work by Ahmed Alharthy and Vladimir V. Kassandrov [48] which overlaps with main theme of our work. The work by these authors is very interesting and based on the old works by F. Edjo Ovono, V. Kassandrov and Ya. Terletsy which develops the works and ideas of Natan Rosen.

Although it seems that these works are very far from the KN electron model, we find that the introduced by Rosen scalar potential is prototype of the considered in our work the Higgs field. The authors come to similar conclusions with our own that gravity has a strong influence on the formation of elementary particles, and although authors do not come to our categorical statement that gravity works on the Compton scale, it seems natural, because they do not take into account the influence of spin, i.e. the frame-dragging of the KN solution.

The fact that the core of an electron takes the form of a relativistic string is very important, as it can explain some of the striking properties of the electron that are known from scattering experiments, where the electron exhibits its seemingly point-like structure.

7 Acknowledgments

I would like to thank first of all our colleagues in Nuclear Safety Institute, L.V. Matveev, Yu.N. Obukhov, P.S. Kondratenko and V.N. Ponomariov, whose permanent discussions supported me in this process of searching for the truth.

References

- [1] J. Schwarz, *The Early History of String Theory and Supersymmetry*, CALT-68-2858 [arXiv:1201.0981]
- [2] N. Arkani-Hamed, Y-t. Huang and D. O'Connell, Kerr Black Holes as Elementary Particles, *J. High Energ. Phys.*, 46 (2020)[arXiv:1906.10100].
- [3] B.S. Schmekel, Quasi-Local Energy of a Charged Rotating Object Described by the Kerr-Newman Metric, *Phys. Rev. D* **100**, 124011 (2019), [arxiv:1811.03551].
- [4] A. Burinskii, Spinning Particle as KerrNewman Black Hole, *Phys.Part.Nuclei Lett.*, 17, 724 (2020).
- [5] A. Burinskii, The KerrNewman Black Hole Solution as Strong Gravity for Elementary Particles, *Gravit. Cosmol.* **26**, 87 (2020).
- [6] B. Carter, "Global Structure of the Kerr Family of Gravitational Fields", *Phys. Rev.* **174** 1559 (1968).
- [7] G.C. Debney, R.P. Kerr and A. Schild, *J. Math. Phys.*, **10** 1842 (1969).

- [8] W. Israel, “Source of the Kerr metric”, *Phys. Rev. D.* **2** 641 (1970).
- [9] V. Hamity, “An interior of the Kerr metric”, *Phys. Lett. A* **56**, 77 (1986).
- [10] C. A. López, *Extended model of the electron in general relativity*, *Phys. Rev. D* **30** 313 (1984).
- [11] O. Grön, “New derivation of Lopez’s source of the Kerr-Newman field”, *Phys. Rev. D* **32** 1588(1985).
- [12] A. Ya. Burinskii, “Microgeons with spin”, *Sov. Phys. JETP* **39** 193 (1974).
- [13] D. D. Ivanenko and A. Ya. Burinskii “Gravitational strings in the models of elementary particles” *Izv. Vuz. Fiz.* **5** 135 (1975).
- [14] M. Gürses and F. Gürsey, “Lorentz covariant treatment of the Kerr-Schild geometry”, *J. Math. Phys.* **16** 2385 (1975).
- [15] A. Burinskii, “Gravitating lepton bag model”, *JETP (Zh. Eksp. Teor. Fiz.)* **148**, 228 (2015).
- [16] A. Burinskii, “Stability of the lepton bag model based on the Kerr-Newman solution ”, *JETP (Zh. Eksp. Teor. Fiz.)* **148** (2015) 937.
- [17] A. Burinskii, “Source of the Kerr-Newman solution as a supersymmetric domain-wall bubble: 50 years of the problem”, *Phys. Lett B* **754**, 99 (2016).
- [18] A. Chodos et al. “New extended model of hadrons”, *Phys. Rev. D* **9**, 3471 (1974).
- [19] W. A. Bardeen et al., “Heavy quarks and strong binding: A field theory of hadron structure.” *Phys. Rev. D* **11**, 1094 (1974).
- [20] R.C. Giles, Semiclassical dynamics of the ”SLAC bag”, *Phys. Rev. D* **70** 1670 (1976).
- [21] K. Johnson K and C.B. Thorn, Stringlike solutions of the bag model *Phys. Rev. D* **13** 1934 (1976),
- [22] P. Fendley, S. Mathur, C. Vafa and N.P. Warner, *Phys. Lett. B* **243**, 257 (1990).
- [23] E.R.C. Abraham and P.K. Townsend, *Nucl. Phys. B* **351**, 313 (1991).
- [24] M. Cvetič, F. Quevedo and S. J. Rey, *Phys. Rev. Lett.* **67**, 1836, (1991).
- [25] G.W. Gibbons and P.K. Townsend, *Phys.Rev.Lett.* **83** 1727 (1999), [hep-th/9905196].
- [26] Xinrui Hou, A. Losev, M. Shifman, *Phys. Rev. D* **61**,085005 (2000), hep-th/9910071,
- [27] B. Chibisov and M. Shifman, *Phys. Rev. D* **56**, 7990 (1997).
- [28] A. Burinskii, Orientifold D-String in the Source of the Kerr Spinning Particle, *Phys. Rev. D* **68**, 105004 (2003).
- [29] R. Penrose, Twistor Algebra, *J. Math. Phys.* **8** 345 (1967).

- [30] A. Burinskii, Wonderful Consequences of the Kerr Theorem, *Grav.&Cosmology*, **11**, 301 (2005), hep-th/0506006.
- [31] A. Burinskii, Complex Kerr geometry and nonstationary Kerr solutions, *Phys. Rev. D* **67** 124024 (2003).
- [32] A. Burinskii Kerr-Newman electron as spinning soliton, *Int. J. of Mod.Phys. A* **29** 1450133 (2014),(Preprint arXiv:1410.2888) .
- [33] A. Burinskii, Int. J. Mod. Phys. *Source of the Kerr-Newman solution as a gravitating bag model: 50 years of the problem of the source of the Kerr solution*, Int. J. Mod. Phys. **A31**, 1641002 (2016).
- [34] A. Burinskii, “Features of spinning gravity in particle physics: supersymmetric core of the Kerr-Newman electron”, *Journal of Physics: Conf. Series* **1275**, 012031 (2019).
- [35] H. B. Nielsen and P. Olesen, “Vortex-line models for dual strings”, *Nucl. Phys. B* **61** 45 (1973).
- [36] L.D. Landau and E.M. Lifshitz, *Electrodynamics of Continuous Media (Volume 8 of A Course of Theoretical Physics)*, Pergamon Press, 1960.
- [37] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, Princeton Univ. Press, New Jersey, 1983.
- [38] P.S. Lomdahl, O.H. Olsen and M.R. Samuelson, *Phys. Rev. A* **29**, 350 (1984).
- [39] B. Zwiebach, *A First Course in String Theory*, Massachusetts Institute of Technology, 2009.
- [40] S. Einstein and R. Finkelstein, Lorentz covariance and the Kerr-Newman geometry, *Phys. Rev. D* **15** 2721 (1977).
- [41] S. Chandrasekhar, “The Mathematical Theory of Black Holes”, Oxford University Press, London, 1992.
- [42] F. Finster, Local $U(2, 2)$ Symmetry in Relativistic Quantum Mechanics, *J. Math. Phys.* **39**, 6276 (1998);
- [43] F. Finster, J. Smoller, and S.-T. Yau, Particlelike solutions of the Einstein-Dirac equations, gr-qc/9801079, *Phys. Rev. D* **59** 104020 (1999).
- [44] F. Finster, J. Smoller, and S.-T. Yau, Non-Existence of Time-Periodic Solutions of the Dirac Equation in a Reissner-Nordstrom Black Hole Background, *J. Math. Phys.* **41** 2173 (2000).
- [45] A. Burinskii, “Emergence of the Dirac Equation in the Solitonic Source of the Kerr Spinning Particle”, *Grav. Cosmol.* **21** 28 (2014), [arXiv:1404.5947].
- [46] Ch. W. Misner, K. S. Thorne, J.A. Wheeler, *Gravitation*, Part 3, San Francisco: W. H. Freeman, ISBN 978-0-7167-0344-0

- [47] A.I. Akhiezer and V. B. Berestetskii, *Quantum Electrodynamics*, Interscience Publishers, 1965.
- [48] A. Alharthy and V. Kassandrov, On a Crucial Role of Gravity in the Formation of Elementary Particles. *Universe*. **6**, 193, 2020.