

Metric Dimension in fuzzy(neutrosophic) Graphs-III

Henry Garrett

Independent Researcher

DrHenryGarrett@gmail.com

Twitter's ID: @DrHenryGarrett | ©DrHenryGarrett.wordpress.com

Abstract

In this study, the term dimension is introduced on fuzzy(neutrosophic) graphs. The classes of these specific graphs are chosen to obtain some results based on dimension. The types of crisp notions and fuzzy(neutrosophic) notions are used to make sense about the material of this study and the outline of this study uses some new notions which are crisp and fuzzy(neutrosophic).

Keywords: Fuzzy Graphs, Neutrosophic Graphs, Dimension

AMS Subject Classification: 05C17, 05C22, 05E45

1 Background

To clarify about the definitions, I use some examples and in this way, exemplifying has key role to make sense about the definitions and to introduce new ways to use on these models in the terms of new notions. The concept of complete is used to classify specific graph in every environment. To differentiate, I use an adjective or prefix in every definition. Two adjectives “fuzzy” and “neutrosophic” are used to distinguish every graph or classes of graph or any notion on them.

$G : (V, E)$ is called a **crisp graph** where V is a set of objects and E is a subset of $V \times V$ such that this subset is symmetric. A crisp graph $G : (V, E)$ is called a **fuzzy graph** $G : (\sigma, \mu)$ where $\sigma : V \rightarrow [0, 1]$ and $\mu : E \rightarrow [0, 1]$ such that $\mu(xy) \leq \sigma(x) \wedge \sigma(y)$ for all $xy \in E$. A crisp graph $G : (V, E)$ is called a **neutrosophic graph** $G : (\sigma, \mu)$ where $\sigma = (\sigma_1, \sigma_2, \sigma_3) : V \rightarrow [0, 1]$ and $\mu = (\mu_1, \mu_2, \mu_3) : E \rightarrow [0, 1]$ such that $\mu(xy) \leq \sigma(x) \wedge \sigma(y)$ for all $xy \in E$. A crisp graph $G : (V, E)$ is called a **crisp complete** where $\forall x \in V, \forall y \in V, xy \in E$. A fuzzy graph $G : (\sigma, \mu)$ is called **fuzzy complete** where it's complete and $\mu(xy) = \sigma(x) \wedge \sigma(y)$ for all $xy \in E$. A neutrosophic graph $G : (\sigma, \mu)$ is called a **neutrosophic complete** where it's complete and $\mu(xy) = \sigma(x) \wedge \sigma(y)$ for all $xy \in E$. A crisp graph $G : (V, E)$ is called a **crisp strong**. A fuzzy graph $G : (\sigma, \mu)$ is called **fuzzy strong** where $\mu(xy) = \sigma(x) \wedge \sigma(y)$ for all $xy \in E$. A neutrosophic graph $G : (\sigma, \mu)$ is called a **neutrosophic strong** where $\mu(xy) = \sigma(x) \wedge \sigma(y)$ for all $xy \in E$. A distinct sequence of vertices v_0, v_1, \dots, v_n in a crisp graph $G : (V, E)$ is called **crisp path** with length n from v_0 to v_n where $v_i v_{i+1} \in E, i = 0, 1, \dots, n - 1$. If one edge is incident to a vertex, the vertex is called **leaf**. A path v_0, v_1, \dots, v_n is called **fuzzy path** where $\mu(v_i v_{i+1}) > 0, i = 0, 1, \dots, n - 1$. A path v_0, v_1, \dots, v_n is called **neutrosophic path** where $\mu(v_i v_{i+1}) > 0, i = 0, 1, \dots, n - 1$. A path v_0, v_1, \dots, v_n with exception of v_0 and



v_n in a crisp graph $G : (V, E)$ is called **crisp cycle** with length n for $v_0 = v_n$.
 A cycle v_0, v_1, \dots, v_0 is called **fuzzy cycle** where there are two edges xy and uv such that $\mu(xy) = \mu(uv) = \bigwedge_{i=0,1,\dots,n-1} \mu(v_i v_{i+1})$. A cycle v_0, v_1, \dots, v_0 is called **neutrosophic cycle** where there are two edges xy and uv such that $\mu(xy) = \mu(uv) = \bigwedge_{i=0,1,\dots,n-1} \mu(v_i v_{i+1})$. A set is **n-set** if its cardinality is n . A **fuzzy**

Table 1. Crisp-fying, Fuzzy-fying and Neutrosophic-fying

Crisp Graphs	Fuzzy Graphs	Neutrosophic Graphs
Crisp Complete	Fuzzy Complete	Neutrosophic Complete
Crisp Strong	Fuzzy Strong	Neutrosophic Strong
Crisp Path	Fuzzy Path	Neutrosophic Path
Crisp Cycle	Fuzzy Cycle	Neutrosophic Cycle

vertex set is the subset of vertex set of (neutrosophic) fuzzy graph such that the values of these vertices are considered. A **fuzzy edge set** is the subset of edge set of (neutrosophic) fuzzy graph such that the values of these edges are considered. Let \mathcal{G} be a family of fuzzy graphs or neutrosophic graphs. This family have **fuzzy(neutrosophic) common** vertex set if all graphs have same vertex set and its values but edges set is subset of fuzzy edge set. A (neutrosophic) fuzzy graph is called **fixed-edge fuzzy(neutrosophic) graph** if all edges have same values. A (neutrosophic) fuzzy graph is called **fixed-vertex fuzzy(neutrosophic) graph** if all vertices have same values. A couple of vertices x and y is called **crisp twin** vertices if either $N(x) = N(y)$ or $N[x] = N[y]$ where $\forall x \in V, N(x) = \{y | xy \in E\}, N[x] = N(x) \cup \{x\}$. Two vertices t and t' are called **fuzzy(neutrosophic) twin** vertices if $N(t) = N(t')$ and $\mu(ts) = \mu(t's)$, for all $s \in N(t) = N(t')$. For using material look at [1-15].

Table 2. Crisp-fying, Fuzzy-fying and Neutrosophic-fying

	Crisp Vertex Set	Fuzzy Vertex Set	Neutrosophic Vertex Set
Crisp Edge Set	Fuzzy Edge Set	Neutrosophic Edge Set	
Crisp Common	Fuzzy Common	Neutrosophic Common	
Crisp Fixed-edge	Fuzzy Fixed-edge	Neutrosophic Fixed-edge	
Crisp Fixed-vertex	Fuzzy Fixed-vertex	Neutrosophic Fixed-vertex	
Crisp Twin	Fuzzy Twin	Neutrosophic Twin	

2 Definitions

We use the notion of vertex in fuzzy(neutrosophic) graphs to define new notions which state the relation amid vertices. In this way, the set of vertices are distinguished by another set of vertices.

Definition 2.1. Let $G = (V, \sigma, \mu)$ be a fuzzy(neutrosophic) graph. A vertex m **fuzzy(neutrosophic)-resolves** vertices f_1 and f_2 if $d(m, f_1) \neq d(m, f_2)$. A set M is **fuzzy(neutrosophic) resolving set** if for every couple of vertices $f_1, f_2 \in V \setminus M$, there's a vertex $m \in M$ such that m fuzzy(neutrosophic)-resolves f_1 and f_2 . $|M|$ is called **fuzzy(neutrosophic) metric number** of G and $\min_M \sum_{m \in M} \sigma(m)$ is called **fuzzy(neutrosophic) metric dimension** of G and if fuzzy(neutrosophic) metric number of set M equals fuzzy(neutrosophic) metric dimension, then M is called **fuzzy(neutrosophic) metric set** of G .

Example 2.2. Let G be a fuzzy(neutrosophic) graph as figure (1). By applying Table (3), the 1-set is explored which its cardinality is minimum. $\{f_6\}$ and $\{f_4\}$ are 1-set which has minimum cardinality amid all sets of vertices but $\{f_4\}$ isn't

fuzzy(neutrosophic) resolving set and $\{f_6\}$ is fuzzy(neutrosophic) resolving set. Thus there's no fuzzy(neutrosophic) metric set but $\{f_6\}$. f_6 fuzzy(neutrosophic)-resolves all given couple of vertices. Therefore one is fuzzy(neutrosophic) metric number of G and 0.13 is fuzzy(neutrosophic) metric dimension of G . By using Table (3), f_4 doesn't fuzzy(neutrosophic)-resolve f_2 and f_6 . f_4 doesn't fuzzy(neutrosophic)-resolve f_1 and f_5 , too.




Figure 1. Black vertex $\{f_6\}$ is only fuzzy(neutrosophic) metric set amid all sets of vertices for fuzzy(neutrosophic) graph G .

Table 3. Distances of Vertices from sets of vertices $\{f_6\}$ and $\{f_4\}$ in fuzzy(neutrosophic) Graph G .

Vertices	f_1	f_2	f_3	f_4	f_5	f_6
f_6	0.22	0.26	0.39	0.24	0.13	0
Vertices	f_1	f_2	f_3	f_4	f_5	f_6
f_4	0.11	0.24	0.37	0	0.11	0.24

Definition 2.3. Consider \mathcal{G} as a family of fuzzy(neutrosophic) graphs on a common vertex set V . A vertex m *simultaneously fuzzy(neutrosophic)-resolves* vertices f_1 and f_2 if $d_G(m, f_1) \neq d_G(m, f_2)$, for all $G \in \mathcal{G}$. A set M is *simultaneously fuzzy(neutrosophic) resolving set* if for every couple of vertices $f_1, f_2 \in V \setminus M$, there's a vertex $m \in M$ such that m resolves f_1 and f_2 , for all $G \in \mathcal{G}$. $|M|$ is called *simultaneously fuzzy(neutrosophic) metric number* of \mathcal{G} and $\min \sigma_{m \in V} \sigma(m)$ is called *simultaneously fuzzy(neutrosophic) metric dimension* of \mathcal{G} and if the simultaneously fuzzy(neutrosophic) cardinality of set M equals simultaneously fuzzy(neutrosophic) metric dimension, then M is called *simultaneously fuzzy(neutrosophic) metric set* of \mathcal{G} .

Example 2.4. Let $\mathcal{G} = \{G_1, G_2, G_3\}$ be a collection of fuzzy(neutrosophic) graphs with common fuzzy(neutrosophic) vertex set and a subset of fuzzy(neutrosophic) edge set as figure (2). By applying Table (4), the 1-set is explored which its cardinality is minimum. $\{f_2\}$ and $\{f_4\}$ are 1-set which has minimum cardinality amid all sets of vertices. $\{f_4\}$ is as fuzzy(neutrosophic) resolving set as $\{f_6\}$ is. Thus there's no fuzzy(neutrosophic) metric set but $\{f_4\}$ and $\{f_6\}$. f_6 as fuzzy(neutrosophic)-resolves all given couple of vertices as f_4 . Therefore one is fuzzy(neutrosophic) metric number of \mathcal{G} and 0.13 is fuzzy(neutrosophic) metric dimension of \mathcal{G} . By using Table (4), f_4 fuzzy(neutrosophic)-resolves all given couple of vertices.




Figure 2. Black vertex $\{f_4\}$ and the set of vertices $\{f_2\}$ are simultaneously fuzzy(neutrosophic) metric set amid all sets of vertices for family of fuzzy(neutrosophic) graphs \mathcal{G} .

Table 4. Distances of Vertices from set of vertices $\{f_6\}$ in Family of fuzzy(neutrosophic) Graphs \mathcal{G} .

Vertices of G_1	f_1	f_2	f_3	f_4
f_4	0.37	0.26	0.13	0
Vertices of G_2	f_1	f_2	f_3	f_4
f_4	0.11	0.22	0.13	0
Vertices of G_3	f_1	f_2	f_3	f_4
f_4	0.24	0.26	0.13	0

3 General Relationships

Proposition 3.1. Let G be a fuzzy(neutrosophic) path. Then every leaf is fuzzy(neutrosophic) resolving set.

Proof. Let l be a leaf. For every given a couple of vertices f_i and f_j , we get $d(l, f_i) \neq d(l, f_j)$. Since if we reassign indexes to vertices such that every vertex f_i and l have i vertices amid themselves, then $d(l, f_i) = \sum_{j \leq i} \mu(f_j f_i) \leq i$. Thus $j \leq i$ implies

$$\sum_{t \leq j} \mu(f_t f_j) + \sum_{s \leq i} \mu(f_s f_i) > \sum_{j \leq i} \mu(f_j f_i) \equiv d(l, f_j) + c = d(l, f_i) \equiv d(l, f_j) < d(l, f_i).$$

Therefore, by $d(l, f_j) < d(l, f_i)$, we get $d(l, f_i) \neq d(l, f_j)$. f_i and f_j are arbitrary so l fuzzy(neutrosophic)-resolves any given couple of vertices f_i and f_j which implies $\{l\}$ is a fuzzy(neutrosophic) resolving set. \square

Corollary 3.2. Let G be a fixed-edge fuzzy(neutrosophic) path. Then every leaf is fuzzy(neutrosophic) resolving set.

Proof. Let l be a leaf. For every given couple of vertices, f_i and f_j , we get $d(l, f_i) = c_i \neq d(l, f_j) = c_j$. It implies l fuzzy(neutrosophic)-resolves any given couple of vertices f_i and f_j which implies $\{l\}$ is a fuzzy(neutrosophic) resolving set. \square

Corollary 3.3. Let G be a fixed-vertex fuzzy(neutrosophic) path. Then every leaf is fuzzy(neutrosophic) metric set, fuzzy(neutrosophic) metric number is one and fuzzy(neutrosophic) metric dimension is c where $c = \sigma(f)$, $f \in V$.

Proof. By Proposition (3.1), every leaf is fuzzy(neutrosophic) resolving set. By $c = \sigma(f)$, $\forall f \in V$, every leaf is fuzzy(neutrosophic) metric set, fuzzy(neutrosophic) metric number is one and fuzzy(neutrosophic) metric dimension is c . \square

Proposition 3.4. Let G be a fuzzy(neutrosophic) path. Then a set including every couple of vertices is fuzzy(neutrosophic) resolving set.

Proof. Let f and f' be a couple of vertices. For every given a couple of vertices f_i and f_j , we get either $d(f, f_i) \neq d(f, f_j)$ or $d(f', f_i) \neq d(f', f_j)$. \square

Corollary 3.5. Let G be a fixed-edge fuzzy(neutrosophic) path. Then every set containing couple of vertices is fuzzy(neutrosophic) resolving set. 103
104

Proof. Consider G is a fuzzy(neutrosophic) path. Thus by Proposition (3.4), every set containing couple of vertices is fuzzy(neutrosophic) resolving set. So it holds for any given fixed-edge path fuzzy(neutrosophic) graph. 105
106
107

Proposition 3.6. Let G be a fuzzy(neutrosophic) graph. An $(k-1)$ -set from an k -set of fuzzy(neutrosophic) twin vertices is subset of a fuzzy(neutrosophic) resolving set. 108
109

Proof. If t and t' are fuzzy(neutrosophic) twin vertices, then $N(t) = N(t')$ and $\mu(ts) = \mu(t's)$, for all $s \in N(t) = N(t')$. 110
111

Corollary 3.7. Let G be a fuzzy(neutrosophic) graph. The number of fuzzy(neutrosophic) twin vertices is $n-1$. Then fuzzy(neutrosophic) metric number is $n-2$. 112
113
114

Proof. Let f and f' be two vertices. By supposition, the cardinality of set of fuzzy(neutrosophic) twin vertices is $n-2$. Thus there are two cases. If both are fuzzy(neutrosophic) twin vertices, then $N(f) = N(f')$ and $\mu(fs) = \mu(f's)$, $\forall s \in N(f)$, $\forall s' \in N(f')$. It implies $d(f, t) = d(f', t)$ for all $t \in V$. Thus suppose if not, then let f be a vertex which isn't fuzzy(neutrosophic) twin vertices with any given vertex and let f' be a vertex which is fuzzy(neutrosophic) twin vertices with any given vertex but not f . By supposition, it's possible and this is only case. Therefore, any given distinct vertex fuzzy(neutrosophic)-resolves f and f' . Then $V \setminus \{f, f'\}$ is fuzzy(neutrosophic) resolving set. It implies fuzzy(neutrosophic) metric number is $n-2$. 115
116
117
118
119
120
121
122
123

Corollary 3.8. Let G be a fuzzy(neutrosophic) graph. The number of fuzzy(neutrosophic) twin vertices is n . Then G is fixed-edge fuzzy(neutrosophic) graph. 124
125

Proof. Suppose f and f' are two given edges. By supposition, every couple of vertices are fuzzy(neutrosophic) twin vertices. It implies $\mu(f) = \mu(f')$. f and f' are arbitrary so every couple of edges have same values. It induces G is fixed-edge fuzzy(neutrosophic) graph. 126
127
128
129

Corollary 3.9. Let G be a fixed-vertex fuzzy(neutrosophic) graph. The number of fuzzy(neutrosophic) twin vertices is $n-1$. Then fuzzy(neutrosophic) metric number is $n-2$, fuzzy(neutrosophic) metric dimension is $(n-2)\sigma(m)$ where m is fuzzy(neutrosophic) twin vertex with a vertex. Every $(n-2)$ -set including fuzzy(neutrosophic) twin vertices is fuzzy(neutrosophic) metric set. 130
131
132
133
134

Proof. By Corollary (3.7), fuzzy(neutrosophic) metric number is $n-2$. By G is a fixed-vertex fuzzy(neutrosophic) graph, fuzzy metric dimension is $(n-2)\sigma(m)$ where m is fuzzy(neutrosophic) twin vertex with a vertex. One vertex doesn't belong to set of fuzzy(neutrosophic) twin vertices and a vertex from that set, are out of fuzzy metric set. It induces every $(n-2)$ -set including fuzzy(neutrosophic) twin vertices is fuzzy metric set. 135
136
137
138
139
140

Proposition 3.10. Let G be a fixed-vertex fuzzy(neutrosophic) graph such that it's fuzzy(neutrosophic) complete. Then fuzzy(neutrosophic) metric number is $n-1$, fuzzy(neutrosophic) metric dimension is $(n-1)\sigma(m)$ where m is a given vertex. Every $(n-1)$ -set is fuzzy(neutrosophic) metric set. 141
142
143
144

Proof. In fuzzy(neutrosophic) complete, every couple of vertices are twin vertices. By G is a fixed-vertex fuzzy(neutrosophic) graph and it's fuzzy(neutrosophic) complete, every couple of vertices are fuzzy(neutrosophic) twin vertices. Thus by Proposition (3.6), the result follows. 145
146
147
148

Proposition 3.11. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with common vertex set. Then simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} is $n - 1$.*

Proof. Consider $(n - 1)$ -set. Thus there's no couple of vertices to be fuzzy(neutrosophic)-resolved. Therefore, every $(n - 1)$ -set is fuzzy(neutrosophic) resolving set for any given fuzzy(neutrosophic) graph. Then it holds for any fuzzy(neutrosophic) graph. It implies it's fuzzy(neutrosophic) resolving set and its cardinality is fuzzy(neutrosophic) metric number. $(n - 1)$ -set has the cardinality $n - 1$. Then it holds for any fuzzy(neutrosophic) graph. It induces it's simultaneously fuzzy(neutrosophic) resolving set and its cardinality is simultaneously fuzzy(neutrosophic) metric number. \square

Proposition 3.12. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with common vertex set. Then simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} is greater than the maximum fuzzy(neutrosophic) metric number of $G \in \mathcal{G}$.*

Proof. Suppose t and t' are simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} and fuzzy(neutrosophic) metric number of $G \in \mathcal{G}$. Thus t is fuzzy(neutrosophic) metric number for any $G \in \mathcal{G}$. Hence, $t \geq t'$. So simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} is greater than the maximum fuzzy(neutrosophic) metric number of $G \in \mathcal{G}$. \square

Proposition 3.13. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with common vertex set. Then simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} is greater than simultaneously fuzzy(neutrosophic) metric number of $\mathcal{H} \subseteq \mathcal{G}$.*

Proof. Suppose t and t' are simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} and \mathcal{H} . Thus t is fuzzy(neutrosophic) metric number for any $G \in \mathcal{G}$. It implies t is fuzzy(neutrosophic) metric number for any $G \in \mathcal{H}$. So t is simultaneously fuzzy(neutrosophic) metric number of \mathcal{H} . By applying Definition about being the minimum number, $t \geq t'$. So simultaneously fuzzy(neutrosophic) metric number of \mathcal{G} is greater than simultaneously fuzzy(neutrosophic) metric number of $\mathcal{H} \subseteq \mathcal{G}$. \square

Theorem 3.14. *Fuzzy(neutrosophic) twin vertices aren't resolved in any given fuzzy(neutrosophic) graph.*

Proof. Let t and t' be fuzzy(neutrosophic) twin vertices. Then $N(t) = N(t')$ and $\mu(ts) = \mu(t's)$, for all $s, s' \in V$ such that $ts, t's \in E$. Thus for every given vertex $s' \in V$, $d_G(s', t) = d_G(s', t')$ where G is a given fuzzy(neutrosophic) graph. It means that t and t' aren't resolved in any given fuzzy(neutrosophic) graph. t and t' are arbitrary so fuzzy(neutrosophic) twin vertices aren't resolved in any given fuzzy(neutrosophic) graph. \square

Proposition 3.15. *Let G be a fixed-vertex fuzzy(neutrosophic) graph. If G is fuzzy(neutrosophic) complete, then every couple of vertices are fuzzy(neutrosophic) twin vertices.*

Proof. Let t and t' be couple of given vertices. By G is fuzzy(neutrosophic) complete, $N(t) = N(t')$. By G is a fixed-vertex fuzzy(neutrosophic) graph, $\mu(ts) = \mu(t's)$, for all edges $ts, t's \in E$. Thus t and t' are fuzzy(neutrosophic) twin vertices. t and t' are arbitrary couple of vertices, hence every couple of vertices are fuzzy(neutrosophic) twin vertices. \square

Theorem 3.16. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with common vertex set and $G \in \mathcal{G}$ is a fixed-vertex fuzzy(neutrosophic) graph such that it's fuzzy(neutrosophic) complete. Then simultaneously fuzzy(neutrosophic) metric number*

is $n - 1$, simultaneously fuzzy(neutrosophic) metric dimension is $(n - 1)\sigma(m)$ where m is a given vertex. Every $(n - 1)$ -set is simultaneously fuzzy(neutrosophic) metric set for \mathcal{G} . 195
196

Proof. G is fixed-vertex fuzzy(neutrosophic) graph and it's fuzzy(neutrosophic) 197
complete. So by Theorem (3.15), we get every couple of vertices in fuzzy(neutrosophic) 198
complete are fuzzy(neutrosophic) twin vertices. So every couple of vertices, by Theorem 199
(3.14), aren't resolved. 200 \square

Corollary 3.17. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with 201
fuzzy(neutrosophic) common vertex set and $G \in \mathcal{G}$ is a fuzzy(neutrosophic) 202
complete. Then simultaneously fuzzy(neutrosophic) metric number is $n - 1$, simultaneously 203
fuzzy(neutrosophic) metric dimension is $(n - 1)\sigma(m)$ where m is a given vertex. Every 204
($n - 1$)-set is simultaneously fuzzy(neutrosophic) metric set for \mathcal{G} . 205*

Proof. By fuzzy(neutrosophic) graphs with fuzzy(neutrosophic) common vertex set, G 206
is fixed-vertex fuzzy(neutrosophic) graph. It's fuzzy(neutrosophic) complete. So by 207
Theorem (3.16), we get intended result. 208 \square

Theorem 3.18. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with common vertex 209
set and for every given couple of vertices, there's a $G \in \mathcal{G}$ such that in that, they're 210
fuzzy(neutrosophic) twin vertices. Then simultaneously fuzzy(neutrosophic) metric 211
number is $n - 1$, simultaneously fuzzy(neutrosophic) metric dimension is $(n - 1)\sigma(m)$ 212
where m is a given vertex. Every $(n - 1)$ -set is simultaneously fuzzy(neutrosophic) 213
metric set for \mathcal{G} . 214*

Proof. By Proposition (3.11), simultaneously fuzzy(neutrosophic) metric number is 215
 $n - 1$. By Theorem (3.14), simultaneously fuzzy(neutrosophic) metric dimension is 216
 $(n - 1)\sigma(m)$ where m is a given vertex. Also, every $(n - 1)$ -set is simultaneously 217
fuzzy(neutrosophic) metric set for \mathcal{G} . 218 \square

Theorem 3.19. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with common vertex 219
set. If \mathcal{G} contains three fixed-vertex fuzzy(neutrosophic) stars with different center, then 220
simultaneously fuzzy(neutrosophic) metric number is $n - 2$, simultaneously 221
fuzzy(neutrosophic) metric dimension is $(n - 2)\sigma(m)$ where m is a given vertex. Every 222
($n - 2$)-set is simultaneously fuzzy(neutrosophic) metric set for \mathcal{G} . 223*

Proof. The cardinality of set of fuzzy(neutrosophic) twin vertices is $n - 1$. Thus by 224
Corollary (3.9), the result follows. 225 \square

Corollary 3.20. *Let \mathcal{G} be a family of fuzzy(neutrosophic) graphs with 226
fuzzy(neutrosophic) common vertex set. If \mathcal{G} contains three fuzzy(neutrosophic) 227
stars with different center, then simultaneously fuzzy(neutrosophic) metric number is $n - 2$, 228
simultaneously fuzzy(neutrosophic) metric dimension is $(n - 2)\sigma(m)$ where m is a given 229
vertex. Every $(n - 2)$ -set is simultaneously fuzzy(neutrosophic) metric set for \mathcal{G} . 230*

Proof. By fuzzy(neutrosophic) graphs with fuzzy(neutrosophic) common vertex set, G 231
is fixed-vertex fuzzy(neutrosophic) graph. It's fuzzy(neutrosophic) complete. So by 232
Theorem (3.19), we get intended result. 233 \square

References

1. M. Akram, and G. Shahzadi, *Operations on Single-Valued Neutrosophic Graphs*, Journal of uncertain systems 11 (1) (2017) 1-26. 235
236
2. K. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets Syst. 20 (1986) 87-96. 237

3. S. Broumi, M. Talea, A. Bakali and F. Smarandache, *Single-valued neutrosophic graphs*, Journal of New Theory 10 (2016) 86-101. 238
239
4. N. Shah, and A. Hussain, *Neutrosophic soft graphs*, Neutrosophic Set and Systems 11 (2016) 31-44. 240
241
5. Henry Garrett, *Big Sets Of Vertices*, Preprints 2021, 2021060189 (doi: 10.20944/preprints202106.0189.v1). 242
243
6. Henry Garrett, *Locating And Location Number*, Preprints 2021, 2021060206 (doi: 10.20944/preprints202106.0206.v1). 244
245
7. Henry Garrett, *Metric Dimensions Of Graphs*, Preprints 2021, 2021060392 (doi: 10.20944/preprints202106.0392.v1). 246
247
8. Henry Garrett, *New Graph Of Graph*, Preprints 2021, 2021060323 (doi: 10.20944/preprints202106.0323.v1). 248
249
9. Henry Garrett, *Numbers Based On Edges*, Preprints 2021, 2021060315 (doi: 10.20944/preprints202106.0315.v1). 250
251
10. Henry Garrett, *Matroid And Its Outlines*, Preprints 2021, 2021060146 (doi: 10.20944/preprints202106.0146.v1). 252
253
11. Henry Garrett, *Matroid And Its Relations*, Preprints 2021, 2021060080 (doi: 10.20944/preprints202106.0080.v1). 254
255
12. A. Shannon and K.T. Atanassov, *A first step to a theory of the intuitionistic fuzzy graphs*, Proceeding of FUBEST (Lakov, D., Ed.) Sofia (1994) 59-61. 256
257
13. F. Smarandache, *A Unifying field in logics neutrosophy: Neutrosophic probability, set and logic*, Rehoboth: American Research Press (1998). 258
259
14. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, *Single-valued neutrosophic sets*, Multispace and Multistructure 4 (2010) 410-413. 260
261
15. L. A. Zadeh, *Fuzzy sets*, Information and Control 8 (1965) 338-353. 262