

Article

Copper-Catalyzed Homocoupling of Boronic Acids: A focus on B-to-Cu and Cu-to-Cu Transmetalations

Aude Salamé,^{1,†} Jordan Rio ^{2,†} Ilaria Ciofini,^{3,*} Lionel Perrin,^{2,*} Laurence Grimaud^{1,*} and Pierre-Adrien Payard^{2,*}

¹ Laboratoire des Biomolécules (LBM), Département de Chimie, Ecole Normale Supérieure, PSL University, Sorbonne Université, CNRS, 75005 Paris, France.

² Université de Lyon, Université Claude Bernard Lyon I, CNRS, INSA, CPE, UMR 5246, ICBMS, 1 rue Victor Grignard, F-69622 Villeurbanne, France.

³ École nationale supérieure de chimie de Paris, Centre national de la recherche scientifique, Institute of Chemistry for Life and HealthSciences, PSL Research University, Paris, France.

* Correspondence: ilaria.ciofini@chimieparistech.psl.eu; lionel.perrin@univ-lyon1.fr; laurence.grimaud@ens.psl.eu; pierre-adrien.payard@univ-lyon1.fr


Abstract: Controlling and understanding Cu-catalyzed homocoupling reaction is crucial to prompt the development of efficient Cu-catalyzed cross-coupling reactions. The presence of a coordinating base (hydroxide, methoxide) enables the B-to-Cu(II) transmetalation from aryl boronic acid to Cu^{II}Cl₂ in methanol, through formation of a mixed Cu-(μ -OH)-B intermediates. A second B-to-Cu transmetalation to form bis-aryl Cu(II) complexes is disfavored. Instead, organocopper(II) dimers undergo a coupled transmetalation-electron-transfer (TET) allowing the formation of a bis-organocopper(III) complexes readily promoting reductive elimination. Based on this mechanism some guidelines are suggested to control the undesired formation of homocoupling product in Cu-catalyzed cross-coupling reactions.

Keywords: coupling reactions; copper; transmetalation; DFT; Reaction Mechanism; cyclic voltammetry

1. Introduction

Metal-catalyzed coupling reactions are a staple of organic synthesis as they allow the controlled extension of the carbon backbone of molecules. One of the first historically described coupling reaction was reported by Ullman at the beginning of the last century (1901) to couple two identical aryl halides (homocoupling) in the presence of a stoichiometric amount of Cu(0).[1-4] While Pd-catalyzed cross-coupling reactions are among the most used nowadays, Cu catalysis proved to be a nice complement[5] and is particularly interesting to catalyze C-C and C-heteroatom bond formations.[5-20]

In the last decade, several protocols have been disclosed describing that mild conditions and low catalyst-loadings can perform the homocoupling of boronic acids using Cu salts (**Scheme 1.A.**).[6-9] Cu-catalyzed homocoupling of boronic acids takes place at room temperature in the presence of a weak base such as K₂CO₃ (**Scheme 1.A.**) using Cu^{II}Cl₂ or Cu^ICl as catalyst precursors.[7]



Scheme 1. A. Cu-catalyzed homo- and cross-coupling of aryl-boronic acids. B. Currently proposed mechanisms of the Cu-catalyzed homocoupling of arylboronic acid ($\text{ArB}(\text{OH})_2$) in the presence of a base and current state-of-art catalytic cycle involving two successive B-to-Cu transmetalations and subsequent reductive elimination.[6-9,21] See **Scheme 3** for the updated mechanism including the conclusion of the present work.

Controlling and understanding Cu-catalyzed homocoupling reaction is of high interest i) to enable efficient formation of symmetrical biphenyl compounds under mild conditions but also ii) to disfavor this process in the case of Cu-catalyzed cross-coupling reactions. Deciphering the mechanism of homocoupling reactions is thus crucial to prompt the development of efficient Cu-catalyzed Cham-Lan [22-26] or Cu-catalyzed Suzuki-like cross-couplings (**Scheme 1.A**).[10,11] By analogy with the established mechanism for Pd-catalyzed oxidative coupling of boronic acids,[27,28] it was proposed that Cu-catalyzed homocoupling of boronic acid takes place through two successive transmetalations on Cu(II) followed by i) direct reductive elimination to yield Cu(0) or ii) one-electron oxidation into Cu(III) and subsequent reductive elimination to yield Cu(I) (**Scheme 1.B**).[6-9,21] However there is no evidence to support these mechanisms. Moreover, the role played by the base was not fully clarified. Indeed, starting from Cu(I), no base is required for the reaction to proceed[15] but the addition of a base is essential starting from Cu(II).[7] Herein we report evidences for an alternative homocoupling reaction pathway involving successive B-to-Cu and Cu-to-Cu transmetalation steps.

The mechanism of the $\text{Cu}^{\text{II}}\text{Cl}_2$ -catalyzed homocoupling reaction of arylboronic acids in methanol under inert atmosphere was investigated experimentally using a combination of NMR, cyclic voltammetry, conductometry pH-metry and kinetics monitoring. The experimental insights were complemented by the exploration of reaction pathways computed at the DFT level. In particular, the effect of the added base and the involvement of higher order Cu intermediates are clarified.


2. Results and Discussion

2.1. Cu-Catalyzed Homocoupling Reaction: Effect of Cu on Selectivity

The stoichiometric reaction of $\text{Cu}^{\text{II}}\text{Cl}_2$ (10 mM) with 2 equiv of *para*-fluorophenylboronic acid ($p\text{-F-C}_6\text{H}_4\text{B}(\text{OH})_2$) (**I**) was monitored using ^{19}F NMR in MeOH (**Scheme 2**). The NMR tube was prepared under inert atmosphere to avoid the oxidative regeneration of Cu(II) from reduced Cu complexes and thereof the subsequent oxidative coupling (**Figure S1**). In MeOH, in the absence of added base, $(p\text{-F-C}_6\text{H}_4\text{B}(\text{OH})_2$ (-112.2 ppm) is stable and no decomposition reaction was observed after 5 h at r.t (**Figure S2**).

Tetrabutylammonium hydroxide (TBAOH) was used as a model base for this reaction since it is soluble in methanol and allows to separate the effect of the base from the potential ones related to a coordinating counter-cation. In the presence of 20 equiv of $(p\text{-F-C}_6\text{H}_4\text{B}(\text{OH})_2$ and 12 equiv of TBAOH, the signal of the boronic acid reagent (**I**) decreases rapidly and several new signals appeared at -117.7, -115.1, -126.5 and -127.9 ppm. They correspond to the homocoupling product **II**, the product of the boronic acid protodemetalation **III**, the phenol **IV** and the ether **V** oxidation products of the boronic

acid, respectively (**Scheme 2** and **Figure S3**). All signals were attributed by addition of a small amount of each pure product in the NMR tube.



Scheme 2. Stoichiometric reaction of $[\text{Cu}^{\text{II}}\text{Cl}_2]$ with $(p\text{-F-C}_6\text{H}_4)\text{B}(\text{OH})_2$ (**I**) in methanol in the presence of tetrabutylammonium hydroxide (TBAOH). The reaction yields to the formation of the homocoupling product **II**, the proto-demetalation product **III** and the oxidation products **IV** and **V**.

Since the boron derivative is stable in the absence of $\text{Cu}^{\text{II}}\text{Cl}_2$, even in the presence of a base (**Figure S4**), protodemetalation is expected to be mediated by Cu. Consequently, the formation of protodemetalation product is indicative of B-to-Cu transmetalation while the formation of homocoupling product comes from transmetalation and subsequent reductive elimination.

The ratio of protodemetalation *vs* homocoupling products was shown to be highly dependent on the $\text{Cu}^{\text{II}}\text{Cl}_2$ and added base concentrations. At high $\text{Cu}^{\text{II}}\text{Cl}_2$ concentration (20 mM), no proto-demetalation product was detected while at low concentration around 20% of proto-demetalation product was observed (**Figure S5**). This indicates that the kinetics of the homocoupling reaction follows a higher reaction order with respect to $\text{Cu}(\text{II})$ than the kinetics of the proto-demetalation reaction. This suggests the possible involvement of multinuclear $\text{Cu}(\text{II})$ complexes. The concentration of TBAOH was shown to impact the final product distribution: high base concentration (0.8 equiv with respect to $\text{PhB}(\text{OH})_2$) favor the proto-demetalation (**Figure S6**).

2.2. Cyclic voltammetry monitoring of the homocoupling reaction

Electrochemical analysis methods were used to monitor the structure of ligandless Cu species under reaction conditions. Cyclic voltammetry is a powerful tool to access the oxidation state of a metal in solution,[29] and consequently to scrutinize the oxidation state of a catalyst during oxidative or reductive chemical steps.[27] $\text{Cu}^{\text{II}}\text{Cl}_2$ (2 mM) is characterized in methanol (**Figure 1.A.**) by two reduction peaks at 0.25 and -0.75 V *vs* the standard calomel electrode (SCE). They respectively correspond to the reduction to $\text{Cu}(\text{I})$ and subsequent deposition of $\text{Cu}(0)$ at the electrode. The deposited $\text{Cu}(0)$ is re-oxidized to $\text{Cu}(\text{I})$ at 0 and 0.2 V *vs* SCE and $\text{Cu}(\text{I})$ to $\text{Cu}(\text{II})$ at 0.5 V *vs* SCE. Addition of phenylboronic acid does not modify the shape of the voltammogram (**Figure S7**).

Upon addition of TBAOH (6 equiv) in the presence of $\text{PhB}(\text{OH})_2$ (10 equiv), the reduction peak related to $\text{Cu}(\text{II})$ disappears progressively (**Figure S8**) while the oxidation peak Ox3 , proportional to the concentration of $\text{Cu}(\text{I})$, is increasing (**Figure 1.B.**). This highlights the formation of $\text{Cu}(\text{I})$ and the consumption of $\text{Cu}(\text{II})$. In addition, since the reduction of $\text{Cu}(\text{I})$ can only originate from reductive elimination, the variation of the concentration of $\text{Cu}(\text{I})$ with time relates to the kinetics of the coupling reaction.




Figure 1. A. Cyclic voltammogram toward reduction potentials of a solution of $\text{Cu}^{\text{II}}\text{Cl}_2$ (2 mM) in MeOH (see **Figure S9** for the voltammogram toward oxidation). B. Cyclic voltammogram toward oxidation potentials monitoring the reduction of Cu(II) to Cu(I) by PhB(OH)₂ in the presence of TBAOH. $\text{Cu}^{\text{II}}\text{Cl}_2$ (2 mM, dark green). Addition of PhB(OH)₂ (10 mM, light green). Addition of TBAOH (5 mM, apple green) at $t = 0$. After 20 minutes (light orange). After 30 minutes (orange). After 45 minutes (red). Working electrode: glassy carbon ($\varnothing = 3$ mm); scan rate: 0.5 V s⁻¹; supporting electrolyte: $n\text{Bu}_4\text{NBF}_4$ (0.3 M); recorded at ambient temperature starting at the Open Circuit Potential (OCP).

To gain more insights into the role played by Cu and the base in the coupling mechanism, we monitored the kinetics of Cu(I) formation using a rotating disk electrode (RDE).[30] Using this set-up, the potential of the working electrode can be imposed at 0.7 V vs SCE to enable Cu(I) oxidation. In this condition, the current intensity measured is proportionated to the concentration of Cu(I) in solution. In the presence of an excess of phenylboronic acid and of base, all kinetic profiles could be fitted by a pseudo first order law in Cu(I) (**Figure S10**).




Figure 2. Variation of the pseudo first order rate constant (k , s^{-1}) of the formation of Cu(I) as estimated by fitting the measured electric current intensity at the polarized rotating electrode (+0.7 V/SCE) as a function of the initial concentration of TBAOH. For each point of the bell-shaped distribution, 4 experiments were repeated and the standard deviation was estimated (error bars).

The variation of the apparent rate constant for the formation of Cu(I) with increasing pH is depicted on **Figure 2**. At low base concentration, the rate of the coupling reaction increased linearly with the concentration of added base, while at higher concentration (6 to 8 equiv) the rate of the reaction decreased sharply. By analogy with the proposed

transmetalation mechanism this bell-shape behavior could be indicative of the formation of a μ -OH-bridged Cu-(μ -OH)-B pre-transmetalation intermediate.[31] In the absence of base, the reaction is apparently limited due to the absence of OH groups while at high concentration the predominance of Cu hydroxide and boronate species likely hamper the formation of mixed complexes between Cu and B.

2.3. Electronic Effects on the Kinetics of the reaction: Hammett Plot

To get more information related to the electronic effects over the rate determining state (RDS) of the overall coupling process, the kinetics of the Cu(I) formation was monitored for a series of *para*-substituted phenylboronic acid derivatives. The variations of the pseudo first-order rate constant were obtained at pH = 11 and 12. In both cases, electron-rich boronic acid derivatives react faster than electron poor ones (Figures 3 and S11). The negative slope of the Hammett plot for the reaction is thus indicative of a decrease of the partial charge on the aromatic ring during the RDS. This is consistent with transmetalation being the rate determining step. The non-linearity of the Hammett plot drawn can be indicative of a shift in pre-equilibrium or a change of the RDS step.[32]




Figure 3. Hammett plot for the reduction of Cu(I) (1.0 mM) by *para*-substituted phenylboronic derivatives (*p*-X-C₆H₄)B(OH)₂ (10 mM) (X = MeO, Me, F, Cl, CN and NO₂) in the presence of TBAOH (8 mM), see Figure S11 for the fit of the associated kinetics; k_H pseudo first order rate constant for X = H, k_X pseudo first order rate constant for a substituent X.

2.4. Structure of Cu(II) in methanol

To interpret the effect of the base concentration on the kinetics of the coupling reaction, we attempted to gather more information regarding the structure of Cu^{II}Cl₂ in MeOH solution as a function of the pH. The conductivity of the Cu^{II}Cl₂ electrolyte in MeOH was studied as a function of the square root of the concentration (Figure S12). The nonlinear plot obtained is typical of a weak electrolyte resulting from a limited dissociation of Cu^{II}Cl₂ into CuCl⁺ and Cl⁻ in MeOH, in contrast to the behavior reported in water.[33] This agrees with the much lower permittivity of MeOH compared to water.[34]

As PhB(OH)₂ is a weak acid in methanol solution accordingly to equation (1) ($pK_{a(\text{MeOH})}(\text{PhB(OH)}_2/\text{PhB(OH})^{3-}) = 11.2$, see Figure S13), the pH of the methanol solution can be controlled by adding sub-stoichiometric amounts of TBAOH to PhB(OH)₂. Thereof, the resulting mixture of boronate PhB(OH)₃₋ and boronic acid acts as a pH buffer.



The $[\text{PhB}(\text{OH})_3]/[\text{PhB}(\text{OH})_2]$ concentration ratios and the associated pH range that enable the coupling reaction to proceed (Figures 2 and S13) were estimated between pH 10.5 and 12.5. Under these conditions, the number of anionic ligands on Cu was determined using the Pourbaix diagram of $\text{Cu}^{\text{II}}\text{Cl}_2$ in MeOH. This diagram was constructed by measuring the open circuit potential (OCP) of i) an equimolar solution of $\text{Cu}^{\text{I}}\text{Cl}$ and $\text{Cu}^{\text{II}}\text{Cl}_2$ (2 mM / 2 mM, top plot) at a Pt working electrode and ii) a $\text{Cu}(0)$ working electrode immersed in a 2 mM solution of $\text{Cu}(\text{I})$, as a function of the pH of the solution which was varied by addition of small quantity of a 1.0 M solution of TBAOH in MeOH (Figure 4).

For pH ranging from 4 to 8, the potentials of the $\text{Cu}(\text{II})/\text{Cu}(\text{I})$ and $\text{Cu}(\text{I})/\text{Cu}(0)$ couple show very little variation, this indicates that counter anions are not coordinated to either $\text{Cu}(\text{I})$ or $\text{Cu}(\text{II})$. Above pH 8, the $\text{Cu}(\text{II})/\text{Cu}(\text{I})$ curve shows a nonlinear transition that is indicative of the partial formation of either $[\text{Cu}^{\text{II}}(\text{OR})\text{Cl}]$ and/or $[\text{Cu}^{\text{II}}(\text{OR})]^+$ complexes ($\text{pK}_{\text{a}1}(\text{CuCl}_2/\text{Cu}(\text{OH})\text{Cl}) = 9$). This transition is also indicated by the formation of green complex. Above pH 10, the slope of the later increases to a value of 113 mV / pH, this corresponds to the coordination of anionic ligands to $\text{Cu}(\text{II})$ to form $[\text{Cu}^{\text{II}}(\text{OR})_2]$ ($\text{R} = \text{H}, \text{Me}$) alkoxide complexes that partially precipitate ($\text{pK}_{\text{a}2}(\text{CuCl}_2/\text{Cu}(\text{OR})_2) = 16.3$). Of note, such a precipitate was never observed under the reaction conditions, even at higher base concentration. For pH above 11, $[\text{Cu}^{\text{I}}(\text{OH})]$ become the predominant species for $\text{Cu}(\text{I})$.




Figure 4. Pourbaix diagram of $\text{Cu}(\text{I})$ and $\text{Cu}(\text{II})$ in MeOH ($\text{Cu}(\text{I})$ and $\text{Cu}(\text{II})$ concentration 2 mM).

2.5. Computational mechanistic investigation: B-to-Cu(II) transmetalation

Based on the results described in section 2.4, $\text{Cu}(\text{II})$ is likely to exist under the form of a $[\text{Cu}^{\text{II}}(\text{OR})_2]$ complex, with $\text{OR}^- = \text{HO}^-, \text{MeO}^-, \text{PhB}(\text{OH})_3^-$ depending on the reaction conditions, for pH ranging between 9.5 and 12.5. However, at low pH, the formation of $[\text{Cu}^{\text{II}}\text{Cl}(\text{OR})]$ or $[\text{Cu}^{\text{II}}(\text{OR})]^+$ cannot be excluded. Thereof, the ability of all these species to promote B-to-Cu(II) transmetalation was evaluated by means of a computational mechanistic investigation performed at the DFT level using the B3LYP hybrid functional.[35-38] Hereafter, energies refer to Gibbs energies ($\Delta G, \text{ kcal mol}^{-1}$) estimated at 298 K and 1 atm upon full geometry optimization. Solvation by methanol has been modeled by means of

a micro-solvation approach[39] that combines an explicit description of the first solvation sphere and an implicit representation of the bulk solvation effects by a polarization continuum model.[40] The fully detailed computational approach is available in **section 4**, Materials and Methods. In the following sections, DFT-optimized structures are tagged by Arabic numeral. If the complex is cationic a + is added (e.g. 5^+). For neutral complexes, the counter anion is specified by a superscript (e.g. 4^{0H}). The presence and the number of explicit solvent molecule is indicated by a superscript n = (e.g. $4^{+,n=2}$), ranging from 0 to 2. Finally, when spin intercrossing occurs, *i.e.* in section 2.6 - the spin state (singlet s, doublet d, triplet t) of the complex is specified in superscript brackets (e.g. $9^{MeO(t),n=0}$). Transition states are indicated using the prefix **TS**-.




Figure 5. Reaction profile of the Cu-to-B transmetalation in the heterobimetallic cationic intermediate $[(S)_2\text{Cu}(\mu\text{-OH})_2\text{B}(\text{Ph})(\text{OH})]^+ (3^{+,n=2})$ ($S = \text{MeOH}$) computed at the DFT level. ΔG (ΔH) are given in kcal mol^{-1} . See the **Figures S14-16** for alternative pathways.

The transmetalation mechanism starting from variously solvated heterobimetallic adducts $[\text{Cu}^{II}(\text{PhB}(\text{OH})_3)(\text{MeOH})_n]^+ (3^{+,n})$ between phenylboronate (**1**) and $[\text{Cu}^{II}\text{MeOH}_2]^{2+}$ (**2**) was investigated. **Figure 5** depicts the most relevant Gibbs energy profile in the case of the di-solvated $[\text{Cu}^{II}(\text{PhB}(\text{OH})_3)(\text{MeOH})_2]^+ (3^{+,n=2})$ heterobimetallic adduct. The influence of solvation on the stabilities of $3^{+,n}$ ($n = 0, 1$ or 2) and on the B-to-Cu transmetalation energy barriers is discussed in the SI (**Figure S15**).[39]

Complex $3^{+,n=2}$ features two bridging $\mu\text{-OH}$ groups between Cu(II) and B atoms and the 4-membered ring atoms are nearly co-planar ($\Phi_{\text{Cu-O-B-O}} = 1.5^\circ$, **Figure 6.A.**). The formation of such a mixed Cu-($\mu\text{-OH}$)-B dimer as a key pre-transmetalation species mirrors previous reports on Pd[41-43] or Ni complexes.[44] Starting from complex $3^{+,n=2}$, one of the bridging hydroxy groups is exchanged with the phenyl group *via* a torsion along the non-exchanging B- $\mu\text{-O}$ bond. This isomerism is endergonic by 11.6 kcal mol^{-1} and gives the pre-transmetalation intermediate $4^{+,n=2}$. In $4^{+,n=2}$, the interaction between Cu and the Ph moiety has a pronounced σ character, as indicated by the open $\text{B-C}_{ipso}\text{-C}_{para}$ angle of 171° and the short Cu- C_{ipso} distance (2.22 \AA) (**Figure 6.A.**). This pre-activation allows the easy transfer of the Ph group *via* the early transition state $\text{TS-}4^{+,n=2}$. Relative to $3^{+,n=2}$, transmetalation *via* $\text{TS-}4^{+,n=2}$ requires to overcome an overall Gibbs energy barrier of 16.2 kcal mol^{-1} to yield the transmetalated complex *trans*- $[\text{Cu}^{II}(\text{Ph})(\text{MeOH})_2(\mu\text{-OH})\text{B}(\text{OH})_2]^+ (5^+)$. This is in good agreement with previous calculations related to boronate transmetalation reported on a parent Cu system.[45] From $3^{+,n=2}$ to $\text{TS-}4^{+,n=2}$ the main contribution to the energy barrier is the formation of pre-complex $4^{+,n=2}$. From cationic complex 5^+ , coordination

of a second phenylboronate molecule $[\text{PhB}(\text{OH})_3]$ (**1**) along with the release of boronic acid leads to intermediate **6**. The large exergonicity of the reaction and the overall Gibbs energy barrier of ca. 16 kcal mol⁻¹ are in line with experimental data.




Figure 6. **A.** 3D structures of optimized intermediates $4^{X,\text{n}}$ ($X = +, \text{Cl}$ or MeO , $n = 1$ or 2). H in white, C in grey, O in red, B in pink, Cu in orange and Cl in green. **B.** Structures of intermediates and transition states involved in the B-to-Cu transmetalation (see **Table 1** for the associated energies). See **Figures S15-16** for alternative solvation states.

Table 1. Transmetalation paths starting from neutral heterobimetallic Cu(II) / B complexes $[\text{Cu}^{\text{II}}(\text{MeOH})_n(\text{X})(\mu\text{-OH})_2\text{B}(\text{Ph})(\text{OH})]$ ($3^{X,\text{n}}$), $\text{X} = \text{Cl}$, MeO , $n = 0$ or 1 . ΔG (ΔH) in kcal mol⁻¹.

| X, n | 3^X | 4^X | $\text{TS-}4^X$ | 5^X |
|---------------------|-----------|-------------|-----------------|------------|
| Cl, n = 1 | 0.0 (0.0) | 9.9 (11.4) | 17.2 (18.5) | 0.0 (3.7) |
| MeO, n = 1 | 0.0 (0.0) | 11.9 (11.8) | 23.7 (22.7) | 6.5 (8.5) |
| MeO, n = 0 | 0.0 (0.0) | 9.5 (19.1) | 18.0 (16.9) | 0.6 (-3.8) |

Alternative pathways of transmetalation have been explored starting from neutral monosolvated Cu(II) complexes $[\text{Cu}^{\text{II}}(\text{MeOH})(\text{X})(\mu\text{-OH})_2\text{B}(\text{Ph})(\text{OH})]$, $\text{X} = \text{Cl}$ ($3^{\text{Cl},\text{n}=1}$), MeO ($3^{\text{MeO},\text{n}=1}$). The Gibbs energies of the computed profiles are gathered in **Table 1**. The presence of a chloride bounded to Cu merely affects the overall transmetalation barrier (17.2 kcal mol⁻¹) compared to the cationic system (16.2 kcal mol⁻¹). By comparison, methoxide is slowing down transmetalation as indicated by an overall energy barrier of 23.7 kcal mol⁻¹. This could be one possible explanation for the observed pH dependence of the homocoupling kinetics.

The lower ability to transmetalate to hydroxo- or methoxy-copper complexes is also reflected in the structure of the pre-transmetalation intermediate $4^{X,\text{n}=1}$. Both $4^{\text{OH},\text{n}=1}$ and $4^{\text{MeO},\text{n}=1}$ show very little σ -character of the Cu-C bond with bond length of 2.4 Å and flat B-Cipso-C par angle of 175°. This is related to a lower donation from Cu to Ph in agreement with the higher partial charge at the Cu in $\text{TS-}4^{\text{MeO},\text{n}=1}$ (1.326 |e⁻|) compared to $\text{TS-}4^{\text{Cl},\text{n}=1}$ (1.256 |e⁻|) (**Table S1**). Finally, the formation of transmetalation products 5^X is either endergonic ($\text{X} = \text{Cl}$) or thermoneutral ($\text{X} = \text{OH}$ and MeO).

Based on these considerations, we have assessed the influence of solvent coordination on the energy barrier of the transmetalation step (**Figures S15-S16**). Mono-solvated $3^{\text{MeO},\text{n}=1}$ is merely more stable than the solvent-free $3^{\text{MeO},\text{n}=0}$ ($\Delta G = -0.7$ kcal mol⁻¹), both are likely in equilibria in MeOH solution. The formation of the pre-complex $4^{\text{MeO},\text{n}=0}$ is facilitated compared to the mono-solvated case (+9.5 kcal mol⁻¹) and the energy of the subsequent transmetalation transition state $\text{TS-}4^{\text{MeO},\text{n}=0}$ is 4.7 kcal mol⁻¹ lower in energy than the one computed for the mono-solvated transition state $\text{TS-}4^{\text{MeO},\text{n}=1}$ ($\Delta G^\ddagger = 18.0$ vs. 22.7 kcal mol⁻¹, **Figure S15**). These specific solvent effects on transmetalation make the use of explicit solvent model mandatory for the description of such systems. Indeed, the dynamics

of solvation plays an important role in slowing down or facilitating the transmetalation as suggested for other related systems.[46-48]

The mechanism presented above accounts for the experimental data regarding the transmetalation but does not rationalize the formation of homocoupling products. One possible mechanism to explain the homocoupling involves a second B-to-Cu transmetalation of the phenyl group followed by a reductive elimination (**Scheme 1**). The energy profile of this reaction pathway were computed starting from $[\text{Cu}^{\text{II}}(\text{MeOH})(\text{PhB}(\text{OH})_3)\text{Ph}]$ (**6**) (see **Figure S17**). Activation energies related to a second transmetalation have been computed to at least 25.1 kcal mol⁻¹. Based on the experimental conditions and the kinetic measurements, this reaction sequence cannot thereof account for the formation of homocoupling products though it is commonly suggested.

2.6. Computational mechanistic investigation: Cu(II)-Cu(II) transmetalation and reductive elimination

To rationalize the formation of homocoupling products, we have considered alternative reaction pathways in which the second transmetalation takes place between two copper centers. Such a mechanism would support the higher kinetic order with respect to Cu experimentally inferred for the homocoupling reaction. In the following section, we will only discuss the reaction sequence starting from monomeric $\text{PhCu}(\text{S})\text{OMe}$ (**7^{MeO}**) that results from the first transmetalation (**Figure 7**).

The second transmetalation starts with the dimerization of **7^{MeO(d)}** in its doublet state ($S = 1/2$) to form the high-spin di-solvated Cu^{II}-Cu^{II} dimer **8^{MeO(t),n=2}** ($S = 1$), whose formation is computed exergonic by 6.3 kcal mol⁻¹. The release of two solvent molecules from **8^{MeO(t),n=2}** leads to an even more stable dimer triplet complex **8^{MeO(t),n=0}** ($\Delta G = -16.1$ kcal mol⁻¹ relative to **7^{MeO(d)}**). Torsion along the non-reactive Cu^{II}-(μ -OH) bond enables the exchange between one bridging μ -MeO group and the Ph ring to yield **9^{MeO(t),n=0}** in its triplet state. This step is endergonic by 8.1 kcal mol⁻¹. More details on the speciation of these intermediates can be found in **Tables S2-3**. For this set of dimers (**8** to **9^{MeO(t),n=0}**), the closed-shell singlet state corresponding to the mixed-valence dimer Cu(III)-Cu(I) is always computed higher in energy.

The triplet Cu(II)-Cu(II) complex **9^{MeO(t),n=0}** can then undergo Cu-Cu transmetalation to yield the mixed-valence (Cu(III)-Cu(I)) singlet complex **10^{MeO(s),n=0}**. This later is not a true minimum on the potential energy surface and spontaneously undergoes reductive elimination to yield Cu(I)-Cu(I) complex **11^{MeO(s),n=0}** and biphenyl (**Figure S18**). Since this transformation requires a spin transition, the associated singlet and triplet energy surface were scanned to estimate the activation energy of this process. The electronic energy of activation for the transmetalation and coupled electron transfer (TET) was estimated to 5 and 7 kcal mol⁻¹. In the case of the monosolvated **10^{MeO,n=1}** the barrier for the TET process is nearly doubled and the product of transmetalation complex **11^{MeO(s),n=1}** is an energy minimum (**Figures S18**). Note that direct reductive elimination at monomeric species Cu(II) is kinetically prohibited (**Figure S17**). This mechanistic picture for reductive elimination is in agreement with results recently reported by Casares on the role of electron transfer between organocupper complexes in transmetalation reactions.[49]




Figure 7. Gibbs energy profile of the Cu-to-Cu second transmetalation and subsequent reductive elimination computed at the DFT level (see the computational details). ΔG (ΔH) are given in kcal mol^{-1} .


3. Conclusion

The presence of a coordinating base (hydroxide, methoxide) enables the first B-to-Cu(II) transmetalation from aryl boronic acid to $\text{Cu}^{\text{II}}\text{Cl}_2$ in methanol, while an excess of base is detrimental. This suggests the formation of heterobimetallic Cu-(μ -OH)-B intermediates as previously suggested[45,50] and as confirmed by our DFT calculations. In contrast with previous suggestions from the literature, a second B-to-Cu transmetalation to form bis-aryl Cu(II) complexes is kinetically prevented.[6,7,9] Alternatively, the formation of organocupper(II) dimers is favored and the later can undergo a coupled transmetalation-electron-transfer (TET) allowing the formation of a mixed-valence organocupper(III)/copper(I) complex able to perform reductive elimination.

The possibility to catalyze homocoupling reactions using Cu(I) in the absence of a base suggests that transmetalation proceeds either *via* Cu(I) oxidation by O_2 and the associated generation of OH^- : [51,52]



or through a mechanism involving a Cu(I) oxo complex as proposed in the case of Pd(0) or *via* another oxidation intermediate.[27,28] This point is currently under study within our group.



Scheme 3. Updated mechanism proposed for Cu(II)-catalyzed homocoupling of arylboronic acids. See **Figure S19** for the most favored pathway for the complete homocoupling process.

Based on mechanistic insights herein reported, we can tentatively propose some guidelines to control the undesired formation of homocoupling product in the context of Cu-catalyzed cross-coupling reactions. The pH should be controlled and optimized for each catalytic system to favor transmetalation and prevent the side reactions. The use of bidentate ligands is expected to disfavor the formation of dimeric Cu(II) species responsible for homocoupling. Alternatively, using a supported Cu-catalysts or working at low catalyst loadings may also be helpful. Finally, the control of the oxygen concentration in the media is crucial to limit the formation of Cu(II) species in Cu-catalyzed non-oxidative cross-coupling reactions.

4. Materials and Methods

Structure optimizations, energy estimation and NBO 6 analyses[53] were performed with the gaussian 09 software.[54] Molecules have been optimized using an ultrafine grid as implemented in gaussian. The Becke hybrid functional B3LYP functional was used for all calculations.[35-38] The 6-31+G augmented double- ζ Pople's basis sets were used for B, C and H atoms, and complemented with an extra polarization function for O atom. Quasi relativistic Stuttgart-Dresden 10MWB electron core pseudopotentials and their associated basis sets were used for Cu and Cl atoms.[55] Since the reaction occurs in highly polar solvent, particular attention was paid to model the solvation. The most favorable solvation states of key intermediates have been verified (see the supporting information), and bulk solvation effect has been represented by continuum model for all calculations.[56] Unless specified, optimized transition states were checked by an analytical frequency calculations and their connectivity were verified by IRC following.

Supplementary Materials: Additional NMR spectra, kinetics, cyclic voltammograms and energy profiles are provided in supporting information (doc). Cartesian coordinates of all the optimized structures and their associated energies are provided (cartesians.xyz).

Author Contributions: A.S. performed the experimental part under the guidance of L.G. and P.-A.P., A.S. and J.R. contributed to the computational part under the guidance of I.C. P.-A.P. and L.P. P.-A.P. designed and lead the project. All authors contributed to writing the manuscript.

Funding: J.R. thanks the MESR for a PhD fellowship. The authors are grateful to the CNRS, Chimie-Paristech-PSL, ENS-PSL, ICBMS, Université Lyon 1, and the Region Auvergne Rhône Alpes for financial support.

Acknowledgments: The authors would like to thank Dr. J. Vantourout and Dr L. A. Perego for fruitful discussions, and P. Pialot for her contribution to the computational part. The authors are grateful to the CCIR of ICBMS, PSMN, GENCI-TGCC (Grants A0100812501 and A0120813435) for providing computational resources and technical support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Ullmann, F.; Bielecki, J. Ueber Synthesen in der Biphenylreihe. *Ber. Dtsch. Chem. Ges.* **1901**, *34*, 2174-2185, doi:<https://doi.org/10.1002/cber.190103402141>.
2. Ullmann, F. Ueber eine neue Bildungsweise von Diphenylaminderivaten. *Ber. Dtsch. Chem. Ges.* **1903**, *36*, 2382-2384, doi:<https://doi.org/10.1002/cber.190303602174>.
3. Ullmann, F.; Sponagel, P. Ueber die Phenyllirung von Phenolen. *Ber. Dtsch. Chem. Ges.* **1905**, *38*, 2211-2212, doi:<https://doi.org/10.1002/cber.190503802176>.
4. Goldberg, I. Ueber Phenyllirungen bei Gegenwart von Kupfer als Katalysator. *Ber. Dtsch. Chem. Ges.* **1906**, *39*, 1691-1692, doi:<https://doi.org/10.1002/cber.19060390298>.
5. Beletskaya, I.P.; Cheprakov, A.V. The Complementary Competitors: Palladium and Copper in C-N Cross-Coupling Reactions. *Organometallics* **2012**, *31*, 7753-7808, doi:10.1021/om300683c.
6. Kirai, N.; Yamamoto, Y. Homocoupling of Arylboronic Acids Catalyzed by 1,10-Phenanthroline-Ligated Copper Complexes in Air. *Eur. J. Org. Chem.* **2009**, *2009*, 1864-1867, doi:<https://doi.org/10.1002/ejoc.200900173>.
7. Cao, Y.-N.; Tian, X.-C.; Chen, X.-X.; Yao, Y.-X.; Gao, F.; Zhou, X.-L. Rapid Ligand-Free Base-Accelerated Copper-Catalyzed Homocoupling Reaction of Arylboronic Acids. *Synlett* **2017**, *28*, 601-606.
8. Kaboudin, B.; Haruki, T.; Yokomatsu, T. CuSO₄-Mediated Homocoupling of Arylboronic Acids under Ligand- and Base-Free Conditions in Air. *Synthesis* **2011**, *2011*, 91-96.
9. Cheng, G.; Luo, M. Homocoupling of Arylboronic Acids Catalyzed by CuCl in Air at Room Temperature. *Eur. J. Org. Chem.* **2011**, *2011*, 2519-2523, doi:<https://doi.org/10.1002/ejoc.201001729>.
10. Yang, C.-T.; Zhang, Z.-Q.; Liu, Y.-C.; Liu, L. Copper-Catalyzed Cross-Coupling Reaction of Organoboron Compounds with Primary Alkyl Halides and Pseudohalides. *Angew. Chem. Int. ed.* **2011**, *50*, 3904-3907, doi:<https://doi.org/10.1002/anie.201008007>.
11. Villalobos, J.M.; Srogl, J.; Liebeskind, L.S. A New Paradigm for Carbon-Carbon Bond Formation: Aerobic, Copper-Templated Cross-Coupling. *J. Am. Chem. Soc.* **2007**, *129*, 15734-15735, doi:10.1021/ja074931n.
12. Kang, S.-K.; Kim, J.-S.; Choi, S.-C. Copper- and Manganese-Catalyzed Cross-Coupling of Organostannanes with Organic Iodides in the Presence of Sodium Chloride. *The Journal of Organic Chemistry* **1997**, *62*, 4208-4209, doi:10.1021/jo970656j.
13. Miyake, Y.; Wu, M.; Rahman, M.J.; Kuwatani, Y.; Iyoda, M. Efficient Construction of Biaryls and Macrocyclic Cyclophanes via Electron-Transfer Oxidation of Lipshutz Cuprates. *The Journal of Organic Chemistry* **2006**, *71*, 6110-6117, doi:10.1021/jo0608063.
14. Strieter, E.R.; Bhayana, B.; Buchwald, S.L. Mechanistic Studies on the Copper-Catalyzed N-Arylation of Amides. *J. Am. Chem. Soc.* **2009**, *131*, 78-88, doi:10.1021/ja0781893.
15. Lefèvre, G.; Franc, G.; Tlili, A.; Adamo, C.; Taillefer, M.; Ciofini, I.; Jutand, A. Contribution to the Mechanism of Copper-Catalyzed C-N and C-O Bond Formation. *Organometallics* **2012**, *31*, 7694-7707, doi:10.1021/om300636f.
16. Mansour, M.; Giacovazzi, R.; Ouali, A.; Taillefer, M.; Jutand, A. Activation of aryl halides by Cu0/1,10-phenanthroline: Cu0 as precursor of CuI catalyst in cross-coupling reactions. *Chem. Commun.* **2008**, 6051-6053, doi:10.1039/B814364A.

17. Stollenz, M.; Meyer, F. Mesitylcopper – A Powerful Tool in Synthetic Chemistry. *Organometallics* **2012**, *31*, 7708-7727, doi:10.1021/om3007689.

18. Piers, E.; Yee, J.G.K.; Gladstone, P.L. CuCl-Mediated Intramolecular Oxidative Coupling of Aryl- and Alkenyltrimethylstannane Functions. *Org. Lett.* **2000**, *2*, 481-484, doi:10.1021/ol9903988.

19. Demir, A.S.; Reis, Ö.; Emrullahoglu, M. Role of Copper Species in the Oxidative Dimerization of Arylboronic Acids: Synthesis of Symmetrical Biaryls. *The Journal of Organic Chemistry* **2003**, *68*, 10130-10134, doi:10.1021/jo034680a.

20. Zhou, Y.; You, W.; Smith, K.B.; Brown, M.K. Copper-Catalyzed Cross-Coupling of Boronic Esters with Aryl Iodides and Application to the Carboboration of Alkynes and Allenes. *Angew. Chem. Int. ed.* **2014**, *53*, 3475-3479, doi:<https://doi.org/10.1002/anie.201310275>.

21. Goj, L.A.; Blue, E.D.; Delp, S.A.; Gunnoe, T.B.; Cundari, T.R.; Petersen, J.L. Single-Electron Oxidation of Monomeric Copper(I) Alkyl Complexes: Evidence for Reductive Elimination through Bimolecular Formation of Alkanes. *Organometallics* **2006**, *25*, 4097-4104, doi:10.1021/om060409i.

22. Vantourout, J.C.; Miras, H.N.; Isidro-Llobet, A.; Sproules, S.; Watson, A.J.B. Spectroscopic Studies of the Chan–Lam Amination: A Mechanism-Inspired Solution to Boronic Ester Reactivity. *J. Am. Chem. Soc.* **2017**, *139*, 4769-4779, doi:10.1021/jacs.6b12800.

23. West, M.J.; Fyfe, J.W.B.; Vantourout, J.C.; Watson, A.J.B. Mechanistic Development and Recent Applications of the Chan–Lam Amination. *Chem. Rev.* **2019**, *119*, 12491-12523, doi:10.1021/acs.chemrev.9b00491.

24. Vantourout, J.C.; Law, R.P.; Isidro-Llobet, A.; Atkinson, S.J.; Watson, A.J.B. Chan–Evans–Lam Amination of Boronic Acid Pinacol (BPin) Esters: Overcoming the Aryl Amine Problem. *The Journal of Organic Chemistry* **2016**, *81*, 3942-3950, doi:10.1021/acs.joc.6b00466.

25. Vantourout, J.C.; Li, L.; Bendito-Moll, E.; Chabarra, S.; Arrington, K.; Bode, B.E.; Isidro-Llobet, A.; Kowalski, J.A.; Nilson, M.G.; Wheelhouse, K.M.P.; et al. Mechanistic Insight Enables Practical, Scalable, Room Temperature Chan–Lam N-Arylation of N-Aryl Sulfonamides. *ACS Catal.* **2018**, *8*, 9560-9566, doi:10.1021/acscatal.8b03238.

26. Bell, N.L.; Xu, C.; Fyfe, J.W.B.; Vantourout, J.C.; Brals, J.; Chabarra, S.; Bode, B.E.; Cordes, D.B.; Slawin, A.M.Z.; McGuire, T.M.; et al. Cu(OTf)2-Mediated Cross-Coupling of Nitriles and N-Heterocycles with Arylboronic Acids to Generate Nitrilium and Pyridinium Products**. *Angew. Chem. Int. ed.* **2021**, *60*, 7935-7940, doi:<https://doi.org/10.1002/anie.202016811>.

27. Adamo, C.; Amatore, C.; Ciofini, I.; Jutand, A.; Lakmini, H. Mechanism of the Palladium-Catalyzed Homocoupling of Arylboronic Acids: Key Involvement of a Palladium Peroxo Complex. *J. Am. Chem. Soc.* **2006**, *128*, 6829-6836, doi:10.1021/ja0569959.

28. Lakmini, H.; Ciofini, I.; Jutand, A.; Amatore, C.; Adamo, C. Pd-Catalyzed Homocoupling Reaction of Arylboronic Acid: Insights from Density Functional Theory. *J. Phys. Chem. A* **2008**, *112*, 12896-12903, doi:10.1021/jp801948u.

29. Jutand, A. Contribution of electrochemistry to organometallic catalysis. *Chem. Rev.* **2008**, *108*, 2300-2347, doi:10.1021/cr068072h.

30. Opekar, F.; Beran, P. Rotating disk electrodes. *Journal of Electroanalytical Chemistry and Interfacial Electrochemistry* **1976**, *69*, 1-105, doi:[https://doi.org/10.1016/S0022-0728\(76\)80129-5](https://doi.org/10.1016/S0022-0728(76)80129-5).

31. Amatore, C.; Le Duc, G.; Jutand, A. Mechanism of Palladium-Catalyzed Suzuki–Miyaura Reactions: Multiple and Antagonistic Roles of Anionic “Bases” and Their Countercations. *Chem. Eur. J.* **2013**, *19*, 10082-10093, doi:<https://doi.org/10.1002/chem.201300177>.

32. Schreck, J.O. Nonlinear Hammett relationships. *Journal of Chemical Education* **1971**, *48*, 103, doi:10.1021/ed048p103.

33. Salmon, P.S.; Neilson, G.W.; Enderby, J.E. The structure of Cu²⁺ aqueous solutions. *Journal of Physics C: Solid State Physics* **1988**, *21*, 1335-1349, doi:10.1088/0022-3719/21/8/010.

34. Reichardt, C.; Welton, T. *Solvents and solvent effects in organic chemistry*; John Wiley & Sons: 2010.

35. Becke, A.D. Density-functional thermochemistry. III. The role of exact exchange. *The Journal of Chemical Physics* **1993**, *98*, 5648-5652, doi:10.1063/1.464913.

36. Lee, C.; Yang, W.; Parr, R.G. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Physical Review B* **1988**, *37*, 785-789, doi:10.1103/PhysRevB.37.785.

37. Vosko, S.H.; Wilk, L.; Nusair, M. Accurate spin-dependent electron liquid correlation energies for local spin density calculations: a critical analysis. *Canadian Journal of Physics* **1980**, *58*, 1200-1211, doi:10.1139/p80-159.

38. Stephens, P.J.; Devlin, F.J.; Chabalowski, C.F.; Frisch, M.J. Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields. *The Journal of Physical Chemistry* **1994**, *98*, 11623-11627, doi:10.1021/j100096a001.

39. Norjmaa, G.; Ujaque, G.; Lledós, A. Beyond Continuum Solvent Models in Computational Homogeneous Catalysis. *Top. Catal.* **2022**, *65*, 118-140, doi:10.1007/s11244-021-01520-2.

40. Tomasi, J.; Mennucci, B.; Cammi, R. Quantum Mechanical Continuum Solvation Models. *Chem. Rev.* **2005**, *105*, 2999-3094, doi:10.1021/cr9904009.

41. Payard, P.-A.; Bohn, A.; Tocqueville, D.; Jaouadi, K.; Escoude, E.; Ajig, S.; Dethoor, A.; Gontard, G.; Perego, L.A.; Vitale, M.; et al. Role of dppf Monoxide in the Transmetalation Step of the Suzuki–Miyaura Coupling Reaction. *Organometallics* **2021**, *40*, 1120-1128, doi:10.1021/acs.organomet.1c00090.

42. Thomas Andy, A.; Denmark Scott, E. Pre-transmetalation intermediates in the Suzuki–Miyaura reaction revealed: The missing link. *Science* **2016**, *352*, 329-332, doi:10.1126/science.aad6981.

43. Thomas, A.A.; Zahrt, A.F.; Delaney, C.P.; Denmark, S.E. Elucidating the Role of the Boronic Esters in the Suzuki–Miyaura Reaction: Structural, Kinetic, and Computational Investigations. *J. Am. Chem. Soc.* **2018**, *140*, 4401-4416, doi:10.1021/jacs.8b00400.

44. Payard, P.-A.; Perego, L.A.; Ciofini, I.; Grimaud, L. Taming Nickel-Catalyzed Suzuki–Miyaura Coupling: A Mechanistic Focus on Boron-to-Nickel Transmetalation. *ACS Catal.* **2018**, *8*, 4812-4823, doi:10.1021/acscatal.8b00933.

45. Bose, S.; Dutta, S.; Koley, D. Entering Chemical Space with Theoretical Underpinning of the Mechanistic Pathways in the Chan–Lam Amination. *ACS Catal.* **2022**, *12*, 1461-1474, doi:10.1021/acscatal.1c04479.

46. del Pozo, J.; Pérez-Iglesias, M.; Álvarez, R.; Lledós, A.; Casares, J.A.; Espinet, P. Speciation of ZnMe₂, ZnMeCl, and ZnCl₂ in Tetrahydrofuran (THF), and Its Influence on Mechanism Calculations of Catalytic Processes. *ACS Catal.* **2017**, *7*, 3575-3583, doi:10.1021/acscatal.6b03636.

47. Peltzer, R.M.; Eisenstein, O.; Nova, A.; Casella, M. How Solvent Dynamics Controls the Schlenk Equilibrium of Grignard Reagents: A Computational Study of CH₃MgCl in Tetrahydrofuran. *J. Phys. Chem. B* **2017**, *121*, 4226-4237, doi:10.1021/acs.jpcb.7b02716.

48. Rio, J.; Perrin, L.; Payard, P.-A. Structure – Reactivity Relationship of Organo-zinc and -zincate Reagents: Key Elements towards Molecular Understanding. *Eur. J. Org. Chem.* **2022**, doi:10.1002/ejoc.202200906.

49. Lozano-Lavilla, O.; Gómez-Orellana, P.; Lledós, A.; Casares, J.A. Transmetalation Reactions Triggered by Electron Transfer between Organocopper Complexes. *Inorg. Chem.* **2021**, *60*, 11633-11639, doi:10.1021/acs.inorgchem.1c01595.

50. King, A.E.; Ryland, B.L.; Brunold, T.C.; Stahl, S.S. Kinetic and Spectroscopic Studies of Aerobic Copper(II)-Catalyzed Methoxylation of Arylboronic Esters and Insights into Aryl Transmetalation to Copper(II). *Organometallics* **2012**, *31*, 7948-7957, doi:10.1021/om300586p.

51. Gray, R.D. Kinetics of oxidation of copper(I) by molecular oxygen in perchloric acid-acetonitrile solutions. *J. Am. Chem. Soc.* **1969**, *91*, 56-62, doi:10.1021/ja01029a012.

52. Nord, H.; Sæland, E.; Morpurgo, A.; Cagliari, A. Kinetics of the Autoxidation of Cuprous Chloride in Hydrochloric Acid Solution. *Acta Chemica Scandinavica* **1955**, *9*, 430-437, doi:10.3891/acta.chem.scand.09-0430.

53. NBO 6.0. E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales, C. R. Landis, and F. Weinhold (Theoretical Chemistry Institute, University of Wisconsin, Madison, WI, 2013; <http://nbo6.chem.wisc.edu/>

54. Gaussian 09, Revision D.01, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2013.

55. Bergner, A.; Dolg, M.; Küchle, W.; Stoll, H.; Preuß, H. Ab initio energy-adjusted pseudopotentials for elements of groups 13–17. *Mol. Phys.* **1993**, *80*, 1431–1441, doi:10.1080/00268979300103121.

56. Marenich, A.V.; Cramer, C.J.; Truhlar, D.G. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* **2009**, *113*, 6378–6396, doi:10.1021/jp810292n.