

Article

Not peer-reviewed version

Naringin Protects Against Colistin-Induced Sciatic Nerve Damage by Reducing Oxidative Stress, Apoptosis and Inflammation Damage

[Selçuk Yılmaz](#)*, Sefa Küçükler, Hasan Şimşek, Serpil Aygörmüş, Fatih Mehmet Kandemir

Posted Date: 14 September 2023

doi: 10.20944/preprints202309.0979.v1

Keywords: apoptosis; colistin; inflammation; naringin; oxidative stress; sciatic nerve



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Naringin Protects Against Colistin-Induced Sciatic Nerve Damage by Reducing Oxidative Stress, Apoptosis and Inflammation Damage

Selçuk Yılmaz ^{1,*}, Sefa Küçükler ², Hasan Şimşek ³, Serpil Aygörmez ⁴
and Fatih Mehmet Kandemir ⁵

¹ Department of Orthopaedics and Traumatology, Faculty of Medicine, Kutahya Health Sciences University, Kütahya, Türkiye. ORCID ID: 0000-0003-2508-8558; E-mail: selcuk.yilmaz@ksbu.edu.tr

² Department of Biochemistry, Faculty of Veterinary, Atatürk University, Erzurum, Türkiye. ORCID ID: 0000-0002-8222-5515; E-mail: sefa.kucukler@atauni.edu.tr

³ Department of Physiology, Faculty of Medicine, Aksaray University, Aksaray, Türkiye. ORCID ID: 0000-0001-5573-4923; E-mail: hasansimsek@aksaray.edu.tr

⁴ Department of Biochemistry, Faculty of Veterinary, Kafkas University, Kars, Türkiye. ORCID ID: 0000-0002-5675-5096; E-mail: serpilaygormez@hotmail.com

⁵ Department of Medical Biochemistry, Faculty of Medicine, Aksaray University, Aksaray, Türkiye. ORCID ID: 0000-0002-8490-2479; E-mail: fmkandemir@aksaray.edu.tr

* Correspondence: selcuk.yilmaz@ksbu.edu.tr; Tel.: +905336651976.

Abstract: Colistin is used in treatment against gram-negative bacteria. It causes toxic effects due to oxidative stress in a dose-dependent manner. Naringin is a flavonoid with strong antioxidant properties found naturally in many plants, especially citrus fruits. This study aimed to determine the protective effects of NRG in CLS-induced sciatic nerve injury. Thirty-five male Sprague rats were randomly divided into five groups: control, NRG, CLS, CLS+NRG50, and CLS+NRG100. CLS was administered orally at a 15mg/kg for seven days. NRG was administered orally at 50 or 100 mg/kg for seven days. CLS caused toxic damage in sciatic nerve tissue by increasing MDA and decreasing antioxidant capacity (SOD, CAT, GPx, and GSH); increasing NF- κ B, TNF- α , nNOS release; increasing apoptotic factor (Caspase-3); decreasing HO-1 and nCAM; and inducing 8-OHdG, MAPK and GFAP. NRG, on the other hand, reversed all these changes in sciatic nerve tissue and reduced CLS-induced oxidative stress damage, inflammatory damage and apoptotic damage. These effects were more effective at the 100mg/kg dose of NRG than at the 50mg/kg dose. In sciatic nerve tissue, CLS-induced sciatic nerve toxicity with increased oxidative stress, inflammation, and apoptotic damage. NRG showed a protective effect against CLS-induced sciatic nerve toxicity.

Keywords: apoptosis; colistin; inflammation; naringin; oxidative stress; sciatic nerve

1. Introduction

Colistin (Polymyxin E, CLS) is an antibiotic drug used in the treatment of infections caused by gram-negative bacteria such as *Escherichia coli* since the 1960s [1]. When CLS acts on bacteria, it interacts with anionic lipopolysaccharides in the bacterial membrane, translocates calcium and magnesium, and neutralizes the bacteria [2]. The use of CLS also produces toxic effects. Among these toxic effects, two important side effects such as neurotoxicity and nephrotoxicity lead to dose limitation or discontinuation of treatment [3]. In previous studies in mice, CLS (15mg/kg, 7 days, i.p.) was reported to cause damage to brain and sciatic nerve tissues [4,5]. In studies of CLS, there is increasing evidence that oxidative stress and mitochondrial damage underlie the mechanism of neurotoxicity [6].

Compounds found naturally in plants, which stand out with their antioxidant properties, protect tissues in the body against damage [7]. Flavonoids are ingredients that are taken daily in the diet and are of interest in research due to their key effects such as anti-oxidant and anti-inflammatory [8].

Naringin (NRG) is a flavanone glycoside derived from the flavanone of naringenin. It occurs naturally in many plants, especially citrus fruits [9]. NRG have gained importance in traditional medicine due to their non-addictive and non-toxic nature [10]. Studies have reported that NRG exhibits antioxidant properties by increasing the activities of antioxidants in different tissues. In addition, NRG is also protective against nitrosative stress by inhibiting the activity of nitric oxide synthase enzyme (eNOS) and removing nitric oxide-derived free radicals [11]. It has also been found to have antioxidant and anti-inflammatory effects in studies on neurodegenerative disorders [12]. The amount of NRG in grapefruit averages 17 mg/100 g of edible fruit. However, studies have reported that taking NRG in the amount of 400 mg/day increases antioxidant capacity in the body [13].

The aim of this study was to investigate the effects of CLS on oxidative stress, inflammatory and apoptotic damage pathways in rat sciatic nerve tissues and the protective properties of NRG against these effects of CLS.

2. Results

2.1. Oxidant and Antioxidant Status Findings

It was found that SOD, CAT and GPx were inhibited and antioxidant activity decreased by decreasing GSH in CLS compared to the control ($p < 0.05$). In addition, lipid peroxidation was manifested by an increase in MDA ($p < 0.05$). Antioxidant enzyme activities and GSH increased and MDA decreased in CLS+NRG50 and CLS+NRG-100 compared to CLS ($p < 0.05$). NRG showed more pronounced effects at a dose of 100mg/kg ($p < 0.05$). (Figure 1).




Figure 1. Effects of CLS and NRG administrations on oxidant and antioxidant markers in sciatic nerve tissues of rats. Values are given as mean \pm SE. Different letters indicate statistical difference: * $p < 0.05$.

2.2. Inflammation Markers findings

Inflammation-related NF- κ B, TNF- α , and nNOS in sciatic nerve tissues were analyzed by ELISA. There was an increase in NF- κ B, TNF- α , and nNOS in the CLS compared to the control ($p < 0.05$). CLS+NRG50 and CLS+NRG100 showed a decrease in all these parameters compared to CLS ($p < 0.05$). When different doses of NRG were compared, 100 mg/kg was found to be more effective ($p < 0.05$) (Figure 2).




Figure 2. Effects of CLS and NRG administrations on NF-κB, TNF-α, and nNOS levels in sciatic nerve tissues of rats. Values are given as mean ± SE. Different letters indicate statistical difference: *p<0.05.

2.3. Findings of HO-1 and 8-OHdG Levels

Compared to the control, HO-1 level decreased in the CLS, while 8-OHdG level increased (p<0.05). With NRG administration, this situation was reversed in CLS+NRG50 and CLS+NRG100 compared to CLS (p<0.05). When different doses of NRG were compared, 100 mg/kg was found to be more effective (p<0.05) (Figure 3).




Figure 3. Effects of CLS and NRG administrations on HO-1 and 8-OHdG levels in sciatic nerve tissues of rats. Values are given as mean ± SE. Different letters indicate statistical difference: *p<0.05.

2.4. Findings of MAPK14, nCAM and GFAP Levels

In CLS, nCAM was inhibited and MAPK and GFAP were increased compared to control (p<0.05). In CLS+NRG50 and CLS+NRG100, nCAM levels increased and MAPK and GFAP levels decreased compared to CLS (p<0.05). NRG showed more pronounced effects at 100mg/kg dose (p<0.05) (Figure 4).




Figure 4. Effects of CLS and NRG administrations on MAPK, nCAM and GFAP levels in sciatic nerve tissues of rats. Values are given as mean \pm SE. Different letters indicate statistical difference: * $p < 0.05$.

2.5. Apoptotic Markers Findings

In CLS group, there was an increase in Caspase-3 ($p < 0.05$). In CLS+NRG50 and CLS+NRG100 groups, the change in Caspase-3 was reversed ($p < 0.05$). NRG showed more pronounced effects at a dose of 100mg/kg (Figure 5).




Figure 5. Effects of CLS and NRG administrations on Caspase-3 levels in sciatic nerve tissues of rats.

3. Discussion

CLS is a drug used to treat drug-resistant gram-negative bacteria but has a dose-dependent toxic effect. It is hypothesized that oxidative stress plays a role in its toxic effects on tissues [14]. There are many antioxidants that reduce oxidative stress by inhibiting ROS production [15]. Flavonoids are naturally occurring compounds found in many plants such as fruits, vegetables, cereals, teas, etc. and have numerous biological effects, including neuroprotective effects [16]. NRG is an active flavanone glycoside found in plants, especially in citrus group plants, and stands out with its antioxidant properties [17]. Therefore, in the present study, the effects of NRG on CLS-induced sciatic nerve tissue toxicity in rats were investigated.

While the balance disrupted as a result of ROS increase causes oxidative stress to damage tissues, antioxidant defense systems protect tissues against this damage [18]. MDA, an indicator of lipid peroxidation, occurs as a result of peroxidation of fatty acids containing three or more double bonds and affects ion exchange across cell membranes [19,20]. Antioxidant enzymes SOD, CAT and GPx are the first line of defense against oxidative stress damage caused by ROS [21]. Another line of defense, GSH, is a tripeptide that acts as the main redox buffer in the cell in oxidative stress damage [22]. In this study, CLS increased oxidative stress by increasing MDA levels and decreasing SOD, CAT and GPx activity and GSH levels in sciatic nerve tissues. NRG administration, on the other hand, increased antioxidant activity by decreasing CLS-induced increased MDA levels and increasing antioxidant enzyme activities and GSH levels. This antioxidant activity was stronger, especially at a 100 mg/kg dose.

If oxidative stress is prolonged, inflammation also occurs [23]. During inflammation, NF- κ B is the first transcription factor to be induced in tissues and accelerates the inflammatory process by stimulating pro-inflammatory cytokines such as TNF- α [24]. The triggering of NO production due to alterations in NOS isoforms activates pathological processes such as shock and pain associated with neurotoxicity. nNOS is one of the 3 isoforms of NOS [13]. In the current study, CLS caused inflammatory damage in sciatic nerve tissues by increasing NF- κ B, TNF- α and nNOS. NRG administration, on the other hand, reduced CLS-induced inflammatory damage by reducing NF- κ B, TNF- α and nNOS. NRG could emerge as an effective therapeutic agent in sciatic nerve tissue toxicity caused by inflammation due to CLS exposure.

MAPK14 is an important member of the MAPK family that is stimulated in cells as a result of increased ROS [13]. MAPK14 induces I κ B to become phosphorylated and subsequently causes I κ B dissociation from NF- κ B [25]. nCAM is a glycoprotein released in the central nervous system and is involved in cell-cell and cell-extracellular matrix interactions [26]. GFAP is an intermediate filament protein located in glial cells (astrocytes, ependymal cells, etc.). GFAP increases with the occurrence of damage and toxic conditions in neuronal cells [27]. In the present study, CLS increased MAPK14 and GFAP levels, while decreasing nCAM levels. NRG administration reversed this situation, decreased MAPK14 and GFAP levels and provided support to the nCAM activity.

8-OHdG is one of the indicators of oxidative DNA damage caused by increased free radicals [28]. HO-1 activity has been shown to regulate key cellular activities such as inflammation and apoptosis [29]. HO-1 plays an active role in cellular defense against cellular damage and shows protective properties, especially by preventing ROS-induced damage [13]. In this study, CLS increased the level of 8-OHdG, an indicator of oxidative DNA damage, and decreased the level of protective HO-1 in sciatic nerve tissues. NRG administration reversed this situation, decreased DNA damage and provided support to the defense system.

Intracellular increase in ROS may trigger cellular aging and apoptosis by damaging macromolecules [27]. Apoptosis (programmed cell death) is an essential physiological process that destroys and removes damaged or dangerous cells in the body. In healthy tissue cells, cellular stress or injury is required to trigger apoptosis [30]. The increase of oxygen radicals negatively affects cellular activities related to intracellular signaling, such as the apoptotic pathway [31]. While caspase family members are inactive in normal healthy cells, they are activated when the death signal arrives. Caspase-3, the most important caspase, is also known as executioner caspase and is pro-apoptotic [32]. In the this study, levels of Caspase-3, which is an apoptotic factor, increased with CLS exposure in sciatic nerve tissues. When NRG was administered together with CLS, the opposite effect was observed and NRG exhibited antiapoptotic properties. Therefore, NRG may be an effective agent against apoptosis in CLS-induced sciatic nerve tissue toxicity.

4. Materials and Methods

4.1. Chemicals

CLS was purchased from Koçak Pharmaceuticals (Colimycin, 150 mg/flakon, Istanbul/Türkiye). NRG and all other chemicals (analytical purities) were obtained from Sigma Chemical Co. (St. Louis, USA).

Experimental Procedure

Thirty-five male rats (Sprague dawley, 220-250g, 10-12 weeks) obtained from Atatürk University Experimental Animal Center (Erzurum, Türkiye) were used in the experiments. Rats were housed under standard laboratory conditions (12-h light and dark cycle, ventilation, 23±2°C, standard cage). Unlimited access to food and drinking water was provided. Rats were randomly divided into 5 groups (n=7). CLS and NRG doses were determined from the literature [3,33].

- 1- Control (CNT): Saline was given orally and intraperitoneally (i.p.) for seven days
- 2- Naringin (NRG): NRG was given orally at a dose of 100 mg/kg once daily for 7 days.
- 3- Colistin (CLS): CLS was given intraperitoneally (i.p.) at a dose of 15 mg/kg once daily for 7 days.
- 4- Colistin+Naringin 50 (CLS+NRG50): For 7 days, CLS was given i.p. at a dose of 15 mg/kg followed by NRG 50 mg/kg orally.
- 5- Colistin+Naringin 100 (CLS+NRG100): For 7 days, CLS was given i.p. at a dose of 15 mg/kg followed by NRG 100 mg/kg orally.

4.2. Collection of samples

The sciatic nerve tissue was removed 24 hours after the last drug administration (day 8), separated from the surrounding tissues, washed in physiologic saline solution and stored.

4.3. Lipid peroxidation analysis

To determine the level of malondialdehyde (MDA) in sciatic nerve tissues, 532 nm absorbance was measured after reaction with thiobarbituric acid. For MDA analysis of sciatic nerve tissues, homogenization was performed according to the previous method [34]. For the analysis of MDA levels, the method in the literature was used [35].

4.4. Antioxidant analysis

Catalase (CAT), superoxide dismutase (SOD), glutathione peroxidase (GPx) activities and glutathione (GSH) levels were analyzed to analyze the antioxidant status of sciatic nerve tissue. For antioxidants markers analysis of sciatic nerve tissues, homogenization was performed according to the previous method [7]. SOD [36], CAT [37], GPx [38], and GSH [39] were determined using the literature. Lowry et al. method was used for protein analysis [40].

4.5. Analysis of Inflammatory Markers

Cytokine production in sciatic nerve tissue was determined by ELISA using commercial kits in accordance with the procedure. Supernatants obtained from homogenates prepared with phosphate buffer (pH 7.4, 0.1 M) were used in the analysis. Nuclear Factor kappa B (NF-κB), Tumor necrosis factor alpha (TNF-α), neuronal nitric oxide synthase (nNOS) levels were determined from sciatic nerve tissue using a rat ELISA kit (NF-κB and TNF-α: Sunred, China; nNOS: YL Biont, China).

4.6. Analysis of MAPK14, nCAM and GFAP Levels

Mitogen-activated protein kinase 14 (MAPK14), neural cell adhesion molecule (nCAM), and Glial fibrillary acidic protein (GFAP) in sciatic nerve tissue was determined by ELISA using commercial kits in accordance with the procedure. Supernatants obtained from homogenates prepared with phosphate buffer (pH 7.4, 0.1 M) were used in the analysis. MAPK14, nCAM, and GFAP levels were determined from sciatic nerve tissue using a rat ELISA kit (Sunred, China).

4.7. Analysis of HO-1 and 8-OHdG Levels

Heme oxygenase-1 (HO-1), and 8-hydroxy-2'-deoxyguanosine (8-OHdG) in sciatic nerve tissue was determined by ELISA using commercial kits in accordance with the procedure. Supernatants obtained from homogenates prepared with phosphate buffer (pH 7.4, 0.1 M) were used in the analysis. HO-1 and 8-OHdG levels were determined from sciatic nerve tissue using a rat ELISA kit (Sunred, China).

4.8. Analysis of Apoptotic Marker

Supernatants obtained from homogenates prepared with phosphate buffer (pH 7.4, 0.1 M) were used for apoptotic damage level analysis in sciatic nerve tissue. Caspase-3 levels were determined from sciatic nerve tissue using a rat ELISA kit (YL Biont, China).

4.9. Statistical analysis

Statistical analysis of the data obtained from sciatic nerve tissues was performed with SPSS 20.0 (IBM, NY) program. One-way ANOVA and Tukey's post hoc tests were used for comparison between groups. Data are presented as mean \pm SE. Statistical significance was accepted: $p < 0.05$.

5. Conclusions

In conclusion, CLS caused toxic effects by increasing inflammation, oxidative stress, and apoptotic damage levels in sciatic nerve tissue. On the other hand, NRG was found to reduce the toxic effect by decreasing all these damages. It can be concluded that administration of NRG, especially at the level of 100mg/kg, will be much more effective in preventing sciatic nerve tissue damage in terms of all these pathways and improving the quality of life of the patient.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was approved by Atatürk University Experimental Animal Ethics Committee (2023/08/119).

Informed Consent Statement: Not applicable.

Data Availability Statement: None.

Acknowledgments: None.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Aksu EH, Kandemir FM, Küçükler S, Mahamadu A. Improvement in colistin-induced reproductive damage, apoptosis, and autophagy in testes via reducing oxidative stress by chrysin. *J Biochem Mol Toxicol* 2018 ; 32(11) : e22201
2. Aksu EH, Kandemir FM, Küçükler S. The effects of hesperidin on colistin-induced reproductive damage, autophagy, and apoptosis by reducing oxidative stress. *Andrologia* 2021 ; 53(2) : e13900
3. Çelik H, Kandemir FM, Çağlayan C, Özdemir S, Çomaklı S, Kucukler S, et al. Neuroprotective effect of rutin against colistin-induced oxidative stress, inflammation and apoptosis in rat brain associated with the CREB/BDNF expressions. *Mol Biol Rep* 2020 ; 47(3) : 2023-2034
4. Dai C, Li J, Lin W, Li G, Sun M, Wang F, et al. Electrophysiology and ultrastructural changes in mouse sciatic nerve associated with colistin sulfate exposure. *Toxicol Mech Methods* 2012 ; 22(8) : 592-6
5. Dai C, Li J, Li J. New insight in colistin induced neurotoxicity with the mitochondrial dysfunction in mice central nervous tissues. *Exp Toxicol Pathol* 2013 ; 65(6) : 941-8
6. Dai C, Tang S, Biao X, Xiao X, Chen C, Li J. Colistin induced peripheral neurotoxicity involves mitochondrial dysfunction and oxidative stress in mice. *Mol Biol Rep* 2019 ; 46(2) : 1963-72
7. Çağlayan C, Kandemir FM, Ayna A, Gür C, Küçükler S, Darendelioğlu E. Neuroprotective effects of 18 β -glycyrrhetic acid against bisphenol A-induced neurotoxicity in rats: involvement of neuronal apoptosis, endoplasmic reticulum stress and JAK1/STAT1 signaling pathway. *Metab Brain Dis* 2022 ; 37(6) : 1931-40

8. Varışlı B, Çağlayan C, Kandemir FM, Gür C, Ayna A, Genç A, et al. Chrysin mitigates diclofenac-induced hepatotoxicity by modulating oxidative stress, apoptosis, autophagy and endoplasmic reticulum stress in rats. *Mol Biol Rep* 2023 ; 50(1) : 433-42
9. Ghanbari-Movahed M, Jackson G, Farzaei MH, Bishayee A. A Systematic Review of the Preventive and Therapeutic Effects of Naringin Against Human Malignancies. *Front Pharmacol* 2021 ; 12: 639840
10. Akintunde JK, Akintola TE, Adenuga GO, Odugbemi ZA, Adetoye RO, Akintunde OG. Naringin attenuates Bisphenol-A mediated neurotoxicity in hypertensive rats by abrogation of cerebral nucleotide depletion, oxidative damage and neuroinflammation. *Neurotoxicology* 2020 ; 81: 18-33
11. Kaur G, Prakash A. Involvement of the nitric oxide signaling in modulation of naringin against intranasal manganese and intracerebroventricular β -amyloid induced neurotoxicity in rats. *J Nutr Biochem* 2020 ; 76 : 108255
12. Garabadu D, Agrawal N. Naringin Exhibits Neuroprotection Against Rotenone-Induced Neurotoxicity in Experimental Rodents. *Neuromolecular Med.* 2020 ; 22(2) : 314-30
13. Semis HS, Kandemir FM, Çağlayan C, Kaynar O, Genc A, Arıkan SM. Protective effect of naringin against oxaliplatin-induced peripheral neuropathy in rats: A behavioral and molecular study. *J Biochem Mol Toxicol* 2022 ; 36(9) : e23121
14. Hanedan B, Ozkaraca M, Kirbas A, Kandemir FM, Aktas MS, Kilic K, et al. Investigation of the effects of hesperidin and chrysin on renal injury induced by colistin in rats. *Biomed Pharmacother* 2018 ; 108 : 1607-16
15. Akaras N, Abuc OO, Koc K, Bal T, Geyikoglu F, Atılay H, et al. (1 \rightarrow 3)- β -d-glucan enhances the toxicity induced by Bortezomib in rat testis. *J Food Biochem* 2020 ; 44(3) : e13155
16. Thayumanavan G, Jeyabalan S, Fuloria S, Sekar M, Ravi M, Selvaraj LK, et al. Silibinin and naringenin against bisphenol a-induced neurotoxicity in Zebrafish model-potential flavonoid molecules for new drug design, development, and therapy for neurological disorders. *Molecules* 2022 ; 27(8) : 2572
17. Kandemir FM, Kucukler S, Çağlayan C, Gur C, Batil AA, Gülçin İ. Therapeutic effects of silymarin and naringin on methotrexate-induced nephrotoxicity in rats: Biochemical evaluation of anti-inflammatory, antiapoptotic, and anti-autophagic properties. *J. Food Biochem* 2017; 41(5) , e12398
18. Sağlıyan A, Benzer F, Kandemir FM, Gunay C, Han MC, Ozkaraca M. Beneficial effects of oral administrations of grape seed extract on healing of surgically induced skin wounds in rabbits. *Revue Méd. Vét* 2012 ; 163(1) : 11-17
19. Ömür AD, Kandemir FM, Yıldırım BA, Akman O, Şenocak EA, Eldutar E, et al. Protective effect of dandelion (*Taraxacum officinale*) extract Against gentamicin-induced reproductive damage in male rats. *Kafkas Univ. Vet. Fak* 2016 ; 22 : 929-36
20. İleriturk M, Kandemir O, Akaras N, Simsek H, Genc A, Kandemir FM. Hesperidin has a protective effect on paclitaxel-induced testicular toxicity through regulating oxidative stress, apoptosis, inflammation and endoplasmic reticulum stress. *Reprod Toxicol* 2023 ; 118 : 108369
21. İleriturk M., Kandemir Ö. Protective Effect of Rutin on Malathion-induced Gastric Toxicity: Evaluation of Oxidative Stress, Inflammation and Apoptosis. *F.U. Vet. J. Health Sci* 2023 ; 37(2) : 139-45
22. Şimşek H, Akaras N, Gür C, Küçükler S, Kandemir FM. Beneficial effects of Chrysin on Cadmium-induced nephrotoxicity in rats: Modulating the levels of Nrf2/HO-1, RAGE/NLRP3, and Caspase-3/Bax/Bcl-2 signaling pathways. *Gene* 2023 ; 875 : 147502
23. Gur C, Akarsu SA, Akaras N, Tuncer SC, Kandemir FM. Carvacrol reduces abnormal and dead sperm counts by attenuating sodium arsenite-induced oxidative stress, inflammation, apoptosis, and autophagy in the testicular tissues of rats. *Environ Toxicol* 2023; 38(6) : 1265-76
24. Simsek H, Akaras N. Acacetin ameliorates acetylsalicylic acid-induced gastric ulcer in rats by interfering with oxidative stress, inflammation, and apoptosis. *Int J Med Biochem* 2023; 6(2) : 96-103
25. Akaras N, Kandemir FM, Şimşek H, Gür C, Aygörmez S. Antioxidant, Anti-inflammatory, and Antiapoptotic Effects of Rutin in Spleen Toxicity Induced by Sodium Valproate in Rats. *TJNS* 2023 ; 12(2) : 138-44
26. Yardım A, Kucukler S, Özdemir S, Çomaklı S, Çağlayan C, Kandemir FM, et al. Silymarin alleviates docetaxel-induced central and peripheral neurotoxicity by reducing oxidative stress, inflammation and apoptosis in rats. *Gene* 2021 ; 769 : 145239
27. Celik H, Kucukler S, Ozdemir S, Comakli S, Gur C, Kandemir FM, et al. Lycopene protects against central and peripheral neuropathy by inhibiting oxaliplatin-induced ATF-6 pathway, apoptosis, inflammation and oxidative stress in brains and sciatic tissues of rats. *Neurotoxicology* 2020 ; 80 : 29-40
28. Yardım A, Kandemir FM, Çomaklı S, Özdemir S, Çağlayan C, Kucukler S, et al. Protective Effects of Curcumin Against Paclitaxel-Induced Spinal Cord and Sciatic Nerve Injuries in Rats. *Neurochem Res* 2021 ; 46(2) : 379-95
29. Gur C, Kandemir FM, Çağlayan C, Satıcı E. Chemopreventive effects of hesperidin against paclitaxel-induced hepatotoxicity and nephrotoxicity via amendment of Nrf2/HO-1 and caspase-3/Bax/Bcl-2 signaling pathways. *Chem Biol Interact* 2022 ; 365 : 110073

30. Akcılar R, Akcılar A, Koçak C, Koçak FE, Bayat Z, Şimşek H, et al. Effects of Ukrain on intestinal apoptosis caused by ischemia-reperfusion injury in rats. *Int J Clin Exp Med* 2015 ; 8(12) : 22158-66
31. Akaras N, Bal T, Atilay H, Selli J, Halici MB. Protective effects of agomelatine on testicular damage caused by bortezomib. *Biotech Histochem* 2017 ; 92(8) : 552-59
32. Şimşek H, Demiryürek Ş, Demir T, Atabay HD, Çeribası AO, Bayraktar R, et al. Assessment of expressions of Bcl-XL, b-FGF, Bmp-2, Caspase-3, PDGFR- α , Smad1 and TGF- β 1 genes in a rat model of lung ischemia/reperfusion. *Iran J Basic Med Sci* 2016 ; 19(2) : 209-14
33. Temel Y, Çağlayan C, Ahmed BM, Kandemir FM, Çiftci M. The effects of chrysin and naringin on cyclophosphamide-induced erythrocyte damage in rats: biochemical evaluation of some enzyme activities in vivo and in vitro. *Naunyn Schmiedebergs Arch Pharmacol* 2021 ; 394(4) : 645-54
34. Semis HS, Kandemir FM, Kaynar O, Dogan T, Arikan SM. The protective effects of hesperidin against paclitaxel-induced peripheral neuropathy in rats. *Life Sci* 2021; 287 : 120104
35. Placer ZA, Cushman LL, Johnson BC. Estimation of product of lipid peroxidation (malonyl dialdehyde) in biochemical systems. *Anal Biochem* 1966 ; 16(2) : 359-64
36. Sun Y, Oberley LW, Li Y. A simple method for clinical assay of superoxide dismutase. *Clin Chem* 1988 ; 34(3) : 497-500
37. Aebi H. Catalase in vitro. *Methods Enzymol* 1984 ; 105 : 121-26
38. Lawrence RA, Burk RF. Glutathione peroxidase activity in selenium-deficient rat liver. *Biochem Biophys Res Commun* 1976 ; 71(4) : 952-58
39. Sedlak J, Lindsay RH. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem.* 1968; 25(1):192-205.
40. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951; 193(1):265-75.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.