

Article

Not peer-reviewed version

A Variational Approach to Excited Fermions on Kinks

[Herbert Weigel](#)^{*} and Danial Saadatmand

Posted Date: 29 November 2023

doi: 10.20944/preprints202311.1824.v1

Keywords: Kinks; fermions; Dirac sea; vacuum polarization energies; variational methods



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

A Variational Approach to Excited Fermions on Kinks

Herbert Weigel ^{1,*}  and Danial Saadatmand ^{1,2} 

¹ Institute for Theoretical Physics, Physics Department, Stellenbosch University, Matieland 7602, South Africa; weigel@sun.ac.za

² National Institute for Theoretical and Computational Sciences (NITheCS) South Africa; saadatmand@sun.ac.za

* Correspondence: weigel@sun.ac.za

Abstract: We study the back-reaction of fermion fields on the kink solution in one space and one time dimension. We employ a variational procedure to determine an upper limit for the minimum of the total energy. This energy has three contributions: the classical kink energy, the energy of valence fermions and the fermion vacuum polarization energy. The latter arises from the interaction of the kink with the Dirac sea and is required for consistency of the semi-classical expansion for the fermions. Earlier studies only considered the valence part and observed a substantial back-reaction. This was reflected by a sizable distortion of the kink profile. We find that this distortion is strongly mitigated when the Dirac sea is properly accounted for. As a result the back-reaction merely produces a slight squeeze or stretch of the kink profile.

Keywords: Kinks, fermions, Dirac sea, vacuum polarization energies, variational methods

1. Introduction and Motivation

Theories in one time and one space dimension ($D = 1 + 1$) of scalar fields with degenerate vacua often lead to static solutions that connect different vacua at the two spatial infinities. We call them solitons (or solitary waves) when the corresponding energy density is localized. Solitons in $D = 1 + 1$ models serve as role models for higher dimensional systems but can also be embedded therein. Thus they have numerous applications on all scales ranging from cosmic strings [1] in the electro-weak theory via hadron [2], nuclear [3] and condensed matter physics [4,5] even to cosmology [6]. A comprehensive summary of applications of solitons in $D = 1 + 1$ has been compiled in the introduction of Ref. [7].

Kink-fermion systems always have a fermion zero mode. Numerous additional fermion bound states emerge when the Yukawa is sufficiently large [8,9]. Though not kinematically stable against decays into free fermions, it is possible to construct local minima (or saddle points) of the static energy functional in which a (valence) fermion resides in an excited bound state. Not too long ago, soliton configurations were constructed that accounted for the back-reaction from such a higher energy valence level [10,11]. We reconsidered those studies and found that the energy of the fermion vacuum, *i.e.* the Dirac sea, which is of the same order as that of the valence fermion in the semi-classical expansion, contributes largely to the total energy [12]. Based on that study we now attempt to identify soliton like minima of the energy functional with an excited valence fermion when all contributions to the fermion energy are included that are leading order in the semi-classical expansion. This extension is also important because we know from the bosonized NJL model that, while coupling to a valence quark strongly binds the chiral soliton, the Dirac sea has a destabilizing effect in the sense that the energy of the polarized sea significantly increases the total energy [13]. However, that model does not fall into the class of renormalizable theories that we explore here.

Some time ago self-consistent configurations from the binding to a single fermion bound state omitting the Dirac sea were also considered in the context of chiral quark [14] and electro-weak [15] models; though only for the lowest energy bound state. For models with fermion couplings the necessity of including the Dirac sea was later pointed out for one [16] and three space dimensions [17]. However, cases in which the coupling goes to an excited level are still interesting and may cause major deformations of the kink even when the Dirac sea is included. This is a major objective of the present

study. We note that indeed excited fermion levels play their roles in physics. For example, in the MIT bag model [18] the Roper (1440) resonance is associated with a radially excited quark level [19,20].

This short report is organized as follows. In Sect. 2 we introduce the model and discuss the classical energy of the kink as well the coupling between the kink and a single fermion mode. In Sect. 3 we examine the fermion contribution to the energy with emphasis on the Dirac sea contribution in the context of the semi-classical expansion. Our numerical results are contained and discussed in Sect. 4 while we briefly conclude in Sect. 5.

2. The Model

In $D = 1 + 1$ the scalar field Φ is dimensionless and the fermion spinors Ψ have canonical energy dimension $\frac{1}{2}$. We take the Yukawa coupling constant g to be dimensionless and write the Lagrangian as

$$\mathcal{L} = \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \frac{\lambda}{4} \left(\Phi^2 - \frac{M^2}{2\lambda} \right)^2 + i \bar{\Psi} \not{\partial} \Psi - g \sqrt{\frac{\lambda}{2}} \bar{\Psi} \Phi \Psi. \quad (1)$$

The scalar (or Higgs) coupling constant λ has dimension energy squared and $m = \frac{gM}{2}$ is the fermion mass which arises from spontaneous symmetry breaking that generates the vacuum expectation value $\langle \Phi \rangle = \frac{\pm M}{\sqrt{2\lambda}}$. Scalar fluctuations about $\langle \Phi \rangle$ have mass M .

In order to find the most generic, *i.e.* parameter independent, formulation and also for numerical practicality it is appropriate to introduce dimensionless quantities:

$$\Phi(t, x) = \frac{M}{\sqrt{2\lambda}} \phi(\tau, \xi) \quad \text{and} \quad \Psi(t, x) = \sqrt{\frac{M}{2}} \psi(\tau, \xi), \quad \text{where} \quad (\tau, \xi) = \frac{M}{2} (t, x). \quad (2)$$

We have introduced the factor $\frac{1}{2}$ in the dimensionless coordinate so that the kink, $\phi_K(\xi) = \tanh(\xi)$, is the soliton solution to the field equation for ϕ when $g = 0$. Choosing $\gamma^0 = \sigma_1$ and $\gamma^1 = i\sigma_3$ as the representation for the Dirac matrices, the stationary Dirac equation then is an eigenvalue equation for the dimensionless energy¹ $\epsilon = \frac{2E}{M}$

$$\epsilon \psi(\xi) = h \psi(\xi) \quad \text{with} \quad h = -i\sigma_2 \partial_\xi + g \phi(\xi) \sigma_1. \quad (3)$$

The normalization condition is $\int d\xi \psi^\dagger(\xi) \psi(\xi) = 1$. Fortunately, in our approach we will not need to construct these spinors; but only the eigenvalues ϵ which can be discrete and, above threshold g , continuous. We try to keep the notation simple and write the labels for these energies (and the corresponding eigen-spinors) only when necessary. In terms of upper (u) and lower (v) spinor components in $\psi = \begin{pmatrix} u \\ v \end{pmatrix}$, Eq. (3) reads (primes denote derivatives with respect to ξ)

$$u' = \epsilon v - g \phi u \quad \text{and} \quad v' = -\epsilon u + g \phi v. \quad (4)$$

The fermion quantum effects on $\phi(\xi)$ are non-local and when they are implemented the field equation $\phi(\xi)$ is not a (simple) differential equation. However, for a given profile we have the classical energy

$$E_{\text{cl}} = \frac{M^3}{4\lambda} \epsilon_{\text{cl}} \quad \text{with} \quad \epsilon_{\text{cl}} = \int_0^\infty d\xi \left[\phi'^2(\xi) + \left(\phi^2(\xi) - 1 \right)^2 \right]. \quad (5)$$

Any legitimate soliton profile connects the vacuum expectations values $\langle \phi \rangle = \pm 1$ between negative and positive spatial infinity and is anti-symmetric under spatial reflection. Then the solutions to Eq. (4) separate into two channels: the one with positive intrinsic parity has even u and odd v , while the

¹ In physical variables the separation leading to the stationary equation is $\Psi(t, x) = e^{-iEt} \Psi(x)$.

negative intrinsic parity channel has it the other way round. As in Ref. [10] we refer to these channels as A - and B -type solutions or configurations. Additional integer labels on A and B count the number of zero-crossings of u on the half-line $x \geq 0$, including the one at $x = 0$ for the B -type solution. In this notation the zero mode of the kink is an A_0 solution.

3. The Fermion Energy Functional

Here we detail the treatment of the fermion contribution to the energy starting from the effective action for fermions interacting with a static background potential. In our case that potential is generated by the soliton $\phi(\xi)$.

3.1. Formal Considerations

Fermions that interact with a static background are subject to a Dirac equation of the form $(i\partial_t - h)\Psi = 0$ and their effective action formally, *i.e.* ignoring the important regularization, is [13,21]

$$\mathcal{A} = \frac{T}{2} \sum_{\nu} |E_{\nu}| - i \ln \sum_{\{\eta_{\nu}\}} \exp \left[-iT \sum_{\nu} \eta_{\nu} |E_{\nu}| \right], \quad (6)$$

where the E_{ν} are again the eigenvalues of the Dirac Hamiltonian h and T is an arbitrarily large time interval discretizing the eigenvalues of $i\partial_t$. The outer sum in the second term runs over all possible sets of occupation numbers $\eta_{\nu} = \pm 1$ for the single fermion levels. When singling out a particular set of occupation numbers, say $\{\bar{\eta}_{\nu}\}$ this outer sum is omitted and we extract the (unregularized) fermion energy functional

$$E_F(\bar{\eta}_{\nu}) = -\frac{1}{2} \sum_{\nu} |E_{\nu}| + \bar{\eta}_{\nu} |E_{\nu}|. \quad (7)$$

The two contributions on the right hand side are the vacuum and valence energies, respectively. Upon comparison with the free case without a static background the vacuum energy turns into the vacuum polarization energy that we will regularize and renormalize utilizing spectral methods [22] in Subsect. 3.3. We furthermore note that the conserved fermion number is

$$N_F(\bar{\eta}_{\nu}) = \sum_{\nu} \left(\bar{\eta}_{\nu} - \frac{1}{2} \right) \text{sign}(E_{\nu}). \quad (8)$$

For a prescribed fermion number therefore the global energy minimum corresponds to a specific set of occupation numbers. Unless the vacuum energy of a self-consistent configuration varies strongly with the selection of occupation numbers, this specific set fills levels starting from the most strongly bound one. Recently, however local energy minima (or at least saddle points) have been discussed for which the non-zero occupation numbers concern the first, second or third excited single particle levels [10–12]. This scenario is the central objective of our project.

These formal considerations clearly show that fermion valance and vacuum energies must be treated on equal level. Yet, Eq. (7) only contains the fermion one-loop contribution to the energy. The loop-counting parameter is the inverse of the ratio of the scales for the classical energy and the fermion energy eigenvalues, *i.e.* $\frac{M^2}{\lambda}$. Hence for this approach to produce reliable results we focus on $M^2 > \lambda$. Furthermore we will omit quantum corrections originating from the scalar field. That is, we assume that the fermion quantum corrections dominate the scalar ones. This is reliable when the fermion energy eigenvalues are strongly skewed by the background potential, which happens to be the case when the Yukawa coupling is large: $g \gg 1$.

3.2. Bound States

The Dirac equation has discrete, normalizable solutions with $|\epsilon| < g$. The stronger the Yukawa coupling, the more of these solutions exist [8,9]. When $\phi(\xi)$ is odd under spatial reflection they have

definite parity. We already mentioned that in the notation of Refs. [10,11] these parity channels are called A and B -type. An additional, integer label on the capital letters counts the bound states in a given parity channel. Then A_0 is the configuration with the most strongly bound fermion mode occupied. This mode has energy eigenvalue zero and is always present, no matter what the Yukawa coupling is. The second most bound fermion mode has opposite parity and its explicit occupation defines the B_1 configuration, followed by a bound state with the same parity as the zero mode. Its occupation defines the A_1 configuration, etc. [10–12]. We will maintain that notation for the particular choices of $\bar{\eta}_\nu$.

In the numerical simulation we find the energy eigenvalues by integrating the differential equations (4) from the origin with initial conditions suitable for either the A or B configurations to some intermediate coordinate, ζ_m . Furthermore we integrate from a large distance, say $\zeta_{\max} \gg \zeta_m$ with initial conditions for exponentially decaying spinor components u and v to ζ_m as well. Only for certain parameters ϵ in Eq. (4) it is possible to match these solutions at ζ_m . These ϵ values are the searched-for energy eigenvalues. We verify that the resulting bound state energies are not sensitive to the choice of ζ_m (ζ_{\max}) when it is taken not too large (small).

3.3. Fermion VPE

In this section we renormalize the divergent vacuum part of the fermion energy in Eq. (7) and express it in terms of scattering data. First we measure this energy relative to the $\Phi \equiv \langle \Phi \rangle$ case so that it turns into the vacuum polarization energy (VPE)

$$E_{\text{VPE}} = -\frac{1}{2} \sum_\nu \left(|E_\nu| - |E_\nu^{(0)}| \right) \Big|_{\text{ren.}} \quad (9)$$

The sum contains bound and scattering states. The latter is expressed as a momentum integral over continuum energies weighted by the change in the density of states generated by $\Phi \neq \langle \Phi \rangle$. That change is computed from scattering data according to the Friedel-Krein formalism [23]. The main ingredient is the momentum derivative of the phase shift that describes the fermion scattering about the background potential. Renormalization (indicated by the subscript) is then accomplished by subtracting sufficiently many terms the Born series, which is an expansion in the strength of the potential, from the integrand and adding those pieces back in as Feynman diagram, which also combine to an expansion in that strength. Those Feynman diagrams are combined with the counterterms of the chosen renormalization scheme. The counterterm Lagrangian may only contain terms that arise from varying parameters (or scaling fields) in Eq. (1) and the coefficients of these terms may not depend on the peculiarities of the fields.

In the present application we still have to account for the fact that with $\phi(\pm\infty) = \pm 1$ the fermion mass terms, as induced by spontaneous symmetry breaking, have opposite signs at positive and negative spatial infinity. To this end we note that for a static system which is invariant under charge conjugation we can formally write the effective action, from which the VPE is extracted, as

$$\int \frac{d\omega}{2\pi} \text{TrLog} [\omega - h] = \frac{1}{2} \int \frac{d\omega}{2\pi} \text{TrLog} [\omega^2 - h^2] . \quad (10)$$

Thus the VPE of this fermion system can be obtained from the average VPE of two scalar systems [24] associated with Eq. (4) since

$$h^2 = \begin{pmatrix} -\partial_\xi^2 - g\phi' + g^2\phi^2 & 0 \\ 0 & -\partial_\xi^2 + g\phi' + g^2\phi^2 \end{pmatrix} . \quad (11)$$

The two potentials are straightforwardly read off as

$$V_S = g^2 (\phi^2 - 1) - g\phi' \quad \text{and} \quad \tilde{V}_S = g^2 (\phi^2 - 1) + g\phi' . \quad (12)$$

They are invariant under spatial reflection. For any scalar potential σ in $D = 1 + 1$ dimensions with that invariance the renormalized VPE is computed as

$$\epsilon_{\text{VPE}}[\sigma] = \int_0^\infty \frac{d\tau}{2\pi} \left\{ \ln \left[G(t,0) \left(G(t,0) - \frac{1}{t} G'(t,0) \right) \right] - \frac{\langle \sigma \rangle}{t} \right\}_{t=\sqrt{\tau^2+g^2}}. \quad (13)$$

Let us outline the connection of Eq. (13) with scattering data: For real momenta k the scattering phase shift is the phase of the Jost function, $f(k, \xi)$. The Jost function is the solution of the wave-equation with potential $\sigma(\xi)$ that obeys the boundary condition $\lim_{\xi \rightarrow \infty} e^{-ik\xi} = 1$. Factorizing the plane wave part $f(k, \xi) = e^{ik\xi} \overline{G}(k, \xi)$ defines $G(t, \xi) = \overline{G}(ik, \xi)$ by analytic continuation. It solves the ordinary differential equation

$$G''(t, \xi) = 2tG'(t, \xi) + \sigma(\xi)G(t, \xi), \quad (14)$$

subject to the boundary condition $\lim_{\xi \rightarrow \infty} G(t, \xi) = 1$. Eq. (13) applies to systems with $\sigma(-\xi) = \sigma(\xi)$ and thus decouples into parity channels. The first factor under the logarithm stems from odd parity (wave-function vanishes at the origin) while the second one originates from even parity (derivative of the wave-function vanishes at the origin).

As few further remarks are in order to explain Eq. (13). First, the last term under the integral proportional to $\langle \sigma \rangle = \int_0^\infty d\xi \sigma(\xi)$ is the Born approximation to the logarithm. The corresponding Feynman diagram is fully canceled by a counterterm within the no-tadpole renormalization scheme. Second, the physical momentum has been analytically continued into the upper half complex plane. Evaluating the momentum integral as a contour integral has contributions from the branch cut along $k = it$ in the dispersion relation $\epsilon = \sqrt{k^2 + g^2}$ with $t > g$ and the poles arising from the logarithmic derivative of the Jost function since the Jost function vanishes at the complex momenta of the bound state energies. Then the third important feature is that the latter contributions exactly cancel the bound state part in the sum of Eq. (9). More details are given in the reviews [22,25].

Restoring physical dimensions finally yields

$$E_{\text{VPE}} = -\frac{M}{4} \left(\epsilon_{\text{VPE}}[V_S] + \epsilon_{\text{VPE}}[\tilde{V}_S] \right). \quad (15)$$

We also observe that the Born subtraction in Eq. (13) involves the integral

$$\langle V_S + \tilde{V}_S \rangle = 2g^2 \int_0^\infty d\xi \left(\phi^2 - 1 \right).$$

It also arises from a counterterm which compensates changes of the vacuum expectation value $\langle \Phi \rangle^2 = \frac{M^2}{2\lambda}$ in Eq. (1). Hence Eq. (15) indeed implements the no-tadpole renormalization condition which requires that the (fermion) quantum corrections do not alter this expectation value.

4. Numerical Results

The model parameters are the mass M , the Higgs coupling λ and the Yukawa coupling constant g . With the scaling in Eq. (2) we factor out an overall constant M from the energy so that the relevant model parameters are g and the dimensionless ratio of the energy scales: $\alpha = \frac{M^2}{2\lambda}$. This ratio weighs the classical vs. vacuum polarization energies and its inverse plays the role of a loop counting parameter. For the numerical analysis it obviously suffices to choose, e. g. $M = 2$ and scan the g - λ parameter space; or equivalently the g - α space. This becomes obvious from the expression for the total energy

$$E_{\text{tot}}[\Phi] = E_{\text{cl}} + \sum_{\nu} \bar{\eta}_{\nu} |E_{\nu}| + E_{\text{VPE}} = \frac{M}{2} \left\{ \alpha \epsilon_{\text{cl}} + \sum_{\nu} \bar{\eta}_{\nu} |\epsilon_{\nu}| - \frac{1}{2} \left(\epsilon_{\text{VPE}}[V_S] + \epsilon_{\text{VPE}}[\tilde{V}_S] \right) \right\}, \quad (16)$$

as g and α are the only the model parameters entering the factor in curly brackets. An important observation is that the classical and the fermion energies scale differently with the model parameters.

Hence the choice of the particular relation $M = \sqrt{2\lambda}$, that was assumed in Ref. [10], may obscure important information [12].

For the particular parameters $M = \lambda = g = 2$ our model matches the super-symmetric one of Ref. [24]. For this case our numerical simulation yields $E_{\text{VPE}} = 0.3479$ when substituting $\phi = \phi_K$ in V_S and \tilde{V}_S . This agrees well with the analytic result $\frac{M}{\pi} \left(1 - \frac{\pi}{4\sqrt{3}}\right)$ found in that super-symmetric model and confirms the validity of our simulation².

The total energy is solely a functional of the scalar field Φ (or ϕ in dimensionless variables) since for any given scalar profile the fermion contributions are determined by the Dirac equation (4) and/or its downstream scattering equation (14). As already mentioned, this extremal condition cannot be formulated as a set of differential equation due to the non-local structure of the VPE. We therefore consider a parameterization for the scalar profile that is modeled after the kink $\phi_K = \tanh(\xi)$ with a rational function as correction³

$$\phi(\xi) = \frac{\xi^2 + a|\xi| + b}{\xi^2 + d|\xi| + e} \tanh(c\xi). \quad (17)$$

For notational convenience we define the set of parameters as

$$\mathbb{P} = \{a, b, c, d, e\}. \quad (18)$$

For a given set of occupation numbers $\{\bar{\eta}_v\}$ we then compute E_{tot} for numerous values of the five variational parameters in \mathbb{P} and identify the minimal value. To do so, we start with a profile function that is either close to the kink or close to one of the solutions constructed in Refs. [10–12]. We then apply a simple steepest descent algorithm to that first choice. Eventually this will converge to a minimum which then will be an upper bound to the actual minimum of E_{tot} because the variational space is limited.

To gauge the quality of this fitting function we will reconsider the minimization of the reduced energy functional

$$E_{\text{red}} = E_{\text{cl}} + E_{\text{val}} = E_{\text{cl}} + \frac{M}{2} \sum_v \bar{\eta}_v \epsilon_v. \quad (19)$$

That minimum was previously [10–12] constructed by self-consistently solving the Dirac equation (4) together with the differential equation

$$\phi'' = 2\phi(\phi^2 - 1) + \frac{4g\lambda}{M^2} \text{sign}(\epsilon)uv, \quad (20)$$

in which u and v are the spinor components of the level for which $\bar{\eta}_v = 1$ normalized to $\int d\xi (u^2 + v^2) = 1$. Here self-consistent refers to the condition that the profile function in the Dirac equation with that particular energy eigenvalue is also the solution to Eq. (20). Those self-consistent profiles turned out to significantly deviate from the kink. Since the VPE typically mitigates the binding from the occupied levels, we expect the actual solution to lie between the kink and those strongly distorted kink profiles. Hence reproducing the latter by the above fitting function to a high precision will justify the parameterization in Eq. (17). In a first step we therefore consider the case $M = \lambda = 2$ for which we constructed self-consistent solutions that minimize the reduced energy functional E_{red} earlier [12]. Subsequently the VPE is computed for this construction. The numerical results for the

² The computations of Ref. [24] make ample use of scattering data; so do we. Additionally we use the analytic properties of the Jost function to write the VPE as a single integral over imaginary momenta. Note also that Ref. [24] computes the VPE for two widely separated kinks and therefore has an additional factor two in Eq. (21).

³ We write absolute values for the odd powers to maintain the reflection property $\phi(-\xi) = -\phi(\xi)$. The actual calculation is performed on the half-line $\xi \geq 0$ and the absolute value sign can be ignored.

B_1 ($\bar{\eta}_v = 1$ for the first excited level with negative parity, all other $\bar{\eta}_v = 0$) and A_1 ($\bar{\eta}_v = 1$ for the first excited level with positive parity, all other $\bar{\eta}_v = 0$) configurations are listed in Table 1 for the choice $g = 4$.

Table 1. Comparison of energies from the fitting function and the self-consistent solution to the reduced problem, Eqs. (4), (19) and (20) for the B_1 and A_1 configurations using $g = 4.0$ and $M = \lambda = 2$: \mathbb{P}_1 is the fit to the self-consistent solution, \mathbb{P}_2 is the variational solution. The column labeled E_{val} denotes $\frac{M}{2} \sum_v \bar{\eta}_v \epsilon_v$ for the respective configurations.

B_1	E_{cl}	E_{val}	E_{VPE}	E_{tot}
	2.223	1.041	2.029	5.300
	self-consistent minimum of E_{red}			
	2.277	1.041	2.043	5.361
	$\mathbb{P}_1 = \{-0.758, -0.063, 0.903, -1.022, 1.018\}$			
	2.171	1.134	1.987	5.293
	$\mathbb{P}_2 = \{-0.733, -0.013, 0.917, -0.924, 0.964\}$			
A_1	E_{cl}	E_{val}	E_{VPE}	E_{tot}
	2.079	2.071	2.000	6.150
	self-consistent minimum of E_{red}			
	2.095	2.065	2.001	6.163
	$\mathbb{P}_1 = \{-2.163, 1.395, 0.759, -2.304, 1.988\}$			
	2.067	2.091	1.983	6.142
	$\mathbb{P}_2 = \{-2.151, 1.402, 0.760, -2.263, 1.958\}$			

Obviously the fit reproduces the results from the self-consistent approach convincingly well, in particular for the fermion ingredients. As a matter of fact, we have considered two scenarios, the first, labeled \mathbb{P}_1 , is a parameter fit to the self-consistent solution and the second, \mathbb{P}_2 , is the variational minimum to E_{red} . Either variational profile essentially equals the self-consistent one. This is the case for both the B_1 and A_1 configurations. Surprisingly those profiles exceed $\phi = 1$ at some moderate distance and approach the asymptotic vacuum expectation value from above as $\xi \rightarrow \infty$. We conclude that the fitting function, Eq. (17) is indeed a well-suited variational *ansatz* to approximate the scalar profile which minimizes the total energy, Eq. (16).

There is a subtlety with this parameterization, though. Asymptotically the profile behaves as $\phi(\xi) \sim 1 + \frac{a-d}{|\xi|}$ which causes the integral in the Born approximation

$$2g^2 \int_0^{\xi_{\text{max}}} d\xi \left[\phi^2(\xi) - 1 \right]$$

to logarithmically diverge as $\xi_{\text{max}} \rightarrow \infty$. The Born approximation has been introduced to cancel the large t component of the logarithm in Eq. (13). Since higher order terms of the Born series are finite when $\xi_{\text{max}} \rightarrow \infty$, we get a sensible result for Eq. (16) when we integrate the differential equation (14) between zero and the very same ξ_{max} . We have confirmed that once ξ_{max} is taken large enough, the numerical simulation of Eq. (13) is stable against further variations of ξ_{max} . In the no-tadpole scheme that first order contribution is exactly removed and there is no further problem. Although the chosen parameterization may not be ideal asymptotically, the results from Table 1 corroborate that it is nevertheless suitable. There is no such problem for the bound states whose wave-functions decay exponentially⁴.

In Table 2 we present the results from minimizing E_{tot} using same model parameters as in Table 1.

⁴ This is similar to the S -wave bound states for the Coulomb problem in the Schrödinger equation.

Table 2. Results from minimizing the total energy Eq. (16) using the variational *ansatz*, Eq. (17). Model parameters are as in Table 1: $g = 4.0$ and $M = \lambda = 2$.

	E_{cl}	E_{val}	E_{VPE}	E_{tot}
B_1	1.475	2.105	1.182	4.761
	$\mathbb{P} = \{-0.472, 0.181, 1.145, -0.544, 0.438\}$			
A_1	1.404	3.103	1.119	5.625
	$\mathbb{P} = \{-0.264, 0.132, 0.681, -0.282, 0.097\}$			

Obviously there are significant changes when including the VPE into the minimization program. This is not unexpected, as we previously found that the VPE is approximately as large as the valence energy. Compared with the data in Table 1 the total energy decreases by about 10% by lowering the classical part more strongly than increasing the fermion contribution. Yet, we still have $E_{tot} > \frac{Mg}{2}$, which is the mass of a free fermion in this case. Hence the soliton configuration is not kinematically stable against a decay into a free fermion. For the model parameters in Tables 1 and 2 the classical energy of the kink without back-reaction from the fermions is $\frac{4}{3}$ and the results from those tables suggests that the VPE strongly mitigates the back-reaction obtained earlier from only the valence levels. This also shows up in the graphical representation of the scalar profiles in Figure 1. The profiles for which we found the minimal E_{tot} are labeled \mathbb{P} . They do not differ from the kink ϕ_K substantially.




Figure 1. Profiles for $M = \lambda = 2$ and $g = 4$. Left panel: B_1 configuration, right panel: A_1 . The full black line (labeled \mathbb{P}) arises from minimizing the total energy, Eq. (16). The other lines (\mathbb{P}_1 , \mathbb{P}_2 , self-cons.) are configurations obtained from minimizing the reduced energy functional Eq.(19). The legends refer to the notation of Tables 1 and 2. For completeness we also show the kink without any fermion interaction (dashed lines).

We will now elaborate on our numerical analysis of scanning the g - α space in more detail. With the VPE included even the A_0 configuration has a non-zero fermion energy⁵ and thus a deviation $\phi \neq \phi_K$ is expected. We will discuss this case first although a very similar setting has already been considered in Ref. [16].

For the A_0 configuration the kink, $\phi_K(\xi) = \tanh(\xi)$ has $E_{tot}^{(K)} = \frac{M}{2} \left[\frac{4}{3}\alpha + \epsilon_{VPE} \right]$, where $\epsilon_{VPE} = 0.943$ and $\epsilon_{VPE} = 1.360$ for $g = 4$ and $g = 5$, respectively. For the cases shown in Table 3 the variational minimum is only slightly less than the total energy of the kink. Again, this corroborates the assertion that the inclusion of the fermion fields only leads to a moderate back-reaction. We also see that $E_{tot} < \frac{Mg}{2}$ for the A_0 case which is thus kinematically stable against a decay into a free fermion. However, since the zero mode does not have definite fermion charge, this configuration cannot be asserted a particle number, in contrast to a free fermion.

⁵ The A_0 bound state by itself is a zero mode with either u or v vanishing so that $\bar{\psi}\psi|_{A_0} = 0$ and thus does not couple to the scalar field.

Table 3. Variational approach for the A_0 configuration. Top panel: $g = 4$, bottom panel: $g = 5$. The valence contribution is not listed as $E_{\text{val}} \equiv 0$ for this configuration. For comparison we also list the total energy for the kink profile $\phi_K(\xi) = \tanh(\xi)$.

α	E_{cl}	E_{VPE}	E_{tot}	$E_{\text{tot}}^{(K)}$
1.0	1.385	0.817	2.202	2.277
	$\mathbb{P} = \{1.053, 1.219, 1.104, 1.105, 0.946\}$			
1.2	1.648	0.829	2.477	2.543
	$\mathbb{P} = \{1.053, 1.191, 1.112, 1.094, 0.976\}$			
1.5	2.043	0.843	2.886	2.943
	$\mathbb{P} = \{1.105, 1.159, 1.138, 1.130, 1.032\}$			
2.0	2.703	0.861	3.564	3.610
	$\mathbb{P} = \{1.104, 1.139, 1.120, 1.122, 1.051\}$			
α	E_{cl}	E_{VPE}	E_{tot}	$E_{\text{tot}}^{(K)}$
1.0	1.437	1.079	2.516	2.693
	$\mathbb{P} = \{1.093, 1.243, 1.208, 1.157, 0.945\}$			
1.2	1.699	1.103	2.802	2.960
	$\mathbb{P} = \{1.098, 1.243, 1.169, 1.158, 0.951\}$			
1.5	2.092	1.131	3.222	3.360
	$\mathbb{P} = \{1.110, 1.221, 1.162, 1.149, 0.980\}$			
2.0	2.748	1.167	3.914	4.026
	$\mathbb{P} = \{1.106, 1.202, 1.125, 1.146, 0.997\}$			

Next we turn to the B_1 configuration with the numerical results displayed in Table 4.

Table 4. Variational approach for the B_1 configuration. Top panel: $g = 4$, bottom panel: $g = 5$. The entry $g = 4$ and $\alpha = 1$ is that of Table 2.

α	E_{cl}	E_{val}	E_{VPE}	E_{tot}	$E_{\text{tot}}^{(K)}$
1.0	1.475	2.105	1.182	4.761	4.922
	$\mathbb{P} = \{-0.472, 0.181, 1.145, -0.544, 0.438\}$				
1.2	1.723	2.186	1.143	5.052	5.189
	$\mathbb{P} = \{-0.472, 0.212, 1.140, -0.533, 0.439\}$				
1.5	2.104	2.270	1.103	5.477	5.589
	$\mathbb{P} = \{-0.474, 0.243, 1.136, -0.522, 0.436\}$				
2.0	2.748	2.358	1.064	6.170	6.256
	$\mathbb{P} = \{-0.466, 0.280, 1.112, -0.504, 0.434\}$				
α	E_{cl}	E_{val}	E_{VPE}	E_{tot}	$E_{\text{tot}}^{(K)}$
1.0	1.485	2.390	1.647	5.522	5.693
	$\mathbb{P} = \{-0.479, 0.132, 1.224, -0.544, 0.321\}$				
1.2	1.730	2.481	1.603	5.814	5.960
	$\mathbb{P} = \{-0.474, 0.159, 1.217, -0.533, 0.335\}$				
1.5	2.107	2.576	1.559	6.241	6.360
	$\mathbb{P} = \{-0.473, 0.194, 1.203, -0.525, 0.358\}$				
2.0	2.749	2.675	1.512	6.936	7.026
	$\mathbb{P} = \{-0.471, 0.237, 1.174, -0.516, 0.380\}$				

In this case the valence energy is substantial. However, since a strongly distorted kink comes with large E_{cl} and E_{VPE} , cf. Table 1, minimizing the valence energy does not lower the total energy. Rather configurations that minimize the total energy have significantly larger valence energies than the solutions constructed in Refs. [10–12]. The resulting configuration is similar to the kink as the variational parameters approximately obey $a \approx d$ and $b \approx e$. We find that c is slightly larger than unity which would indicate that profile is a squeezed kink. However, the remaining differences

between a and d as well as between b and e lead to a stretched kink. This is seen in Figure 1 in which we show minimizing profiles for various channels. As we increase α , the classical contribution to the energy becomes more dominant and the minimizing configuration should get even closer to the kink. Indeed the numerical simulations confirm that as we find the variational parameters $\mathbb{P} = \{-0.401, 0.253, 1.070, -0.416, 0.342\}$ and $\mathbb{P} = \{-0.396, 0.274, 1.041, -0.408, 0.339\}$ for $\alpha = 3.0$ and $\alpha = 4.0$, respectively; with $g = 4.0$ in both cases. The respective energies are $E_{\text{tot}} = 7.532$ and $E_{\text{tot}} = 8.879$. The discrepancies to the energy of the kink configuration (7.589 and 8.922) are marginal and decrease as α increases.

As a final example we consider the A_1 configuration when the valence fermion dwells in the third bound state, the second one with even parity. The numerical results are displayed in Table 5.

Table 5. Variational approach for the A_1 configuration. Top panel: $g = 4$, bottom panel: $g = 5$. The entry $g = 4$ and $\alpha = 1$ is that of Table 2.

α	E_{cl}	E_{val}	E_{VPE}	E_{tot}	$E_{\text{tot}}^{(\text{K})}$
1.0	1.404	3.103	1.119	5.625	5.741
	$\mathbb{P} = \{-0.264, 0.132, 0.681, -0.282, 0.097\}$				
1.2	1.669	3.136	1.101	5.905	6.007
	$\mathbb{P} = \{-0.262, 0.133, 0.702, -0.281, 0.102\}$				
1.5	2.065	3.176	1.079	6.319	6.407
	$\mathbb{P} = \{-0.260, 0.139, 0.728, -0.277, 0.109\}$				
2.0	2.724	3.228	1.051	7.003	7.074
	$\mathbb{P} = \{-0.258, 0.141, 0.769, -0.271, 0.116\}$				
α	E_{cl}	E_{val}	E_{VPE}	E_{tot}	$E_{\text{tot}}^{(\text{K})}$
1.0	1.381	3.663	1.562	6.606	6.693
	$\mathbb{P} = \{-0.248, 0.081, 0.749, -0.263, 0.063\}$				
1.2	1.645	3.697	1.541	6.882	6.960
	$\mathbb{P} = \{-0.255, 0.107, 0.766, -0.263, 0.083\}$				
1.5	2.044	3.725	1.523	7.292	7.360
	$\mathbb{P} = \{-0.252, 0.105, 0.781, -0.263, 0.085\}$				
2.0	2.706	3.768	1.496	7.970	8.026
	$\mathbb{P} = \{-0.245, 0.095, 0.814, -0.255, 0.081\}$				

The scenario is pretty much the same as for the previous configurations: The inclusion of the fermion VPE moves the distorted kink back to a slightly stretched kink. We see that even for $\alpha = 1$ the variational solution is not very different from the kink. This implies that the total energy is dominated by the classical part which is another indication that there are sizable cancellations between the gain from binding a valence fermion and the fermion VPE. All our variational searches yield $a > d$ so that the profile approaches the vacuum expectation value from above as $\zeta \rightarrow \infty$.

For a certain parameter set we display the profiles that minimize E_{tot} in the various channels in Figure 2.




Figure 2. Variational solutions of the A_0 , B_1 and A_1 configurations for $\alpha = 1.0$ and $g = 4.0$. For comparison the kink profile is also shown.

When compared to the kink, the solution of the A_0 configuration is squeezed, while the others are stretched. The reason is that squeezing the kink slightly decreases the VPE. To considerably gain energy from binding a non-zero mode, the Yukawa interaction must be strongly attractive which requires an extended kink profile. That is what we observe for the B_1 and A_1 configurations.

We should, however, mention that we find various local minima for the total energy in the space of the chosen variational parameters. Their total energies may be slightly smaller or larger than those reported above; but they turn out to be stationary under the steepest descent algorithm. This is the case for all configurations but we will discuss this only for the A_1 configuration in more detail. The data for some of the alternative solutions are listed in Tables 6 and 7. Looking at the corresponding variational parameters suggests that these solutions would be quite different. In particular those from Table 6 exhibit significant deviations from the kink relation $b \approx e$. However, when plotting them, *cf.* Figure 3, we observe that the profiles are approximately identical and differences get smaller as the Yukawa coupling increases. We conjecture that these solutions are pretty close to the actual solution, but the particular variational ansatz is not able of capturing it exactly. Furthermore the minimum may be very shallow. The ansatz most likely does not allow for a continuous transition between the solutions on a path that can be constructed from a steepest descent procedure. In any event, we have sufficient evidence to state that the actual solution will be similar to the kink but quite distinct from the solutions of Refs. [10–12].

Table 6. Alternative solutions for the A_1 configuration: Top panel $g = 4$, bottom panel $g = 5$.

α	E_{cl}	E_{val}	E_{VPE}	E_{tot}
1.0	1.443	3.035	1.141	5.618
	$\mathbb{P} = \{-1.963, 1.521, 1.345, -2.060, 1.875\}$			
1.2	1.696	3.092	1.113	5.900
	$\mathbb{P} = \{-1.917, 1.530, 1.268, -2.002, 1.848\}$			
1.5	2.073	3.156	1.088	6.317
	$\mathbb{P} = \{-1.832, 1.551, 1.141, -1.896, 1.807\}$			
2.0	2.733	3.220	1.051	7.004
	$\mathbb{P} = \{-1.846, 1.580, 1.154, -1.905, 1.807\}$			
α	E_{cl}	E_{val}	E_{VPE}	E_{tot}
1.0	1.419	3.507	1.644	6.571
	$\mathbb{P} = \{-1.664, 0.995, 1.245, -1.727, 1.120\}$			
1.2	1.660	3.634	1.586	6.879
	$\mathbb{P} = \{-1.833, 1.457, 1.165, -1.918, 1.719\}$			
1.5	2.052	3.692	1.549	7.293
	$\mathbb{P} = \{-1.824, 1.519, 1.116, -1.898, 1.744\}$			
2.0	2.710	3.746	1.514	7.971
	$\mathbb{P} = \{-1.748, 1.490, 1.107, -1.804, 1.666\}$			

Table 7. Second set of alternative solutions for the A_1 configuration: Top panel $g = 4$, bottom panel $g = 5$.

α	E_{cl}	E_{val}	E_{VPE}	E_{tot}
1.0	1.394	3.109	1.138	5.642
	$\mathbb{P} = \{2.600, 0.170, 0.717, 2.581, 0.088\}$			
1.2	1.660	3.144	1.115	5.918
	$\mathbb{P} = \{2.441, 0.207, 0.734, 2.428, 0.117\}$			
1.5	2.057	3.185	1.090	6.332
	$\mathbb{P} = \{2.590, 0.182, 0.766, 2.579, 0.110\}$			
2.0	2.716	3.239	1.058	7.013
	$\mathbb{P} = \{2.558, 0.188, 0.805, 2.575, 0.126\}$			

α	E_{cl}	E_{val}	E_{VPE}	E_{tot}
1.0	1.370	3.667	1.578	6.621
	$\mathbb{P} = \{2.609, 0.143, 0.770, 2.578, 0.074\}$			
1.2	1.637	3.694	1.561	6.892
	$\mathbb{P} = \{2.449, 0.193, 0.780, 2.428, 0.108\}$			
1.5	2.036	3.728	1.536	7.300
	$\mathbb{P} = \{2.442, 0.193, 0.801, 2.425, 0.117\}$			
2.0	2.700	3.771	1.507	7.978
	$\mathbb{P} = \{2.592, 0.155, 0.834, 2.577, 0.103\}$			




Figure 3. Comparison of the different solutions for the A_1 scenario according to the respective tables. We always have $\alpha = 1$ as in Table 1 and Figure 1. Left panel $g = 4$, right panel $g = 5$.

Without the fermion coupling, the boson contribution to the VPE is $\frac{1}{2\sqrt{3}} - \frac{3}{\pi} \approx -0.666$ for $M = 2$ [22,26,27]. With that coupling included the boson VPE is difficult to estimate because there is a linear term in the harmonic approximation for the fluctuations when the profile is not a solution to the classical kink equation. Furthermore imaginary frequency eigenvalues emerge for the boson fluctuations [28]. Yet, we assume that the above number is a useful estimate because the variational profiles are quite similar to the kink. As conjectured earlier, the numerical simulations verify that the fermion VPE becomes significantly larger when the Yukawa coupling increases. In subsection 3.1 we especially argued that for large enough values of the Yukawa coupling constant the fermion VPE should dominate the boson counterpart. And indeed we always find that $E_{VPE} > \frac{M}{2}$ when $g \geq 5$.

5. Conclusions

In this project we have considered a model with fermions coupled to the kink in one space and one time dimension and investigated the fermion back-reaction on the kink profile. In this approach a valence fermion appears as an explicitly occupied bound state level and the back-reaction may be significant when this is not the ground state level. And indeed earlier studies, that only considered the coupling of the kink to a single fermion valence level, found considerable back-reactions. However, it is important to not only consider the energy of the occupied valence level but also to add the energy of the Dirac sea to the functional that determines the back-reaction. The main argument for

its inclusion is the consistency of the semi-classical expansion. This contribution is obtained as the renormalized sum of the changes of the one-particle fermion energies. These changes emerge because the Yukawa coupling between the kink and the fermions polarizes the fermion vacuum. They concern the discrete bound states as well as the continuous scattering states. The boson contribution to the vacuum polarization energy was not included in the energy functional. Rather we have argued that, for large values of the Yukawa coupling constant, the fermion energies would dominate the quantum corrections. We have verified that conjecture *a posteriori*.

We have utilized a variational approach to find (an upper bound) to the minimal total energy. We have only considered a single variational parameterization of the kink profile with five parameters to lessen the numerical efforts. Certainly this leaves space for improvement. Nevertheless, we consider it sufficient to show that the Dirac sea contribution brings back the strongly distorted kink profiles from the self-consistent treatment for the case that omits the Dirac sea, to a slightly squeezed or stretched kink profile. The similarity to the kink is particularly pronounced for the A_0 configuration where we do not have a valence quark contribution to the energy. For that reason, a slightly squeezed kink profile is energetically favorable for the A_0 configuration. In all other cases, some energy is gained from binding the valence level by stretching the kink because it makes the Yukawa interaction more attractive. Since this deformation increases both the classical energy and the vacuum polarization energy, the stretch can only be modest.

The dominant effect for mitigating the kink distortion is the balance between the bound state and vacuum energies. This could well be a consequence of Levinson's theorem [29,30], which relates the number of bound states to the density of continuum states. Essentially it states that any bound state must have emerged from the continuum. So the energy gain from one comes with an energy loss of the other.

We have considered cases in which only a single fermion occupation number was non-zero. The valence energy should become more important when this is true for several occupation numbers. Then we expect stronger distortion of the kink. This should be subject of future projects.

Author Contributions: The authors mutually agree that their contributions warrant co-authorship.

Funding: H. W. is supported in part by the National Research Foundation of South Africa (NRF) by grant 150672.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Nambu, Y. String-Like Configurations in the Weinberg-Salam Theory. *Nucl. Phys. B* **1977**, *130*, 505.
2. Weigel, H. *Chiral Soliton Models for Baryons*; Vol. 743, Lect. Notes Phys, 2008.
3. Feist, D.T.J.; Lau, P.H.C.; Manton, N.S. Skyrmions up to Baryon Number 108. *Phys. Rev. D* **2013**, *87*, 085034.
4. U. Schollwöck, *et al.*. *Quantum Magnetism*; Vol. 645, Lect. Notes Phys, 2004.
5. Nagaosa, N.; Tokura, Y. *Nature Nanotech.* **2013**, *8*, 899.
6. Vilenkin, A.; Shellard, E.P.S. *Cosmic Strings and Other Topological Defects*; Cambridge University Press, 2000.
7. Moradi Marjaneh, A.; Simas, F.C.; Bazeia, D. Collisions of kinks in deformed I. 160 ϕ^4 and I. 160 ϕ^6 models. *Chaos Solitons and Fractals* **2022**, *164*, 112723.
8. Chu, Y.Z.; Vachaspati, T. Fermions on one or fewer kinks. *Phys. Rev. D* **2008**, *77*, 025006.
9. Brihaye, Y.; Delsate, T. Remarks on bell-shaped lumps: Stability and fermionic modes. *Phys. Rev. D* **2008**, *78*, 025014.
10. Klimashonok, V.; Perapechka, I.; Shnir, Y. Fermions on kinks revisited. *Phys. Rev. D* **2019**, *100*, 105003.
11. Perapechka, I.; Shnir, Y. Kinks bounded by fermions. *Phys. Rev. D* **2020**, *101*, 021701.
12. Saadatmand, D.; Weigel, H. Excited fermions on kinks and the Dirac sea. *Phys. Rev. D* **2023**, *107*, 036006.
13. Alkofer, R.; Reinhardt, H.; Weigel, H. Baryons as chiral solitons in the Nambu-Jona-Lasinio model. *Phys. Rept.* **1996**, *265*, 139.
14. Friedberg, R.; Lee, T.D. Fermion Field Nontopological Solitons. 1. *Phys. Rev. D* **1977**, *15*, 1694.
15. Nolte, G.; Kunz, J. The Sphaleron barrier in the presence of Fermions. *Phys. Rev. D* **1993**, *48*, 5905–5916.

16. Farhi, E.; Graham, N.; Jaffe, R.L.; Weigel, H. Heavy fermion stabilization of solitons in (1+1)-dimensions. *Nucl. Phys. B* **2000**, *585*, 443.
17. Farhi, E.; Graham, N.; Jaffe, R.L.; Weigel, H. Searching for quantum solitons in a (3+1)-dimensional chiral Yukawa model. *Nucl. Phys. B* **2002**, *630*, 241–268.
18. Chodos, A.; Jaffe, R.L.; Johnson, K.; Thorn, C.B.; Weisskopf, V.F. A New Extended Model of Hadrons. *Phys. Rev. D* **1974**, *9*, 3471.
19. Bowler, K.C.; Hey, A.J.G. The Roper Resonance: A Problem for the MIT Bag? *Phys. Lett. B* **1977**, *69*, 469.
20. Guichon, P.A.M. A Nonstatic Bag Model for the Roper Resonances. *Phys. Lett. B* **1985**, *164*, 361.
21. Reinhardt, H. The Chiral Soliton in the Proper Time Regularization Scheme. *Nucl. Phys. A* **1989**, *503*, 825.
22. Graham, N.; Quandt, M.; Weigel, H. *Spectral Methods in Quantum Field Theory*; Vol. 777, Lect. Notes Phys, 2009.
23. J. S. Faulkner. Scattering theory and cluster calculations. *J. Phys.* **1977**, *C10*, 4661.
24. Graham, N.; Jaffe, R.L. Fermionic one loop corrections to soliton energies in (1+1)-dimensions. *Nucl. Phys. B* **1999**, *549*, 516.
25. Graham, N.; Weigel, H. Quantum corrections to soliton energies. *Int. J. Mod. Phys. A* **2022**, *37*, 2241004.
26. Dashen, R.F.; Hasslacher, B.; Neveu, A. Nonperturbative Methods and Extended Hadron Models in Field Theory 2. Two-Dimensional Models and Extended Hadrons. *Phys. Rev. D* **1974**, *10*, 4130–4138.
27. Rajaraman, R. *Solitons and Instantons*; North Holland, 1982.
28. Graham, N.; Jaffe, R.L. Unambiguous one loop quantum energies of (1+1)-dimensional bosonic field configurations. *Phys. Lett. B* **1998**, *435*, 145–151.
29. Levison, N. On the uniqueness of the potential in a Schrödinger equation for a given asymptotic phase. *Kgl. Danske Videnskab Selskab; Mat.-fys. Medd.* **1949**, *25*, 9.
30. Barton, G. Levinson's Theorem in One-dimension: Heuristics. *J. Phys. A* **1985**, *18*, 479–494.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.