

Communication

Not peer-reviewed version

---

# Could SP-A and SP-D Serum Levels Predict COVID-19 Severity?

---

[Luca Maddaloni](#) <sup>\*</sup>, [Veronica Zullino](#) , [Ginevra Bugani](#) <sup>\*</sup>, [Alessandro Lazzaro](#) , [Matteo Brisciani](#) , [Claudio Mastroianni](#) , [Letizia Santinelli](#) , Franco Ruberto

Posted Date: 19 April 2024

doi: 10.20944/preprints202402.0739.v2

Keywords: SP-A; SP-D; SARS-CoV-2; collectins; surfactant proteins



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Communication

# Could SP-A and SP-D Serum Levels Predict COVID-19 Severity?

Luca Maddaloni <sup>1,†</sup>, Veronica Zullino <sup>2,†</sup>, Ginevra Bugani <sup>1</sup>, Alessandro Lazzaro <sup>1</sup>,  
Matteo Brisciani <sup>2</sup>, Claudio Maria Mastroianni <sup>1</sup>, Letizia Santinelli <sup>1</sup> and Franco Ruberto <sup>2</sup>

<sup>1</sup> Department of Public Health and Infectious Diseases, Sapienza University of Rome, Italy;  
luca.maddaloni@uniroma1.it (L.M.); ginevra.bugani@uniroma1.it (G.B.);  
alessandro.lazzaro@uniroma1.it (A.L.); claudio.mastroianni@uniroma1.it (C.M.M.);  
letizia.santinelli@uniroma1.it (L.S.)

<sup>2</sup> Department of General and Specialistic Surgery, Sapienza University of Rome, Italy;  
v.zullino@policlinicoumberto1.it (V.Z.); m.brisciani@policlinicoumberto1.it (M.B.);  
f.ruberto@policlinicoumberto1.it (F.R.)

\* Correspondence: luca.maddaloni@uniroma1.it (L.M.); ginevra.bugani@uniroma1.it (G.B.)

† These authors contributed equally to this work.

**Abstract:** **Background:** given the various clinical manifestations that characterize coronavirus disease 2019 (COVID-19), the scientific community is constantly searching for biomarkers with prognostic value. Surfactant protein A (SP-A) and D (SP-D) play a crucial role in ensuring proper alveolar function and an alteration of their serum levels have been reported in several pulmonary diseases characterized by acute respiratory distress syndrome (ARDS) and pulmonary fibrosis. Considering that such clinical manifestations can also occur during severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection, we wondered if these collectins could act as prognostic markers. **Methods:** in this regard, serum levels of SP-A and SP-D were measured by enzyme immunoassay in patients with SARS-CoV-2 infection (n=51) at admission (T0) and after 7 days (T1) and compared with healthy donors (n=11). **Results:** SP-D increased in COVID-19 patients compared to healthy controls during the early phases of infection, while a significant reduction was observed at T1. Stratifying SARS-CoV-2 patients according to disease severity, increased serum SP-D levels were observed in severe compared to mild patients. **Conclusions:** In the light of these results SP-D, but not SP-A, seems to be an eligible marker of COVID-19 pneumonia and the early detection of SP-D serum levels could be crucial for a preventive clinical management.

**Keywords:** SP-A; SP-D; SARS-CoV-2; collectins; surfactant proteins

## 1. Introduction

The continuous global pandemic of coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), represents a public health menace on a worldwide scale<sup>1</sup>. Most of SARS-CoV-2-infected patients are asymptomatic or characterized by mild symptoms (throat, loss of taste and smell, malaise, and joint pain). However, in individuals older (>50 years old) or with comorbidities (hypertension, diabetes, impaired immunity) the risk of severe manifestations of COVID-19 is higher. The most common clinical manifestation is acute respiratory distress syndrome (ARDS) with hypoxic respiratory failure. During this kind of severe critical illness, the overexpression of proinflammatory cytokines leads to endothelial dysfunction and damage. Considering the heterogeneous prognosis of COVID-19 patients, the scientific community has long wondered about potential biomarker candidates for a poorer prognosis. This can enable therapeutic interventions and monitor patients within an appropriate window of action. SARS-CoV-2 has the capacity to affect type I pneumocytes, type II pneumocytes, and alveolar macrophages, leading to impaired lung function<sup>2</sup>. Pulmonary surfactants consist primarily of phospholipids and a smaller fraction of surfactant proteins (SP-A, SP-B, SP-C, SP-D)<sup>3</sup>. These surfactant phospholipids, located within the alveolar space, have a critical role in reducing the surface tension along the alveolar walls during inhalation and prevent alveolar collapse after exhalation<sup>4</sup>. In particular, SP-A and SP-D

are collectins that interact with the surface oligosaccharides of viruses and this phenomenon bolsters viral clearance, primarily through the actions of macrophages and monocytes<sup>4</sup>. Additionally, SP-A, as an immunomodulator, has the capacity to restrain dendritic cell maturation and inhibit the excessive release of IL-8 by eosinophils<sup>5</sup>. Following SARS-CoV-2 invasion, the ensuing viral replication and damage to type II pneumocytes negatively affect surfactant production, giving rise to breathing difficulties and the development of ARDS in COVID-19 patients. To this extent, in more severe instances, the extensive loss of these critical "alveolar defenders" culminates in respiratory failure<sup>2</sup>. Many studies evaluated SP-D levels in bronchoalveolar lavage fluid (BALF). Decreased SP-D levels were reported in children affected by cystic fibrosis with acute lung infection or in children with gastroesophageal reflux disease (GERD) who have continuous acid airway injury and suffer from chronic lung infections. Furthermore, decreased SP-D levels in BALF were observed in patients affected by ARDS with worse oxygenation compared to patients with good oxygenation. In addition, in smokers, BALF SP-D levels are reduced compared to nonsmokers, while increased levels were observed in patients with sarcoidosis, asthma, eosinophilic pneumonia and pulmonary alveolar proteinosis (PAP)<sup>6</sup>. Several studies assessed the value of SP-D serum levels as a disease marker for human lung diseases. This evidence concerns patients with various form of interstitial lung disease, including interstitial pneumonia with collagen disease, idiopathic pulmonary fibrosis, sarcoidosis and other. The mechanisms that explain how pulmonary SP-D comes into the circulation are unclear, but there are several hypotheses: inflammatory conditions, as in ARDS, induce an increased permeability of lung vessels that might result in alveolar to vascular shift of SP-D; the same inflammatory status could be responsible for a loss of epithelial secretory cells integrity and an efflux of SP-D from epithelial cells into alveolar vessels<sup>6-8</sup>.

Considering the prognostic increased of SP-A and SP-D serum concentrations in patients with lung damage (idiopathic pulmonary fibrosis and interstitial pneumonia) with collagen vascular diseases<sup>9</sup>, we hypothesized that in COVID-19 as well, the levels of these collectins could predict the severity of the disease.

## 2. Materials and Methods

### 2.1. Study Participants and Ethical Approvals

Peripheral blood samples were consecutively collected at T0 (at admission) and T1 (seven days post hospitalization) from SARS-CoV-2 infected individuals hospitalized at the Policlinico Umberto I, Sapienza University of Rome (Italy) from January to March 2022. A control group of healthy donors was also enrolled. Patients who were  $\geq 18$  years old, who have received or not vaccination, with a positive SARS-CoV-2 RT-PCR test were enrolled. Patients with any one of the following conditions were excluded: a diagnosis of autoimmune lung disease, pulmonary fibrosis, obstructive pulmonary disease or bacterial pneumonia. Oxygen therapy was delivered via Venturi mask (VMK) in spontaneous breathing patients; if hypoxemia persisted, continuous positive airway pressure (CPAP) was applied. From 51 patients studied, 11 patients had comorbidities (21,6%) and 40 no comorbidities (78,4%). The most common comorbidities were arterial hypertension (37,3%), cardiovascular diseases (25,4%), oncological diseases (17,6%) and controlled lung diseases (5,1%). Among patients with controlled lung diseases: 4,08% had chronic obstructive pulmonary disease (COPD) and 1,02% had pulmonary emphysema. In all cases, patients were under treatment with oral steroid. Hematologic diseases were the most frequent pathological condition in the field of oncological diseases. Furthermore, we used body mass index to identify patients with obesity: 8,16% had an obesity class I while the other had a body mass index less than  $30 \text{ kg/m}^2$ . Finally, patients with smoking habit were not enrolled.

The COVID-19 case definitions of European Centre for Disease Prevention and Control (ECDC) were adopted<sup>7</sup> and severity stratification was based on World Health Organization (WHO) criteria<sup>8</sup>. The study was approved by the ethics committee of the Policlinico Umberto I Hospital, "Sapienza" University of Rome and informed consent was obtained from both COVID-19 patients and healthy

individuals. Finally, patients' data were anonymized and a progressive alphanumeric code was assigned to them.

## 2.2. Serological Evaluation of SP-A and SP-D Levels

Twenty milliliters of whole blood were collected; serum was separated by centrifugation at 1620 RCF for 10' within one hour after collection and stored at -80°C for further analysis. Serum SP-A and SP-D levels at T0 and T1 were evaluated using a commercially available enzyme-linked immunosorbent assay (ELISA) kit (Catalog numbers MBS167144 and ELH-SPD, respectively), according to the manufacturers' protocol. The assay was based on an indirect assay with an antigen-antisera bound and a conjugate antibody using HRP-Streptavidin as a substrate for the detection. Absorptions were measured at 450 nm with a Microplate Multimode Reader (Biorad). Concentrations in each sample were calculated from the standard curve, obtained by plotting absorbance versus known standard concentrations. SP-A and SP-D concentrations were expressed as ng/L and ng/mL, respectively.

## 2.3. RT-qPCR Detection of SARS-CoV-2 RNA

Viral RNA was extracted from nasopharyngeal swabs using Versant SP 1.0 Kit (Siemens Healthcare Diagnostics). Briefly, 10 µl of extracted RNA was reverse-transcribed and simultaneously amplified by a real-time RT-PCR system (RealStar SARS-CoV-2 RT-PCR, Altona Diagnostics), targeting E and S viral genes.

## 2.4. Statistical Analysis

Patients' data were expressed as median (interquartile range) or number (percentage). Demographic and clinical patients' characteristics were analyzed using "N-1" Chi-squared test. Cross-sectional data between patients and healthy donors were analyzed using the Mann-Whitney U test, while Wilcoxon signed-rank test for paired samples was used to evaluate longitudinal data between T0 and T1. A p-value less than 0.05 was considered statistically significant. Statistical analyses were performed using GraphPad Prism software, version 9.4 (GraphPad Software Inc., La Jolla, California, USA).

# 3. Results

## 3.1. Study Participants

A total of 51 SARS-CoV-2-infected patients and 11 healthy donors were enrolled in this study. Demographic and clinical characteristics of the study population are reported in Table 1.

**Table 1.** Demographic and clinical characteristics of SARS-CoV-2 infected patients and healthy donors.

| Parameters   | SARS-CoV-2 infected patients (n=51) (A)       | Healthy donors (n=11) (B) | A vs B p-values |
|--|---|---------------------------|-----------------|
| Age (years) <sup>a</sup>                               | 68 (56.5-78)                                  | 39 (30-52)                | < 0.0001        |
| Male <sup>b</sup>                                      | 24 (47%)                                      | 6 (54.5%)                 | 0.7465          |
| SARS-CoV-2 vaccination (at least 2 doses) <sup>b</sup> | 14 (27%)                                      | 11 (100%)                 | < 0.0001        |
| Respiratory support <sup>b</sup>                       | VMK 12 (23%)<br>A.A. 9 (18%)<br>CPAP 30 (59%) | NA                        | -               |
| D-dimer (µg/l) <sup>a</sup>                            | 1092 (631.5-1621)                             | NA                        | -               |
| PCR (mg/dl) <sup>a</sup>                               | 4.17 (1.39-6.5)                               | NA                        | -               |
| Ferritin (ng/ml) <sup>a</sup>                          | 823 (357-1244)                                | NA                        | -               |
| LDH (Ui/L) <sup>a</sup>                                | 242 (209.5-252)                               | NA                        | -               |

|  |                 |    |   |
|--|-----------------|----|---|
| Fibrinogen (mg/dl) <sup>a</sup>                | 407 (325-440)   | NA | - |
| Lymphocytes (10 <sup>3</sup> /ml) <sup>a</sup> | 1285 (815-1795) | NA | - |


NA = not applicable, VMK = Venturi Mask, CPAP = Continuous Positive Airway Pressure, A.A. = Ambient Air.

<sup>a</sup> Data are expressed as median (interquartile ranges). <sup>b</sup> Data are expressed as number (percentage). Differences in demographic characteristics between SARS-CoV-2 infected patients at admission (T0) and healthy individuals were evaluated using Mann-Whitney U and Chi-squared tests.  $P < 0.05$  was considered statistically significant.

### 3.1. Serum SP-A and SP-D Levels Evaluation

At baseline, COVID-19 patients displayed higher median levels of SP-D protein ( $p = 0.0063$ ) compared to healthy donors, while no differences were recorded for SP-A levels between both groups ( $p = 0.0824$ ).


At T1, serum SP-A and SP-D levels were similar among COVID-19 patients and healthy donors ( $p = 0.3451$  and  $p = 0.0569$ , respectively). When comparing T0 and T1, a significant reduction of serum SP-A and SP-D levels were observed at T1 among COVID-19 patients ( $p = 0.0123$  for both proteins) (Figure 1).



**Figure 1.** Comparison of serum SP-A (A) and SP-D (B) between SARS-CoV-2 infected patients at hospitalization (T0) and 7 days after (T1) and healthy controls. Data were analyzed using the Mann-Whitney U-test for unpaired samples and the Wilcoxon signed-rank test for paired samples. \* $p < 0.05$ ; \*\* $p < 0.01$ .

To investigate the possible association between surfactant protein expression and disease severity, patients were stratified according to respiratory support requirement: the mild group were in ambient air or needed a VMK support ( $n=21$ ), while the severe group was under CPAP ( $n=30$ ) (Table 2). A lower vaccination rate against SARS-CoV-2 was found in the severe compared to the mild patient group. None of these patients developed ARDS. Serum SP-A levels were comparable between healthy controls, mild and severe COVID-19 patients at both timepoints ( $p > 0.05$ ), while the only difference observed was a longitudinally reduction in the mild group ( $p = 0.0153$ ). Regarding SP-D, no differences were recorded for serum levels of this protein among healthy donors and mild patients at both time points; by contrast, severe patients exhibited higher SP-D levels at T0 ( $p = 0.0006$ )

and T1 ( $p = 0.0063$ ) compared to the control group. In addition, the serum levels of this protein were higher in the COVID-19 severe group compared to the mild one at T1 ( $p = 0.0369$ ) (Figure 2).



**Figure 2.** Comparison of serum SP-A (A) and SP-D (B) levels between mild and severe SARS-CoV-2 infected patients at hospitalization (T0) and 7 days after (T1) and healthy controls. Data were analyzed using the Mann-Whitney U-test for unpaired samples and the Wilcoxon signed-rank test for paired samples. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ .

**Table 2.** Demographic and clinical characteristics of mild and severe SARS-CoV-2 infected patients.

| Parameters   | Mild SARS-CoV-2 infected patients (n=21) (A) | Severe SARS-CoV-2 infected patients (n=30) (B) | A vs B p-values |
|--|--|--|-----------------|
| Age (years) <sup>a</sup>                               | 75 (62-83)                                   | 65 (56-74)                                     | 0,0739          |
| Male <sup>b</sup>                                      | 8 (38%)                                      | 66 (53%)                                       | 0.2955          |
| SARS-CoV-2 vaccination (at least 2 doses) <sup>b</sup> | 12 (57%)                                     | 2 (7%)   | <b>0.0001</b>   |
| Respiratory support <sup>b</sup>                       | VMK 12 (57%)<br>A.A. 9 (43%)                 | CPAP 30 (100%)                                 | <b>0.0001</b>   |
| ARDS development <sup>b</sup>                          | 0 (0%)                                       | 0 (0%)   | -               |
| D-dimer ( $\mu\text{g/l}$ ) <sup>a</sup>               | 1209 (669-1621)                              | 1007 (535-1920)                                | 0.6906          |
| PCR (mg/dl) <sup>a</sup>                               | 4.63 (0.83-6.5)                              | 4.17 (1.4-7.35)                                | 0.6814          |
| Ferritin (ng/ml) <sup>a</sup>                          | 821 (239-1199)                               | 823 (511-1294)                                 | 0,5183          |
| LDH (Ui/L) <sup>a</sup>                                | 284 (199-390)                                | 265 (229-284)                                  | 0.9007          |
| Fibrinogen (mg/dl) <sup>a</sup>                        | 514 (384-555)                                | 444 (362-555)                                  | 0.2244          |
| Lymphocytes ( $10^3/\text{ml}$ ) <sup>a</sup>          | 910 (630-1345)                               | 1055 (763-1465)                                | 0.1622          |

VMK = Venturi Mask, CPAP = Continuous Positive Airway Pressure, A.A. = Ambient Air. <sup>a</sup> Data are expressed as median (interquartile ranges). <sup>b</sup> Data are expressed as number (percentage). Differences in demographic characteristics between mild and severe SARS-CoV-2 infected patients at admission (T0) were evaluated using Mann-Whitney U and Chi-squared tests.  $P < 0.05$  was considered statistically significant.

#### 4. Discussion

Several studies associated SARS-CoV-2 infection with pulmonary fibrosis and ARDS<sup>12,13</sup>. Previous evidence from literature in a plethora of pulmonary diseases reported that decrease of SP-A concentration in BALF was associated with ARDS, while both SP-A and SP-D concentrations were reduced in BALF fluids of patients with pulmonary fibrosis. By contrast, their serum concentration increased in these patients<sup>9,14,15</sup>. Afterwards, elevated plasma SP-D levels were observed in patients experiencing pneumonia after SARS-CoV infection<sup>16</sup> and more recently high serum levels of this protein was associated with severe COVID-19, suggesting its potential utility for the early identification of patients who can face worsening conditions<sup>8,17</sup>. In the light of these considerations, we measured the levels of these two collectins in patients' serum and evaluated their association with the clinical outcome. SP-D, but not SP-A, increased among COVID-19 patients compared to healthy controls during the early phases of infection, while a significant reduction was observed at T1. To better clarify the role of the two soluble lectin members family, we stratified SARS-CoV-2 patients according to mild and severe disease and we observed increased serum levels of SP-D in patients under CPAP compared to patients supported with VMK or without respiratory support at T1. This may be due to an earlier recovery of the latter than the former as they are presumably characterized by less extended lung injury. During COVID-19, the previously SARS-CoV-2 vaccination plays a relevant role in the fight against infection. In line with this, our group of severe patients, requiring more invasive respiratory support, was characterized by a higher frequency of unvaccinated individuals. Our findings suggest a role for SP-D as a predictive marker of COVID-19 outcome, as proposed in other studies, in which this collectin levels were associated with ARDS severity and clinical outcome<sup>7,18,19</sup>. On the other hand, no differences were observed for SP-A serum levels according to COVID-19 severity, as previously observed also by Takenaka et al.<sup>20</sup> The lower validity of SP-A as a marker might be due to its reduced serum levels compared to SP-D, both under physiological and pathological conditions. Despite these findings, the limitations of this study are the small sample size, significant age and vaccination differences between COVID-19 patients and controls and the lacking quantification of these proteins' levels in BALF, to assess their likely reduction corresponding to their serum increase. Further studies might focus on the correlation between such collectins and more objective clinical indicators, such as lung imaging and haemogasanalysis parameters. In conclusion, our study suggests SP-D, but not SP-A, as an eligible marker of COVID-19 pneumonia and the early detection of SP-D serum levels could be determinant for a preventive treatment of COVID-19.

**Author Contributions:** Conceptualization, L.M., V.Z. and L.S.; methodology, L.M., G.B. and L.S.; formal analysis, L.M. and G.B.; investigation, A.L., V.Z. and M.B.; writing—original draft preparation, L.M. and G.B.; writing—review and editing, L.M. and V.Z.; supervision, L.S.; project administration, F.R. and C.M.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board of the Department of Public Health and Infectious Diseases, University of Rome "Sapienza" and Ethics Committee Azienda Policlinico Umberto I of Rome (Approval code: Rif. 6484 Prot. 1100/2021; approval date: 15/12/2021).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The data that support the findings of this study are available from the corresponding authors, L.M. and G.B., upon reasonable request.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Majumder, J.; Minko, T. Recent Developments on Therapeutic and Diagnostic Approaches for COVID-19. *AAPS J.* **2021**;23(1):14.
2. Mason, RJ. Pathogenesis of COVID-19 from a cell biology perspective. *Eur Respir J.* **2020**;55(4):2000607.
3. Chroneos, Z.; Sever-Chroneos, Z.; Shepherd, V. Pulmonary Surfactant: An Immunological Perspective. *Cell Physiol Biochem.* **2010**;25(001):013-026.

4. Ghati, A.; Dam, P.; Tasdemir, D.; et al. Exogenous pulmonary surfactant: A review focused on adjunctive therapy for severe acute respiratory syndrome coronavirus 2 including SP-A and SP-D as added clinical marker. *Curr Opin Colloid Interface Sci.* **2021**;51:101413.
5. Han, S.; Mallampalli, R.K. The role of surfactant in lung disease and host defense against pulmonary infections. *Ann Am Thorac Soc.* **2015**;12:765–774.
6. Hartl, D.; Grieser Ludwig-Maximilians, M. Surfactant protein D in human lung diseases *Eur J Clin Invest.* **2006**;36(6):423-35.
7. Agustama, A.; Surgean Veterini, A.; Utariani, A. Correlation of Surfactant Protein-D (SP-D) Serum Levels with ARDS Severity and Mortality in Covid-19 Patients in Indonesia. *Acta medica academica.* **2022**;51(1), 21–28.
8. Kerget, B.; Kerget, F.; Koçak, A. O.; Kızıltunç, A.; Araz, Ö.; Uçar, E. Y.; Akgün, M. Are Serum Interleukin 6 and Surfactant Protein D Levels Associated with the Clinical Course of COVID-19? *Lung.* **2020**;198(5), 777–784.
9. Wang, K.; Ju, Q.; Cao, J.; Tang, W.; Zhang, J. Impact of serum SP-A and SP-D levels on comparison and prognosis of idiopathic pulmonary fibrosis: A systematic review and meta-analysis. *Medicine (Baltimore).* **2017** Jun;96(23):e7083.
10. European Centre for Disease Prevention and Control. Case Definition for Coronavirus Disease 2019 (COVID-19). 2020. Available online: <https://www.ecdc.europa.eu/en/covid-19/surveillance/case-definition> (accessed on 1 February 2021).
11. World Health Organization. Clinical Management of COVID-19; WHO: Geneva, Switzerland, 2020; pp. 1–62.
12. Polak, S. B.; Van Gool, I. C.; Cohen, D.; von der Thüsen, J. H.; van Paassen, J. A systematic review of pathological findings in COVID-19: A pathophysiological timeline and possible mechanisms of disease progression. *Modern pathology.* **2020**;33(11), 2128–2138.
13. Krynytska, I.; Marushchak, M.; Birchenko, I.; Dovgalyuk, A.; Tokarskyy, O. COVID-19-associated acute respiratory distress syndrome versus classical acute respiratory distress syndrome (a narrative review). *Iranian journal of microbiology.* **2021**;13(6), 737–747.
14. Park, J.; Pabon, M.; Choi, A. M. K.; Siempos, I. I.; Fredenburgh, L. E.; Baron, R. M.; Jeon, K.; Chung, C. R.; Yang, J. H.; Park, C. M.; Suh, G. Y. Plasma surfactant protein-D as a diagnostic biomarker for acute respiratory distress syndrome: Validation in US and Korean cohorts. *BMC pulmonary medicine.* **2017**;17(1), 204.
15. Wu, Y. P.; Liu, Z. H.; Wei, R.; Pan, S. D.; Mao, N. Y.; Chen, B.; Han, J. J.; Zhang, F. S.; Holmskov, U.; Xia, Z. L.; de Groot, P. G.; Reid, K. B.; Xu, W. B.; Sorensen, G. L. Elevated plasma surfactant protein D (SP-D) levels and a direct correlation with anti-severe acute respiratory syndrome coronavirus-specific IgG antibody in SARS patients. *Scandinavian journal of immunology.* **2009**;69(6), 508–515.
16. Tong, M.; Xiong, Y.; Zhu, C.; Xu, H.; Zheng, Q.; Jiang, Y.; Zou, L.; Xiao, X.; Chen, F.; Yan, X.; Hu, C.; Zhu, Y. Serum surfactant protein D in COVID-19 is elevated and correlated with disease severity. *BMC infectious diseases.* **2021**;21(1), 737.
17. Tiezzi, M.; Morra, S.; Seminerio, J.; Van Muylem, A.; Godefroid, A.; Law-Weng-Sam, N.; Van Praet, A.; Corbière, V.; Orte Cano, C.; Karimi, S.; Del Marmol, V.; Bondu, B.; Benjelloun, M.; Lavis, P.; Mascart, F.; van de Borne, P.; Cardozo, A. K. SP-D and CC-16 Pneumoproteins' Kinetics and Their Predictive Role During SARS-CoV-2 Infection. *Frontiers in medicine.* **2022**;8, 761299.
18. Salvioni, L.; Testa, F.; Sulejmani, A.; Pepe, F.; Giorgio Lovaglio, P.; Berta, P.; Dominici, R.; Leoni, V.; Prosperi, D.; Vittadini, G.; Colombo, M.; Fiandra, L. Surfactant protein D (SP-D) as a biomarker of SARS-CoV-2 infection. *Clinica chimica acta; international journal of clinical chemistry.* **2022**;537, 140–145.
19. Jayadi; Airlangga, P. S.; Kusuma, E.; Waloejo, C. S.; Salinding, A.; Lestari, P. Correlation between serum surfactant protein-D level with respiratory compliance and acute respiratory distress syndrome in critically ill COVID-19 Patients: A retrospective observational study. *International journal of critical illness and injury science.* **2022**;12(4), 204–210.
20. Takenaka, H.; Saito, A.; Kuronuma, K.; Moniwa, K.; Nishikiori, H.; Takahashi, S.; Chiba, H. The Soluble Lectin Families as Novel Biomarkers for COVID-19 Pneumonia. *In vivo (Athens, Greece),* **2023**;37(4), 1721–1728.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.