
The Work Ability Index (WAI) in the Healthcare Sector. A Cross-Sectional/Retrospective Assessment of the Questionnaire

[Nicola Magnavita](#)*, [Igor Meraglia](#), Giacomo Viti, Lorenzo Borghese

Posted Date: 19 February 2024

doi: 10.20944/preprints202402.0997.v1

Keywords: disability management; total worker health; psychometrics; medical surveillance; health promotion; musculoskeletal disorders; psychiatric disorders; ageing; gender differences; night work



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

The Work Ability Index (WAI) in the Healthcare Sector—A Cross-Sectional/Retrospective Assessment of the Questionnaire

Nicola Magnavita ^{1,2,*}, Igor Meraglia ¹, Giacomo Viti ¹ and Lorenzo Borghese ²

¹ Department of Life Sciences and Public Health; Università Cattolica del Sacro Cuore, Roma, Italy ; meraglia.igor93@gmail.com; giacomo.viti.med@gmail.com;

² Association of Italian Knights of the Sovereign Military Order of Malta, S. Giovanni Battista Hospital, Rome, Italy; presidente@acismom.it

* Correspondence: nicolamagnavita@gmail.com; Tel.: +393473300367

Abstract: The Work Ability Index (WAI) is the most widely used questionnaire for the self-assessment of working ability. Because of its different applications, shorter versions, and widespread use in healthcare activities, it is worthwhile assessing its characteristics. The WAI was distributed online among the workers of a healthcare company; the results were compared with data contained in the workers' personal health records and with absence registers. 340 out of 575 workers (59.1%) participated; 6.5% of them reported poor work ability. Exploratory factor analysis indicated that the one-factor version best described the characteristics of the WAI. The scores of the complete WAI, the shorter form without the list of diseases, and the minimal one-item version (WAS) had equal distribution and were significantly correlated. The WAI score was inversely related to age and significantly lower in women than in men, but it was higher in night workers than in colleagues due to the probable effect of selective factors. The WAI score was also correlated with absenteeism, but no differences were found between males and females in the average number of absences, suggesting that cultural or emotional factors influence the self-rating of the WAI. Workers tended to overreport illnesses in the online survey compared to data collected during occupational health checks. Musculoskeletal disorders were the most frequently reported illnesses (53%). Psychiatric illnesses affected 21% of workers and had the greatest impact on work ability. Multilevel ergonomic and human factor intervention seems to be needed to recover the working capacity of healthcare workers.

Keywords: disability management; total worker health; psychometrics; medical surveillance; health promotion; musculoskeletal disorders; psychiatric disorders; ageing; gender differences; night work

1. Introduction

The Work Ability Index (WAI) [1] is a questionnaire for the self-assessment of working capacity. It is used in occupational healthcare to reveal how well a worker can perform his or her work. It was developed in the late 1990s by Finnish researchers and is widely used as one of the methods for assessing work ability during medical surveillance and workplace surveys. This tool enables the occupational doctor who is beginning risk prevention activities in a company to rapidly collect information on the occupational health status of workers and subsequently create a well-targeted health plan. Over 97% of Finnish doctors and a substantial share of occupational physicians, epidemiologists, psychologists, and sociologists from all over the world are currently using it for health promotion, population health trends analysis, individual case management, and for calculating work ability across age groups [2]. An abundant quantity of literature bears witness to the diffusion of the questionnaire. As of February 15, 2024, digitizing the term "Work Ability Index" on the PubMed search engine yielded 5,535 articles. The professional category most frequently

investigated with the WAI are healthcare workers, and a meta-analysis indicates that low working capacity patterns are frequent among these workers [3]. Despite the abundance of contributions in the literature, some important elements concerning the structure and application of the questionnaire still need to be clarified.

The first aspect to clarify concerns the psychometric characteristics of the questionnaire. The authors conceived the questionnaire as a unitary instrument that would provide a single score, while some studies have observed that a two- or three-factor solution could be more advantageous [4]. The second relevant aspect regards the diffusion of shortened versions of the WAI, which could be particularly useful in investigations conducted in the workplace, but which require a careful evaluation of the instrument. The third characteristic needing clarification is related to the observation that the WAI score is influenced by age, gender, and type of job. The last important aspect concerns the level of reliability of self-assessments.

Despite the amount of research conducted with this tool, few critical evaluations have associated the administration of the questionnaire with objective data (e.g., the registration of absences due to illness or clinical assessments of workers) to verify the correspondence between subjective answers and objective information. This aspect is very important for the company's occupational doctor who must take the necessary measures to protect the health of workers and must therefore know whether subjective declarations correspond with objective data. Moreover, although several authors have used shorter versions of the questionnaire (e.g., without a list of diseases or with just one component), there are very few studies that have verified the usefulness of such tools. Lastly, it is also important for the occupational doctor to know whether the questionnaire functions in a particular way in specific work sectors, such as the healthcare field, where the greatest number of applications have been made to date.

The WAI includes a list of diseases and seven occupational health characteristics. The final WAI score is calculated as the total unweighted score covering the WAI's seven dimensions, or indicators. These are:

- I. Current work ability compared with lifetime best;
- II. Work ability in relation to the demands of the job;
- III. Number of current diseases diagnosed by a physician;
- IV. Estimated work impairment due to illness;
- V. Sick leave during the past 12 months;
- VI. Personal prognosis of work ability 2 years from now;
- VII. Mental resources.

The combination of dimension values results in a total WAI score that can range from 7 (unable to work) to 49 (full work ability). Based on this score, an individual is classified into standard work ability categories of excellent (WAI 44–49), good (WAI 37–43), moderate (WAI 28–36), and poor (WAI ~27).

The list of diseases is the aspect of the questionnaire that has posed most problems in its application. A lengthy list of 51 medical problems was included in the original WAI, requiring employees to note any illnesses for which they had received a diagnosis. Filling out the questionnaire was too complicated to perform in the workplace where health surveillance must not interfere with production needs. A shortlist of 15 diseases was included in the WAI version introduced by Nubling et al. [5]. This turned out to be comparable to the extended version and is nowadays commonly used in occupational medicine. Despite this simplification, internal consistency studies have shown that most missing responses on the WAI relate to the list of diseases, and that the coherence of the WAI is not significantly influenced by self-reported symptoms and illnesses [6]. Some researchers have therefore proposed solving the question of ambiguous WAI scores resulting from missing answers by using a version of the WAI that does not include the list of diseases [7]. This shortened version, without the list of diseases (WAI_{nodis}), has values ranging from 6 to 42.

An even shorter version of the questionnaire is based on the first question "Assume that your work ability at its best has a value of 10 points. How many points would you give your current work

ability?". This single-item questionnaire, also called Work Ability Score (WAS), has been proposed for use in extensive population surveys [7].

To obtain a more thorough knowledge of the questionnaire, the questions we wanted to answer were:

1. Is the WAI score significantly influenced by the age, gender, or occupational category of the workers?
2. Does the questionnaire have a unifactorial structure, or is it better to recognize two distinct factors within it?
3. Does the simplified questionnaire, without the list of diseases, have an acceptable correlation with the complete version?
4. Does the shortest version of the questionnaire (limited to the first dimension) have an acceptable correlation with the final score obtained from the complete questionnaire?
5. Is the assessment of the demands of the job reliable, i.e., do workers who carry out the same job consider it equally demanding from a physical or psychological point of view?
6. Is there a correspondence between the diseases reported in the questionnaire and those reported to the occupational doctor during the medical examination carried out at the workplace?
7. Are illnesses not diagnosed by a doctor, and which therefore do not contribute to the WAI score, irrelevant for the purposes of assessing work capacity?
8. Is there an association between the WAI and sickness absence?
9. Which group of pathologies determines the greatest impairment of the WAI? And which determines the highest burden of sickness absence?
10. Are mental resources correlated with the presence of mental illnesses?

To answer these questions, we administered the WAI to all the workers of a healthcare company who were exposed to occupational risks and compared the answers with the data collected during medical surveillance and in sickness absence records.

2. Materials and Methods

2.1. Population

In Italy, according to Council Directive 89/391/EEC of 12 June 1989 [8], concerning the implementation of measures aimed at promoting improvements in the safety and health of workers at work, and the Legislative Decree 81/2008, which incorporated the EEC Directive into national legislation [9], workers who are exposed to occupational risks must be subjected to health surveillance in the workplace. During this surveillance, workers of a healthcare company were invited to complete the WAI questionnaire using the SurveyMonkey® online platform. The estimated completion time of the questionnaire was 5 minutes.

All workers exposed to occupational risks and included in the health surveillance register (575) were confidentially contacted by email and invited to participate in the survey. Participation was completely voluntary, no economic incentive was provided for response, and no sanctions were adopted for non-responders. Participants were enrolled between 1st and 30th November 2023. Two weeks after commencing the investigation, a reminder mail was sent to non-responders.

The research was conducted in accordance with the Helsinki Declaration. Participants signed an informed consent and, by signing their personal health file, authorized the analysis of their personal data and the dissemination of the results in a collective anonymous form, according to the principles of confidentiality of occupational medicine and the ICOH code of ethics [10]. Ethics approval was obtained from the Catholic University Ethics Committee (ID 3008). Given the cross-sectional study design, no imputation was applied for missing data and the results were based on completed survey responses.

2.2. Questionnaire and objective data

Via the online platform, workers were invited to provide their name, age, gender, occupational category, and to state whether they worked night shifts. They were then invited to self-assess their working capacity using the Italian version of the WAI [11].

The WAI is made up of a series of questions that consider the resources and health status of the worker as well as the physical and mental demands of work. Answers are evaluated in the seven dimensions mentioned in the Introduction, yielding a score ranging from 7 to 49. The value assigned represents the employee's perception of their own work capacity and, consequently, the work capacity level.

In addition to the score resulting from the complete WAI questionnaire, we calculated the score of the shortened version (with values ranging from 6 to 42), which does not take into account diseases diagnosed by a doctor (WAI_{nodis}), and the shortest version, also known as Work Ability Scale (WAS), that corresponds to the first dimension of the questionnaire, with values ranging from 0 to 10.

For the period 2021 to October 2023, the healthcare company administration provided us with the absence registers for all their employees, including those who freely chose not to participate in the survey. It was the I.M.'s task to reconcile the absences with the employees who responded to the questionnaire.

We also analyzed the workers' personal health files containing the results of medical examinations and health checks conducted in the workplace over the years to evaluate the correspondence between subjective indications and pathologies objectively identified by the occupational doctor.

2.4. Statistics

Firstly, using Cronbach's α , the internal consistency of the WAI and its shortened versions was assessed. Using Kolmogorov-Smirnova and Shapiro-Wilk tests, distribution of the variables of interest (WAI, WAI_{nodis} and WAS) was studied to ascertain whether it was normal. However, for non-normally distributed variables, we preferred to use nonparametric tests since, as reported by Lumley et al. [12], the assumption of normality is only required for small samples, due to the central limit theorem. With sample sizes exceeding 30 (as occurred in our case), violations of the normality assumptions are not a problem, and they become less so as the sample size increases, even when they are extreme.

To estimate the adequacy of substituting the WAI_{nodis} or the WAS for the WAI when assessing work ability, the convergent validity between the two instruments was tested by evaluating the Spearman correlation between the shortened forms and the total WAI score. Then, the construction of a linear regression model enabled us to investigate the concurrent validity of the WAI and its shorter versions. In particular, we looked at whether the scores of the shorter WAI versions predicted the overall WAI score to a reasonable extent.

Exploratory factor analysis, principal components analysis with direct oblimin rotation, and the maximum likelihood technique were used to assess the construct validity of the questionnaire.

Distribution of the WAI score between the different occupational categories was studied using analysis of variance (One-way ANOVA) and Bonferroni post-hoc comparisons.

Simple linear regression was used to evaluate the relationship between the independent variables (gender, age, job category, night shifts, days of absence) and the WAI score.

The IBM/SPSS Statistics for Windows, Version 28.0. (Armonk, NY, USA: IBM Corp) was used for the analyses.

3. Results

An invitation to participate was sent via email to all workers of the healthcare company who were exposed to occupational risks (n=575). 412 workers (71.7% of those eligible) responded to the invitation, but 62 of them were excluded because their responses were incomplete. Of the 340 workers

(59.1% of the cohort) who finally took part, 133 were male (39.1%) and 207 were female (60.9%). The mean age was 46.3 ± 11.6 years.

The average score of the WAI questionnaire was equal to 37.0 ± 5.6 points; median value 38. The distribution of scores was significantly non-normal (Kolmogorov-Smirnov test 0.105, $p < 0.001$; Shapiro-Wilk test 0.951, $p < 0.001$) (Figure 1). The reliability coefficient (Cronbach's α) of the questionnaire was 0.73.




Figure 1. Distribution of the total WAI total score (WAI_tot).

In accordance with the authors' indications [1,13], we classified the workers by dividing them into four groups of increasing ability based on the WAI score. 22 workers (6.5%) reported poor work ability (score 7-27); 119 (35.0%) had moderate work ability (score 28-36); 157 (46.2%) reported good work ability (score 37-43), and 42 (12.4%) showed excellent work ability (score 44-49) (Figure 2).




Figure 2. Classification of workers according to work ability.

3.1. Distribution of work ability by age, gender and occupational category

The WAI was inversely correlated with age (Spearman's rho = -0.294 $p < 0.001$). The WAI score decreased by approximately one point every three years of age.

Females had a significantly worse WAI score than males (36.3±6.0 in female vs. 37.9±4.9 in male workers, Mann-Whitney U Test $p < 0.05$).

The distribution of the WAI score varied significantly between the different occupational categories (ANOVA $p < 0.001$). Nurses reported the lowest work ability, while assistant nurses had the highest average rating of work ability. The Bonferroni test showed that there was a significant difference between nurses and assistant nurses ($p < 0.001$), between nurses and employees ($p < 0.01$), and between assistant nurses and technicians ($p < 0.05$) (Table 1).

Table 1. Work ability in different categories of workers.

Category	WAI (mean±s.d.)	Significant Comparisons ¹
1. Physician (n=38)	37.1±4.7	none
2. Nurse (n=97)	35.3±6.2	2 vs. 3 $p < 0.001$; 2 vs. 4 $p < 0.01$
3. Assistant nurse (n=75)	38.8±5.6	3 vs. 2 $p < 0.001$; 3 vs. 5 $p < 0.05$
4. Clerk (n=50)	38.6±4.8	4 vs. 2 $p < 0.01$
5. Technician (n=80)	36.2±5.4	5 vs. 4 $p < 0.05$

¹ Bonferroni test.

Workers performing night shifts (n=90) reported a significantly better WAI score than others (38.2±5.8 vs. 36.5±5.5, $p < 0.01$). Night workers were significantly younger than other workers (42.4±9.9 vs. 47.7±11.9 years, Student's t test $p < 0.001$) and included a larger proportion of nurses than the others. Limiting the comparison to nurses and assistant nurses (n=172 workers), night workers had a better WAI than those who did not perform night shifts (37.9±6.0 vs. 35.8±6.2, $p < 0.05$) and were also younger (42.3±9.3 vs. 48.9±10.9 years, Student's t test $p < 0.001$).

3.2. Factorial structure of the work ability index

Principal components analysis suggested that the presence of a single factor explained over 42% of the variance. All items showed acceptable loadings (>0.49) within the model. The least variance

was explained by item 5, which accounted for 49% of variance, whilst the most explanatory item was item 2 (74%). An alternative two-factor model was examined to provide further assurance that the one-factor structure was the most appropriate. The two-factor model explained over 55% of the variance. In the two-factor model, the first factor was associated with six indicators, while the second factor (which explained 13% of the variance) was associated with sickness absences (Table 2).

Table 2. Exploratory factor analysis of the work ability index administered to healthcare workers.

Items	Principal component analysis	Principal Component Analysis ¹	
	Factor 1	Factor 1	Factor 2
1. Current work ability compared with lifetime best	0.71	0.73	
2. Current work ability in relation to its demands	0.74	0.75	
3. Number of current diseases diagnosed by a physician	0.62	0.59	
4. Estimated work impairment due to disease	0.70	0.68	
5. Sick leave during the past year (12 months)	0.49		0.83
6. Own prognosis of work ability two years from now	0.55	0.71	
7. Mental resources (feelings of joy, alertness, or optimism)	0.68	0.67	
Variance of the component (%)	42.09	42.09	13.57

¹ Rotation method: Oblimin with Kaiser normalization. Rotation converged in 13 iterations.

3.3. Relationships between the complete WAI and the shortened version.

The shortened version of the questionnaire without the list of diseases (WAI_{nodis}) had a similar distribution to that of the total score (Figure 3). The two scores were strongly correlated with each other (Spearman's $\rho=0.96$ $p<0.001$) (Table 3). Using simple linear regression to calculate the WAI score and taking the WAI_{nodis} score as a predictor, we obtained a standardized coefficient beta = 0.962, and a coefficient of determination (R square) of 0.93.




Figure 3. Distribution of the shortened WAI without the list of diseases (WAI_{nodis}).

Table 3. Spearman's bivariate correlations between the shortened WAI without the list of diseases (WAI_{nodis}), the one-item form (WAS) and the total score of the complete version (WAI_{tot}).

	WAI _{nodis}	WAS	WAI
WAI _{nodis}	1.000	0.832**	0.958**
WAS		1.000	0.789**
WAI			1.000

** . Correlation is significant at the 0.01 level (2-tailed).

3.4. Relationships between the complete WAI and the one-item version.

The first dimension of the WAI, or work ability score (WAS), demonstrating current working capacity compared to lifetime best, produced a similar distribution to the total score obtained from the seven indicators (Figure 4). To interpret the WAS results, the inventors of this method [14] suggested using the same type of categorization as for the WAI, i.e., poor (0–5 points), moderate (6,7), good (8,9), excellent (10).




Figure 4. Distribution of the one-item version of the WAI (WAS).

The correlation between the WAS and the WAI was highly significant (Table 3). In a simple linear regression model, the WAS, inserted as a predictor of the WAI, yielded a standardized coefficient beta = 0.752, with a coefficient of determination (R square) of 0.57.

3.5. Reliability of the question on job characteristics

The second indicator of the WAI, "Work ability in relation to the demands of the job" is based on three questions: the first asks whether the job is psychologically demanding, physically demanding, or both; the other two questions ask workers to evaluate the physical and psychological commitment required by the job. Depending on the answer to the first question, the scores obtained from the last two questions must be corrected, either by increasing them by 50% or by reducing them by 50%. It is therefore important to check whether the answer to the first question is consistent with the work profile of the respondents.

In our sample, most workers (64.4%) described their job as psychologically and physically demanding; 30.6% believed it was psychologically demanding. Only 5% (three nurses, three physiotherapists and 11 assistant nurses) indicated that the work was physically demanding but

without any psychological burden. The tasks assigned to the aforementioned workers were not unlike those performed by other members of the same occupational category.

3.6. Reliability of the reported diseases

The third indicator of the WAI is “Number of current diseases diagnosed by a physician”. The availability of health records made it possible to compare the WAI responses with data collected in personal health files.

First, we verified that the workers who reported suffering from one or more illnesses had declared these conditions during medical examinations carried out in the workplace to assess work fitness. In most cases, the reported disease groups roughly corresponded with the pathologies reported to the occupational doctor. However, we noticed that many workers indicated a greater number of classes of disorders than declared in the routine health checks they undergo to establish fitness for work. In detail, two workers declared suffering from 14 groups of diseases diagnosed by the doctor, one worker declared suffering from 10 groups of diseases and 3 other subjects reported more than 5 groups of diseases; in all these cases, personal health records indicated fewer illnesses. The tendency not to report illnesses during routine medical examinations seemed to depend on the type of illness. Although musculoskeletal disorders were almost always reported during health surveillance, respondents reported other types of diseases less frequently. For example, cancers had a fair degree of under-reporting. In this sample, 4 workers reported a benign or malignant tumor diagnosed by the doctor, but in their personal health records we found only non-cancerous pathologies. Another 12 workers declared that “in their opinion” they suffered from benign or malignant tumors. In the medical records we found a tumor history in 7 cases (2 breast tumors, one thyroid tumor, a neurofibroma, a mastocytosis, a giant cell tumor, and a lymphoma), but no reports of neoplasia in the remaining 5. The workers who had reported the presence of tumors were all declared fit for work, without any limitations.

Another aspect that we were able to compare with the health record data was the first dimension, concerning current work capacity compared to maximum work capacity (WAS). Although five workers declared that they were completely unable to work (WAS=0), their health records enabled us to ascertain that they did not have serious chronic diseases and had never expressed problems in carrying out work tasks in the past. A more thorough investigation, conducted by examining the absence registers and questioning the workers, revealed that they had completed the questionnaire while they were on sick leave but had regularly resumed work after recovering from illness.

Finally, we analyzed the situation of the 22 workers with a WAI score <27. This group, which was classified as “poor” and in need of recovering work ability, was composed of 2 doctors, 12 nurses, 4 assistant nurses, 3 technicians, and 1 white collar worker. For 6 of them, the occupational doctor had expressed limited fitness for work, while all the others had been declared fit for work.

3.7. Relevance of diseases not confirmed by a doctor

In the disease section, the WAI questionnaire provides three response options, relating respectively to absence of disease, the presence of self-diagnosed disease and disease certified by a doctor. We wanted to evaluate whether there was a difference in work capacity between subjects who reported illnesses and colleagues. To make comparisons without being influenced by the number of diseases reported, we used the WAI_{nodis} score (Table 4). The ANOVA analysis of variance and post-hoc comparisons using the Bonferroni test showed that difference in scores was generally significant between the absence and presence of diseases, regardless of whether they had been diagnosed by a doctor or not. Across nearly all disease groups, there was no difference in work ability levels in doctor-diagnosed cases compared to workers’ self-diagnosed cases. Only musculoskeletal disorders were associated with worse work ability in cases diagnosed and treated by a doctor, compared to self-diagnosed disorders. In four disease groups (endocrine and metabolic diseases, diseases of the blood, abortions or neonatal defects in children, and tumors), no significant difference in WAI_{nodis} was found between healthy and sick people.

Table 4. Distribution of work ability (shortened WAldis version without the disease-related score) in relation to the different disease classes. Analysis of variance and Bonferroni post-hoc comparisons.

Class of Disease	0	1	2	Anova <i>p</i>	Bonferroni <i>p</i>		
	No disease N (mean ± s.d.)	Personal diagnosis N (mean ± s.d.)	Medical diagnosis N (mean ± s.d.)		0 vs.1	0 vs.2	1 vs.2
Injury	256 (32.5±4.0)	49 (27.9±5.0)	35 (28.7±4.8)	<0.001	<0.001	<0.001	n.s.
Musculoskeletal	160 (33.8±3.4)	82 (28.4±4.7)	98 (30.1±4.5)	<0.001	<0.001	<0.001	<0.05
Cardiovascular	276 (32.0±4.3)	51 (29.1±5.2)	13 (28.9±5.5)	<0.001	<0.001	<0.05	n.s.
Respiratory	312 (31.7±4.5)	17 (28.2±5.1)	11 (30.1±6.1)	<0.01	<0.01	n.s.	n.s.
Psychiatric	267 (32.7±3.7)	15 (27.9±5.1)	58 (26.5±4.7)	<0.001	<0.001	<0.001	n.s.
Neurological	271 (32.4±4.2)	32 (28.1±4.5)	37 (27.3±4.4)	<0.001	<0.001	<0.001	n.s.
Gastroenteric	264 (32.1±4.5)	38 (29.5±4.3)	38 (28.9±4.7)	<0.001	<0.01	<0.001	n.s.
Genitourinary	302 (31.7±4.7)	24 (29.8±4.0)	14 (29.3±4.6)	<0.05	n.s.	n.s.	n.s.
Skin	288 (31.9±4.4)	28 (28.5±4.7)	24 (28.6±5.3)	<0.001	<0.001	<0.01	n.s.
Cancer	324 (31.5±4.7)	12 (30.3±2.5)	4 (32.5±3.4)	n.s.	n.s.	n.s.	n.s.
Endocrine	277 (31.7±4.6)	48 (30.0±5.1)	15 (31.4±4.3)	n.s.	n.s.	n.s.	n.s.
Blood	324 (31.5±4.6)	11 (30.1±4.2)	5 (28.2±7.8)	n.s.	n.s.	n.s.	n.s.
Abortion	320 (31.5±4.7)	16 (29.5±4.4)	4 (33.5±2.5)	n.s.	n.s.	n.s.	n.s.
Other	316 (31.7±4.5)	16 (27.2±5.4)	8 (30.9±5.9)	<0.01	<0.001	n.s.	n.s.

3.8. Association between diseases and impairment

An analysis of the questionnaire answers enabled us to evaluate the prevalence of different groups of diseases. Most workers (180 out of 340, 53%) reported being affected by musculoskeletal disorders in the back, neck, limbs, or other parts of the body (e.g., neck pain, back pain, sciatica, recurrent pain, rheumatism, arthritis). The third category of problems in order of frequency, but the first in order of impairment, was the presence of mental health disorders (depression, anxiety, sleep problems and burnout), which were reported by 73 respondents (21.5%).

In terms of frequency and impact, other very important illnesses were neurological and sensory diseases (migraine, epilepsy, stroke, neuralgia, eye and hearing diseases, other neurological diseases, 20.3%), gastroenteric diseases (gastritis, gastroduodenal ulcer, gallbladder stones, liver or pancreatic disease, colitis, constipation, 22.4%), cardiovascular diseases (high blood pressure, coronary heart disease, angina, heart attack, heart failure, 18.8%) and skin diseases (eczema, allergies, other skin rashes or varicose veins, 15.3%).

The total number of sick days recorded in the observation period were strongly correlated with age (Pearson's $r = 0.269$, $p < 0.001$). Over the three-year period, no differences were observed in the number of days of absence among the different categories of workers (ANOVA $F = 0.256$, $p = 0.91$). Furthermore, no significant difference was found in the number of days of sickness absence in males (33.6 ± 45.6) and females (30.6 ± 36.5) (Student's $t = 0.544$, $p = 0.59$).

In a simple linear regression model that included as independent variables occupational category, working night shifts, gender, age, and days of absence over the three-year period, only the last three variables were predictors of the WAI score (Table 5).

Table 5. Effect of sex, age, and days of absence on work ability. Simple linear regression.

	Beta	t	p
(Constant)		16,187	,000
Age	-,142	-2,304	,022
Sex	-,123	-2,125	,034
Night shift	,086	1,312	,191
Category	,071	1,124	,262
Absence	-,259	-4,317	,000

3.9. Reliability of reported sickness absences

To verify the reliability of the answers given to the question concerning the number of absences in the previous year, we correlated the value of the corresponding WAI indicator with the days of absence in the period 2021-2023. The two quantities were correlated in a very significant way (Spearman's $\rho = -0.553$, $p < 0.001$).

In Table 6 we have reported the mean days of sick leave taken by workers who reported a medical diagnosis for one of the envisaged classes of disease. The longest mean absence period observed between 2021 and 2023 was in workers reporting a blood disease. This was followed by mean absences for skin and other diseases.

Table 6. Days of absence of workers who indicated they had a medical diagnosis for one of the 14 classes of pathologies.

Medical diagnosis Class of Disease	Days of absence 2021 range (mean \pm s.d.)	Days of absence 2022 range (mean \pm s.d.)	Days of absence 2023 range (mean \pm s.d.)	Days of absence 2021-23 range (mean \pm s.d.)
Injury	0-21 (5.0 \pm 5.2)	0-36 (11.0 \pm 10.1)	0-31 (7.9 \pm 8.3)	1-83 (23.9 \pm 18.5)
Musculoskeletal	0-54 (7.5 \pm 10.4)	0-120 (11.8 \pm 16.8)	0-83 (9.0 \pm 13.4)	1-172 (28.3 \pm 31.5)
Cardiovascular	0-14 (7.0 \pm 4.9)	0-33 (12.6 \pm 10.5)	0-134 (2.7 \pm 39.7)	1-179 (40.4 \pm 49.1)
Respiratory	0-172 (21.7 \pm 53.1)	0-24 (9.7 \pm 8.2)	0-21 (7.4 \pm 7.7)	2-196 (39.8 \pm 52.3)
Psychiatric	0-172 (14.8 \pm 30.6)	0-64 (11.3 \pm 12.6)	0-134 (15.2 \pm 27.0)	1-179 (37.0 \pm 47.0)

Neurological	0-36 (7.3±9.2)	0-45 (13.0±12.7)	0-134 (15.2±27.0)	1-179 (35.5±39.7)
Gastroenteric	0-172 (12.1±29.7)	0-45 (11.4±12.3)	0-83 (10.3±17.3)	2-196 (33.8±42.6)
Genitourinary	0-172 (22.4±47.4)	0-40 (10.9±12.7)	0-83 (10.2±21.7)	1-196 (34.5±60.6)
Skin	0-172 (23.2±45.5)	0-45 (11.3±11.0)	0-41 (8.1±9.2)	2-196 (42.6±48.9)
Cancer	0-10 (4.7±4.2)	0-16 (7.3±8.5)	2-19 (7.5±7.8)	2-35 (19.5±15.6)
Endocrine	0-10 (4.9±4.5)	0-24 (11.0±8.1)	0-66 (11.7±20.7)	2-80 (27.6±23.2)
Blood	0-172 (37.2±75.4)	0-24 (9.2±10.0)	0-19 (7.2±7.6)	2-196 (53.6±80.6)
Abortion	0-6 (2.2±2.8)	0-13 (3.2±6.5)	2-19 (7.2±7.9)	2-35 (12.7±15.4)
Other	0-172 (29.6±63.4)	0-24 (7.0±8.8)	0-19 (6.0±6.6)	2-196 (42.6±69.2)

3.10. Relationship between mental resources and psychiatric diseases

It is reasonable to expect that mental resources, which are the seventh dimension on which the WAI is calculated, are not equally distributed in workers with or without mental disorders.

Analysis of variance indicates that workers who do not have mental disorders have greater mental resources. No difference was observed in resources between those who declared disorders treated by a doctor and self-diagnosed disorders; in both these categories resources were significantly lower than in healthy people (Table 7).

Table 7. Distribution of mental resources among workers with psychiatric disease. Analysis of variance and Bonferroni post-hoc comparisons.

Class of Disease	0	1	2	Anova <i>p</i>	Bonferroni <i>p</i>		
	No psychiatric disease N = 267	Personal diagnosis N = 15	Medical diagnosis N = 58		0 vs.1	0 vs.2	1 vs.2
Mental resources (mean ± s.d.)	3.57±0.61	2.93±0.70	2.83±0.75	<0.001	<0.001	<0.001	n.s.

4. Discussion

Application of the WAI in a healthcare company not only provides the company doctor with an initial picture of the occupational health conditions of the employees, but it also offers useful indications concerning the use of the questionnaire in healthcare activities.

In our experience, the WAI decreases significantly as the age of the workers increases. This association was reported in most previous studies [15–24], although there is no shortage of studies that failed to observe this association [25,26]. Previous authors reported that the decrease in the WAI with ageing is strictly dependent on the type of task assigned [27]. In our sample, however, the role of age is more prevalent than the occupational category, which nevertheless has a certain relevance. Nurses had significantly lower WAI levels than other occupational categories; this result was observed in previous studies [28] and is certainly significant due to the possible consequences on healthcare activities [29]. The worsening of the WAI with age is accompanied by an increase in absences due to illness. Other studies that observed the same phenomenon demonstrated that the worsening of the WAI is followed by an increase in absenteeism [30,31]. The WAI is a recognized predictor of absenteeism [32–34], especially if it is accompanied by stress [35], burnout [36], or depression [37].

Data previously reported in the literature [27,28,38] led us to expect the greater impairment of the self-rated working ability of women compared to men observed in our sample. We also expected to have a greater number of absences for females, given that women have a higher WAI than men. Interestingly, this phenomenon did not occur, since there was no difference in the average number

of days of sickness absence for males and females. This finding suggests that socio-cultural or emotional factors are involved in the self-assessment of work ability. We have planned a further study to evaluate this possibility by analyzing, in addition to the WAI, levels of occupational stress, anxiety and depression in male and female workers. In fact, the literature tells us that the WAI score is associated with occupational stress [39–42] and emotional factors [43–45]. The relationship between stress, psychiatric problems and the WAI is complex and probably cyclical: according to the Stockholm County longitudinal study, after 4 years, compared to their colleagues, workers with a low WAI had an almost two-fold rate of distress [46].

In our sample, night workers were younger and reported greater work ability than those who did not work at night. The finding we obtained was probably the result of selective factors that determined the removal of workers with health problems from night work, while the younger and more capable employees carried out this type of service willingly. Previous studies on the relationship between shift work and work ability had not contradicted this hypothesis. When compared to day work, no undue adverse health effects of shift schedules were observed in German workers [47] and in the Nurses' Early Exit Study conducted in seven European countries [48]. In a Brazilian study, impairment of the WAI was observed only in workers who performed a disproportionate number of night shifts or had a temporary employment status [49]. A more recent study has shown that employees who work a greater number of night shifts are healthier than their colleagues since they have fewer chronic musculoskeletal and cardiorespiratory pathologies, a lower level of oxidation indices (DNA adducts and telomere length) and a higher WAI [50]. Therefore, selective factors appear to be predominant in the relationship between the WAI and the night work of healthcare workers.

An issue frequently addressed in the literature is the mono-factorial or multifactorial nature of the questionnaire. The authors of the WAI conceived this instrument as unitary. In fact, the calculation of a total score derived from the seven components of the WAI assumes the unidimensionality of the questionnaire. The calculation of Cronbach's alpha is also based on this assumption. In the various studies conducted with this tool it ranged from 0.54 to 0.83 [6,51–55], and was 0.73 in our own study. However, it is easy to claim that work ability is a multidimensional concept and that dividing the seven dimensions into different factors could theoretically provide better results. In previous research, the underlying structure of the WAI was studied with a variety of techniques, including sampling strategies, statistical analyses, and the administration of the questionnaire in various languages and cultural contexts. These analyses produced inconsistent findings, making it difficult to interpret the WAI. Some studies confirmed the unifactorial structure of the questionnaire [52], while others recognized the existence of two [54,56,57] or three factors [51,53,55]. Country-specific or language-specific variations in the factor structure of the WAI were demonstrated in a study that involved about 40,000 nurses from several European nations [6]. Furthermore, different occupational categories can cause the heterogeneity observed in the WAI factor structure results. Two psychometric studies [4,57] were concordant in recognizing two factors, one composed of indicators 1,2,6 and 7, the other of indicators 3,4, and 5, referring respectively to "subjective work ability and resources", and "ill-health". In our sample we recognized the unifactorial nature of the questionnaire and did not obtain significant advantages by subdividing into two factors. This result can probably be attributed to the highly sectorial nature of our investigation which was carried out on a single healthcare facility. For many years, the company in which we conducted our study had implemented an active disability management policy according to which health problems were solved by changes in the work plan to adapt the work to the man rather than by terminating employment. This policy is not uncommon in healthcare companies. Under these conditions, the two possible factors underlying the WAI merged into a single concept: the difficulty of performing as effectively as in the past. Summing up, we concluded that researchers who apply the WAI in a healthcare company can consider this tool to be unifactorial.

A phenomenon observed in our sample, and expected in other healthcare companies, was the failure to find any difference between illnesses diagnosed by a doctor and self-diagnosed diseases, both of which have a significant impact on working ability. Healthcare professionals usually have

sufficient competence to diagnose their own pathologies without the need to turn to a physician for confirmation, however, since the instructions of the authors of the questionnaire indicate that only diagnoses made by a doctor contribute to the formation of the final score, this introduces an inherent weakness in the assessment of the work ability of healthcare workers. Based on our experience, the shortened form of the questionnaire without the list of diseases (WAI_{nodis}) could be more effective than the full questionnaire because it contains fewer errors. However, a further study to compare the results of the WAI and the WAI_{nodis} with an external construct is needed to verify this hypothesis.

The shortened forms of the questionnaire (WAI_{nodis} and WAS) were shown to have the same distribution as the total score and to be strongly correlated with it. This suggests that they are acceptable substitutes for the full version and can be used when epidemiological needs require workers to be given a more limited number of questions. Numerous studies confirm the feasibility of using shortened versions to replace the WAI, at least in occupational health research and employee surveys [7,58–60]. We applied the one-item questionnaire in previous investigations on post-Covid [61] and on the return to work of women with breast cancer [62], as well as in previous studies [63].

Without careful monitoring of responses, the online administration of the questionnaire probably favors the over-reporting of pathologies. We therefore paid particular attention to verifying the reliability of the workers' answers by comparing responses obtained from the online questionnaire with data recorded in individual health records. Another critical aspect emerged when some workers declared themselves completely unable to work, but the condition was not confirmed in their health surveillance records. An in-depth investigation revealed that these workers suffered from transitory morbid conditions. A certain ambiguity therefore emerges in the first indicator of the WAI (also known as the WAS) which is interpreted by some as "at this precise moment", by others as "in your current working condition". These problems may occur mainly in online administration of the questionnaire, while misunderstandings can be avoided if the questionnaire is administered immediately before the health check and the doctor can analyze it during the worker's medical examination.

A further critical aspect concerned the definition of the work as being predominantly physical, psychological or both, since this often led to disparity in the answers provided by workers who had the same job tasks. In our opinion, the mechanism devised by the authors to determine the second dimension of the WAI, i.e., attributing a different psychological and physical load according to whether the work is classified as psychologically or physically demanding, is certainly useful for comparing different work sectors, but may introduce a bias if applied to workers from the same healthcare company.

A careful analysis of the responses of the employees in this healthcare company revealed that more than half of them (53%) complained of musculoskeletal disorders that impaired work ability. Mental disorders were reported by over 20% of workers and were associated with the greatest reduction in work ability. It is well known that both disorders are very frequent in healthcare activities, and may be correlated [64,65]. Studies showed that the association of musculoskeletal problems and stress are responsible for a considerable reduction in the WAI in healthcare workers [66]. The high prevalence of both diseases demonstrates the need to address this problem with environmental and organizational solutions, rather than with restrictions on individual fitness for work which can be decided by the occupational doctor. The association between mental resources and psychiatric disease leads us to believe that individual intervention aimed at improving resilience and coping strategies must be combined with structural and administrative ergonomic interventions to improve the quality of work. A systematic review with meta-analysis of multilevel interventions in the workplace has shown that this type of combined action can bring about a significant improvement in work ability [67].

The present study has some limitations, mainly because it was conducted in a single healthcare company, so the results cannot be applied to other occupational sectors. Nevertheless, the method used in this company can be applied elsewhere and lead to confirmation or improvement of our results. Online administration of the questionnaire allows for immediate assessment of the cohort's conditions but may lead to over-reporting or inaccurate reporting by workers in the absence of the

interaction with the occupational doctor that normally occurs when the questionnaire is completed just prior to a medical examination. Our findings have led to new research initiatives to clarify the impact of socio-cultural and personal factors on the WAI and its relationship to job satisfaction and work engagement. In addition, they have encouraged health interventions aimed at improving work from an ergonomic and interpersonal point of view.

5. Conclusions

This study confirms the efficacy of using the WAI for health surveillance since it enables the occupational health practitioner to quickly find out how workers assess their working ability and consequently to draw up a health plan that considers the workers' handicaps and the best ways to remedy them. For healthcare workers, this instrument functions very well as a unitary indicator and there is no need to divide it into sub-scales. Shortened versions (without the list of diseases or limited to a single question) can be useful in epidemiological investigations, although the reporting of diseases can undoubtedly be advantageous for the physician in charge of workers' health. When the WAI is used to study healthcare workers, it should be borne in mind that uncertainty may arise over doctor-diagnosed and self-diagnosed diseases, and it is therefore advisable to consider both conditions. The lower work ability reported by women is not associated with a greater number of absences due to illness. This should lead us to examine the socio-cultural and work-related factors that influence reporting in both genders. We hope that this article can contribute to the evaluation of the questionnaire and its application in healthcare activities.

Author Contributions: Conceptualization, N.M.; methodology, N.M and I.M.; investigation, I. M. and G. V.; data curation, N.M.; writing—original draft preparation, N.M.; writing—review and editing, I.M. and. N.M.; supervision, L.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Catholic University of the Sacred Heart of Rome (protocol code 3008; date of approval 16 April 2020).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Anonymized data can be obtained upon request.

Acknowledgments: We thank the workers who participated in the survey, the Human Resources Service of the Italian Association of the Knights of the Sovereign Military Order of Malta who provided the data on absences and Ms. E.A. Wright who revised the English language.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tuomi, K.; Ilmarinen, J.; Jahkola, A.; Katajarinne, L.; Tulkki, A. *Work Ability Index*, 2nd ed. Finnish Institute of Occupational Health: Helsinki, Finland, 1998.
2. Coomer, K.; Houdmont, J. Occupational health professionals' knowledge, understanding and use of work ability. *Occup Med (Lond)* **2013**, *63*, 405-409. doi: 10.1093/occmed/kqt070.
3. Romero-Sánchez, J.M.; Porcel-Gálvez, A.M.; Paloma-Castro, O.; García-Jiménez, J.; González-Domínguez, M.E.; Palomar-Aumatell, X.; Fernández-García, E. Worldwide prevalence of inadequate work ability among hospital nursing personnel: A systematic review and meta-analysis. *J Nurs Scholarsh* **2022**, *54*, 513-528. doi: 10.1111/jnu.12749.
4. Mateo Rodríguez, I.; Knox, E.C.L.; Oliver Hernández, C.; Daponte Codina, A.; The esTAR Group. Psychometric Properties of the Work Ability Index in Health Centre Workers in Spain. *Int J Environ Res Public Health* **2021**, *18*, 12988. doi: 10.3390/ijerph182412988.
5. Nubling, M.; Hasselhorn, H.M.; Seitsamo, J.; Ilmarinen, J. Comparing the use of the short and the long disease list in the Work Ability Index Questionnaire. In: *Assessment and Promotion of Work Ability, Health, and Wellbeing of Ageing Workers*, Costa, G., Goedhard, W., Ilmarinen, J., Eds.; Elsevier: Amsterdam, 2004, 292–295.

6. Radkiewicz, P.; Widerszal-Bazyl, M. Psychometric properties of Work Ability Index in the light of comparative survey study. *Int Congr Ser* **2005**, *1280*, 304–309. <https://doi.org/10.1016/j.ics.2005.02.089>. 23.
7. Schouten, L.S.; Bültmann, U.; Heymans, M.W.; Joling, C.I.; Twisk, J.W.; Roelen, C.A. Shortened version of the work ability index to identify workers at risk of long-term sickness absence. *Eur J Public Health* **2016**, *26*, 301-5. doi: 10.1093/eurpub/ckv198.
8. European Union. Council Directive 89/391/EEC of 12 June 1989. Available at: <https://eur-lex.europa.eu/legal-content/IT/TXT/?uri=CELEX:31989L0391>. Accessed January 24, 2024.
9. Legislative Decree 9 April 2008, n. 81 Implementation of article 1 of law 3 August 2007, n. 123, regarding the protection of health and safety in the workplace. Gazzetta Ufficiale della Repubblica Italiana. Serie Generale n.101 del 30-04-2008 - Suppl. Ordinario n. 108. Available at: <https://www.gazzettaufficiale.it/eli/gu/2008/04/30/101/so/108/sg/pdf>. Accessed 24 January, 2024.
10. ICOH. International Commission on Occupational Health. Code of ethics. Available at: <https://www.icohweb.org/site/code-of-ethics.asp>. Accessed January 24, 2024.
11. Converso, D.; Sottimano, I.; Garzaro, G.; Guidetti, G.; Pira, E.; Viotti, S. Contribution to validation of the Italian version of work ability personal radar. *Med Lav* **2022**, *113*, e2022051. doi: 10.23749/mdl.v113i6.13470.
12. Lumley, T.; Diehr, P.; Emerson, S.; Chen, L. The importance of the normality assumption in large public health data sets. *Annu Rev Public Health* **2002**, *23*, 151-169. doi:10.1146/annurev.publhealth.23.100901.140546.
13. Ilmarinen, J. The work ability index (WAI). *Occup Med* **2007**, *57*, 160.
14. Gould, R.; Ilmarinen, J.; Järvisalo, J. Dimensions of work ability. Results from the Health 2000 Survey. Helsinki (Finland): Finnish Centre for Pensions, **2008**, 25–34. *Occup Environ Health* **2008**, *81*, 495–501. doi: 10.1007/s00420-007-0222-z.].
15. Gajewski, P.D.; Rieker, J.A.; Athanassiou, G.; Bröde, P.; Claus, M.; Golka, K.; Hengstler, J.G.; Kleinsorge, T.; Nitsche, M.A.; Reinders, J.; Tisch, A.; Watzl, C.; Wascher, E.; Getzmann, S. A Systematic Analysis of Biological, Sociodemographic, Psychosocial, and Lifestyle Factors Contributing to Work Ability Across the Working Life Span: Cross-sectional Study. *JMIR Form Res* **2023**, *7*, e40818. doi: 10.2196/40818.
16. Suorsa, K.; Mattila, V.M.; Leskinen, T.; Heinonen, O.J.; Pentti, J.; Vahtera, J.; Stenholm, S. Work ability and physical fitness among aging workers: the Finnish Retirement and Aging Study. *Eur J Ageing* **2022**, *19*, 1301-1310. doi: 10.1007/s10433-022-00714-1.
17. Ezzatvar, Y., Calatayud, J., Andersen, L.L., Vieira, E.R., López-Bueno, R., Casaña, J. Muscular Fitness and Work Ability among Physical Therapists. *Int J Environ Res Public Health* **2021**, *18*, 1722. doi: 10.3390/ijerph18041722.
18. Rypicz, Ł.; Wiczak, I.; Rosińczuk, J.; Karniej, P.; Kołcz, A. Factors affecting work ability index among Polish nurses working in hospitals - A prospective observational survey. *J Nurs Manag* **2021**, *29*, 468-476. doi: 10.1111/jonm.13192.
19. Firoozeh, M.; Saremi, M.; Kavousi, A.; Maleki, A. Demographic and occupational determinants of the work ability of firemen. *J Occup Health* **2017**, *59*, 81-87. doi: 10.1539/joh.15-0296-FS. Erratum in: *J Occup Health*. 2017;59(2):e1.
20. Cotrim, T.P.; Ribeiro, C.; Teles, J.; Reis, V.; Guerreiro, M.J.; Janicas, A.S.; Candeias, S.; Costa, M. Monitoring Work Ability Index During a Two-Year Period Among Portuguese Municipality Workers. *Int J Environ Res Public Health* **2019**, *16*, 3674. doi: 10.3390/ijerph16193674.
21. Cotrim, T.; Soares, G.; Ferreira, P.; Barnabé, R.; Teles, J.; Prata, N. Measuring psychosocial factors and predicting work ability among cemetery workers. *Work* **2020**, *65*, 111-119. doi: 10.3233/WOR-193063.
22. Kim, J.W.; Yang, S.; Chung, I.; Lee, M.Y. Factors that determine the Work Ability Index of street cleaners. *Yeungnam Univ J Med* **2019**, *36*, 219-224. doi: 10.12701/yujm.2019.00199.
23. Garzaro, G.; Sottimano, I.; Di Maso, M.; Bergamaschi, E.; Coggiola, M.; Converso, D.; Iavicoli, S.; Pira, E.; Viotti, S. Work Ability among Italian Bank Video Display Terminal Operators: Socio-Demographic, Lifestyle, and Occupational Correlates. *Int J Environ Res Public Health* **2019**, *16*, 1653. doi: 10.3390/ijerph16091653.
24. Thanapop, S.; Thanapop, C. Work ability of Thai older Workers in Southern Thailand: a comparison of formal and informal sectors. *BMC Public Health* **2021**, *21*, 1218. doi: 10.1186/s12889-021-10974-8.
25. Padula, R.S.; da Silva Valente Ldo, S.; de Moraes, M.V.; Chiavegato, L.D.; Cabral, C.M. Gender and age do not influence the ability to work. *Work* **2012**, *41*, 4330-4332. doi: 10.3233/WOR-2012-0727-4330.
26. Žmauc, T.; Železnik, D.; Težak, O. Work ability index in Slovenian hospital nurses aged over fifty years. *Arh Hig Rada Toksikol* **2019**, *70*, 265-272. doi: 10.2478/aiht-2019-70-3291.
27. Garzaro, G.; Clari, M.; Ciocan, C.; Albanesi, B.; Guidetti, G.; Dimonte, V.; Sottimano, I. Physical Health and Work Ability among Healthcare Workers. A Cross-Sectional Study. *Nurs Rep* **2022**, *12*, 259-269. doi: 10.3390/nursrep12020026.
28. Ronchese, F.; Ricci, F.; Peccolo, G.; Persechino, B.; Rondinone, B.M.; Buresti, G.; Negro, C.; Bovenzi, M.; Miani, A. Relation of the work ability index to fitness for work in healthcare and public employees in a region of Northeastern Italy. *Int Arch Occup Environ Health* **2023**, *96*, 1203-1210. doi: 10.1007/s00420-023-02001-7.

29. Abbasi, M.; Zakerian, A.; Kolahdouzi, M.; Mehri, A.; Akbarzadeh, A.; Ebrahimi, M.H. Relationship between Work Ability Index and Cognitive Failure among Nurses. *Electron Physician* **2016**, *8*, 2136-43. doi: 10.19082/2136.
30. Ohta, M.; Higuchi, Y.; Kumashiro, M.; Yamato, H.; Sugimura, H. Decrease in Work Ability Index and sickness absence during the following year: a two-year follow-up study. *Int Arch Occup Environ Health* **2017**, *90*, 883-894. doi: 10.1007/s00420-017-1251-x.
31. Palmlöf, L.; Skillgate, E.; Talbäck, M.; Josephson, M.; Vingård, E.; Holm, L.W. Poor work ability increases sickness absence over 10 years. *Occup Med (Lond)* **2019**, *69*, 359-365. doi: 10.1093/occmed/kqz083.
32. Lundin, A.; Leijon, O.; Vaez, M.; Hallgren, M.; Torgén, M. Predictive validity of the Work Ability Index and its individual items in the general population. *Scand J Public Health* **2017**, *45*, 350-356. doi: 10.1177/1403494817702759
33. Reeuwijk, K.G.; Robroek, S.J.; Niessen, M.A.; Kraaijenhagen, R.A.; Vergouwe, Y.; Burdorf, A. The Prognostic Value of the Work Ability Index for Sickness Absence among Office Workers. *PLoS One* **2015**, *10*, e0126969. doi: 10.1371/journal.pone.0126969.
34. Notenbomer, A.; Groothoff, J.W.; van Rhenen, W.; Roelen, C.A. Associations of work ability with frequent and long-term sickness absence. *Occup Med (Lond)* **2015**, *65*, 373-9. doi: 10.1093/occmed/kqv052.
35. Oberlinner, C.; Yong, M.; Nasterlack, M.; Pluto, R.P.; Lang, S. Combined effect of back pain and stress on work ability. *Occup Med (Lond)* **2015**, *65*, 147-53. doi: 10.1093/occmed/kqu190.
36. Schouteten, R. Predicting absenteeism: screening for work ability or burnout. *Occup Med (Lond)* **2017**, *67*, 52-57. doi: 10.1093/occmed/kqw161.
37. Pranjic, N.; Males-Bilic, L. Work ability index, absenteeism and depression among patients with burnout syndrome. *Mater Sociomed* **2014**, *26*, 249-52. doi: 10.5455/msm.2014.249-252.
38. La Torre, G.; Grima, D.; Romano, F.; Polimeni, A. Perceived work ability and work-family conflict in healthcare workers: An observational study in a teaching hospital in Italy. *J Occup Health* **2021**, *63*, e12271. doi: 10.1002/1348-9585.12271.
39. Li, H.; Liu, Z.; Liu, R.; Li, L.; Lin, A. The relationship between work stress and work ability among power supply workers in Guangdong, China: a cross-sectional study. *BMC Public Health* **2016**, *16*, 123. doi: 10.1186/s12889-016-2800-z.
40. Yong, M.; Nasterlack, M.; Pluto, R.P.; Lang, S.; Oberlinner, C. Occupational stress perception and its potential impact on work ability. *Work*, **2013**, *46*, 347-54. doi: 10.3233/WOR-121556.
41. Amirmahani, M.; Hasheminejad, N.; Tahernejad, S.; Reza Tohidi Nik, H. Evaluation of work ability index and its association with job stress and musculoskeletal disorders among midwives during the Covid-19 pandemic. *Med Lav* **2022**, *113*, e2022031. doi: 10.23749/ml.v113i4.12834.
42. Kazemi, R.; Zamanian, Z.; Khalifeh, M.; Hemmatjo, R. The Effects of Noise and Heat Strain on the Work Ability Index (WAI) among Rubber Factory Workers. *Ann Glob Health* **2019**, *85*, 96. doi: 10.5334/aogh.2515.
43. Bernburg, M.; Vitzthum, K.; Groneberg, D.A.; Mache, S. Physicians' occupational stress, depressive symptoms and work ability in relation to their working environment: a cross-sectional study of differences among medical residents with various specialties working in German hospitals. *BMJ Open* **2016**, *6*, e011369. doi: 10.1136/bmjopen-2016-011369.
44. Abdelrehim, M.G.; Eshak, E.S.; Kamal, N.N. The mediating role of work-family conflicts in the association between work ability and depression among Egyptian civil workers. *J Public Health (Oxf)* **2023**, *45*, e175-e183. doi: 10.1093/pubmed/fdac061.
45. Björkelund, C.; Saxvik, A.; Svenningsson, I.; Petersson, E.L.; Wiegner, L.; Larsson, M.; Törnbom, K.; Wikberg, C.; Ariai, N.; Nejati, S.; Hensing, G.; Hange, D. Rehabilitation cooperation and person-centred dialogue meeting for patients sick-listed for common mental disorders: 12 months follow-up of sick leave days, symptoms of depression, anxiety, stress and work ability - a pragmatic cluster randomised controlled trial from the CO-WORK-CARE project. *BMJ Open* **2023**, *13*, e074137. doi: 10.1136/bmjopen-2023-074137.
46. Onell, C.; Holm, L.W.; Bohman, T.; Magnusson, C.; Lekander, M.; Skillgate, E. Work ability and psychological distress in a working population: results from the Stockholm Public Health Cohort. *Scand J Public Health* **2023**, *51*, 595-601. doi: 10.1177/14034948211033692
47. Yong, M.; Nasterlack, M.; Pluto, R.P.; Elmerich, K.; Karl, D.; Knauth, P. Is health, measured by work ability index, affected by 12-hour rotating shift schedules? *Chronobiol Int* **2010**, *27*, 1135-48. doi: 10.3109/07420528.2010.490111.
48. Camerino, D.; Conway, P.M.; Sartori, S.; Campanini, P.; Estryn-Béhar, M.; van der Heijden, B.I.; Costa, G. Factors affecting work ability in day and shift-working nurses. *Chronobiol Int* **2008**, *25*, 425-442. doi: 10.1080/07420520802118236.
49. Rotenberg, L.; Griep, R.H.; Fischer, F.M.; Fonseca Mde, J.; Landsbergis, P. Working at night and work ability among nursing personnel: when precarious employment makes the difference. *Int Arch Occup Environ Health* **2009**, *82*, 877-85. doi: 10.1007/s00420-008-0383-4.
50. Pavanello, S.; Stendardo, M.; Mastrangelo, G.; Casillo, V.; Nardini, M.; Mutti, A.; Campisi, M.; Andreoli, R.; Boschetto, P. Higher Number of Night Shifts Associates with Good Perception of Work Capacity and

- Optimal Lung Function but Correlates with Increased Oxidative Damage and Telomere Attrition. *Biomed Res Int* **2019**, 8327629. doi: 10.1155/2019/8327629.
51. Abdolalizadeh, M.; Arastoo, A.A.; Ghsemzadeh, R.; Montazeri, A.; Ahmadi, K.; Azizi, A. The psychometric properties of an Iranian translation of the Work Ability Index (WAI) questionnaire. *J Occup Rehabil* **2012**, *22*, 401–408. <https://doi.org/10.1007/s10926-012-9355-3>.
 52. Bethge, M.; Radoschewski, F.M.; Gutenbrunner, C. The Work Ability Index as a screening tool to identify the need for rehabilitation: longitudinal findings from the Second German Sociomedical Panel of Employees. *J Rehabil Med* **2012**, *44*, 980–987. <https://doi.org/10.2340/16501977-1063>.
 53. Martinez, M.C.; Latorre Mdo, R.; Fischer, F.M. Validity and reliability of the Brazilian version of the Work Ability Index questionnaire. *Rev Saude Publica* **2009**, *43*, 525–532. <https://doi.org/10.1590/S0034-89102009005000017>.
 54. Martus, P.; Jakob, O.; Rose, U.; Seibt, R.; Freude, G. A comparative analysis of the Work Ability Index. *Occup Med* **2010**, *60*, 517–524. <https://doi.org/10.1093/occmed/kqq093>.
 55. Peralta, N.; Godoi Vasconcelos, A.G.; Härter Griep, R.; Miller, L. Validity and reliability of the Work Ability Index in primary care workers in Argentina. *Salud Colectiva* **2012**, *8*, 163–173.
 56. Alexopoulos, E.; Merekoulias, G.; Gnardellis, C.; Jelastopulu, E. Work Ability Index: validation of the Greek version and descriptive data in heavy industry employees. *Br J Med Med Res* **2013**, *3*, 608–621. <https://doi.org/10.9734/bjmmr/2013/2552>.
 57. Freyer, M.; Formazin, M.; Rose, U. Factorial Validity of the Work Ability Index Among Employees in Germany. *J Occup Rehabil* **2019**, *29*, 433–442. doi: 10.1007/s10926-018-9803-9.
 58. Ebener, M.; Hasselhorn, H.M. Validation of Short Measures of Work Ability for Research and Employee Surveys. *Int J Environ Res Public Health* **2019**, *16*, 3386. doi: 10.3390/ijerph16183386.
 59. Mokarami, H.; Cousins, R.; Kalteh, H.O. Comparison of the work ability index and the work ability score for predicting health-related quality of life. *Int Arch Occup Environ Health* **2022**, *95*, 213–221. doi: 10.1007/s00420-021-01740-9.
 60. El Fassi, M.; Bocquet, V.; Majery, N.; Lair, M.L.; Couffignal, S.; Mairiaux, P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health* **2013**, *13*, 305. doi: 10.1186/1471-2458-13-305.
 61. Magnavita, N.; Arnesano, G.; Di Prinzio, R.R.; Gasbarri, M.; Meraglia, I.; Merella, M.; Vacca, M.E. Post-COVID Symptoms in Occupational Cohorts: Effects on Health and Work Ability. *Int J Environ Res Public Health* **2023**, *20*, 5638. doi: 10.3390/ijerph20095638
 62. Magnavita, N.; Di Prinzio, R.R.; Meraglia, I.; Vacca, M.E.; Arnesano, G.; Merella, M.; Mauro, I.; Iuliano, A.; Terribile, D.A. Supporting Return to Work after Breast Cancer: A Mixed Method Study. *Healthcare* **2023**, *11*, 2343. <https://doi.org/10.3390/healthcare11162343>
 63. Magnavita, N. Workplace Health Promotion Embedded in Medical Surveillance: The Italian Way to Total Worker Health Program. *Int J Environ Res Public Health* **2023**, *20*, 3659. doi: 10.3390/ijerph20043659.
 64. Bernal, D.; Campos-Serna, J.; Tobias, A.; Vargas-Prada, S.; Benavides, F.G.; Serra, C. Work-related psychosocial risk factors and musculoskeletal disorders in hospital nurses and nursing aides: a systematic review and meta-analysis. *Int J Nurs Stud* **2015**, *52*, 635–48. doi: 10.1016/j.ijnurstu.2014.11.003.
 65. Del Campo, M.T.; Romo, P.E.; de la Hoz, R.E.; Villamor, J.M.; Mahillo-Fernández, I. Anxiety and depression predict musculoskeletal disorders in health care workers. *Arch Environ Occup Health* **2017**, *72*, 39–44. doi: 10.1080/19338244.2016.1154002
 66. Lindegård, A.; Larsman, P.; Hadzibajramovic, E.; Ahlborg, G. Jr. The influence of perceived stress and musculoskeletal pain on work performance and work ability in Swedish health care workers. *Int Arch Occup Environ Health* **2014**, *87*, 373–9. doi: 10.1007/s00420-013-0875-8.
 67. Oakman, J.; Neupane, S.; Proper, K.I.; Kinsman, N.; Nygård, C.H. Workplace interventions to improve work ability: A systematic review and meta-analysis of their effectiveness. *Scand J Work Environ Health* **2018**, *44*, 134–146. doi: 10.5271/sjweh.3685.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.