

Review

Not peer-reviewed version

---

# The Fossil Record and Diversity of Pycnodondiform Fishes in Non-Marine Environments

---

[Jürgen Kriwet](#)<sup>\*</sup> and John Joseph Cawley

Posted Date: 20 March 2024

doi: 10.20944/preprints202403.1176.v1

Keywords: Pycnodonts, freshwater, brackish, Jurassic, Cretaceous, extinction, refuge.



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

# The Fossil Record and Diversity of Pycnodondiform Fishes in Non-Marine Environments

John J. Cawley<sup>1</sup> and Jürgen Kriwet<sup>1,2,\*</sup>

<sup>1</sup> Department of Palaeontology, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

<sup>2</sup> Vienna Doctoral School of Ecology and Evolution (VDSEE), University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria

\* Correspondence: juergen.kriwet@univie.ac.at

**Abstract:** Pycnodont fishes were a successful clade of neopterygian fishes that are predominantly found from shallow marine deposits. However, throughout their long 180 million years reign (Late Triassic - end Eocene), they have made multiple incursions into both brackish and freshwater environments. This fossil record mostly consists of fragmentary dental material but articulated specimens are known from Early Cretaceous lacustrine localities in Spain. This review article aims to document all non-marine occurrences of Pycnodontiformes throughout most of the Mesozoic and early Palaeogene. This review highlights two interesting trends in the history of non-marine habitat colonization by pycnodonts: (1) a huge spike in non-marine occurrences during the Cretaceous, (2) the most occurrences in non-marine localities occurred at the latest Cretaceous, the Maastrichtian. The high number of colonization events within the Cretaceous lines up with extreme climatic events such as high temperatures resulting in high sea levels which regularly flooded continental masses, allowing pycnodonts easier access to non-marine habitats. The increased presence of pycnodonts in brackish and freshwater habitats during the Maastrichtian might have played a role in their survival through the K/Pg extinction event. Freshwater habitats are not as vulnerable as marine ecosystems to environmental disturbance as the base of their food chain relies on detritus. Pycnodonts might have used such environments as a refuge and began to occupy marine waters after the K/Pg extinction event.

**Keywords:** Pycnodonts; freshwater; brackish; Jurassic; Cretaceous; extinction; refuge

## 1. Introduction

Despite making up less than 0.3% of available global water masses, freshwater environments contain a staggering diversity of fish species comprising approximately 25% of all known vertebrates [1] and 51% of all extant fishes. This high diversity in such a small area of habitat is due to high productivity, geographical isolation and various physiographic processes [2,3] such as light [4], thermal zonation [5] and oxygen concentration [6]. However, the fossil record of freshwater fishes is of substantially poorer quality than their marine counterparts and thus, far less is known about the evolutionary history of many freshwater fish lineages. This relative paucity of fossil material is due to various environmental conditions such as strong currents and greater risk of drying out during arid seasons. Additionally, the freshwater fish fossil record is biased towards fish in lentic habitats (e.g., stillwater such as lakes and ponds) contrary to those from lotic habitats (moving waters such as rivers and torrents). This is powerfully demonstrated by the fossil record of two freshwater teleost lineages from the Neotropical region, i.e., Characiformes and Siluriformes. The fossil record of characiforms consists of fragmentary material such as isolated teeth and scales testifying their occurrence in sediments representing faster moving water, whilst fossil siluriforms are more commonly represented by complete fossils and are typically found in stillwater environments [7]. However, a factor to also consider is that siluriforms have denser and thicker bones than characiforms, making them more amenable to fossilization. Many neotropical siluriforms are also armoured in some manner whether they possess pectoral and dorsal spines or dermal plates, increasing preservation chances further.

While many fish clades today are strictly freshwater [see 8] there are many predominantly marine clades that have made incursions into and adapted to freshwaters such as members of Belontiidae, Tetraodontidae, Acipenseridae, Mugilidae, and Gobiidae. Many other clades include diadromous taxa, i.e., moving in between marine and freshwater environments during their life history (9). Pertinent examples are members of Salmoniformes and Clupeidae moving from marine environments into freshwaters to breed (anadromy) and freshwater eels of the family Anguillidae that migrate from marine environments as larvae and juveniles to rivers and lakes where they spend their adult life (catadromy).

Considering the history of any particular freshwater fish clade, it is rare that the clade in question originated in freshwater and it has been noted by Cavin [10] that most basal taxa for any freshwater clade occurred in marine waters. Among 15 present day families that had their origins in the Mesozoic, Patterson [11] only identified hiodontids, which are members of Osteoglossomorpha, as having originated and stayed in freshwater throughout their evolutionary history. During the course of their evolutionary history, many invasions of fishes into freshwater environments from lineages originating in marine waters have occurred. Some examples include Amiiformes [12], Ginglymodi [13], Acipenseriformes [14] and Coccolepididae [15].

One of the most successful lineages of fishes that lived during the Mesozoic were the pycnodontiforms or pycnodonts [16]. These fishes were considered as stem teleosts by Nursall [17], but a more recent phylogenetic analysis by Poyato-Ariza [18] showed pycnodonts to be rather positioned on the stem of Neopterygii.

Pycnodonts are most commonly associated with shallow marine environments such as reefs and appear to be specialized for such complex, structured habitats due to the deep bodied morphology with large medial fins (Figure 1) that are common to this order and are morphologically convergent with many extant coral reef fish families such as chaetodontids and balistids [19]. Cawley et al. [16] hypothesized that pycnodont fish diversity may be correlated with reef areas as their diversity was low when area occupied by reefs was minimal, but was high when reef area occupation increased. However, such a body plan also can be useful for other complex environments such as well vegetated freshwater biotopes and pycnodonts seemingly have repeatedly invaded brackish and freshwater environments multiple times throughout their history. Poyato-Ariza [20] consequently advised not to use pycnodonts as indicator taxa for any particular palaeoenvironment such as reefs due to occurrence of some in freshwater environments. Nevertheless, the true nature of these alleged freshwater pycnodonts has been since discussed. The review presented here aims to (1) compile all available pycnodont occurrences in brackish or freshwater deposits, (2) how this can inform about their palaeobiogeography through geological time and (3) discuss the occurrence of fully freshwater adapted pycnodonts.

## 2. Nonmarine occurrences

### 2.1 Middle Jurassic

#### 2.1.1 Thailand

The earliest records of pycnodonts occurring in non-marine environments are difficult to accurately determine but can be traced back to the Middle Jurassic. The only definite record from this time interval to our knowledge is an isolated prearticular dentition from the Khlong Min Formation in Thailand that was presented by Cavin et al. [21] and identified as cf. *Gyrodus*. The Mab Ching locality has only been dated to be Middle to Late Jurassic in age based on charophytes [22,23] and fossil turtles [24] but unfortunately, more precise dating has not been inferred for this locality. A Middle Jurassic age for the pycnodont remain, nevertheless, seems likely since the Khlong Min Fm. is at the base of the stratigraphic sequence.

### 2.2 Late Jurassic

#### 2.2.1 China

The Asian pycnodont record is very poor. So far, the only Late Jurassic Asian pycnodont that was reported from freshwater deposits is the little known *Tibetodon gyroides* from Changtu in the Tibetan highlands [25,26]. This taxon only is known by an isolated vomerine dentition exposing five

rows of regularly arranged teeth and its age cannot be determined any further than being of Late Jurassic age. Young and Liu [25] assumed this taxon to be a member of the family Gyrodontidae due to the dental similarities with *Gyrodus* such as the depressed pit in the occlusal surface of the teeth. However, they also argue that due to its widely separated geographic position from most other pycnodonts around this time, it is also possible for this taxon to belong to a different family. The morphology of the teeth actually differs from that of typical members of Gyrodontidae supporting the latter interpretation.



**Figure 1.** Holotype of *Ocloedus subdiscus* (MNHN MSE 301G) from the Early Cretaceous (Berriasian-Valanginian) El Montsec, Spain. This species represents one of the few articulated specimens of a pycnodont from a freshwater environment. Collection abbreviations: MNHN - Muséum national d'Histoire naturelle, Paris Scale bar is 1 cm.

### 2.2.2. North America

The only Late Jurassic records of freshwater pycnodonts from North America include a single broken tooth belonging to an undescribed pycnodontiform from the Morrison Formation (USA), which is of Oxfordian-Tithonian age [27]. It was assigned to Pycnodontoidea due to its highly ornamented crown surface and oval shape. The preserved portion of the tooth displays at least seven small cusplets surrounding a median occlusal depression. Such ornate teeth are found in pycnodontiforms such as *Turbomesodon arcuatus* from the Berriasian (Early Cretaceous) of South England and NW Germany [28,29], as well as *Nonophalagodus trinitiensis* and *Texasensis coronatus* from the Albian of North America [30,31] (see below for more information on these pycnodontiforms).

### 2.2.3. Africa

The only Late Jurassic African pycnodont non-marine record is the dubious *Congopycnodus cornutus* from the lacustrine Stanleyville Formation in the Democratic Republic of Congo. This taxon was based on the occipital region of the skull and a nuchal horn [32]. Previously, this formation was assumed to be Aalenian-Bathonian in age, which would have made this fish the oldest freshwater occurrence for pycnodonts. However, there is strong evidence that the Stanleyville Formation is actually Late Jurassic to Early Cretaceous (Kimmeridgian to Barremian-Valanginian) in age see [33,34] for arguments. While the material of *C. cornutus* is fragmentary, the association of such bones together indicated that these remains represent the earliest known representative of the Coccodontoidea [35], a pycnodont clade containing highly derived armored pycnodonts such as Coccodontidae [36,37] Gladiopycnodontidae [35] and Gebrayelichthyidae [38], which are otherwise only found in the

famous Cenomanian (Upper Cretaceous) marine limestone deposits of Lebanon. Whether this bizarre clade had its origins in freshwater environments or *Congopycnodus* was an outlier within the early horned pycnodonts remains ambiguous momentarily requiring more and better-preserved material including the characteristic dentition. We consider the material not indisputably enough to unambiguously attribute it to a pycnodontiform at all.

An isolated vomerine dentition referred to as cf. *Pycnodus* was reported from the Mughar Mudstone of the Abay (Blue River) Gorge, northwestern Ethiopian plateau, which is of ?Kimmeridgian–Tithonian age [39]. The depositional environment of the Mughar Mudstone is interpreted as representing meandering river deposits on a coastal plain [40,41]. The tentative assignment to *Pycnodus* has to be considered incorrect since members of *Pycnodus* are exclusively restricted to the Palaeogene [42]. This occurrence within a fluvial environment on a coastal plain does not support this pycnodont to be a true freshwater fish, but also could represent a sporadic marine invader.

### 2.3. Early Cretaceous


In contrast to the Jurassic, the records of freshwater pycnodonts increased during the Early Cretaceous with a wide geographic distribution including some taxa, which have fully preserved articulated specimens. This pattern is in good accordance with the fossil record of other fishes, for which the Early Cretaceous seemingly was an important time interval for adapting to freshwater environments e.g., [10,43].

#### 2.3.1. Thailand

Small incomplete vomerine dentitions and isolated teeth assigned to cf. *Anomoeodus* were found from the Phu Phan Thong locality in the Sao Khua Formation in Thailand [21]. The teeth are of Hauterivian-Barremian age. Cavin et al. [21] assigned this material to cf. *Anomoeodus* due to the similarities of the shape and ornamentation of the teeth shared with *Anomoeodus* which was widespread throughout the Cretaceous [44]. Of particular interest is that all specimens of cf. *Anomoeodus* found in this locality are tiny, suggesting that this pycnodont travelled into freshwater environments to breed and their young would have developed in freshwater habitats before returning to sea as an adult [45].

#### 2.3.2. United Kingdom

Pycnodont records from non-marine deposits of the United Kingdom come exclusively from England. Hornung [29] presented a revision of Early Cretaceous pycnodonts not only from Germany (see below), but also from England, which includes various fragmentary prearticulars and vomers previously referred to as *Coelodus*. According to this author all specimens can either be assigned to *Turbomesodon arcuatus* (Figure 2) or *T. laevidens*, both occurring in the Middle Purbeck limestone, which is Berriasian and Berriasian-Valanginian age, respectively, to *T. microdon* from the Grinstead Clay Formation of Valanginian age, and *T. multidentis* from the Sevenoaks locality of the Weald Clay Group of Hauterivian-Barremian age. Hornung [29] moreover infers that "*Pycnodus*" *mantelli* and *T. arcuatus* may be synonymous. If that is true, then the location of *T. arcuatus* in southern England and "*Pycnodus*" *mantelli* in Germany could be an indication that this taxon might be amphidromous due to the lack of a hydrographic connection between the continental Wessex and the Lower Saxony basins. However, both a better quality and higher quantity of specimens from both localities will be necessary to confirm such a hypothesis.



**Figure 2.** Examples of jaw fragments of pycnodont taxa from nonmarine localities. (A) cf. *Coelodus* sp. (MTM V 2010.139.1), modified from [200] Szabo et al. 2016. (B) Pycnodontiformes incertae sedis (UFRJ-DG 423-PD), modified from [151]. (C) *Agassizilia erfoudina* (FSAC-KK 5073), modified from [47]. (D) *Neoprosclinetes africanus* (FSAC-KK 5070), modified from [47]. (E) *Turbomesodon* cf. *arcuatus* (NHM PV P.45763), modified from [29]. Collection abbreviations: FSAC - Faculté des Sciences Aïn Chock; MTM - Hungarian Natural History Museum (Magyar Természettudományi Múzeum); NHM - Natural History Museum, London; UFRJ - Universidade Federal do Rio de Janeiro. Scale bar for A, C, D and E is 10 mm. Scale bar for B is 5 mm.

Sweetman et al. [46] reported premaxillary, dentary, vomer and prearticular bones with teeth from the Wessex Formation (Barremian) from southern England and assigned them to *Coelodus* sp. However, Cooper and Martill [47] observed that the morphology of the teeth resemble more that seen in *Ocloedus* [48]. An additional, still unidentified pycnodont referred only to as Pycnodontiformes indet. in the same locality is represented by fragmentary vomers and prearticulars along with isolated teeth.

The occurrence of the pycnodont *Coelodus* was reported from two layers in the Berriasian Middle Purbeck beds of Durlston Bay in southern England [26]. These remains were obtained from the Cherty freshwater layer of the Lulworth Formation and the inter-marine layer of the Durlston Formation, the palaeoenvironment of the former being a series of freshwater ponds and marshes and the latter being a freshwater-brackish lake based on salinity measures inferred from mollusc shells and evaporite data [49].

However, the taxonomic assignment to *Coelodus* or to a pycnodont at all remains dubious, because no description nor any figure of this material has been provided up to now, enabling any unambiguous identification of this pycnodont fish; it only was listed in a table. Thus, unless the specimens related to this taxonomic assignment can be found or further material recovered from this locality, these specimen/specimens should be considered dubious at the best.

Another Early Cretaceous freshwater pycnodont record based on isolated teeth comes from the Pevensey Pit of the Ashdown Brickworks quarry of Bexhill in southern England, which is Valanginian in age [50,51]. Even though these authors did not provide any taxonomic assignment, we tentatively assign them here to *Ocloedus*, because of the characteristic tooth morphologies. These teeth alongside material belonging to a diverse fauna of sharks, teleosts, amphibians and dinosaurs would have been deposited in a rapidly flowing river due to such material being reworked together with sediments along pebble shorefaces [52,53]. Additional evidence for this environment being freshwater can also be seen in the predominance of certain genera of ostracods and bivalves such as *Cypridea* and *Neomiodon*, respectively [54].

### 2.3.3 Germany

The non-marine Early Cretaceous fossil record of Germany is according to our current knowledge restricted mainly to the Berriasian of northern and north-western Germany. Due to the lack of complete or taxonomically informative fossil material, most of these specimens are referred to as Pycnodontiformes indet. Isolated teeth were recovered from various lacustrine facies of the Berriasian Bückberg Group such as Sehnde and Wätzum [Fuhse Fm.] [55,56], Egestorf in Lower Saxony [Isterberg Fm.] (late Berriasian) that originally were referred to *Pycnodus mantelli* by Struckmann [57], but lack any description or illustration, Barenberg near Borgholzhausen in

Northrhine-Westphalia [either Oesede or Isterberg formations], previously referred to as *Gyrodus mantelli*, which only were mentioned but not illustrated by Dunker [58], and finally Lobber Ort on the Island of Rügen [59]. These remains have to be considered ambiguous as long as no further information is available.

Prearticulars assigned to “*Pycnodus*” *mantellii* (aka *T. arcuatus*) also were reported from various Berriasian sites in Lower Saxony [58] and North-Rhine Westphalia, northern Germany [60]. Vomers and isolated teeth belonging to *Turbomesodon* cf. *arcuatus* also were discovered in North-Rhine Westphalia [29,60–62]. Another specimen of this taxon represented by a vomerine dentition was reported from an unknown locality in the Isterberg Formation of the Bückenberg Group [29]. Kriwet [63] also figured a specimen of this taxon as “*Coelodus*” sp. from northern Germany [63; fig 28c].

While the palaeoenvironment of all these Berriasian localities are interpreted as lacustrine, occasional marine influences cannot be completely rejected.

#### 2.3.4. Belgium

While the locality of Bernissart in Belgium is most famous for the *Iguanodon* remains found at the bottom of a pit, a staggering number of actinopterygian specimens were unearthed from this locality, amounting to nearly 3000 specimens containing 17 species belonging to 12 genera [64,65]. The fossil remains were found in lacustrine clays, which are defined as the Sainte-Barbe Clays Formation [66,67]. The environment of Bernissart would have been lacustrine to swampy [68–71] and using palynological dating is estimated to be of late Barremian to early Aptian age [72–74]. The only pycnodont taxa found so far in this locality are some complete but imperfectly preserved specimens of *Turbomesodon bernissartensis* [75,76].

#### 2.3.5. France

An assortment of isolated teeth of various morphotypes from the Berriasian locality of Champblanc Quarry in Cherves-de-Cognac, France can be referred to Pycnodonformes [77]. These teeth are commonly found in the upper levels of the Champblanc section, which are characterized by lacustrine limestones [78,79]. The most common morphotype found is similar to the morphology seen in *Gyrodus* with a papilla-like structure present in the middle of an occlusal pit and Pouech et al. [77] subsequently considered this morphotype as aff. *Gyrodus*.

The only other morphotype that could be identified with a particular genus are teeth with two rounded tubercles on each end of the tooth crown with a groove in between. Such a tooth morphology is similar to that seen in *Arcodonichthys* [80]. Every other morphotype from this locality cannot be assigned to any genus or species with the most granular assignment given being that of the family Pycnodontidae.

The large amount of different and diverse morphotypes in the Berriasian locality of Champblanc Quarry nevertheless indicate that a diverse pycnodont fauna occupied this habitat that probably underwent niche partitioning in order to exploit different resources efficiently [16,47,81]. Another interesting observation by Pouech et al. [77] was the distribution of pycnodonts within the sedimentary series of Cherves-de-Cognac compared to ginglymodians, another actinopterygian clade with many durophagous representatives. Pycnodonts are more commonly represented in the upper levels of the sedimentary series, which are dominated by lacustrine limestones leading the authors to propose a lacustrine life environment for pycnodonts. Lentic environments such as lakes might have been more ideal for the typical laterally flattened and dorso-ventrally high body of pycnodonts as their morphology already permits them to manoeuvre in shallow complex environments without the need for expending energy fighting the current. In contrast, isolated teeth of ginglymodians such as *Scheenstia mantelli* dominate the actinopterygian assemblage in the lower levels. The environment of this lower series is confirmed by geochemical analyses to be upstream waters close to the lagoon where these teeth were deposited [82]. This pattern of pycnodonts and ginglymodians occupying different habitats mirrors that seen in marine environments as shown by the separation of these two groups from the same locality in a body shape morphospace occupation [16].

#### 2.3.6. Spain

The fossil record of non-marine pycnodont occurrences of Spain consists of not only abundant isolated dental remains but also many articulated skeletal remains (Figure 1) including holomorphic specimens.

A large abundance of isolated vomerine and prearticular tooth plates belonging to the pycnodont *Arcodonichthys pasiegae* were recovered from the Vega de Pas Formation (Hauterivian-Barremian) in the Cantabrian basin, northern Spain [80]. Sedimentation of this formation occurred in a lacustrine setting. *Arcodonichthys pasiegae* is easily identifiable due to the strongly curved anterior and posterior border of the vomerine teeth with swollen lateral ends. Due to the large number of specimens found of this taxon (over 100), some interesting aspects of its ecology can be inferred. The measurements of the tooth plates [see table in 80] suggest a small pycnodont close in size to taxa such as *Stemmatodus* [83] with an estimated size range of 4-7 cm. Tooth wear is extensive even in smaller specimens and the inner layer of dentine can be exposed in larger specimens [80; fig 1c]. This dental evidence along with the abundance of bivalves and gastropods found within the same fossil layers [84] as *Arcodonichthys* suggest that this species was particularly specialized to feed on hard shelled prey in comparison to many other pycnodonts, which were hypothesized to be more generalized durophages with a broad range of potential prey items as advocated by Poyato-Ariza [20] and Darras [85]. However, Kriwet [86] already hypothesized that pycnodonts were highly specialized feeders.

Several Lower Cretaceous localities in Eastern Spain yielded additional pycnodont fish material including complete skeletons. All these localities are situated in the Mesozoic Iberian Basin, an intracratonic basin situated at the eastern margin of the Iberian plate. The development of the Lower Cretaceous sub-basins of the Iberian Basin is related to the anticlockwise rotation of the Iberian Plate during the Mesozoic and extensional tectonic events [87].

Two of these famous localities are La Pedrera de Santa Maria de Meià (= La Pedrera de Rúbies) near Vilanova de Meià and La Cubrua in the Sierra de Montsec, Province of Lérida, north-eastern Spain [e.g., 88,89]. Originally assumed to be of Kimmeridgian age [90], Peybèrnes and Oertli [91] and Brenner [92] attributed a late Berriasian to early Valanginian age to the fossiliferous strata. The fish fauna includes chondrichthyans [only hybodonts; 93-95], an actinistian, and many actinopterygians [96-100]. Pycnodonts are abundant in the Sierra del Montsec and are represented by adult and juvenile specimens of a single species, *Ocloedus subdiscus* [48,89,101]. The depositional area of Montsec is subject to dispute. While some authors [102,103] interpret the depositional area as freshwater environment based on terrestrial faunal elements such as plants, insects, and tetrapods, the occurrence of agglutinating foraminifers, acritarchs, and brackish ostracodes indicate a marine environment [104]. This interpretation also is supported by the composition of the fish fauna [105]. The palaeogeographic situation of the fossiliferous strata also supports a coastal, intertidal environment, which is advocated here.

The syncline of Galve in the Province of Teruel in north-eastern Spain belongs structurally to the Lower Cretaceous sub-basin of Aliaga. The basin is built up of about 1000 m of Jurassic and Lower Cretaceous sediments and yielded ca. 60 vertebrate localities ranging from the Jurassic to Early Cretaceous in age [for reviews of the vertebrate fauna see 106,107]. The Early Cretaceous Aliaga sub-basin can be regarded as one of the most productive vertebrate localities of Early Cretaceous age on the Iberian Peninsula. Pycnodonts are represented by abundant isolated dentitions including both large and small specimens of the same taxon indicating the presence of juveniles and adults. Indeterminate pycnodontiforms were reported by Sánchez-Hernández et al. [107] from the upper part of the Berriasian-Barremian El Castellar Formation, which is interpreted as being an open lacustrine environment with distributary channels and a lagoonal system near the coast based on the presence of bioclastic limestones, sand beds, charophyte remains and root marks [108]. Soría and Meléndez [109] reported the presence of cf. *Proscinetes* sp. and *Coelodus* sp., as well as unidentifiable forms in the lower Barremian Camarillas Formation, which was interpreted by these authors as a fluvial environment. The presence of *Anomoeodus* or a similar taxon is indicated by the presence of an isolated premaxilla, which is characterized by three incisiform teeth (JK, pers. obser.). Lagoonal conditions in the syncline of Galve are supposed to have been established during the upper Barremian Artoles Formation, in which pycnodont dental remains are common (JK, pers. obser.). In contrast to this environmental interpretation, the selachians and associated actinopterygian fauna seems to favour an environment with at least sporadic connections to the open marine realm in the East [110].

Additional isolated dental remains of three different pycnodonts including two different pycnodontids of uncertain affinities (probably representing *Ocloedus*) and *Anomoeodus nursalli* were described by Kriwet [44] from coaly limestones (carbonates and clastics interbedded with coal seams of 60 to 65 cm thickness) of Uña in the Province of Cuenca, eastern Spain. The coals were described as brown coals or lignites [111–113]. They belong to the Huérguina Formation, which is early Barremian in age [114]. The depositional area generally is interpreted as freshwater. In addition to the pycnodonts, still un-described actinopterygians and a diverse tetrapod fauna were derived from one of these coals. Kriwet [44] hypothesized that the coals of Uña were accumulated at the margin of the Bay of Cuenca, which was connected to the marine realm in the east. Accordingly, continental faunal and floral contents were washed into this fast-changing area of paralic coal formation or lived at the edge of the bay. The pycnodonts thus probably were not primary inhabitants of this restricted, probably fast changing environment.

Although the vast majority of the Spanish Early Cretaceous non-marine pycnodont fossil record seemingly consists of fragmentary material such as isolated teeth and jaws, some complete pycnodont specimens from freshwater deposits nevertheless were recovered [89,115–117]. The most notable in addition to *Ocloedus subdiscus* from Montsech (see above) are *Stenamara mia* and *Turbomesodon praeclarus* from the lacustrine deposits of Las Hoyas in the Province of Cuenca, eastern Spain, which also is of Barremian age [76,118]. This locality was shown to represent a freshwater environment without any marine influence based on many factors including palaeoecology, palaeogeography, sedimentology, stratigraphy, taphonomy, and extensive isotope analysis of the fish fossils [119]. Both co-occurring pycnodonts are members of the Pycnodontidae, the most speciose lineage of pycnodonts and are considered the first unambiguous record of non-marine pycnodonts that were discovered [119]. *Stenamara* is both the rarest and smallest of the two Las Hoyas pycnodonts with only two specimens known [117] and its standard length being 73 mm. While *Stenamara* is comparatively poorly preserved it can still be ascertained to include an adult specimen due to the presence of bifurcation in the lepidotrichia [120] and its extremely deep body (maximum body height/standard length being 124.5%) indicates it to have been a slow swimmer but well adapted to manoeuvre in complex environments such as the vegetated shallows of the lake of Las Hoyas. *Turbomesodon praeclarus* originally identified as *Macromesodon* aff. *M. bernissartensis* [121], conversely, was obviously much more common in Las Hoyas with 25 or more complete or partially preserved specimens along with isolated dentition [76]. Like *Stenamara*, it was a deep-bodied fish well adapted to manoeuvre in structured habitats. The discovery of small specimens of *T. praeclarus* from the Las Hoyas sediments indicate the presence of juveniles, which is considered additional evidence that these fishes were fully freshwater adapted as they evidently also reproduced in the lake [76,121].

Poyato-Ariza [117] speculated that the Las Hoyas pycnodonts might have been more herbivorous than what is typically assumed for the group as the usual pycnodont prey such as molluscs is not present in the Las Hoyas locality and crustaceans such as crayfish are too large for the small gaped pycnodonts to feed on and break apart. The charophyte alga *Clavatoraxis robustus* was in great abundance in Los Hoyas [117] and was reinforced with spine-cell rosettes, which Martín-Closas and Diéguez [122] interpreted as a defence mechanism against herbivory. Both *Stenamara* and *Turbomesodon* with their powerful molariform teeth and nipping incisiform teeth might have been able to tackle such well defended prey. This herbivory hypothesis is more plausible than initially assumed due to adaptations to plant eating being found in other pycnodonts including bicuspid teeth in the Late Triassic *Gibbodon cenensis* [123], *Nursallia tethysensis* [124] and *Thiollierepycnodus wagneri* [125]. A large premaxillary tooth with multiple cusps similar to that seen in modern herbivorous teleosts such as marine acanthurids and freshwater cichlids also can be seen in the Palaeocene *Pycnodus multicuspidatus* from Moroccan deposits [126]. A tooth microwear analysis by Baines [127] on the jaws of the Jurassic pycnodont *Gyrodus planidens* revealed that the orientation of dental wear patterns on the teeth show that it was capable of propalinal or horizontal jaw movements, which is similar to that seen in mammals. These jaw movements would have been effective for processing plant material in contrast to the vertical crushing actions more typically used to access hard shelled prey items. The comparatively high number of *T. praeclarus* specimens in particular could be used to test such dietary hypotheses in the future.

Some interesting trace fossils at Las Hoyas possibly can be attributed to pycnodonts. These horizontal trails consisting of a sinusoidal wave are referred to as the ichnospecies *Undichna unisulca*

[128] and are preserved as epichnial grooves or hypichnial ridges. These trace fossils are more commonly preserved in freshwater environments and are a decent indicator of palaeoenvironmental setting [128]. The producer of these trace fossils would need to possess a large caudal fin with no overlap with the anal fin to produce such traces in the sediment. There are currently no fishes known from Las Hoyas that have such a caudal fin morphology without another fin leaving occasional traces. Pycnodonts typically have large anal fins that would fit in with the form of the trace fossils left behind and so were posited by Gibert et al. [128] to have been the most likely producers of *U. unisulca*. *Turbomesodon praeclarus*, being the more common of the two pycnodont taxa found at Las Hoyas is the most likely producer of these traces. Nevertheless, the palaeogeographic situation of Las Hoyas close to the marine realm during the Barremian does not preclude occasional marine incursions.

In addition to these well-known vertebrate sites, many other Early Cretaceous localities on the Iberian Peninsula, which are supposed to represent freshwater settings yielded isolated dental remains of pycnodonts, such as from the Berriasian of Lérida, eastern Spain (*Proscinetes* sp. cf. *P. bernardi*), the Barremian of Aguilón near Zaragoza, eastern Spain (*Coelodus* sp.), the Aptian of Cintorres near Valencia and Mirambel in Teruel, eastern Spain (*Anomoeodus complanatus*), the Aptian of Morella, Valencia, eastern Spain (*Coelodus* sp. aff. *C. soleri*), the Albian of Caceda, Asturias (undetermined pycnodont), the Albian of Condemios de Abajo, Castilla-La-Mancha, central Spain (*Anomoeodus* sp. aff. *A. muensteri*), and the Albian-Cenomanian of the area of Girona in Catalonia (*Coelodus soleri*) [93,129–132]. No pycnodont fish remains from non-marine deposits of Early Cretaceous age have been reported from western or southern Spain, or even Portugal to date. Nevertheless, this review emphasizes the incomplete knowledge of these fishes from the Early Cretaceous of the Iberian Peninsula, both in terms of taxonomy and environmental distribution.

### 2.3.8. North America

The currently known non-marine Early Cretaceous record of pycnodonts from North America seemingly is restricted to the Aptian-Albian. Oreska et al. [133] reported five partial vomers which they assigned to aff. Pycnodontidae indet. This material was collected from two vertebrate microfossil beds, at the Little Sheep Mudstone Member of the Cloverly Formation of Wyoming which has been dated to Aptian [134]. These microfossils were accumulated in concentrations within clay and mudstones and the lack of wear on these fossils indicate low transport and a low energy depositional environment [135] which, along with fine grained sediments and weathering volcanic ash, can be interpreted as a lacustrine environment [136].

Pycnodont remains also were reported from the Holly Creek Fm., which is part of the Trinity Group in Arkansas and which is dated to the Albian [137]. The depositional environment is suggested to be a coastal plain, being a mixed water to brackish environment based on stable isotope work performed on turtle shells [138]. This formation contains two pycnodonts: Pycnodontiformes indet. and *Anomoeodus caddoi* [137]. *Anomoeodus caddoi* is represented by one isolated tooth and two prearticular toothplates. A singular branchial and oral tooth are the only material representing Pycnodontiformes indet.

The greatest number of North American non-marine pycnodont occurrences during the Early Cretaceous can be observed in northern Texas [30,139–142] and consists exclusively of isolated dental remains. Based on these remains, several taxa were identified such as *Texasensis coronatus* from the Paluxy Fm. of the Huggins and Pecan Valley Estates sites of northern Texas (Albian), which consists of a series of sandstones and shales representing fluvial meanderbelt to coastal barrier and deltaic facies [143]. This species originally was referred to as *Callodus coronatus* by, e.g., Thurmond [30] and Winkler et al. [140,141] but transferred to *Texasensis* by Özdikmen [31], because the name of *Callodus* is preoccupied by a beetle.

Another pycnodont that also occurs in the Pecan Valley site is "*Macromesodon*" *dumblei* (most likely a member of *Turbomesodon*), while *Nonaphalgotodus trinitiensis* was described from the Albian deposits of Butler Farm [144], which also is considered to represent a freshwater pycnodont. The pycnodont *Thurmondella estesi*, which originally was ascribed to the genus *Paramicrodon* that, however, represents a synonym of a genus of flies [145], is known from two Texas localities spanning two different geological stages, i.e., Albian and Aptian, both belonging to the Paluxy Fm. [141]. The fossils of this species were recovered from deposits, which are in close association with thin, variably developed bituminous coal and plant fragment-bearing mudstones indicating continental influence

[141]. Other taxa recovered from Albian non-marine localities of northern Texas include "*Palaeobalistum*" *geiseri*, "*Proscinetes*" *texanus* and "*Coelodus*" sp. [141].

### 2.3.9. South America

South American Early Cretaceous non-marine pycnodont records come from a fossil fish assemblage of the Açú Formation, which is located in the Potiguar Basin in Ceará, north-eastern Brazil. The deposits of this formation were interpreted as a meandering riverine environment with some tidal influence [146]. Precise dating of this formation has been unsuccessful so far due to the lack of stratigraphically useful and diagnostic fossils, but an Albian-Cenomanian age was suggested based on palynological analyses [147,148]. More recent stratigraphic data provided by by Arai [149,150] specified the age of the Açú Formation as Albian in age.

Isolated teeth and a partially preserved vomerine and prearticular dentition of pycnodonts were recovered from the Açú Formation [151] and assigned to "Pycnodontiformes incertae sedis" due to the lack of more complete and taxonomically diagnosable material. These specimens are likely to represent a new genus as it differs from all other known Brazilian pycnodont taxa in the shape and ornamentation of the teeth, which are present as a series of crenulations on the margin of the tooth (Figure 2) [151].

### 2.3.10. Africa

A fragmentary prearticular and isolated teeth from the Albian Jebel Boulouha North locality in Tunisia tentatively were assigned to aff. *Gyrodus* sp. by Cuny et al. [152]. Sedimentation of the fossiliferous layers were interpreted as a terrestrial carbonate rich environment [153]. The teeth display an apical depression, which is surrounded by crenulated margins similar to what is seen in European specimens also assigned to aff. *Gyrodus* sp. from the Hauterivian by Kriwet and Schmitz [154]. This similarity prompted Cuny et al. [152] to assign this material to a similar taxon, pending better preserved material.

Another Tunisian specimen from the Albian described by Cuny et al. [152] is an isolated tooth identified as Pycnodontiformes indet., which was recovered from the Oued el Khil assemblage. The depositional area of this assemblage was interpreted as a brackish environment due to the mix of freshwater and marine influences found in deposition [153,155].

The Kem Kem Group of Morocco is one of the most famous Cretaceous freshwater localities, being Albian-Cenomanian in age. While better known for hosting a multitude of giant vertebrate taxa such as the theropod dinosaurs *Spinosaurus* [156–158] and *Carcharodontosaurus* [159], the sawfish *Onchopristus* [160,161] and the coelacanth *Mawsonia* [162], the Kem Kem also provided a diverse pycnodont fauna [47]. All of the known pycnodont taxa from this locality are represented by fragmentary dentitions and isolated teeth, which are poorly preserved probably as result of the high energy riverine environment that characterizes the sedimentary setting of the Kem Kem. The corresponding pycnodont fauna is of note for two things in particular: 1) it is the oldest record of pycnodonts from Morocco and 2) it is the most diverse record of freshwater pycnodont assemblages known in the fossil record with at least four genera being represented. This assemblage also allowed the identification of a second species of *Neoproscinetes*, *N. africanus*. This species shares with *N. penalvai* from the Early Cretaceous of Brazil similar circular to subcircular medial vomerine teeth and similar morphologies, numbers and arrangement of teeth on the prearticular (Figure 2). *Agassizilia erfoudina*, an endemic taxon for the Kem Kem group, is similar to *Iemanja* from the Early Cretaceous of Brazil in that the prearticular teeth are small and irregularly arranged, but they are only restricted to the lateral margin in *Agassizilia* (Figure 2). This suggests it may have also fed on soft shelled crustaceans as was suggested for *Iemanja* [47]. The other two pycnodont taxa are not well enough preserved to allow a definitive taxonomic assignment but were identified as cf. *Macromesodon* and cf. *Coelodus* due to sharing many similarities with the aforementioned genera of pycnodont fishes. Cooper and Martill [47] hypothesized that the high diversity of pycnodonts in the Kem Kem group could be due to niche partitioning as each taxon found has a unique arrangement of teeth, which, combined with the few stomach contents of pycnodonts that are preserved being monospecific [86], suggests high specialization of each genus to different prey items. Additional evidence for such niche

partitioning was found through the results of a morphospace analysis concerning pycnodont lower jaws [16].

#### 2.4. Late Cretaceous

##### 2.4.1. China

Zhou et al. [163] documented a hitherto unknown pycnodont, *Xinjiangodus gyrodooides*, which is represented by a fragmentary vomerine and prearticular dentition coming from the sandy mudstones of the Donggou Formation at the southern margin of the Junggar Basin in Xinjiang, China. The exact age of this formation is problematic with some authors indicating a Coniacian-Santonian age [164] and others restricting it to the Maastrichtian [165]. More recent work based on fossil ostracods attributes a Coniacian-Campanian age to the Donggou Formation [166,167]. The sedimentary environment of the fossiliferous site of this pycnodont is indicative of shallow lake deposits. A characteristic feature of *X. gyrooides* distinguishing it from other pycnodonts is the large spacing between teeth, and the presence of an apical depression on the tooth crowns with radiating wrinkles [163: table 1). An isolated opercular bone also was assigned to this taxon. However, the shape and size of the fragmentary opercular bone is far too wide and large for pycnodonts and it is more probable that it belongs to a teleost. Alternatively, it could be the remains of a preopercular bone of a pycnodont as that bone is typically more expanded in pycnodonts, but this specimen is too fragmented to genuinely identify this bony element to any pycnodontiform let alone *X. gyrodooides*.

##### 2.4.2. India

Complete pycnodont skeletons representing a single species, "*Pycnodus lametae*" have long been known from the Late Cretaceous Lameta Formation of central India (Province of Maharashtra) [168,169]. The Lameta Fm. is part of the infratrappean beds, which were deposited below the Deccan Traps. These traps represent the deposits of active volcanic flows during the latest Cretaceous (Maastrichtian) found in association with sedimentary beds of continental origin. The infratrappean beds of the Lameta Fm. were interpreted as palustrine/lacustrine depositional environments in an alluvial-plain setting under semiarid climatic conditions based on palaeontological and sedimentological evidence [170–173].

An additional incomplete specimen of "*Pycnodus lametae*" consisting of a crushed skull and missing caudal region was described from laminated clays of the same area from which the specimen of Woodward [168] originated, at Dongargaon by Mohabey and Udhoji [174]. These authors mistakenly assigned this specimen as the holotype despite the holotype being the specimen described by Woodward [168]. The assignment of this pycnodont to *Pycnodus* is problematic due to the lack of a post-parietal process [reported in 42], so revision of this material is necessary for taxonomic purposes.

Isolated teeth of various tetraodontiforms, which were initially assigned to *Stephanodus libycus*, *Indotrigonodon ovatus*, *Pisdurodon spatulatus* and *Eotrigonodon wardhaensis* also were reported from the Lameta Fm. [175,176]. Prasad [169] provided a detailed review of the vertebrate fauna of the Deccan Traps and reassigned these teeth to an indeterminate pycnodontid. The teeth of *S. libycus* were interpreted as branchial teeth, whereas the teeth of *I. ovatus*, *P. spatulatus* and *E. wardhaensis* were interpreted as oral teeth [169]. Shome and Chandel [177] reported the presence of branchial teeth, which were assigned to *Pycnodus* sp. from the Papro Fm. of Lalitpur District, Uttar Pradesh, which is dated to the Maastrichtian based on the vertebrate microfauna fossil record. Oral teeth of "*Pycnodus*" sp. and branchial teeth named *Stephanodus libycus* were also discovered from predominantly freshwater, lacustrine and palustrine (swamp) deposits of the Maastrichtian Lameta Fm. in the Province of Madhya Pradesh [178,179].

Many additional records of pycnodonts are known from intertrappean beds. In contrast to infratrappean beds, intertrappean beds were deposited during the dormancy stages of volcanic activity and are thus intercalated within the lava flows. These depositions are generally considered to have been freshwater lacustrine palaeoenvironments that would have occupied low-lying areas on the surface of lava flows when volcanic flow had ceased.

Khosla et al. [180] reports the presence of a branchial tooth assigned to an indeterminate pycnodontid from an intertrappean section of Kisalपुरi, which also contains freshwater ostracods. Teeth named *Pycnodus bicresta* were reported from the Naskal locality of the Andhra Pradesh Fm., which was interpreted as an alkaline lake in a floodplain setting based on sedimentological, palaeontological and taphonomic evidence [181,182]. Since these authors considered pycnodonts to be an exclusively marine group, they assumed that *P. bicresta* had migrated into this lake through river channels and became subsequently entrapped.

Also, from Naskal, a specimen of "*Pycnodus lametae*" was reported to have been recovered, which, unfortunately, was lost during preparation [183]. It seems that "*Pycnodus lametae*" is a common freshwater taxon in India as Rana [184] has reported specimens from several localities including Asifabad, Nagpur and Rangapur. As well as "*Pycnodus lametae*" [185], two other pycnodont taxa, *Pycnodus bicresta* [169,186] which was originally assigned as an indeterminate pycnodontid [176,185] and *Pycnodus* cf. *P. praecursor* [184] are known from the Asifabad Fm., the latter restricted to infratrappian beds while the other pycnodont specimens also are found in intertrappean beds.

Another unidentified pycnodontid, represented by a prearticular bone, branchial, incisiform and molariform oral teeth, is known from the Maastrichtian Piplanarayanwar of the Chhindwara District, Madhya Pradesh [186]. This locality was interpreted to have a nearshore, coastal plain environment due to the abundance of the myliobatid batomorph *Igdabatis indicus*. Diverse mixes of nonmarine and marine taxa at the Asifabad and Nagpur sites also indicate some marine influences [185,187,188].

Two other Indian localities of Late Cretaceous to Palaeogene age provided additional isolated pycnodont teeth, which all were identified as *Pycnodus* sp. These include the estuarine deposits of the Maastrichtian-Selandian Fatehgarh Fm. of Rajasthan [189,190] and the Maastrichtian-Danian intertrappean beds at the Jhilmili locality, Chhindwara District [191], in which the presence of *Pycnodus* together with *Lepisosteus* remains indicate a fluvio-lacustrine environment. It nevertheless should be noted that all Late Cretaceous pycnodont remains assigned to *Pycnodus* have to be considered as different taxon since *Pycnodus* is not known before the Cenozoic [compare 42].

#### 2.4.3. Austria

Isolated teeth and a fragmentary right prearticular were reported from the Turonian Schönleiten Formation in SE Austria (Steiermark) and assigned to *Coelodus plethodon* by Schultz and Paunović [192]. This species originally was described by Arambourg and Joleaud [193] from the marine Campanian of Niger (Africa). Ósi et al. [194] argued that the Austrian material is too fragmentary to assign it unambiguously to any particular genus, with which we agree here. Consequently, these remains have to be considered as Pycnodontidae or even Pycnodontiformes indet. The specimen was recovered from grey shales which represents a brackish, estuarine environment [195–197].

#### 2.4.4 Hungary

The pycnodontid cf. *Coelodus* was reported from the Santonian Csehbánya Fm. of Iharkút, Hungary based on isolated dental remains [198–200] (Figure 2). The prearticulars were initially considered to represent probably two different morphotypes by Ósi et al. [201], but Szabó et al. [200] nevertheless assigned all the Iharkút material to a single taxon until more complete material is available for a definite identification.

Sedimentation of the Csehbánya Fm. was determined to have been a floodplain environment [202]. Work on the pycnodont dentitions from multiple authors has revealed quite a lot about the palaeoecology of these pycnodonts. Stable isotope composition performed on their teeth revealed low concentrations of strontium and that migration from marine or brackish environments thus were unlikely [198]. Additionally, different ontogenetic stages are represented evidencing that these pycnodonts inhabited these non-marine environments for their entire life [200]. Botfalvai et al. [199] revealed that these pycnodonts were comparatively abundant in the vertebrate assemblage being the third most frequent group represented by 28 individuals. An abundance of potential molluscan prey such as bivalves and gastropods are known from the Csehbánya Formation and Szabó et al. [200] hypothesized that these would have formed the main food source of the pycnodonts. However, so far, no direct evidence of a predator-prey relationship between these pycnodonts and molluscs has been observed.

Another Hungarian pycnodont represented by isolated molariform oral and hook-like branchial teeth, which currently is assigned to Pycnodontiformes indet. was recovered from the Santonian Ajka Coal Fm. [203]. This pycnodont also can be determined to be a freshwater inhabitant as the Ajka Formation was dominated by swampy and lacustrine environments in multiple sub basins [204–206].

#### 2.4.5. France

Fragmentary vomerine and prearticular dentitions identified as cf. *Phacodus* were recovered from the Campanian Villeveyrac - L'Olivet locality in southern France [45]. The fossil assemblage of this site, particularly the remains of plants and molluscs such as gastropods and bivalves correspond to taxa that are typically known from lacustrine environments [207]. As with cf. *Anomoedus* from the Barremian Sao Khua Fm. of Thailand, Cavin et al. [45] also observed that all specimens of cf. *Phacodus* from this locality were small (e.g., the vomer was estimated to be roughly 5 mm long if complete). Using the proportions of a typical pycnodont [63] then it corresponds to a total length of just over 5 cm. All other known records of *Phacodus* show it to be a marine fish with specimens 20 times larger than the vomer from Villeveyrac – L'Olivet. Assuming that the French specimens represent juveniles, *Phacodus* might have represented an anadromous fish, with adults returning to freshwater to spawn and juveniles travelling to marine waters once they reached adulthood.

#### 2.4.6. Spain

Isolated teeth belonging to unidentified pycnodonts were reported by many authors from the Campanian-Maastrichtian locality of Lo Hueco of the Province of Cuenca in eastern Spain [208–211], which was a swamp or wetland type environment based on palynological analysis [210]. During the Maastrichtian, a number of freshwater and estuarine environments from north-eastern Iberia [212] were home to several taxa of pycnodonts but unfortunately, all of these are represented by isolated teeth only and thus, are rarely assigned to genus level. These environments include the brackish coastal plain Fontllonga-6 locality (Pycnodontiformes indet., ?Pycnodontiformes indet.), the coastal lagoonal site of Els Nerets (cf. *Coelodus* sp.), which is interpreted to be predominantly freshwater due to the richness of cypriniforms and freshwater zygospores in the lower beds from the site but also contain some marine influence due to the presence of dinoflagellate cysts in the upper beds, the estuarine wetland L'Espinau site (?Pycnodontiformes indet., cf. *Coelodus* sp.), the floodplains of Serrat del Rostiar-1 (Pycnodontiformes indet.), which contain ephemeral ponds and fluvial channels with some marine influence, the riverine environment of the Camí del Soldat locality of the Tremp Fm. (Pycnodontiformes indet.), which had a stable salinity due to tidal currents accumulating seawater in shallow pools at the margins of the river [213].

#### 2.4.7. North America

Jaws and teeth belonging to two indeterminate species of pycnodontids were uncovered from the Mussentuchit Member of the Cenomanian Cedar Mountain Fm. in Emery County, Utah [214,215], which represents a broad alluvial floodplain with no evidence of marine or brackish water facies [216]. The Cenomanian Dakota Fm. of the Grand Staircase–Escalante National Monument region of Utah, which was a floodplain environment yielded vomerine and prearticular tooth plates, branchial teeth and contour scales that were assigned to *Coelodus* sp. by Brinkman et al. [217]. Fragmentary vomers, isolated vomerine teeth, branchial teeth and contour scales belonging to *Coelodus* sp. are also uncovered from the Turonian Smoky Hollow Member of the Straight Cliffs Formation in Utah, which contains sediments indicating a lacustrine/floodplain palaeoenvironment from the middle Turonian [217]. Additionally, isolated vomerine and branchial teeth of *Coelodus* sp. were found in the Coniacian John Henry Member of the Straight Cliffs Formation. These Coniacian specimens also are considered to be non-marine as they co-occur with teeth of the freshwater shark *Lonchidion* [218,219]. Brinkman et al. [217] also reported isolated oral and branchial teeth of *Micropycnodon* sp. from two brackish localities, the Santonian Straight Cliffs and Campanian Wahweap formations, both localities also from southern Utah.

#### 2.4.8. South America

Just as in the Early Cretaceous, South America provided a wide assortment of non-marine Late Cretaceous pycnodonts. The most diverse continental vertebrate fauna of Late Cretaceous age of South America comes from the Cenomanian Alcântara Formation of Brazil [220,221]. Here, the Laje do Coringa outcrop is the richest fossil site of the Alcântara Fm. and shows high taxonomic similarities with the vertebrate fauna of the Kem Kem beds, which is related to South America and Africa drifting apart in the late Aptian-Albian [222]. Pycnodonts of the Laje do Coringa assemblage are represented by isolated remains of a single indeterminate pycnodontiform [223]. The palaeoenvironment of Laje do Coringa was inferred by Medeiros et al. [222] as an estuary located in a forested coastline in a semi-arid to arid climate.

In Argentina, isolated vomerine and prearticular dentitions of at least two pycnodontiforms were recovered from non-marine sediments of the Maastrichtian Yacoraite Fm. Several dental remains identified as *Coelodus toncoensis* were reported from gray and black shale levels of the Amblayo Member in the Salta Basin [224–228]. Sedimentation of the Amblayo Member was predominantly marine but becomes increasingly mixed with freshwater and decreases in sea levels near the boundary with the overlying Güemes Member as can be seen in the low values of oxygen and carbon isotopes obtained from the deposited limestone [229]. This environmental change can also be observed in the increase in terrigenous content and the accumulation of silty red sandstone and mudstone-wackestone facies [229]. Both types of evidence indicate that *C. toncoensis* occupied a shallow water environment with varying salinity. The gray and black shales from which the remains of *C. toncoensis* were recovered are indicators of a profundal setting within the center of a lacustrine basin [230].

Additionally, an indeterminate pycnodontiform was reported from the Tres Cruces sub-basin of the Yacoraite Fm. [231]. Marquillas et al. [229] conducted an isotope analysis of the limestones of this sub-basin and found a similar decrease in oxygen and carbon isotope values as was found for the upper Amblayo Member suggesting a similar brackish environment for this pycnodont as for *C. toncoensis*.

The only other South American country where freshwater pycnodonts were recorded is Bolivia. Here, a tooth of an indetermined pycnodont was described from the Campanian Chaunaca Fm. representing a continental fluvial deposit [232,233]. Three pycnodont taxa were recovered from the brackish/lagoonal Paja Patcha locality of the Lower El Morino Formation, which is Maastrichtian in age: branchial teeth assigned to an indeterminate pycnodontid [234,235], a fragmentary prearticular of *Coelodus* sp. [235] and fragmentary prearticulars, vomers and isolated teeth belonging to *Coelodus toncoensis* [234–236]. An indeterminate pycnodontid also was reported by de Muizon et al [237] from the marshy to palustral Vila Vascarra locality, which also belongs to the El Morino Fm.

#### 2.4.9. Africa

Taxonomically undiagnostic pycnodont teeth (Pycnodontiformes indet.) from the Cenomanian “Continental Intercalaire” of the Guir Basin of Algeria were reported by Benyoucef et al. [238]. One tooth is more rounded than the other and unornamented while the other tooth is ornamented by a series of crenulations surrounding an apical depression. They were recovered from the Menaguir site, which is interpreted to be a near shore environment based on sedimentological and ichnological features implying brackish water conditions [238].

Two taxa of Maastrichtian non-marine pycnodonts, *Pycnodus* sp. and *Pycnodus jonesae*, were reported from the Ménaka Formation in the Iullemeden Basin, Mali [239,240]. Both *Pycnodus* taxa were deposited in a phosphate conglomerate, which is indicative of shallow marine to brackish environmental conditions rather than being freshwater [241]. *Pycnodus* sp. is represented by fragmentary tooth plates, which are insufficiently preserved to assign to a species level. Specimens of *Pycnodus jonesae* are represented by isolated teeth, prearticular dentitions and indeterminate tooth plates. However, the assignment of these remains to *Pycnodus* are dubious and they certainly represent other, probably new pycnodontiforms [compare 42].

#### 2.4.10. Madagascar

Only one possible freshwater pycnodont represented by an incomplete vomerine tooth plate assigned to *Coelodus* sp. has been reported from Madagascar up to now, which is Maastrichtian in

age [242]. This specimen comes from the Anembalemba Member of the Maevarano Formation in Boeny, which is interpreted as a coastal alluvial plain that was subject to strong seasonality with intense rainfall followed by periods of drought based on sedimentary features in the facies deposits showcasing alternating episodes of turbulent flow and mass flow [243–245]. The Maevarano Formation being coastal and being overlain by the Maastrichtian marine Berivotra Formation could explain why *Coelodus* and other marine fish taxa were present in this continental setting as marine transgression could allow for easier access to such environments [243].

## 2.5. Palaeogene

Pycnodont diversity in the Palaeogene drops significantly compared to what is seen in the prior Late Cretaceous [16,19] and as a result, less non-marine occurrences of pycnodonts were identified in the fossil record.

### 2.5.1. India

Palaeogene non-marine pycnodontiforms of India were reported from the intertrappean Andhra Pradesh Formation at Rajahmundry including two taxa, *Pycnodus* sp. and *Pycnodus* cf. *P. praecursor* [184]. These fossiliferous layers were deposited just after the K/Pg extinction event during the early Danian according to <sup>40</sup>Ar/<sup>39</sup>Ar radioisotopic dating and palaeomagnetism [246]. Branchial teeth originally identified as *Eotrigonodon* aff. *E. jonesi* and also found in Rajahmundry are now considered to belong to a pycnodontid [169,247]. Additional teeth of unidentified pycnodontids were additionally uncovered from this site [248]. All of these fishes were deposited in silty claystones, which reflect supratidal to terrestrial environments followed by a return to restricted brackish conditions [249].

### 2.5.2. North America

Isolated molariform teeth of *Pycnodus* sp. have been collected from both the Tallahatta (Ypresian-Lutetian) and Lisbon formations (Lutetian) of the Claiborne Group in Alabama, USA [250]. The palaeoenvironment of the Claiborne Group was most likely a marine environment with a soft/muddy bottom based on an analysis of otoliths of various teleosts [250]. However, the presence of teleost families in this locality known to occupy brackish waters [250], as well as fossils of turtle taxa that occur in estuaries today being found in the same locality [251] indicate that this pycnodont might have been an inhabitant of an estuarine environment.

### 2.5.3. South America

Benedetto and Sanchez [225] report *Coelodus toncoensis* not only from the Maastrichtian Amblayo Member of the Yacoraite Formation but also from the limestones of the Yacoraite Formation close to the boundary of the Santa Barbara Formation. This description correlates well with the Alemania Member of the Yacoraite Formation, which is of Danian age [229,230] indicating that this species might have survived the K-Pg extinction event. In the Palaeogene part of the Yacoraite Fm., a return to more marine conditions can be observed based on a change from negative to positive values in carbon and oxygen isotopes [229]. However, palaeontological evidence such as the presence of euryhaline invertebrates including ostracodes, bivalves and gastropods indicate a shallow marine environment with some localised mixing of fresh and brackish waters [252], which matches the mixed environment of the upper Amblayo Member (Maastrichtian) of the Yacoraite Fm. where *C. toncoensis* was also present.

### 2.5.4. Africa

Like with the *Pycnodus* specimens from the Maastrichtian, the Palaeogene pycnodonts from Mali also were found in phosphatic conglomerates indicating shallow marine to brackish conditions. An earlier Malian representative of *Pycnodus* from the Palaeocene (Selandian-Thanetian), *P. jonesae*, might have survived the K-Pg extinction event and was recovered from the Teberemt Formation of


the Gao Trench and Iullemeden basins. Additionally, *P. jonesae* was found in the late Palaeocene (Thanetian) Teberemt Fm. of the Iullemeden Basin [239,240]. *Pycnodus* sp. was also reported from the Teberemt Fm. The most famous of these localities is located in the Tamaguélelt Fm. of the Taoudenit Basin, Mali, which yielded three taxa of *Pycnodus*, *Pycnodus* sp., *P. zeaformis* and *P. maliensis*, which are all Eocene (Ypresian) in age [239,240]. It is noteworthy that *P. maliensis* is represented by prearticular remains, while *P. zeaformis* is known by vomers indicating that both could be synonymous.

### 3. Discussion

One clear pattern arises when the entire non-marine pycnodont fossil record is considered: The incredible rise in such occurrences throughout the Cretaceous. While Jurassic occurrences are sporadic with low generic diversity, that changes dramatically when one enters the Early Cretaceous with 14 known genera observed during this time period and the number of occurrences is eleven times higher than the preceding period (Tables 1 and 2). A map of these occurrences shows that pycnodonts were initially present in far eastern Asia, North America and central Africa during the Late Jurassic and have extended their range into Western Europe, South America and northern Africa in the Early Cretaceous (Figure 3). Number of occurrences and known genera both decrease in the Late Cretaceous with the latter being especially affected by a poorer fossil record with six known genera. Late Cretaceous non-marine occurrences also see the first appearance of pycnodonts in both Madagascar and India. There is a westward shift throughout the late Cretaceous in the distribution of non-marine South American pycnodonts from Brazil in the Cenomanian to Bolivia and Argentina in the Campanian-Maastrichtian.

It is only in the Cretaceous that fully articulated fossil non-marine specimens are found such as *Stenamara mia* and *Turbomesodon praeclarus* [48,76] from the Barremian Las Hoyas Fm. in Spain and *Pycnodus lametae* [168,169,174] from the Maastrichtian Lameta Fm. In India, which has greatly helped researchers in the accurate classification of specimens whereas the Jurassic and Palaeogene non-marine pycnodonts are represented by far more fragmentary material. Also of note is that all articulated pycnodont specimens are found in lacustrine deposits but are a significant minority of non-marine pycnodont specimens discovered so far 5 out of 129 listed occurrences or 3.88% (see Appendix A). Although the preservation quality of the freshwater pycnodont fossil record is generally poor, with the few exceptions mentioned above, Guinot and Cavin [253] detected a major decrease in body size among pycnodonts when they moved into freshwater habitats. When all occurrences are presented for just the Cretaceous, both the earliest (Berriasian) and final (Maastrichtian) stage of the Cretaceous, have the largest number of non-marine pycnodont occurrences by 17.3% and 26.4%, respectively (Figure 4, Table 3).

Why pycnodonts were so common in non-marine habitats during the Cretaceous could be a result of the environmental conditions throughout much of this time. The Early Cretaceous in particular was a time of immense environmental change with not only sea level changes but a shift from an arid to a humid climate [254]. Sea levels were much higher than today due to the absence of polar ice caps. Such variations in sea level increase the possibility of marine fishes colonizing continental environments and was a major factor for why many marine fish lineages made incursions into freshwater environments during the Miocene [255]. Low sea levels, conversely, increased the spread of low-lying areas, which would have allowed for the movement of freshwater fishes from one hydrographic basin to another and which thus could have been an additional factor for why freshwater occurrences in pycnodonts had risen so dramatically from the Late Jurassic to the Early Cretaceous (Table 1-2, Appendix A).



**Figure 3.** Palaeobiogeographic distribution of non-marine pycnodonts through time. Numbered stars refer primarily to geological age. When geological age is unclear, the star will represent the locality the fossil was collected from as well as the currently time span that the locality is believed to represent. All cited localities and localities referring to a particular stage are discussed in further detail within the main text. Maps modified from Scotese [227, 228, 229, 230 2014a, b, c, d]. (A) Late Jurassic, Oxfordian, 158.4 Ma: 1. Changtu, Tibet, Late Jurassic; 2. Khlong Min Formation, Thailand, Mid-Late Jurassic; 3. Morrison Formation, USA, Oxfordian-Tithonian; 4. Mugher Mudstone, Ethiopia, ?Kimmeridgian-Tithonian; 5. Stanleyville Formation, Democratic Republic of Congo, Kimmeridgian-Valanginian. (B) Early Cretaceous, early Aptian, 121.8 Ma: 1. Berriasian; 2. Iberian Basin, eastern Spain, Berriasian-Valanginian, Barremian-Albian; 3. Southern England, Berriasian-Valanginian, Hauterivian-Barremian; 4. Valanginian 5. Phu Phan Thong, Thailand, Hauterivian-Barremian; 6. Vega de Pas Formation, northern Spain, Hauterivian-Barremian; 7. Bernissart, Belgium, Barremian-Aptian; 8. Paluxy Formation, northern Texas, USA, Albian-Aptian; 9. Aptian; 10. Albian; 11. Kem Kem, Albian-Cenomanian. (C) Late Cretaceous, Maastrichtian, 68 Ma: 1. Cenomanian; 2. Turonian; 3. Southern Utah, USA, Turonian-Campanian; 4. Donggou Formation, China, Coniacian-Campanian; 5. Santonian; 6. Campanian; 7. Lo Hueco (Cuenca), Spain, Campanian-Maastrichtian; 8. Maastrichtian; 9. Jhilmili, India, Maastrichtian-Danian; 10. Rajasthan, India, Maastrichtian-Selandian. (D) Palaeogene, Ypresian, 52.2 Ma: 1. Danian; 2. Teberemt and Tamaguélelt Formations, Mali, Selandian-Ypresian; 3. Lutetian.

Epicontinental seas were a common phenomenon during the Cretaceous such as the Western Interior Seaway Albian-Maastrichtian; [256,257] and the Trans-Saharan Seaway Cenomanian-Ypresian; [240]. Due to the increase in sea levels, many brackish water ecosystems as well as interior seas developed and would have encountered extreme variations in salinity. This probably explains why many fossil localities from the Cretaceous contain mixed assemblages of both freshwater and marine fish taxa making habitat interpretations based on fish taxa alone difficult and prone to errors. There also was a period of high sea levels during the Palaeogene such as the Paleocene–Eocene Thermal Maximum event (56 MYA), which coincides with the exclusive presence of pycnodonts in

brackish water environments (Appendix A), e.g., those occurring in Mali [240]. The increase in freshwater habitat occupation during the Cretaceous may reflect a genuine signal as similar movements into these habitats can be observed in other fish groups at the time such as Ginglymodi [13] and Hybodontiformes [11].


**Table 1.** Non-marine pycnodont occurrences and diversity through time. Localities with uncertain stratigraphic age included.

<b>Time Bin</b>	<b>Nonmarine occurrences</b>	<b>No. of known genera</b>
Late Jurassic	4	3
Early Cretaceous	61	15
Late Cretaceous	50	8
Palaeocene	9	2
Eocene	5	1

**Table 2.** Non-marine pycnodont occurrences and diversity through time. Specimens from localities with known stratigraphic ages only included.

<b>Time Bin</b>	<b>Nonmarine occurrences</b>	<b>No. of known genera</b>
Late Jurassic	3	1
Early Cretaceous	60	14
Late Cretaceous	45	5
Palaeocene	7	2
Eocene	5	1

Freshwater habitats being used as a refugia for ancient fish clades is a consistent evolutionary pattern and was noted in Amiiiformes, Lepisosteiformes and Dipnoi [11,258–260]. Cavin [10] stated that freshwater environments appeared to be far less vulnerable to environmental disturbance than marine habitats during the K-Pg extinction event. The majority of fish clades (14 of 15 families) that became extinct at this time were marine forms that were also piscivores at the top of the food chain [261] and no exclusively freshwater clade was wiped out during the K-Pg boundary event. An analysis by Sheehan and Fastovsky [262] revealed that over 80% of actinopterygian species from freshwater deposits in eastern Montana crossed the K-Pg boundary. Similarly, work by Cavin and Martin [263] showed that most actinopterygian families (24 out of 25) crossing the K-Pg boundary had representatives living in freshwater or brackish environments ranging from those living exclusively in freshwater to marine groups that had occasional freshwater/brackish representatives. The reason for this discrepancy in victims of this mass extinction most likely lies in how their food webs are structured. Marine food chains, particularly those of the open ocean, have phytoplankton at the base of the food chain, which was drastically affected by the darkening of the atmosphere from the asteroid impact and which ultimately had a domino effect on the rest of the ecosystems until it hit the apex predators [264]. This is supported by the documented decline in pelagic plankton after the K-Pg boundary [265], which agrees well with the disappearance of pelagic predatory fish.



**Figure 4.** Relative number of non-marine pycnodont occurrences from each geological stage of the (A) Early and (B) Late Cretaceous. Numbers in parentheses represent total number of occurrences. Occurrences used to represent pycnodont presence as fossil material typically too fragmented to assign genera/species.

The food webs of freshwater ecosystems in contrast, rely on detritus, which continues to function regardless of the potential for photosynthesis being diminished. Supporting this further, the two families of fishes that were not strictly marine that perished during the K-Pg extinction event had representatives that would have been planktivorous, e.g., the mawsoniid sarcopterygians and aspidorhynchid actinopterygians [10,266]. Coastal marine and brackish habitats would also be able to weather the extinction event better as they also had access to the detritus from river systems that flowed into the ocean [264]. Since pycnodonts not only occupied numerous freshwater ecosystems at the boundary but were predominantly present in coastal habitats such as reefs, this could be one particular factor for why they survived the K-Pg extinction event while many other marine fish clades were wiped out.

**Table 3.** Nonmarine pycnodont occurrences during each stage in the Cretaceous in total numbers of occurrences and in relation to total non-marine pycnodontiform occurrences throughout the Cretaceous.

Stage	Nonmarine occurrences	% of Cretaceous nonmarine occurrences
-------	-----------------------	---------------------------------------

Berriasian	19	17.3
Berriasian-Valanginian	2	1.8
Valanginian	2	1.8
Hauterivian-Barremian	3	2.7
Berriasian-Barremian	1	0.9
Barremian	13	11.8
Barremian-Aptian	1	0.9
Aptian	8	7.3
Albian	11	10.0
Albian-Cenomanian	5	4.5
Cenomanian	5	4.5
Turonian	2	1.8
Coniacian	1	0.9
Coniacian-Campanian	1	0.9
Santonian	3	2.7
Campanian	3	2.7
Campanian-Maastrichtian	1	0.9
Maastrichtian	29	26.4

#### 4. Conclusions

Pycnodonts were a predominantly marine group that were especially prevalent in structured habitats such as reefs from the Norian (Late Triassic) to the Ypresian (middle Eocene, Mid-Palaeogene) spanning an evolutionary history of ca. 170 Ma [42]. Nevertheless, they seemingly made numerous incursions into brackish and even freshwater habitats throughout their evolutionary history starting in the Jurassic, but particularly during the Cretaceous. The last freshwater occurrences coincide with those from marine deposits [42,250]. The occurrence in the Lutetian Lisbon Formation of the Claiborne Group in Alabama, USA [250] could even indicate that pycnodonts survived longer in freshwater than in seawater, if the stratigraphic assignment is correct. However, despite the repeatedly incursion into continental environments, pycnodontiform seemingly never became permanent inhabitants of freshwater environments like more ancient clades e.g., Lepisosteiformes and Amiiformes.

The marine to freshwater transition of pycnodonts appeared to be more similar to that seen in younger marine derived fish clades today, such as silversides, drums, needlefishes, anchovies and pufferfishes [267–269], where a mostly marine clade made numerous incursions into freshwater habitats throughout their evolution but had the bulk of their diversity present in marine habitats. The freshwater fossil record of pycnodontiforms is, for the most part, very fragmentary, represented by isolated dental remains. Only a few taxa from the Cretaceous are preserved as fully articulated specimens. This is an issue for most freshwater fish fossils as certain habitats, particularly fluvial, are hostile to fossilisation due to the high energy movements of currents. However, despite these limitations, there are certain clues about ecology and life history that can be deduced from certain taxa and their occurrences such as pycnodontiforms from the Early Cretaceous of Spain [121] and Hungary [198,200] being permanent freshwater residents due to the wide range of body and tooth plate sizes, respectively, being present in the same locality indicating the simultaneous presence of juveniles and adults, whereas other pycnodont taxa migrated to freshwaters to reproduce, while otherwise living in marine environments as seen in cf. *Phacodus* [45], and finally, possible niche partitioning as seen in the Kem Kem pycnodont assemblage [47].

Our review of the freshwater pycnodont fossil record is far from being complete assuming it to be rather incomplete due to taphonomic processes and insufficient data. But yet with such limited

data, fascinating and tantalizing glimpses has already been obtained from these specimens. Renewed collection effort in freshwater localities as well as the discovery of new fossil sites combined with isotopic analyses should assist immensely in not only discovering further the extent of pycnodont diversity in non-marine environs but hopefully shed further light on evolutionary questions concerning both extinct and extant clades of freshwater fishes and how they came to dominate the rivers and lakes of the world.

**Author Contributions:** Conceptualization, J.J. Cawley and J. Kriwet; writing—original draft preparation, J.J. Cawley.; writing—review and editing, J. Kriwet. Both authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** All data used for this review is provided in the main body of the text.

**Acknowledgments:** This work is the result of many years of research into pycnodontiform fishes. Many people contributed to the various studies by providing access to collections under their care. It is impossible to name all and we apologize to everybody who is missing in the following list: G. Arratia and H.-P. Schultze (formerly Natural History Museum Berlin, Germany, now Biodiversität Institute & Natural History Museum, The University of Kansas); G. Bergér (Museum Bergér, Eichstätt, Germany); R. Böttcher (now retired) and E. Maxwell (Staatliches Museum für Naturkunde, Stuttgart, Germany); the late P. Forey, A. Longbottom (now retired), Z. Johanson, and E. Bernard (The Natural History Museum London, UK); M.A. Herrero (Museo Paleontológico de Galve, Spain); M. Kölbl-Ebert and C. Ifrim (Jura-Museum Eichstätt, Germany); the late M. Mäuser and O. Wings (Naturkunde Museum Bamberg, Germany); W. Munk (Staatliches Museum für Naturkunde Karlsruhe, Germany); O.W.M. Rauhut, M. Kölbl, W. Werner (now retired), and M. Krings (all Bayerische Staatssammlung für Geologie und Paläontologie), who are all acknowledged for supporting our research. Finally, we would like to thank the reviewers of the present work for the comments and valuable insights. Open Access Funding was provided by the Library of the University of Vienna.

**Conflicts of Interest:** The authors declare no conflicts of interest

#### Appendix A. Taxonomy, affinities and palaeogeographic distribution of nonmarine pycnodontiforms in stratigraphic order from oldest to youngest.

Taxa	Family	Country	Locality	Period	Stage	Ref.
cf. <i>Gyrodus</i>	Gyrodontidae	Thailand	Khlong Min Formation (Mab Ching)	Middle-Late Jurassic	Unknown	[21]
<i>Tibetodus gyroides</i>	Family incertae sedis	China	Tibet	Late Jurassic	Unknown	[25,26]
Pycnodontidea gen et spp. indet	Family incertae sedis	USA	Morrison Formation, Dinosaur National Monument	Late Jurassic	Oxfordian-Tithonian	[27]
cf. <i>Pycnodus</i>	Pycnodontidae	Ethiopia	Abay Gorge, Mughher Mudstone	Late Jurassic	?Kimmeridgian-Tithonian	[39]
<i>Congopycnodus cornutus</i>	Family incertae sedis	Democratic	Stanleyville	Late Jurassic-	Kimmeridgian-	[32]

		Republic of Congo		Early Cretaceous	Valanginia n	
?Pycnodonti formes indet. (tooth morphotype 2)	Family incertae sedis	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77]
aff. <i>Gyrodus</i> (tooth morphotype 6)	Gyrodontidae	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77]
cf. <i>Arcodonichth</i> <i>ys</i> (tooth morphotype 7)	Family incertae sedis	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77,80]
Pycnodonti formes indet. (tooth morphotype 8)	Family incertae sedis	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77]
Pycnodonti dae gen. et spp. Indet (tooth morphotype 9)	Pycnodontidae	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77]
Pycnodonti dae gen. et spp. Indet (tooth morphotype 10)	Pycnodontidae	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77]
Pycnodonti dae gen. et spp. Indet (tooth morphotype 11)	Family incertae sedis	France	Champbla nc Quarry, Cherves- de-Cognac	Early Cretaceous	Berriasian	[77]
<i>Coelodus</i>	Pycnodontidae	England	Lulworth Formation	Early Cretaceous	Berriasian	[26,49]
<i>Coelodus</i> **	Pycnodontidae	England	Durlston Formation	Early Cretaceous	Berriasian	[26,49]

<i>"Pycnodus"</i> <i>mantelli</i>	Pycnodontidae	Germany	Gronau, North Rhine- Westphalia	Early Cretaceous	Berriasian	[29,60]
<i>"Pycnodus"</i> <i>mantelli</i>	Pycnodontidae	Germany	Osterwald/ Otternhage n, Lower Saxony	Early Cretaceous	Berriasian	[29,58]
<i>Turbomesodon</i> <i>cf.</i> <i>arcuatus</i>	Pycnodontidae	Germany	Bückeberg Group, Istenberg Formation	Early Cretaceous	Berriasian	[29,63]
<i>Turbomesodon</i> <i>cf.</i> <i>arcuatus</i>	Pycnodontidae	Germany	North Rhine- Westphalia	Early Cretaceous	Berriasian	[29,60– 62]
Pycnodontif ormes indet.	Family incertae sedis	Germany	Bückeberg Group, Fuhse Formation	Early Cretaceous	Berriasian	[29,55,5 6]
Pycnodontif ormes indet.	Family incertae sedis	Germany	Bückeberg Group, Istenberg Formation	Early Cretaceous	Berriasian	[29,57]
Pycnodontif ormes indet.	Family incertae sedis	Germany	Bückeberg Group, Oesede Formation or Istenberg Formation	Early Cretaceous	Berriasian	[29,58]
Pycnodontif ormes indet.	Family incertae sedis	Germany	Lobber Ort, Island of Rügen	Early Cretaceous	Berriasian	[29,59]
<i>Turbomesodon</i> <i>arcuatus</i>	Pycnodontidae	England	Middle Purbeck Limestone	Early Cretaceous	Berriasian	[29]
<i>Proscinetes</i> sp. cf. <i>P.</i> <i>bernardi</i>	Pycnodontidae	Spain	Lérida	Early Cretaceous	Berriasian	[93,129 –132]
<i>Ocloedus</i> <i>subdiscus</i> **	Pycnodontidae	Spain	El Montsec, Lérida	Early Cretaceous	Berriasian- Valanginia n	[48,89,1 01]

<i>Turbomesodon laevidens</i>	Pycnodontidae	England	Middle Purbeck Limestone	Early Cretaceous	Berriasian-Valanginian	[29]
<i>Ocloedus</i>	Pycnodontidae	England	Pevensey Pit, Ashdown Brickworks quarry (Wealden Group)	Early Cretaceous	Valanginian	[50,51,53]
<i>Turbomesodon microdon</i>	Pycnodontidae	England	Grinstead Clay Formation	Early Cretaceous	Valanginian	[29]
cf. <i>Anomoeodus</i>	Pycnodontidae	Thailand	Sao Khua Formation (Phu Phan Thong)	Early Cretaceous	Hauterivian-Barremian	[21]
<i>Turbomesodon multidentis</i>	Pycnodontidae	England	Weald Clay Group, Sevenoaks, Kent	Early Cretaceous	Hauterivian-Barremian	[29]
<i>Arcodonichthys pasiegae</i>	Family incertae sedis	Spain	Vega de Pas Formation, Basque-Cantabrian Basin	Early Cretaceous	Hauterivian-Barremian	[80]
Pycnodontiformes indet.	Family incertae sedis	Spain	El Castellamar Formation	Early Cretaceous	Berriasian-Barremian	[107]
<i>Stenamaramia</i>	Pycnodontidae	Spain	Las Hoyas	Early Cretaceous	Barremian	[118]
<i>Turbomesodon praeclarus</i>	Pycnodontidae	Spain	Las Hoyas	Early Cretaceous	Barremian	[76]
<i>Coelodus*</i>	Pycnodontidae	England	Wessex (Wealden Group)	Early Cretaceous	Barremian	[46]
Pycnodontiformes indet.	Family incertae sedis	England	Wessex (Wealden Group)	Early Cretaceous	Barremian	[46]

Pycnodonti dae gen. et spp. indet cf. <i>Proscinetes</i> <i>Coelodus</i> <i>Anomoeodus</i>	Pycnodontidae	Spain	Camarillas Formation	Early Cretaceous	Barremian	[109]
	Pycnodontidae	Spain	Camarillas Formation	Early Cretaceous	Barremian	[109]
	Pycnodontidae	Spain	Camarillas Formation	Early Cretaceous	Barremian	[109]
	Pycnodontidae	Spain	Camarillas Formation	Early Cretaceous	Barremian	Kriwet, pers. obser,
Pycnodontif ormes indet.	Family incertae sedis	Spain	Artoles Formation	Early Cretaceous	Barremian	Kriwet, pers. obser,
cf. <i>Ocloedus</i> sp. 1	Pycnodontidae	Spain	Huérgiuna Formation	Early Cretaceous	Barremian	[44]
cf. <i>Ocloedus</i> sp. 2	Pycnodontidae	Spain	Huérgiuna Formation	Early Cretaceous	Barremian	[44]
<i>Anomoeodus</i> <i>nursalli</i>	Pycnodontidae	Spain	Huérgiuna Formation	Early Cretaceous	Barremian	[44]
<i>Coelodus</i>	Pycnodontidae	Spain	Aguilón	Early Cretaceous	Barremian	[93,129 –132]
<i>Turbomesodon</i> <i>bernissartensis</i>	Pycnodontidae	Belgium	Bernissart	Early Cretaceous	Barremian- Aptian	[75,76]
<i>“Palaeobalistum” geiseri</i>	Pycnodontidae	USA	Twin Mountains Formation, Paluxy Church	Early Cretaceous	Aptian	[141]
<i>“Proscinetes” texanus</i>	Pycnodontidae	USA	Twin Mountains Formation, Paluxy Church	Early Cretaceous	Aptian	[30,141]
<i>Coelodus</i>	Pycnodontidae	USA	Twin Mountains Formation, Paluxy Church	Early Cretaceous	Aptian	[141]
<i>Thurmondella estesi</i>	Family incertae sedis	USA	Twin Mountains Formation,	Early Cretaceous	Aptian	[141,14 5]

			Paluxy Church Cloverly Formation, Little Sheep Mudstone Member	Early Cretaceous	Aptian	[133]
Pycnodonti dae indet.	Pycnodontidae	USA				
<i>Anomoeodus complanatus</i>	Pycnodontidae	Spain	Cintorres, Valencia	Early Cretaceous	Aptian	[93,129 -132]
<i>Anomoeodus complanatus</i>	Pycnodontidae	Spain	Mirambel, Tereul	Early Cretaceous	Aptian	[93,129 -132]
<i>Coelodus</i> sp. aff. <i>C. soleri</i>	Pycnodontidae	Spain	Morella, Valencia	Early Cretaceous	Aptian	[93,129 -132]
aff. <i>Gyrodus</i>	Gyrodontidae	Tunisia	Jebel Boulouha	Early Cretaceous	Albian	[152]
<i>Thurmondella a estesi</i>	Family incertae sedis	USA	Paluxy Formation, Stephenville Printing	Early Cretaceous	Albian	[30,141]
<i>Texasensis coronatus</i>	Pycnodontidae	USA	Paluxy Formation, Pecan Valley Estates	Early Cretaceous	Albian	[30,31,1 41]
" <i>Macromesodon</i> " <i>dumblei</i>	Family incertae sedis	USA	Paluxy Formation, Pecan Valley Estates	Early Cretaceous	Albian	[30,141]
<i>Nonaphalgodus trinitensis</i>	Pycnodontidae	USA	Antlers Formation, Butler Farm	Early Cretaceous	Albian	[30,141]
<i>Anomoeodus caddoi**</i>	Pycnodontidae	USA	Holly Creek Formation	Early Cretaceous	Albian	[137]
Pycnodontif ormes indet.**	Family incertae sedis	USA	Holly Creek Formation	Early Cretaceous	Albian	[137]

?Pycnodonti formes indet.**	Family incertae sedis	Tunisia	Oued el Khil	Early Cretaceous	Albian	[152]
Pycnodontif ormes indet.	Family incertae sedis	Brazil	Açu Formation, Potiguar Basin	Early Cretaceous	Albian	[151]
<i>Anomoeodus</i> sp. aff. <i>A.</i> <i>muensteri</i>	Pycnodontidae	Spain	Condemios de Abajo, Castilla- La-Mancha	Early Cretaceous	Albian	[93,129 -132]
Pycnodontif ormes indet.	Family incertae sedis	Spain	Ceceda, Asturias	Early Cretaceous	Albian	[93,129 -132]
<i>Neoproscinetes</i> <i>africanus</i>	Pycnodontidae	Morocco	Kem Kem	Early-Late Cretaceous	Albian- Cenomania n	[47]
cf. <i>Macromesodon</i>	Pycnodontidae	Morocco	Kem Kem	Early-Late Cretaceous	Albian- Cenomania n	[47]
<i>Agassizilia</i> <i>erfoudina</i>	?Pycnodontidae	Morocco	Kem Kem	Early-Late Cretaceous	Albian- Cenomania n	[47]
cf. <i>Coelodus</i>	Pycnodontidae	Morocco	Kem Kem	Early-Late Cretaceous	Albian- Cenomania n	[47]
<i>Coelodus</i> <i>soleri</i>	Pycnodontidae	Spain	Girona, Catalonia	Early-Late Cretaceous	Albian- Cenomania n	[93,129 -132]
Pycnodontif ormes indet.**	Family incertae sedis	Brazil	Alcântara Formation, Laje do Coringa	Late Cretaceous	Cenomania n	[221,22 3]
Pycnodontif ormes indet.**	Family incertae sedis	Algeria	Continenta l Intercalaire , Guir Basin	Late Cretaceous	Cenomania n	[238]
Pycnodonti dae gen. et spp. Indet 1	Pycnodontidae	USA	Cedar Mountain Formation	Late Cretaceous	Cenomania n	[214]
Pycnodonti dae gen. et spp. Indet 2	Pycnodontidae	USA	Cedar Mountain Formation	Late Cretaceous	Cenomania n	[214]

<i>Coelodus</i>	Pycnodontidae	USA	Dakota Formation	Late Cretaceous	Cenomanian	[217]
<i>Coelodus</i>	Pycnodontidae	USA	Straight Cliffs Formation, Smoky Hollow Member	Late Cretaceous	Turonian	[217]
Pycnodontiformes indet./Pycnodontidae indet.**	?Pycnodontidae	Austria	Gams, Schönleiten Formation	Late Cretaceous	Turonian	[192,194]
<i>Coelodus</i>	Pycnodontidae	USA	Straight Cliffs Formation, John Henry Member	Late Cretaceous	Coniacian	[217]
<i>Xinjiangodus gyrodoides</i>	Pycnodontidae	China	Donggou Formation, Junggar Basin	Late Cretaceous	Coniacian-Campanian	[163]
cf. <i>Coelodus</i>	Pycnodontidae	Hungary	Csehbánya Formation, Iharkút	Late Cretaceous	Santonian	[198,201]
<i>Micropynodon</i> **	Family incertae sedis	USA	Straight Cliffs Formation, John Henry Member	Late Cretaceous	Santonian	[217]
Pycnodontiformes indet.	Family incertae sedis	Hungary	Ajka Formation	Late Cretaceous	Santonian	[203]
cf. <i>Phacodus</i>	Family incertae sedis	France	Villeveyrac - L'Olivet	Late Cretaceous	Campanian	[45]
<i>Micropynodon</i> **	Family incertae sedis	USA	Wahweap Formation	Late Cretaceous	Campanian	[217]
Pycnodontidae gen. et spp. indet.	Pycnodontidae	Bolivia	Chaunaca Formation	Late Cretaceous	Campanian	[232]

Pycnodontoida gen. et spp. indet	Family incertae sedis	Spain	Lo Hueco (Cuenca)	Late Cretaceous	Campanian-Maastrichtian	[208–211]
<i>Pycnodus lametae</i>	Pycnodontidae	India	Lameta Formation (Maharashtra)	Late Cretaceous	Maastrichtian	[168,169,174]
cf. <i>Coelodus</i> **	Family incertae sedis	Spain	Els Neret	Late Cretaceous	Maastrichtian	[212]
cf. <i>Coelodus</i> **	Family incertae sedis	Spain	L'Espinau	Late Cretaceous	Maastrichtian	[212]
Pycnodontiformes indet.**	Family incertae sedis	Spain	Serrat del Rostiar-1	Late Cretaceous	Maastrichtian	[212]
Pycnodontiformes indet.**	Family incertae sedis	Spain	Cami del Soldat	Late Cretaceous	Maastrichtian	[212]
Pycnodontiformes indet.**	Family incertae sedis	Spain	Fontllonga-6	Late Cretaceous	Maastrichtian	[212]
?Pycnodontiformes indet.**	Family incertae sedis	Spain	Fontllonga-6	Late Cretaceous	Maastrichtian	[212]
?Pycnodontiformes indet.**	Family incertae sedis	Spain	L'Espinau	Late Cretaceous	Maastrichtian	[212]
Pycnodontidae gen. et spp. indet	Pycnodontidae	India	Madhya Pradesh (Kisalpuri)	Late Cretaceous	Maastrichtian	[180]
Pycnodontidae gen. et spp. indet	Pycnodontidae	India	Lameta Formation (Pisdura)	Late Cretaceous	Maastrichtian	[169,174]
<i>Pycnodus bicresta</i>	Pycnodontidae	India	Naskal, Andhra Pradesh	Late Cretaceous	Maastrichtian	[181,182]
Pycnodontidae gen. et spp. indet**	Pycnodontidae	India	Chhindwara District, Madhya Pradesh	Late Cretaceous	Maastrichtian	[186]
<i>Pycnodus</i>	Pycnodontidae	India	Papro Formation,	Late Cretaceous	Maastrichtian	[177]

			Uttar Pradesh			
<i>"Pycnodus"</i> <i>lametae</i> ***	Pycnodontidae	India	Naskal, Andhra Pradesh	Late Cretaceous	Maastrichtian	[183]
<i>Pycnodus</i>	Pycnodontidae	India	Lameta Formation (Madhya Pradesh)	Late Cretaceous	Maastrichtian	[178,179]
<i>Pycnodus bicresta</i>	Pycnodontidae	India	Asifabad	Late Cretaceous	Maastrichtian	[169,176,186]
<i>Pycnodus lametae</i>	Pycnodontidae	India	Nagpur	Late Cretaceous	Maastrichtian	[184]
<i>Pycnodus lametae</i>	Pycnodontidae	India	Asifabad	Late Cretaceous	Maastrichtian	[184,185]
<i>Pycnodus lametae</i>	Pycnodontidae	India	Rangapur	Late Cretaceous	Maastrichtian	[184]
<i>Pycnodus</i> cf. <i>P. praecursor</i>	Pycnodontidae	India	Asifabad	Late Cretaceous	Maastrichtian	[184]
<i>Coelodus toncoensis</i> **	Pycnodontidae	Argentina	Yacoraite Formation	Late Cretaceous	Maastrichtian	[224–228]
Pycnodontiformes indet.**	Family incertae	Argentina	Yacoraite Formation	Late Cretaceous	Maastrichtian	[231]
<i>Coelodus toncoensis</i> **	Pycnodontidae	Bolivia	El Molino Formation, Paja Patcha	Late Cretaceous	Maastrichtian	[234–236]
<i>Coelodus</i> **	Pycnodontidae	Bolivia	El Molino Formation, Paja Patcha	Late Cretaceous	Maastrichtian	[235]
Pycnodontidae gen. et spp. indet.**	Pycnodontidae	Bolivia	El Molino Formation, Paja Patcha	Late Cretaceous	Maastrichtian	[234,235]
Pycnodontidae gen. et spp. indet.**	Pycnodontidae	Bolivia	El Molino Formation, Vila Vascarra	Late Cretaceous	Maastrichtian	[237]
<i>Coelodus</i>	Pycnodontidae	Madagascar	Maevarano Formation, Anembalemba Member	Late Cretaceous	Maastrichtian	[242]

<i>Pycnodus jonesae</i> **	Pycnodontidae	Mali	Ménaka Formation, Iullemeden Basin	Late Cretaceous	Maastrichtian	[239,240]
<i>Pycnodus</i> **	Pycnodontidae	Mali	Ménaka Formation, Iullemeden Basin	Late Cretaceous	Maastrichtian	[239,240]
<i>Pycnodus</i>	Pycnodontidae	India	Jhilmili, Chhindwara District	Late Cretaceous-Palaeocene	Maastrichtian-Danian	[191]
<i>Pycnodus</i> **	Pycnodontidae	India	Fatehgarh Formation, Rajasthan	Late Cretaceous-Palaeocene	Maastrichtian-Selandian	[189,190]
<i>Coelodus toncoensis</i> **	Pycnodontidae	Argentina	Yacoraite Formation	Palaeocene	Danian	[225]
<i>Pycnodontidae</i> gen. et spp. indet**	Pycnodontidae	India	Rajahmundry	Palaeocene	Danian	[169,247,248]
<i>Pycnodus</i> cf. <i>P. praecursor</i> **	Pycnodontidae	India	Rajahmundry	Palaeocene	Danian	[184]
<i>Pycnodus</i> **	Pycnodontidae	India	Rajahmundry	Palaeocene	Danian	[184]
<i>Pycnodus jonesae</i> **	Pycnodontidae	Mali	Teberemt Formation, Gao Trench Basin	Palaeocene	Selandian-Thanetian	[239,240]
<i>Pycnodus jonesae</i> **	Pycnodontidae	Mali	Teberemt Formation, Iullemeden Basin	Palaeocene	Thanetian	[239,240]
<i>Pycnodus</i> **	Pycnodontidae	Mali	Teberemt Formation, Iullemeden Basin	Palaeocene	Thanetian	[240]
<i>Pycnodus maliensis</i> **	Pycnodontidae	Mali	Tamaguéle It Formation, Taoudenit Basin	Eocene	Ypresian	[239,240]

<i>Pycnodus zeiformis**</i>	Pycnodontidae	Mali	Tamaguéle It Formation, Taoudenit Basin	Eocene	Ypresian	[239,24 0]
<i>Pycnodus**</i>	Pycnodontidae	Mali	Tamaguéle It Formation, Taoudenit Basin	Eocene	Ypresian	[240]
<i>Pycnodus**</i>	Pycnodontidae	USA	Tallahata Formation	Eocene	Ypresian- Lutetian	[250]
<i>Pycnodus**</i>	Pycnodontidae	USA	Lisbon Formation	Eocene	Lutetian	[250]

## References

- Reid, G.M.G. Introduction to freshwater fishes and their conservation. *Int Zoo Yearb* **2013**, *47*, 1–5.
- Oberdorff, T.; Guegan, J.F.; Hugueny, B. Global scale patterns of fish species richness in rivers. *Ecography* **1995**, *18*, 345–352.
- Amarasinghe, U.S.; Welcomme, R.L. An analysis of fish species richness in natural lakes. *Environ Biol Fishes* **2002**, *65*, 327–339.
- Horne, A.J.; Goldman, C. *Limnology*; McGraw-Hill: University of Minnesota, USA, 1994; 480 pp.
- Boyd, C.E. *Water Quality: An Introduction*, 2<sup>nd</sup> ed.; Springer: Auburn, USA, 2015; 440 pp.
- Dodds, W.K.; Whiles, M.R. *Freshwater ecology: concepts and environmental applications of limnology*; 2<sup>nd</sup> ed.; Academic Press: Cambridge, USA, 2010; 829 pp.
- Lundberg, J.G. The temporal context for the diversification of neotropical fishes. In *Phylogeny and Classification of Neotropical Fishes*; Malabarba L.R., Reis R.E., Vari R.P., Lucena, C.A.S., Lucena, Z.M.S., Eds.; EDIPUCRS: University of California, USA, 1998; pp. 49–68.
- Berra, T.M. *An atlas of distribution of the freshwater fish families of the world*; University of Nebraska Press: Lincoln, USA, 1981; 197 pp.
- McDowall, R.M. The occurrence and distribution of diadromy among fishes. *Am Fish Soc Symp* **1987**, *1*, 1–13.
- Cavin, L. *Freshwater fishes: 250 million years of evolutionary history*; ISTE Press Ltd: London, UK, 2017, 214 pp.
- Patterson, C. The distribution of mesozoic freshwater fishes, Biogéographie et liaisons intercontinentales au cours du Mésozoïque. *Mém Mus Natl Hist* **1975**, *11*, 156–174.
- Bogan, S.; Taverne, L.; Agnolin, F. First triassic and oldest record of a South American amiiform fish: *Caturus* sp. from the Los Menucos Group (lower Upper Triassic), Rio Negro province, Argentina. *Geol Belg* **2013**, *16*, 191–195.
- Cavin, L. Diversity of Mesozoic semionotiform fishes and the origin of gars (Lepisosteidae). *Sci Nat* **2010**, *97*, 1035–1040.
- Egerton, P. de M.G. On *Chondrosteus*, an extinct genus of the Sturionidae, found in the Lias Formation at Lyme Regis. *Philos trans R Soc Lond* **1858**, *148*, 871–885.
- Hilton E. J.; Grande L.; Bemis W.E. Morphology of †*Coccolepis bucklandi* Agassiz, 1843 (Actinopterygii, †Coccolepidae) from the Solnhofen Lithographic Limestone deposits (Upper Jurassic, Germany). In *Mesozoic Fishes 3 – Systematics, Paleoenvironments and Biodiversity*; Arratia G., Tintori, A., Eds.; Verlag Dr. Friedrich Pfeil: Munich, Germany, 2004; pp. 209–238.
- Cawley, J.J.; Marramà, G.; Carnevale, G.; Villafañá, J.A.; López-Romero, F.A.; Kriwet, J. Rise and fall of †Pycnodontiformes: Diversity, competition and extinction of a successful fish clade. *Ecol Evol* **2021**, *11*, 1769–1796.
- Nursall, J.R. The case of pycnodont fishes as the fossil sister-group of teleosts. In *Origin and phylogenetic interrelationships of teleosts*; Nelson, J.S., Schultze, H.P., Wilson, M.V.H. Eds.; Verlag Dr. Friedrich Pfeil: Munich, Germany, 2010; pp. 37–60.
- Poyato Ariza, F.J. Studies on pycnodont fishes (I): evaluation of their phylogenetic position among actinopterygians. *Riv Ital Paleontol S* **2015**, *121*, 329–343.

19. Kriwet, J. A comprehensive study of pycnodont fishes (Neopterygii, Pycnodontiformes): morphology, taxonomy, functional morphology, phylogeny, and palaeobiogeography. Unpublished PhD thesis, Humboldt University, Berlin, 2001.
20. Poyato-Ariza, F.J. Pycnodont fishes: morphologic variation, ecomorphologic plasticity, and a new interpretation of their evolutionary history. *Bull Kitakyushu Mus Nat Hist Hum Hist, Ser A* **2005**, *3*, 169–184.
21. Cavin, L.; Deesri, U.; Suteethorn, V. The Jurassic and Cretaceous bony fish record (Actinopterygii, Dipnoi) from Thailand. In *Late Palaeozoic and Mesozoic Ecosystems in SE Asia*; Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V., Eds.; Geological Society: London, UK, 2009; pp. 125–139.
22. Buffetaut, E.; Raksaskulwong, L.; Suteethorn, V.; Tong, H. First post-Triassic temnospondyl amphibians from the Shan-Thai Block: intercentra from the Jurassic of peninsular Thailand. *Geol Mag* **1994**, *131*, 837–839.
23. Buffetaut, E.; Tong, H.; Suteethorn, V.; Raksaskulwong, L. Jurassic vertebrates from the southern peninsula of Thailand and their implications. A preliminary report. In *Proceedings of the International Symposium on Stratigraphic Correlation of Southeast Asia, Bangkok*, Bangkok, Thailand, 15–20 November 1994; Angsuwathana, P., Wongwanich, T., Tansathien, W., Wongsomak, S., Tulyatid, S., Tulyatid, J. Eds.; Department of Mineral Resources: Bangkok, Thailand, 1994; pp. 253–256.
24. Tong, H.; Buffetaut, E.; Suteethorn, V. Middle Jurassic turtles from southern Thailand. *Geol Mag* **2002**, *139*, 687–697.
25. Young, C.C.; Liu, H. *Tibetodus*, a new pycnodont fish from Changtu (in Chinese with English summary). *Acta Palaeo Sin* **1954**, *2*, 95–102.
26. Schaeffer, B.; Patterson, C. Jurassic fishes from the western United States, with comments on Jurassic fish distribution. *Am Mus Novit* **1984**, *2796*, 1–86.
27. Kirkland, J.I. Morrison fishes. *Mod Geol* **1998**, *22*, 503–533.
28. Woodward, A.S. The fossil fishes of the English Wealden and Purbeck formations. Part II. *Palaentogr Soc Monogr* **1918**, *80*, 49–104.
29. Hornung, J. Pycnodonte Fische (Actinopterygii: Pycnodontiformes) in der Unterkreide von Norddeutschland - Diversität und palökologische Beziehungen -- Pycnodont fishes (Actinopterygii: Pycnodontiformes) from the Lower Cretaceous of northern Germany - diversity and paleocological relationships. *Ber Naturwiss Ver Bielef Umgeg* **2021**, *58*, 4–77.
30. Thurmond, J.T. Lower vertebrate faunas of the Trinity Division in north-central Texas. *Geosci Man* **1974**, *8*, 103–129.
31. Özdikmen, H. *Texasensis* nom. nov., a new name for the preoccupied fossil fish genus *Callodus* Thurmond, 1974 (Osteichthyes: Pycnodontiformes). *Mun Ent Zool* **2009**, *4*, 616.
32. Taverne, L. A horny pycnodont fish (Pycnodontiformes) in the continental Middle Jurassic (Stanleyville Formation) of the Democratic Republic of Congo. *Geo-Eco-Trop* **2019**, *43*, 25–34.
33. Myers, T.S.; Tabor, N.J.; Jacobs, L.L. Late Jurassic paleoclimate of central Africa. *Palaeogeogr Palaeoclimatol Palaeoecol* **2011**, *311*, 11–125.
34. Caillaud, A.; Blanpied, C.; Delvaux, D. The Upper Jurassic Stanleyville Group of the eastern Congo Basin: An example of perennial lacustrine system. *J Afr Earth Sci* **2017**, *132*, 80–98.
35. Taverne, L.; Capasso, L. Gladiopycnodontidae, a new family of pycnodontiform fishes from the Late Cretaceous of Lebanon, with the description of three genera. *Eur J Taxon* **2013**, *57*, 1–30.
36. Gayet, M. *Ichthyoceros spinosus* nov. gen., nov. sp., du Cénomaniens inférieur de Hakel (Liban) et ses affinités avec le genre *Trewavasias* (Pisces, Pycnodontiformes, Coccodontidae). *Bull Mus Natl Hist Nat* **1984**, *3*, 287–307.
37. Taverne, L.; Capasso, L. Ostéologie et phylogénie des Coccodontidae, une famille remarquable de poissons Pycnodontiformes du Crétacé supérieur marin du Liban, avec la description de deux nouveaux genres. *Palaeontos* **2014**, *25*, 3–43.
38. Nursall, J.R.; Capasso, L. *Gebrayelichthys* (novum), an extraordinary genus of neopterygian fishes from the Cenomanian of Lebanon. In *Mesozoic Fishes 3: Systematics, Palaeoenvironments and Biodiversity*; Arratia, G., Tintori, A., Eds.; Verlag Dr. Friedrich Pfeil: Munich, Germany, 2004; pp. 317–340.
39. Goodwin, M.B.; Clemens, W.A.; Hutchison, J.H.; Wood, C.B.; Zavada, M.S.; Kemp, A.; Duffin, C.J.; Schaff, C.R. Mesozoic continental vertebrates with associated palynostratigraphic dates from the northwestern Ethiopian plateau. *J Vertebr Paleontol* **1999**, *19*, 728–741.
40. Wolela, A. Sedimentation and depositional environments of the Barremian-Cenomanian Debre Libanose Sandstone, Blue Nile (Abay) Basin, Ethiopia. *Cretac Res* **2009**, *30*, 1133–1145.
41. Getaneh, W.; Atnafu, B. Geochemistry and lithostratigraphy of the mugher mudstone: Insights into the late jurassic-early Cretaceous clastic sedimentation in Ethiopia and its surroundings. *J Afr Earth Sci* **2020**, *164*, 103770.
42. Cawley, J.J.; Marrama, G.; Carnevale, G.; Kriwet, J. A quantitative approach to determine the taxonomic identity and ontogeny of the pycnodontiform fish *Pycnodus* (Neopterygii, Actinopterygii) from the Eocene of Bolca Lagerstätte, Italy. *PeerJ* **2018**, *6*, e4809.

43. Cuny, G. 2002. Requins d'eau douce, actuels et fossiles. *PSci* **2002**, 295, 30–36.
44. Kriwet, J. Pycnodont fishes (Neopterygii, Pycnodontiformes) from the Lower Cretaceous of Uña (E-Spain) with comments on branchial teeth in pycnodontid fishes. In *Mesozoic Fishes 2: Systematics and Fossil Record*; Arratia, G., Schultze, H.P., Eds.; Verlag Dr. Friedrich Pfiel: Munich, Germany, 1999; pp. 215–238.
45. Cavin, L.; Garcia, G.; Valentin, X. A minute freshwater pycnodont fish from the Late Cretaceous of southern France: Palaeoecological implications. *Cretac Res* **2020**, 106, 104242.
46. Sweetman, S.C.; Goedert, J.; Martill, D.M. A preliminary account of the fishes of the Lower Cretaceous Wessex Formation (Wealden Group, Barremian) of the Isle of Wight, southern England. *Biol J Linn Soc* **2014**, 113, 872–896.
47. Cooper, S.L.; Martill, D.M. A diverse assemblage of pycnodont fishes (Actinopterygii, Pycnodontiformes) from the mid-Cretaceous, continental Kem Kem Group of south-east Morocco. *Cretac Res* **2020**, 112, p.104456.
48. Poyato-Ariza, F.J.; Wenz, S. A new insight into pycnodontiform fishes. *Geodiversitas* **2002**, 24, 139–248.
49. Coram, R.A.; Radley, J.D. 2021. Revisiting climate change and palaeoenvironments in the Purbeck Limestone Group (Tithonian–Berriasian) of Durlston Bay, southern UK. *Proc Geol Assoc Lond* **2021**, 132, 392–404.
50. Austen, P.; Brockhurst, D.; Honeysett, K. Vertebrate fauna from Ashdown brickworks, Bexhill, east Sussex. *Wealden News* **2010**, 8, 13–23.
51. Naish, D.; Sweetman, S.C. A tiny maniraptoran dinosaur in the Lower Cretaceous Hastings Group: evidence from a new vertebrate-bearing locality in south-east England. *Cretac Res* **2011**, 32, 464–471.
52. Allen, P. Wealden of the Weald: a new model. *Proc Geol Assoc* **1975**, 86, 389–437.
53. Turmine-Juhel, P.; Wilks, R.; Brockhurst, D.; Austen, P.A.; Duffin, C.J.; Benton, M.J. Microvertebrates from the Wadhurst Clay Formation (Lower Cretaceous) of Ashdown Brickworks, East Sussex, UK. *Proc Geol Assoc* **2019**, 130, 752–769.
54. Hayward, R.J. The Geology of part of the Wadhurst Clay Formation and part of the Tunbridge Wells Formation at the Ashdown Brickworks, Bexhill, Sussex. Unpublished B.Sc. dissertation, University of Greenwich, London, 1996.
55. Struckmann, C. Die Wealdenbildungen von Sehnde bei Lehrte. *Neues Jahrb Mineral Geol Palaeontol* **1891**, 1891, 117–131.
56. Jordan, R. Die Ziegeleitongrube westlich Wätzum, Blatt Sarstedt, an der Ostflanke des Salzstockes von Sarstedt-Lehrte (Dogger, Wealden, marine Unterkreide). *Ber Naturhist Ges Hann* **1959**, 104, 5–24.
57. Struckmann, C. Geognostische Studien am Deister. *Jahresber Naturhist Ges Hann* **1880**, 29/30, 60–75.
58. Dunker, W. *Monographie der Norddeutschen Wealdenbildung. Ein Beitrag zur Geognosie und Naturgeschichte der Vorwelt*. Oehme und Müller: Rome, Italy, 1846; 83 pp.
59. Ansoerge, J. Fischreste (Selachii, Actinopterygii) aus der Wealdentonscholle von Lobber Ort (Mönchgut/Rügen/DDR). *PalZ* **1990**, 64, 133–144.
60. Nyhuis, C.; Herbig, H.G. Ichthyolithen aus dem Berriasium von Gronau/Westfalen (westliches niedersächsisches Kreidebecken) – Rekonstruktion einer trophischen Kette aus sturmkondensierten Bonebeds. *Terra Nostra* **2009**, 2009, 85.
61. Kemper, E. *Geologischer Führer durch die Grafschaft Bentheim und die angrenzenden Gebiete*. 3rd edition. Verlag Heimatverein der Grafschaft Bentheim: Nordhorn, Germany, 1968; 172 pp.
62. Kemper, E. *Geologischer Führer durch die Grafschaft Bentheim und die angrenzenden Gebiete mit Abriss einem der emsländischen Unterkreide*. Verlag Heimatverein der Grafschaft Bentheim: Nordhorn, Germany, 1976; 206 pp.
63. Kriwet, J. A comprehensive study of the skull and dentition of pycnodont fishes. *Zitteliana* **2005**, 45, 135–188.
64. Olive, S.; Taverne, L.; López-Arbarello, A. A new genus of coccolepidid actinopterygian from the Cretaceous *Iguanodon*-bearing locality of Bernissart, Belgium. *Cretac Res* **2019**, 95, 318–335.
65. Cavin, L.; Deesri, U.; Olive, S. *Scheenstia bernissartensis* (Actinopterygii: Ginglymodi) from the Early Cretaceous of Bernissart, Belgium, with an appraisal of ginglymodian evolutionary history. *J Syst Palaeontol* **2020**, 18, 513–527.
66. Cornet, J.; Schmitz, G. Note sur les puits naturels du terrain houiller du Hainaut et le gisement des iguanodons de Bernissart-Bull. *Bull Soc belge Géol Paléontol Hydrol* **1898**, 12, 301–318.
67. Cornet, J. L'époque wealdienne dans le Hainaut. *Bull Soc belge Géol Paléontol Hydrol* **1927**, 50, 89–104.
68. Yans, J. *Lithostratigraphie, minéralogie et diagenèse des sédiments à faciès wealdien du Bassin de Mons (Belgique)*. Académie royale de Belgique: Brussels, Belgium, 2007; 179 pp.
69. Schnyder, J.; Dejax, J.; Keppens, E.; Tu, T.T.N.; Spagna, P.; Boulila, S.; Galbrun, B.; Riboulleau, A.; Tshibangu, J.P.; Yans, J. An Early Cretaceous lacustrine record: organic matter and organic carbon isotopes at Bernissart (Mons Basin, Belgium). *Palaeogeogr Palaeoclimatol Palaeoecol* **2009**, 281, 79–91.
70. Spagna, P. Les faciès wealdiens du Bassin de Mons (Belgique): paléoenvironnements, géodynamique et valorisation industrielle. Unpublished PhD thesis, Faculté Polytechnique de l'Umons, Mons, 2010.

71. Spagna, P.; Yans, J.; Schnyder, J.; Dupuis, C. The Paleoenvironment of the Bernissart Iguanodons: Sedimentological Analysis of the Lower Cretaceous Wealden Facies in the Bernissart Area. In *Bernissart Dinosaurs and Early Terrestrial Ecosystems*; Godefroit, P., Ed.; Indiana University Press: Bloomington, USA, 2012; pp. 87–96.
72. Dejax, J.; Pons, D.; Yans, J. Palynology of the dinosaur-bearing Wealden facies in the natural pit of Bernissart (Belgium). *Rev Palaeobot Palynol* **2007**, *144*, 25–38.
73. Yans, J.; Dejax, J.; Pons, D.; Taverne, L.; Bultynck, P. The iguanodons of Bernissart (Belgium) are middle Barremian to earliest Aptian in age. *Bull Inst R Sci Nat Belg Sci Terre* **2006**, *76*, 91–95.
74. Yans, J.; Dejax, J.; Schnyder, J. On the age of the Bernissart Iguanodons. In *Bernissart Dinosaurs and Early Terrestrial Ecosystems*; Godefroit, P., Ed.; Indiana University Press: Bloomington, USA, 2012; pp. 79–86.
75. Traquair, R.H. Les poissons wealdiens de Bernissart. *Mém Mus R Hist Nat Belg* **1911**, *6*, 1–65.
76. Poyato-Ariza, F.J.; Wenz, S. The new pycnodontid fish genus *Turbomesodon*, and a revision of *Macromesodon* based on new material from the Lower Cretaceous of Las Hoyas, Cuenca, Spain. In *Mesozoic Fishes 3: Systematics, Paleoenvironments and Biodiversity*; Arratia, G., Tintori, A., Eds.; Verlag Dr. Friedrich Pfeil: Munich, Germany, 2004; pp. 341–378.
77. Pouech, J.; Mazin, J.M.; Cavin, L.; Poyato-Ariza, F.J. A Berriasian actinopterygian fauna from Cherves-de-Cognac, France: biodiversity and palaeoenvironmental implications. *Cretac Res* **2015**, *55*, 32–43.
78. Mazin, J.M.; Pouech, J.; Hantzpergue, P.; Lenglet, T. The Purbeckian site of Cherves-de-Cognac (Berriasian, Early Cretaceous, SW France): a first synthesis. In *Mid-Mesozoic Life and Environments Symposium*, Cognac, France, 24-28 June 2008; Mazin, J.M., Pouech, J., Hantzpergue, P., Lacombe, V., Eds.; Documents des Laboratoires de Géologie de Lyon: Lyon: Cognac, France, 2008; pp. 68–71.
79. Pouech, J. Position des mammifères dans les écosystèmes mésozoïques d'Europe Occidentale: le gisement de Cherves-de-Cognac (Berriasien, Charente, France). Unpublished PhD thesis, Claude Bernard University Lyon 1, Lyon, 2008.
80. Poyato-Ariza, F.J.; Bermúdez-Rochas, D.D. New pycnodont fish (*Arcodonichthys pasiegae* gen. et sp. nov.) from the Early Cretaceous of the Basque-Cantabrian Basin, Northern Spain. *J Vertebr Paleontol* **2009**, *29*, 271–275.
81. Cooper, S.L.; Martill, D.M. Pycnodont fishes (Actinopterygii, Pycnodontiformes) from the Upper Cretaceous (lower Turonian) Akrabou Formation of Asfla, Morocco. *Cretac Res* **2020**, *116*, p.104607.
82. Pouech, J.; Amiot, R.; Lecuyer, C.; Mazin, J.M.; Martineau, F.; Fourel, F. Oxygen isotope composition of vertebrate phosphates from Cherves-de-Cognac (Berriasian, France): environmental and ecological significance. *Palaeogeogr Palaeoclimatol Palaeoecol* **2014**, *410*, 290–299.
83. Kriwet, J. Dental morphology of the pycnodontid fish †*Stemmatodus rhombus* (Agassiz 1844) (Neopterygii, {Pycnodontiformes}) from the Early Cretaceous, with comments on its systematic position. *Trans R Soc Edinb Earth Sci* **2004**, *94*, 145–155.
84. Bermúdez-Rochas, D.D.; Delvene, G.; Moratalla, J.; Hernán, J.; De La Fuente, M. Primeros datos paleontológicos del yacimiento del Cretácico Inferior Vega de Pas 1 (Cuenca Vasco-Cantábrica, Cantabria, España). In *II Semana de Jóvenes Investigadores del IGME*; Bermúdez, D.D., Najarro, M., Quesada, C., Eds.; Publicaciones del Instituto Geológico y Minero de España: Madrid, Spain, 2007; pp. 23–28.
85. Darras, L.P.G. The evolution of macroecological consequences of grazing and shell-crushing in fishes. Unpublished PhD thesis, University of Leicester, Leicester, 2012.
86. Kriwet, J. Feeding mechanisms and ecology of pycnodont fishes (Neopterygii, Pycnodontiformes). *Fossil Record* **2001**, *4*, 139–165.
87. Kriwet, J. Late Jurassic elasmobranch and actinopterygian fishes from Portugal and Spain. *Cuad Geol Ibérica* **1998**, *24*, 241–260.
88. Gómez-Alba, J. Catálogo razonado de los vertebrados fósiles de España del Museo de Geología de Barcelona (1882-1982). *Treb Mus Geol Barc* **1997**, *6*, 1–296.
89. Kriwet, J.; Poyato-Ariza, F.J.; Wenz, S. A revision of the pycnodontid fish *Coelodus subdiscus* Wenz 1989, from the Early Cretaceous of Montsec (Lleida, Spain). *Treb Mus Geol Barc* **1999**, *8*, 33–65.
90. Vidal, L.M. Nota sobre la presencia del tramo Kimeridgense en el Montsec (Lérida) y hallazgo de un batracio en sus hiladas. *Mems R Acad Cienc Artes Barcelona* **1902**, *4*, 1–12.
91. Peybernés, B.; Oertli, H. La série de passage du Jurassique au Crétacé dans le Bassin sud-pyrénéen (Espagne). *C R Acad Sci* **1972**, *274*, 3348–3351.
92. Brenner, P. Ostracoden und Charophyten des spanischen Wealden (Systematik, Gkologie, Stratigraphie, Palaogeographie). *Palaeontogr A* **1976**, *152*, 113–201.
93. Gomez Pallerola, J.E. Nuevas aportaciones a la ictiofauna y a la flora del Neocomiense del Montsec de Rubies (Lérida). *Bol Geol Min* **1982**, *93*, 199–213.
94. Gomez Pallerola, J.E. Nuevos Hybodóntidos del Cretácico Inferior de Santa Maria de Meyá (Lérida). *Bol Geol Min* **1985**, *96*, 372–380.
95. Gomez Pallerola, J.E. Nota sobre los tiburones hybodontos de las calizas litográficas del Cretácico Inferior del Motsec (Lérida). *Bol Geol Min* **1992**, *103*, 783–813.

96. Wenz, S. Note préliminaire sur la faune ichthyologique du Jurassique supérieur du Montsec (Espagne). *Bull Soc géol France* **1968**, *7*, 116–119.
97. Wenz, S. Les Amiidés (Pisces, Halecomorphi) du Crétacé inférieur du Montsec (province de Lérida, Espagne): *Amiopsis woodwardi* (Sauvage, 1903). *Quaderns Institut d'Estudis Ilerdencs* **1988**, *1*, 1–50.
98. Wenz, S. Peixos del Cretaci Inferior de la Serra del Montsec (Espanya). In *Les calcàries litogràfiques del Cretaci inferior del Montsec. Deu anys de campanyes paleontològiques*; Martínez-Delclòs, X., Ed.; Institut d'Estudis Ilerdencs: Lleida, Spain, 1991; pp. 111–132.
99. Poyato-Ariza, F.J. Teleósteos primitivos del Cretácico inferior español: órdenes Elopiformes y Gonorhynchiformes. Unpublished PhD Thesis, Universidad Autónoma, Madrid, 1991.
100. Wenz, S.; Poyato-Ariza, F.J. Les actinoptérygiens juvéniles du Crétacé inférieur du Montsec et de Las Hoyas (Espagne). *Geobios* **1994**, *27*, 203–212.
101. Wenz, S. Une nouvelle espèce de *Coelodus* (Pisces, Pycnodontiformes) du Crétacé inférieur du Montsec (Province de Lérida, Espagne): *Coelodus subdiscus* n. sp. *Geobios* **1989**, *22*, 515–520.
102. Martínez-Delclòs, X.; Nel, A. Arthropods. In *Montsec and Montral-Alcover, Two Konservat-Lagerstätten, Catalonia, Spain*; Martínez-Delclòs, X., Ed.; Institut d'Estudis Ilerdencs: Lleida, Spain, 1995; pp. 39–46.
103. De Gibert, J.M.; Fregenal-Martínez, M.A.; Buatois, L.A.; Mangano, M.G. Trace fossils and their palaeoecological significance in Lower Cretaceous lacustrine conservation deposits, El Montsec, Spain. *Palaeogeogr Palaeoclimatol Palaeoecol* **2000**, *156*, 89–101.
104. Schairer, G.; Janicke, V. Sedimentologisch-paläontologische Untersuchungen an den Plattenkalken der Sierra de Montsec (Prov. Lérida. NE-Spanien). *Neues Jahrb Geol Paläont Abh* **1970**, *135*, 171–189.
105. Schultze, H.P. The fossil record of the intertidal zone. In *Intertidal fishes: Life in two worlds*; Horn, M.H., Martin, K.L.M., Chotkowski, M.A. Eds.; Academic Press: San Diego, 1999; pp. 373–392.
106. Cuenca-Bescós, G.; Amo, O.; Aurell, M.; Buscalioni, A.D.; Canudo, J.I.; Laplana, C.; Pérez Oñate, J.; Ruiz Omeñaca, J.I.; Sanz, J.L.; Soria, A.R. Los vertebrados del tránsito Jurásico-Cretácico de Galve (Teruel). In *Comunicaciones de las X Jornadas Paleontología*, Madrid, Spain, 3-5 November 1994; Sociedad Española de Paleontología, Madrid, Spain, 1994; pp. 50–53.
107. Sánchez-Hernández, B.; Benton, M.J.; Naish, D. Dinosaurs and other fossil vertebrates from the Late Jurassic and Early Cretaceous of the Galve area, NE Spain. *Palaeogeogr Palaeoclimatol Palaeoecol* **2007**, *249*, 180–215.
108. Díaz, M.; Yébenes, A. La sedimentación litoral y continental durante el Cretácico Inferior. Sinclinal de Galve, Teruel. *Estud Geol* **1987**, *43*, 3–21.
109. Soria, A.R.; Meléndez, A.; Cuenca-Bescós, G.; Canudo, J.I.; Liesa, C.L. Los sistemas lacustres del Cretácico Inferior de la Cordillera Ibérica Central: la Cubeta de Aliaga. In *XIII Congreso Español de Sedimentología. Guía de Excursiones*, Zaragoza, Spain, 28-30 June 1995; Meléndez, A., Aurell, M., Eds.; Universidad de Zaragoza: Zaragoza, Spain, 1995; pp. 91–141.
110. Kriwet, J. Neoselachier (Pisces, Elasmobranchii) aus der Unterkreide (unteres Barremium) von Galve und Alcaine (Spanien, Provinz Teruel). *Palaeo Ichthyol* **1999**, *9*, 113–142.
111. Henkel, S.; Krebs, B. Zwei Säugetier-Unterkiefer aus der Unteren Kreide von Uña (Prov. Cuenca, Spanien). *Neues Jahrb Geol Paläont Mh* **1969**, *1969*, 449–463.
112. Krebs, B. The search for Mesozoic Mammals in Spain and Portugal. *Mesozoic Vert Life* **1980**, *1*, 23–25
113. Krebs, B. The Barremian vertebrate locality Uña (Province of Cuenca) materials for a comparison with Las Hoyas. In *2nd International Symposium on Lithographic Limestones, Lleida - Cuenca (Spain), Extended Abstracts*, Madrid, Spain, 9-16 July 1995; Meléndez, N., Ed.; Ediciones de la Universidad Autónoma de Madrid: Madrid, Spain, 1995, pp. 95–97.
114. Geyer, O.F., Krautter, M. Die Unterkreide im Grenzgebiet von Aragonien, Neukastilien und Valencia (südwestliches Keltiberikum, Spanien). *Profil* **1998**, *15*, 163–239.
115. Sanz, J. L.; Wenz, S.; Yébenes, A.; Estes, R.; Martínez-Delclòs, X.; Jiménez-Fuentes, E.; Diéguez, C.; Buscalioni, A.D.; Barbadillo, L.J.; Vía, L. An Early Cretaceous faunal and floral continental assemblage: Las Hoyas fossil-site (Cuenca, Spain). *Geobios* **1988**, *21*, 611–631.
116. Poyato-Ariza, F.J.; Wenz, S. Ichthyofauna. In *Las Hoyas. A Lacustrine Konservat-Lagerstätte, Cuenca, Spain. Field trip guide book of the II International Symposium on Lithographic Limestones*; Meléndez, N., Ed.; Universidad Complutense de Madrid: Madrid, Spain, 1995; pp. 43–49.
117. Poyato-Ariza, F.J. Palaeoecology of the fishes from the Early Cretaceous lake of Las Hoyas Cuenca, Spain, with a hypothesis of sexual dimorphism for the Chanidae *Rubiesichthys*. *Bull Kitakyushu Mus Nat Hist Hum Hist, Ser A* **2005**, *3*, 153–168.
118. Poyato-Ariza, F.J.; Wenz, S. A new pycnodontiform fish from the Early Cretaceous of Las Hoyas (Cuenca, Spain). *Bull Soc géol Fr* **2000**, *171*, 251–258.
119. Poyato-Ariza, F.J.; Talbot, M.R.; Fregenal-Martínez, M.A.; Meléndez, N.; Wenz, S. First isotopic and multidisciplinary evidence for nonmarine coelacanths and pycnodontiform fishes: palaeoenvironmental implications. *Palaeogeogr Palaeoclimatol Palaeoecol* **1998**, *144*, 65–84.

120. Wenz, S.; Poyato-Ariza, F.J. Les Actinoptérygiens juvéniles du Crétacé inférieur du Montsec et de Las Hoyas. *Geobios Mem Spec* **1994**, *16*, 203–212.
121. Wenz, S.; Poyato-Ariza, F.J. Pycnodontiform fishes from the early Cretaceous of Las Hoyas (Spain). In *Extended abstracts: II International Symposium on Lithographic Limestones, Cuenca, Spain, July 1995*, Cuenca, Spain, 9-16 July 1995; Ediciones de la Universidad Autónoma de Madrid, Madrid, Spain, 1995; pp. 157–161.
122. Martín-Closas, C.; Diéguez, C. Charophytes from the Lower Cretaceous of the Iberian ranges (Spain). *Palaeontology* **1998**, *41*, 1133–1152.
123. Tintori, A. Two new pycnodonts (Pisces, Actinopterygii) from the Upper Triassic of Lombardy (N. Italy). *Riv Ital Paleontol Stratigr* **1981**, *86*, 795–824.
124. Capasso, L.L.; Abi Saad, P.; Taverne, L. *Nursallia tethysensis* sp. nov., a new pycnodont fish (Neopterygii: Halecostomi) from the Cenomanian of Lebanon. *Bull Inst R Sci Nat Belg Sci Terre* **2009**, *79*, 117–136.
125. Ebert, M. A new genus of Pycnodontidae (Actinopterygii) from the Upper Jurassic of France and Germany, included in a phylogeny of Pycnodontiformes. *Zool J Linn Soc* **2020**, *188*, 434–454.
126. Vullo, R.; Bardet, N.; Gheerbrant, E.; Jalil, N. Multicuspid tooth morphology in a gigantic Palaeocene pycnodont fish: Evolutionary and palaeoecological significance. *Geol Mag* **2019**, *156*, 1618–1622.
127. Baines, D.C. Tooth Microwear in Fishes. Unpublished PhD thesis, University of Leicester, Leicester, 2010.
128. Gibert, J.M. De; Buatois, L.A.; Fregenal-Martínez, M.A.; Gabriela Mangano, M.; Ortega, F.; Poyato-Ariza, F.J.; Wenz, S. The fish trace fossil *Undichna* from the Cretaceous of Spain. *Palaeontology* **1999**, *42*, 409–427.
129. Bauzá Rullán, J. Contribución al conocimiento de la fauna ictiológica de Cataluña. *Estud Geol* **1948**, *8*, 241–242.
130. Bauzá Rullán, J. Contribución a la fauna ictiológica de España. *Brev Geol Asturica* **1957**, *1*, 7–19.
131. Bataller, J.R. Los vertebrados del Cretácico español. *Not Comunic Inst Geol Miner España* **1960**, *60*, 141–164.
132. Soría, A.R. La sedimentación en las cuencas marginales del surco Ibérico durante el Cretácico Inferior y su control estructural. Unpublished PhD thesis, Universidad de Zaragoza, Zaragoza, 1996.
133. Oreska, M.P.; Carrano, M.T.; Dzikiewicz, K.M. Vertebrate paleontology of the Cloverly Formation (Lower Cretaceous), I: faunal composition, biogeographic relationships, and sampling. *J Vertebr Paleontol* **2013**, *33*, 264–292.
134. D'Emic, M.D.; Foreman, B.; Jud, N.A.; Britt, B.B.; Schmitz, M.; Crowley, J.L. Chronostratigraphic revision of the Cloverly Formation (Lower Cretaceous, Western Interior, USA). *Bull Peabody Mus Nat Hist* **2019**, *60*, 3–40.
135. Rogers, R.R.; Brady, M.E. Origins of microfossil bonebeds: insights from the Upper Cretaceous Judith River Formation of north-central Montana. *Paleobiology* **2010**, *36*, 80–112.
136. Moberly, R. Morrison, Cloverly, and Sykes Mountain formations, northern Bighorn Basin, Wyoming and Montana. *Geol Soc Am Bull* **1960**, *71*, 1137–1176.
137. Suarez, C.A.; Frederickson, J.; Cifelli, R.L.; Pittman, J.G.; Nydam, R.L.; Hunt-Foster, R.K.; Morgan, K. A new vertebrate fauna from the Lower Cretaceous Holly Creek Formation of the Trinity Group, southwest Arkansas, USA. *PeerJ* **2021**, *9*, p.e12242.
138. Frucci, M.N. 2018, Oxygen isotopic composition of aquatic reptile phosphate from the Early Cretaceous Holly Creek Formation, southern Arkansas. Unpublished B.S. Honors Thesis, University of Arkansas, Fayetteville, 2018.
139. Thurmond, J.T. *Lower vertebrates and paleocology of the Trinity Group (Lower Cretaceous) in North Central Texas*; Southern Methodist University: Southern Methodist University, USA, 1969; 128 pp.
140. Winkler, D.A.; Murry, P.A.; Jacobs, L.L. Vertebrate paleontology of the Trinity Group, Lower Cretaceous of central Texas. In *Field guide to the vertebrate paleontology of the Trinity Group, Lower Cretaceous of central Texas*; Winkler, D.A., Murry, P.A., Jacobs, L.L., Eds.; Institute for the Study of the Earth and Man: Dallas, USA, 1989; pp.1-22.
141. Winkler, D.A.; Murry, P.A.; Jacobs, L.L. Early Cretaceous (Comanchean) vertebrates of central Texas. *J Vertebr Paleontol* **1990**, *10*, 95–116.
142. Cifelli, R.L.; Gardner, J.D.; Nydam, R.L.; Brinkman, D.L. Additions to the vertebrate fauna of the Antlers Formation (Lower Cretaceous), Southwestern Oklahoma. *Oklahoma Geol Notes* **1997**, *57*, 124–131.
143. Caughey, C.A. *Depositional systems in the Paluxy Formation (Lower Cretaceous), Northeast Texas—oil, gas, and groundwater resources*; Bureau of Economic Geology: Austin, USA, 1977; 59 pp.
144. Thurmond, J.T. Cartilaginous fishes of the Trinity Group and related rocks (Lower Cretaceous) of north central Texas. *Southeast Geol* **1971**, *13*, 207–227.
145. Özdikmen, H.; Akbulut, A. *Thurmondella* nom. nov., A new name for the preoccupied genus *Paramicrodon* Thurmond, 1974 (Osteichthyes: Actinopterygii: Pycnodontiformes). *Mun Ent Zool* **2012**, *7*, 1280–1281.
146. Costa, A.B.C.; Córdoba, V.C.; Netto, R.G.; Lima Filho, F.P. Registro faciológico e paleoambiental da transgressão que marca a passagem do Cenomaniano para o Turoniano na Bacia Potiguar, NE do Brasil. *Comun Geol* **2014**, *101*, 415–420.
147. Araripe, P.T.; Feijó, F.J. Potiguar Basin; Bacia Potiguar. *Bol geociênc Petrobras* **1994**, *8*, 127–141.

148. Maraschin, A.; Mizusaki, A.M.; Vasconcelos, P.M.; Hinrichs, R.; Luiz, D.E.; Dos Anjos, S. Depositional age definition of the Açu Formation (Potiguar Basin, northeastern Brazil) through 40 Ar- 39 Ar dating of early-aauthigenic K-feldspar overgrowths. *Pesqui em Geocienc* **2010**, *37*, 85–96.
149. Arai, M. Paleogeografia do Atlântico Sul no Aptiano: um novo modelo a partir de dados micropaleontológicos recentes. *Bol geociênc Petrobras* **2009**, *17*, e351.
150. Arai, M. Aptian/Albian (early Cretaceous) paleogeography of the South Atlantic: a paleontological perspective. *Braz J Geology* **2014**, *44*, 339–350.
151. Veiga, I.M.M.G.; Bergqvist, L.P.; Brito, P.M. The fish assemblage of the Cretaceous (? Albian-Cenomanian) Açu Formation, Potiguar Basin, Northeastern Brazil. *J South Am Earth Sci* **2019**, *93*, 162–173.
152. Cuny, G.; Cobbett, A.M.; Meunier, F.J.; Benton, M.J. Vertebrate microremains from the Early Cretaceous of southern Tunisia. *Geobios* **2010**, *43*, 615–628.
153. Anderson, P.E.; Benton, M.J.; Trueman, C.N.; Paterson, B.A.; Cuny, G. Palaeoenvironments of vertebrates on the southern shore of Tethys: The nonmarine Early Cretaceous of Tunisia. *Palaeogeogr Palaeoclimatol Palaeoecol* **2007**, *243*, 118–131.
154. Kriwet, J.; Schmitz, L. New insight into the distribution and palaeobiology of the pycnodont fish *Gyrodus*. *Acta Palaeontol Pol* **2005**, *50*, 49–56.
155. Srarfi, D. Biostratigraphie, biodiversité, taphonomie and paléoenvironnement des niveaux à vertébrés du Jurassique–Crétacé du Sud-Est de la Tunisie. Implications paléobiogéographiques. Unpublished PhD thesis, Claude Bernard University Lyon 1, Lyon, 2006.
156. Russell, D.A. Isolated dinosaur bones and teeth from the Middle Cretaceous of the Tafilalt, Morocco. *Bull Mus Natl Hist Nat 4C* **1996**, *18*, 349–402.
157. Dal Sasso, C.; Maganuco, S.; Buffetaut, E.; Mendez, M.A. New information of the skull of the enigmatic theropod *Spinosaurus*, with remarks on its size and affinities. *J Vertebr Paleontol* **2005**, *25*, 888–896.
158. Ibrahim, N.; Sereno, P.C.; Dal Sasso, C.; Maganuco, S.; Fabri, M.; Martill, D.M.; Zouhri, S.; Myhrvold, N.; Lurino, D.A. Semi-aquatic adaptations in a giant predatory dinosaur. *Science* **2014**, *345*, 1613–1616.
159. Sereno, P.C.; Dutheil, D.B.; Iarochene, M.; Larsson, H.C.; Lyon, G.H.; Magwene, P.M.; Sidor, C.A.; Varricchio, D.J.; Wilson, J.A. Predatory dinosaurs from the Sahara and Late Cretaceous faunal differentiation. *Science* **1996**, *272*, 986–991.
160. Martill, D.M.; Ibrahim, N. Aberrant rostral teeth of the sawfish *Onchopristis numidus* from the Kem Kem beds (? early Late Cretaceous) of Morocco and a reappraisal of *Onchopristis* in New Zealand. *J Afr Earth Sci* **2012**, *64*, 71–76.
161. Ibrahim, N.; Sereno, P.C.; Varricchio, D.J.; Martill, D.M.; Dutheil, D.B.; Unwin, D.M.; Baidder, L.; Larsson, H.C.; Zouhri, S.; Kaoukaya, A. Geology and paleontology of the Upper Cretaceous Kem Kem group of eastern Morocco. *ZooKeys* **2020**, *928*, 1–216.
162. Yabumoto, Y.; Uyeno, T. New material of a Cretaceous coelacanth, *Mawsonia lavocati* Tabaste from Morocco. *Bull Natn Sci Mus Tokyo* **2005**, *31*, 39–49.
163. Zhou, X.; You, J.; Jin, J.; Feng, Q.; Luo, Z.; Hu, J.; Li, Y. A minute freshwater pycnodont fish from the Late Cretaceous on the southern margin of the Junggar Basin: Palaeoecological implications. *Hist Biol* **2023**, *35*, 1528–1535.
164. Hao, Y.C.; Su, D.Y.; Yu, J. X.; Li, P.X.; Li, Y.G.; Wang, N.W.; Qi, H.; Guan, S.Z.; Hu, H.G.; Liu, X.; Yang, W.D.; Ye, L.S.; Shou, Z.X.; Zhang, Q.B. *Strata of China 12: Cretaceous*; Geological Publishing House: Beijing, China, 1986; 301 pp.
165. Chen, P. A survey of the non-marine Cretaceous of China. *Cretac Res* **1983**, *4*, 123–143.
166. Zheng, X.L.; Zheng, X.M.; Zheng, X.S.; Li, J.G.; Yang, Y.M.; Weng, Y.X.; Wu, C.X. The late Cretaceous ostracod fossils of the Junggar Basin (in Chinese). *J Stratigr* **2013**, *37*, 206–209.
167. Xi, D.; Wan, X.; Li, G.; Li, G. Cretaceous integrative stratigraphy and timescale of China. *Sci China Earth Sci* **2019**, *62*, 256–286.
168. Woodward, A.S. On some fish-remains from the Lameta Beds at Dongargaon, Central Provinces. *Palaeont Ind* **1908**, *3*, 1–6.
169. Prasad, G.V.R. Vertebrate biodiversity of the Deccan volcanic province of India: A review. *Bull Soc géol Fr* **2012**, *183*, 597–610.
170. Sarkar, A.; Bhattacharya, S.K.; Mohabey, D.M. Stable isotope analyses of dinosaur eggshells: Palaeoenvironmental implications. *Geology* **1991**, *19*, 1068–1107.
171. Ghosh, P.; Bhattacharya, S.K.; Jani, R.A. Paleoclimate and paleovegetation in Central India during the Upper Cretaceous based on the stable isotope composition of the paleosol carbonates. *Palaeogeogr Palaeoclimatol Palaeoecol* **1995**, *114*, 285–296.
172. Tandon, S.K.; Sood, A.; Andrews, J.E.; Dennis, P.F. Palaeoenvironments of dinosaur-bearing Lameta Beds (Maastrichtian), Narmada Valley, Central India. *Palaeogeogr Palaeoclimatol Palaeoecol* **1995**, *117*, 153–184.
173. Mohabey, D.M. *Pycnodus lametae* (Pycnodontidae), a holostean fish from freshwater Upper Cretaceous Lameta Formation of Maharashtra. *J Geol Soc India* **1996**, *47*, 593–598.

174. Mohabey, D.M.; Udhoji, S.G. Fauna and flora from Late Cretaceous (Maastrichtian) non-marine Lameta sediments associated with Deccan volcanic episode, Maharashtra: its relevance to the K-T boundary problem, palaeoenvironment and palaeogeography. *Gondwana Geol Mag* **1996**, *2*, 349–364.
175. Jain, S.L.; Sahni, A. Some Upper Cretaceous vertebrates from central India and their palaeogeographical implications. In *Symposium on Cretaceous of India: palaeoecology, palaeogeography and time boundaries*, Lucknow, India, 24–26 November; Maheshwari, H.K., Ed.; Indian Association Palynostratigraphers: Lucknow, India, 1983; 66–83.
176. Sahni, A. Cretaceous-Paleocene terrestrial faunas of India: lack of endemism during drifting of the Indian plate. *Science* **1984**, *226*, 441–443.
177. Shome, S.; Chandel, R.S. Palaeontological studies of Papro Formation (Infratrappean) of Lalitpur District, Uttar Pradesh-Its age, correlation and palaeoecology. *Indian J Geosci* **2013**, *65*, 49–62.
178. Khosla, A. Dinosaur eggshells from the Late Cretaceous Lameta Formation along the east-central Narbada River region: Biomineralization and morphotaxonomical studies. Unpublished PhD Thesis, Panjab University, Chandigarh, 1996.
179. Khosla, A.; Sahni, A. Late cretaceous (Maastrichtian) ostracodes from the Lameta formation, Jabalpur cantonment area, Madhya Pradesh, India. *J Palaeontol Soc India* **2000**, *45*, 57–78.
180. Khosla, A.; Prasad, G.V.R.; Verma, O.; Jain, A.K.; Sahni, A. Discovery of a micromammal-yielding Deccan intertrappean site near Kisalpuri, Dindori District, Madhya Pradesh. *Curr Sci* **2004**, *87*, 380–383.
181. Prasad, G.V.R.; Khajuria, C.K. Palaeoenvironments of the Late Cretaceous Mammal-Bearing Intertrappean Beds of Naskal, Andhra Pradesh, India. *Mem Geol Soc India* **1996**, *37*, 337–362.
182. Khajuria, C.K.; Prasad, G.V.R. Taphonomy of a Late Cretaceous mammal-bearing microvertebrate assemblage from the Deccan inter-trappean beds of Naskal, peninsular India. *Palaeogeogr Palaeoclimatol Palaeoecol* **1998**, *137*, 153–172.
183. Prasad, G.V.R.; Khajuria, C.K. A record of microvertebrate fauna from the intertrappean beds of Naskal, Andhra Pradesh. *J Palaeontol Soc India* **1990**, *35*, 151–161.
184. Rana, R.S. Palaeontology and Palaeoecology of the Intertrappean (Cretaceous–Tertiary transition) beds of the Peninsular India. *J Palaeontol Soc India* **1990**, *35*, 105–120.
185. Prasad, G.V.R.; Sahni, A. Coastal-plain microvertebrate assemblage from the terminal Cretaceous of Asifabad, Peninsular India. *J Palaeontol Soc India* **1987**, *32*, 5–19.
186. Lourembam, R.S.; Prasad, G.V.; Grover, P. Ichthyofauna (Chondrichthyes, Osteichthyes) from the Upper Cretaceous intertrappean beds of Piplanarayanwar, Chhindwara District, Madhya Pradesh, India. *Isl Arc* **2017**, *26*, p.e12180.
187. Sahni, A. Upper Cretaceous palaeobiogeography of peninsular India and e Cretaceous-Paleocene transition: the vertebrate evidence. In *Symposium on Cretaceous of India: palaeoecology, palaeogeography and time boundaries*, Lucknow, India, 24–26 November; Maheshwari, H.K., Ed.; Indian Association Palynostratigraphers: Lucknow, India, 1983; 128–140.
188. Keller, G.; Adatte, T.; Bajpai, S.; Mohabey, D.M.; Widdowson, M.; Khosla, A.; Sharma, R.; Khosla, S.C.; Gertsch, B.; Fleitmann, D.; Sahni, A. K-T transition in Deccan Traps of central India marks major marine Seaway across India. *Earth Planet Sci Lett* **2009**, *282*, 10–23.
189. Rana, R.S.; Kumar, K.; Singh, H. Palaeocene vertebrate fauna from the Fatehgarh Formation of Barmer district, Rajasthan, western India. In *Micropalaeontology: Application in Stratigraphy and Paleocyanography*; Sinha, D.K., Ed.; Narosa Publishing House: New Delhi, India, 2006; pp. 113–130.
190. Dolson, J.; Burley, S.D.; Sunder, V.R.; Kothari, V.; Naidu, B.; Whiteley, N.P.; Farrimond, P.; Taylor, A.; Direen, N.; Ananthakrishnan, B. The discovery of the Barmer Basin, Rajasthan, India, and its petroleum geology. *AAPG Bull* **2015**, *99*, 433–465.
191. Khosla, A. Palaeoenvironmental, palaeoecological and palaeobiogeographical implications of mixed fresh water and brackish marine assemblages from the Cretaceous-Palaeogene Deccan intertrappean beds at Jhilmili, Chhindwara District, central India. *Rev Mex de Cienc Geol* **2015**, *32*, 344–357.
192. Schultz, O.; Paunović, M. Der Nachweis von *Coelodus* (Osteichthyes, Pycnodontidae) im Turonien (Oberkreide) von Gams bei Hieflau, Steiermark, Österreich, und aus der Oberkreide von Kroatien und Italien. *Ann Naturh Mus Wien, Ser A* **1997**, *98A*, 73–141.
193. Arambourg, C.; Joleaud, L. Vertébrés fossiles du Bassin du Niger. *Bull Dir Mines Af Occid* **1943**, *7*, 27–74.
194. Ósi, A.; Szabó, M.; Kollmann, H.; Wagreich, M.; Kalmar, R.; Makádi, L.; Szentesi, Z.; Summesberger, H. Vertebrate remains from the Turonian (Upper Cretaceous) Gosau Group of Gams, Austria. *Cretac Res* **2019**, *99*, 190–208.
195. Kollmann, H.A. *Megalonoda* n. gen.(Melanopsidae, Gastropoda) aus der Oberkreide der Nördlichen Kalkalpen (Österreich). *Ann Naturh Mus Wien, Ser A* **1984**, *86*, 55–62.
196. Kollmann, H.A.; Sachsenhofer, R.F. Zur Genese des Gagats von Gams bei Hieflau (Oberkreide, Steiermark). *Mitt Ref Geol Paläont Landesmuseum Joanneum* **1998**, *2*, 223–228.

197. Neubauer, T.A.; Harzhauser, M.; Mandic, O.; Georgopoulou, E.; Kroh, A. Paleobiogeography and historical biogeography of the non-marine Caenogastropod family Melanopsidae. *Palaeogeogr Palaeoclimatol Palaeoecol* **2016**, *44*, 124–143.
198. Kocsis, L.; Ósi, A.; Vennemann, T.; Trueman, C.N.; Palmer, M.R. Geochemical study of vertebrate fossils from the Upper Cretaceous (Santonian) Csehbánya Formation (Hungary): Evidence for a freshwater habitat of mosasaurs and pycnodont fish. *Palaeogeogr Palaeoclimatol Palaeoecol* **2009**, *280*, 532–542.
199. Botfalvai, G.; Ósi, A.; Mindszenty, A. Taphonomic and paleoecologic investigations of the Late Cretaceous (Santonian) Iharkút vertebrate assemblage (Bakony Mts, northwestern Hungary). *Palaeogeogr Palaeoclimatol Palaeoecol* **2015**, *417*, 79–405.
200. Szabó, M.; Gulyás, P.; Ósi, A. Late Cretaceous (Santonian) pycnodontid (Actinopterygii, Pycnodontidae) remains from the freshwater deposits of the Csehbánya Formation, (Iharkút, Bakony Mountains, Hungary). *Ann Paleontol* **2016**, *102*, 123–134.
201. Ósi, A.; Rabi, M.; Makádi, L.; Szentesi, Z.; Botfalvai, G.; Gulyás, P. The Late Cretaceous continental vertebrate fauna from Iharkút (western Hungary, Central Europe): a review. In *Bernissart Dinosaurs and Early Terrestrial Ecosystems*; Godefroit, P., Ed.; Indiana University Press: Bloomington, USA, 2012; pp. 532–569.
202. Jocha-Edelényi, E. History of evolution of the Upper Cretaceous Basin in the Bakony Mts at the time of the terrestrial Csehbánya Formation. *Acta Geol Hung* **1988**, *31*, 19–31.
203. Ósi, A.; Bodor, E.R.; Makádi, L.; Rabi, M. Vertebrate remains from the Upper Cretaceous (Santonian) Ajka Coal Formation, western Hungary. *Cretac Res* **2016**, *57*, 228–238.
204. Császár, G.H.; Góczán, F.A. Bakony felső-kréta kőszénkutatás és kőszén lágvizsgálat (Upper Cretaceous coal prospecting and peat bog studies in the Bakony Mts). *Magyar Állami Földtani Intézet Évi Jelentése* **1988**, *1986*, 155–178.
205. Siegl-Farkas, Á. Palynostratigraphy and evolution history of the Ajka Coal Formation, W. Hungary. (Hungarian with English summary). *Magyar Állami Öldtani Intézet Évi Jelentése* **1988**, *1986*, 179–209.
206. Bodrogi, I.; Fogarasi, A.; Yazikova, E.A.; Sztanó, O.; Báldi-Beke, M. Upper Cretaceous of the Bakony Mts. (Hungary): sedimentology, biostratigraphy, correlation. *Zentbl Geol Paläontologie* **1998**, *1*, 1179–1194.
207. Garcia, G.; Bardet, N.; Houssaye, A.; Pereda-Suberbiola, X.; Valentin, X. Mosasauroid (Squamata) discovery in the Late Cretaceous (Early Campanian) continental deposits of Villeveyrac–L'Olivet, southern France. *C R Palevol* **2015**, *14*, 495–505.
208. Torices, A.; Diaz Berenguer, E.; Narvaéz, I.; Ortega, F.; Perez, S.; Serrano, H. Preliminary analysis of the microvertebrate fossils of “Lo Hueco” (Upper Cretaceous, Cuenca, Spain). In *Mésogée-Bulletin du Muséum d'histoire Naturelle de Marseille 66, Proceedings of the 8ème Congrès de l'European Association of Vertebrate Palaeontology (EAVP)*, Aix-en-Provence, France, 7–12 June 2010; Muséum d'histoire naturelle: Marseille, France, 2010; 80.
209. Torices, A.; Barroso-Barcenilla, F.; Cambra-Moo, O.; Perez, S.; Serrano, H. Vertebrate microfossil analysis in the Palaeontological site of “Lo Hueco” (Upper Cretaceous, Cuenca, Spain). In *Abstracts 71<sup>st</sup> Meeting of the Society of Vertebrate Paleontology*, Las Vegas, USA, 2–5 November 2011; Supplement to the Journal of Vertebrate Paleontology: Las Vegas, USA, 2011; 205.
210. Ortega, F.; Bardet, N.; Barroso-Barcenilla, F.; Callapez, P.M.; Cambra-Moo, O.; Daviero-Gómez, V.; Díez Díaz, V.; Domingo, L.; Elvira, A.; Escaso, F.; García-Oliva, M.; Gómez, B.; Houssaye, A.; Knoll, F.; Marcos-Fernández, F.; Martín, M.; Mocho, P.; Narvaéz, I.; Pérez-García, A.; Peyrot, D.; Segura, M.; Serrano, H.; Torices, A.; Vidal, D.; Sanz, J.L. The biota of the upper Cretaceous site of Lo Hueco (Cuenca, Spain). *J Iber Geol* **2015**, *41*, 83–99.
211. Marmi, J.; Blanco, A.; Fondevilla, V.; Dalla Vecchia, F.M.; Sellés, A.G.; Vicente, A.; Martín-Closas, C.; Oms, O.; Galobart, A. The Molí del Baró-1 site, a diverse fossil assemblage from the uppermost Maastrichtian of the southern Pyrenees (north-eastern Iberia). *Cretac Res* **2016**, *57*, 519–539.
212. Blanco, A.; Szabó, M.; Blanco-Lapaz, À.; Marmi, J. Late Cretaceous (Maastrichtian) Chondrichthyes and Osteichthyes from northeastern Iberia. *Palaeogeogr Palaeoclimatol Palaeoecol* **2017**, *465*, 278–294.
213. Díez-Canseco, D.; Arz, J.A.; Benito, M.; Díaz-Molina, M.; Arenillas, I. Tidal influence in redbeds: a palaeoenvironmental and biostratigraphic reconstruction of the Lower Tremp Formation (South-Central Pyrenees, Spain) around the Cretaceous/Paleogene boundary. *Sediment Geol* **2014**, *312*, 31–49.
214. Cifelli, R.L.; Nydam, R.L.; Gardner, J.D.; Weil, A.; Eaton, J.G.; Kirkland, J.I.; Madsen, S.K.; Gillette, D.D. Medial Cretaceous vertebrates from the Cedar Mountain Formation, Emery County, Utah: the Mussentuchit local fauna. In *Vertebrate paleontology in Utah (MP 99-1)*; Gillette, D.D., Ed.; Utah Geological Survey: Salt Lake City, USA, 1999; pp. 219–242.
215. Fiorillo, A.R.; Gillette, D.D. Non-mammalian microvertebrate remains from the Robison Eggshell site, Cedar Mountain Formation (Lower Cretaceous), Emery County, Utah. In *Vertebrate paleontology in Utah (MP 99-1)*; Gillette, D.D., Ed.; Utah Geological Survey: Salt Lake City, USA, 1999; pp. 259–268.
216. Eaton, J.G.; Kirkland, J.I.; Kauffman, E.G. Evidence and dating of mid-Cretaceous tectonic activity in the San Rafael Swell, Emery County, Utah. *Mt Geol* **1990**, *27*, 39–45.

217. Brinkman, D.B.; Newbrey, M.G.; Neuman, A.G.; Eaton, J.G.; Titus, A.L. Freshwater Osteichthyes from the Cenomanian to late Campanian of Grand Staircase-Escalante National Monument, Utah. In *At the Top of the Grand Staircase: The Late Cretaceous of Southern Utah*; Titus, A.L., Loewen, M.A., Eds.; Indiana University Press: Bloomington, USA, 2013; pp. 195-236.
218. Kirkland, J.I.; Eaton, J.G.; Brinkman, D.B. Elasmobranchs from Upper Cretaceous freshwater facies in southern Utah. In *At the Top of the Grand Staircase: The Late Cretaceous of Southern Utah*; Titus, A.L., Loewen, M.A., Eds.; Indiana University Press: Bloomington, USA, 2013; pp. 153-194.
219. Titus, A.; Eaton, J.; Sertich, J. Late Cretaceous stratigraphy and vertebrate faunas of the Markagunt, Paunsaugunt, and Kaiparowits plateaus, southern Utah. *Geol Intermt West* **2016**, *3*, 229-291.
220. Pedrão, E.; Arai, M.; Barrilari, I.M.R.; Carvalho, I.S. *Análise palinológica de uma amostra de superfície de Querru (Formação Itapecuru), Município de Itapecuru-Mirim – MA*. Relatório Técnico Petrobrás, Rio de Janeiro, Brazil, 1993; 11 pp.
221. Candeiro, C.R.A.; Fanti, F.; Therrien, F.; Lamanna, M.C. Continental fossil vertebrates from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation, Brazil, and their relationship with contemporaneous faunas from North Africa. *J Afr Earth Sci* **2011**, *60*, 79-92.
222. Medeiros, M.A.; Lindoso, R.M.; Mendes, I.D.; De Souza Carvalho, I. The Cretaceous (Cenomanian) continental record of the laje do coringa flagstone (Alcântara formation), northeastern South America. *J South Am Earth Sci* **2014**, *53*, 50-58.
223. Sousa, E.P.; Medeiros, M.A.; Bertini, R.J.; Pereira, A.A.; Toledo, C.E.V. *Ocorrência inédita de pycnodontiformes para a Laje do Coringa, Ilha do Cajual (Formação Alcântara), Eo-cenomaniano do Estado do Maranhão*, Simpósio Brasileiro de Paleontologia de Vertebrados IV, UNESP, Rio Claro, Brazil, July 12-16; 2004; 67-68.
224. Benedetto, J.L.; Sanchez, T. El hallazgo de peces Pycnodontiformes (Holostei) en la Formacion Yacoraite (Cretacico superior) de la provincia de Salta (Argentina) y su importancia paleoecologica. *Acta Geol Lilloana* **1971**, *11*, 153-175.
225. Benedetto, J.L.; Sanchez, T. *Coelodus toncoensis* nov. sp. (Pisces, Holostei, Pycnodontiformes) de la Formacion Yacoraite (Cretacico superior) de la provincia de Salta. *Ameghiniana* **1972**, *9*, 59-71.
226. Cione, A.L. Algunas consideraciones sobre los Pycnodontiformes (Pisces, Holostei) procedentes de la Formacion Yacoraite, Cretacico tardio de la provincia de Salta. *Ameghiniana* **1977**, *14*, 315-316.
227. Gasparini, Z.; Buffetaut, E. *Dolichoampsia minima*, n. gen., n. sp., a representative of a new family of eusuchian crocodiles from the late Cretaceous of northern Argentina. *Neues Jahrb Geol Paläont Mh* **1980**, *5*, 257-271.
228. Cione, A.L.; Pereira, S.M.; Alonso, R.; Arias, J. Los peces bagres (Osteichthyes, Siluriformes) de la Formación Yacoraite (Cretacico tardio). Consideraciones biogeograficas y bioestratigraficas. *Ameghiniana* **1985**, *21*, 294-304.
229. Marquillas, R.; Sabino, I.; Sial, A.N.; Del Papa, C.; Ferreira, V.; Matthews, S. Carbon and oxygen isotopes of Maastrichtian-Danian shallow marine carbonates: Yacoraite Formation, northwestern Argentina. *J South Am Earth Sci* **2007**, *23*, 304-320.
230. Deschamps, R.; Rohais, S.; Hamon, Y.; Gasparini, M. Dynamic of a lacustrine sedimentary system during late rifting at the Cretaceous-Palaeocene transition: Example of the Yacoraite Formation, Salta Basin, Argentina. *Depos Rec* **2020**, *6*, 490-523.
231. Arratia, G.; Cione, A. The record of fossil fishes of southern South America. *Münchner Geowiss Abh* **1996**, *30*, 9-72.
232. Gayet, M. Holostean and teleostean fishes of Bolivia. *Rev Téc Yaci Petrol Fisc Bol* **1991**, *12*, 453-494.
233. Gayet, M.; Marshall, L.G.; Sempéré, T. The Mesozoic and Paleocene vertebrates of Bolivia and their stratigraphic context: a review. *Rev Téc Yaci Petrol Fisc Bol* **1991**, *12*, 393-433.
234. Gayet, M.; Sempere, T.; Cappetta, H.; Jaillard, E.; Lévy, A. La présence de fossiles marins dans le Crétacé terminal des Andes centrales et ses conséquences paléogéographiques. *Palaeogeogr Palaeoclimatol Palaeoecol* **1993**, *102*, 283-319.
235. Gayet, M.; Marshall, L.G.; Sempere, T.; Meunier, F.J.; Cappetta, H.; Rage, J.C. Middle Maastrichtian vertebrates (fishes, amphibians, dinosaurs and other reptiles, mammals) from Pajcha Pata (Bolivia). Biostratigraphic, palaeoecologic and palaeobiogeographic implications. *Palaeogeogr Palaeoclimatol Palaeoecol* **2001**, *169*, 39-68.
236. Schultze, H.P. Pycnodont fish (Actinopterygii, Osteichthyes) from the El Molino Formation (Late Cretaceous to Early Paleocene) of Bolivia. *Rev Téc Yaci Petrol Fisc Bol* **1991**, *12*, 449-452.
237. Muizon, C. de; Gayet, M.; Lavenu, A.; Marshall, L.G.; Sigé, B.; Villaroel, C. Late Cretaceous vertebrates, including mammals, from Tiupampa, southcentral Bolivia. *Geobios* **1983**, *16*, 747-753.
238. Benyoucef, M.; Läng, E.; Cavin, L.; Mebarki, K.; Adaci, M.; Bensalah, M. Overabundance of piscivorous dinosaurs (Theropoda: Spinosauridae) in the mid-Cretaceous of North Africa: The Algerian dilemma. *Cretac Res* **2015**, *55*, 44-55.
239. Longbottom, A.E. New Tertiary pycnodonts from the Tilemsi valley, Republic of Mali. *Bull br Mus Nat Hist* **1984**, *38*, 1-26.

240. O'Leary, M.A.; Bouaré, M.L.; Claeson, K.M.; Heilbronn, K.; Hill, R.V.; Mccartney, J.; Sessa, J. A.; Sissoko, F.; Tapanila, L.; Wheeler, E.; Roberts, E.M. Stratigraphy and paleobiology of the upper Cretaceous-lower Paleogene sediments from the trans-Saharan seaway in Mali. *Bull Am Mus Nat Hist* **2019**, *2019*, 1–183.
241. Tapanila, L.; Roberts, E.; Bouaré, M.L.; Sissoko, F.; O'Leary, M.A. Phosphate taxonomy of bone and comprolite conglomerates: a case study from the Eocene of Mali, NW Africa. *Palaios* **2008**, *23*, 139–152.
242. Ostrowski, S.A. The teleost ichthyofauna from the Late Cretaceous of Madagascar: Systematics, distribution, and implications for Gondwanan biogeography. Unpublished PhD thesis, Michigan State University, East Lansing, 2012.
243. Rogers, R.R.; Hartman, J.H.; Krause, D.W. Stratigraphic analysis of Upper Cretaceous rocks in the Mahajanga Basin, northwestern Madagascar: implications for ancient and modern faunas. *J Geol* **2000**, *108*, 275–301.
244. Rogers, R.R. Fine-grained debris flows and extraordinary vertebrate burials in the Late Cretaceous of Madagascar. *Geology* **2005**, *33*, 297–300.
245. Krause, D.W.; Sertich, J.J.W.; Rogers, R.R.; Kast, S.C.; Rasoamiaramanana, A.H.; Buckley, G.A. Overview of the discovery, distribution, and geological context of *Simosuchus clarki* (Crocodyliformes: Notosuchia) from the Late Cretaceous of Madagascar. *J Vertebr Paleontol* **2010**, *30*, 4–12.
246. Fendley, I.M.; Sprain, C.J.; Renne, P.R.; Arenillas, I.; Arz, J.A.; Gilabert, V.; Self, S.; Vanderkluisen, L.; Pande, K.; Smit, J.; Mittal, T. No Cretaceous-Paleogene boundary in exposed Rajahmundry Traps: A refined chronology of the longest Deccan lava flows from  $40\text{Ar}/39\text{Ar}$  dates, magnetostratigraphy, and biostratigraphy. *Geochem Geophys Geosyst* **2020**, *21*, e2020GC009149.
247. Bhalla, S.N. On the occurrence of *Eotrigonodon* in the Eocene of Rajahmundry, Andhra Pradesh. *J Geol Soc India* **1974**, *15*, 335–337.
248. Prasad, G.V.R. Fossil fish remains from the intertrappean beds of Duddukuru, Andhra Pradesh. *Bull Indian Geol Assoc* **1987**, *20*, 33–39.
249. Keller, G.; Adatte, T.; Gardin, S.; Bartolini, A.; Bajpai, S. Main Deccan volcanism phase ends near the K-T boundary: Evidence from the Krishna-Godavari Basin, SE India. *Earth Planet Sci Lett* **2008**, *268*, 293–311.
250. Ebersole, J.A.; Cicimurri, D.J.; Stringer, G.L. Taxonomy and biostratigraphy of the elasmobranchs and bonyfishes (Chondrichthyes and Osteichthyes) of the lower-to-middle Eocene (Ypresian to Bartonian) Claiborne Group in Alabama, USA, including an analysis of otoliths. *Euro J Taxon* **2019**, *585*, 1–274.
251. Holman J.A.; Case G.R. Reptiles from the Eocene Tallahatta Formation of Alabama. *J Vertebr Paleontol* **1988**, *8*, 328–333.
252. Marquillas, R.A.; Del Papa, C.E.; Sabino, I.F. Sedimentary aspects and paleoenvironmental evolution of a rift basin: Salta Group (Cretaceous-Paleogene), northwestern Argentina. *Int J Earth Sci* **2005**, *54*, 94–113.
253. Guinot, G.; Cavin, L. Body size evolution and habitat colonization across 100 million years (Late Jurassic–Paleocene) of the actinopterygian evolutionary history. *Fish Fish* **2018**, *19*, 577–597.
254. Föllmi, K.B. Early Cretaceous life, climate and anoxia. *Cretac Res* **2012**, *35*, 230–257.
255. Lovejoy, N.R.; Albert, J.S.; Crampton, W.G. Miocene marine incursions and marine/freshwater transitions: evidence from Neotropical fishes. *J South Am Earth Sci* **2006**, *21*, 5–13.
256. Kauffman, E.G. Paleobiogeography and evolutionary response dynamic in the Cretaceous Western Interior seaway of North America. In *Jurassic–Cretaceous Biochronology and Paleogeography of North America*; Westermann, G.E.G., Ed.; Geological Association of Canada: St. John's, Canada, 1984; pp. 275–306.
257. Wang, C.; Scott, R.W.; Wan, X.; Graham, S.A.; Huang, Y.; Wang, P.; Wu, H.; Dean, W.E.; Zhang, L. Late Cretaceous climate changes recorded in Eastern Asian lacustrine deposits and North American Epiherc sea strata. *Earth Sci Rev* **2013**, *126*, 275–299.
258. Grande, L.; Bemis, W.E. A comprehensive phylogenetic study of Amiid fishes (Amiidae) based on comparative skeletal anatomy. An empirical search for interconnected patterns of natural history. *J Vertebr Palaeontol* **1998**, *18*, 1–696.
259. Clement, A.M.; Long, J.A. Air-breathing adaptation in a marine Devonian lungfish. *Biol Lett* **2010**, *6*, 509–512.
260. Betancur-R, R.; Ortí, G.; Pyron, R.A. Fossil-based comparative analyses reveal ancient marine ancestry erased by extinction in ray-finned fishes. *Ecol Lett* **2015**, *18*, 441–450.
261. Guinot, G.; Cavin L. Contrasting 'fish' diversity dynamics between marine and freshwater environments. *Curr Biol* **2015**, *25*, 2314–2318.
262. Sheehan, P.M.; Fastovsky, D.E. Major extinctions of land-dwelling vertebrates at the Cretaceous-Tertiary boundary, Eastern Montana. *Geology* **1992**, *20*, 556–560.
263. Cavin, L.; Martin, M. Les actinopterygiens et la limite Cretace-Tertiaire. *Geobios* **1995**, *28*, 183–188.
264. Cavin, L. Effects of the Cretaceous-Tertiary event on bony fishes. In *Geological and Biological Effects of Impact Events*; Buffetaut, E., Koeberl, C., Eds.; Springer-Verlag: Berlin, Germany, 2002; pp. 141–158.
265. D'Hondt, S.; Donaghay, P.; Zachos, J. C.; Luttenberg, D.; Lindinger, M. Organic carbon fluxes and ecological recovery from the Cretaceous-Tertiary mass extinction. *Science* **1998**, *282*, 276–279.

266. Bartholomai, A. The large aspidorhynchid fish, *Richmondichthys sweeti* (Etheridge Jnr and Smith Woodward, 1891) from Albian marine deposits of Queensland, Australia. *Mem Queensl Mus* **2004**, *49*, 521–536.
267. Bloom, D.D.; Weir, J.T.; Piller, K.R.; Lovejoy, N.R. Do freshwater fishes diversify faster than marine fishes? A test using state-dependent diversification analyses and molecular phylogenetics of new world silversides (Atherinopsidae). *Evol* **2013**, *28*, 2689–2690.
268. Santini, F.; Nguyen, M.T.; Sorenson, L.; Waltzek, T.B.; Lynch Alfaro, J.W.; Eastman, J.M.; Alfaro, M.E. Do habitat shifts drive diversification in teleost fishes? An example from the pufferfishes (Tetraodontidae). *J Evol Biol* **2013**, *26*, 1003–1018.
269. Bloom, D.D.; Lovejoy, N.R. On the origins of marine-derived freshwater fishes in South America. *J Biogeogr* **2017**, *44*, 1927–1938.
270. Scotese, C.R. *Atlas of Jurassic Paleogeographic Maps, PALEOMAP Atlas for ArcGIS, Maps 32–42, Vol. 4, The Jurassic and Triassic,, Mollweide Projection*; PALEOMAP Project: Evanston, USA, 2014; 13 pp.
271. cotese, C.R. *Atlas of Early Cretaceous Paleogeographic Maps, PALEOMAP Atlas for ArcGIS, Maps 23–31, Vol. 2, The Cretaceous, Mollweide Projection*; PALEOMAP Project: Evanston, USA, 2014; 12 pp.
272. Scotese, C.R. *Atlas of Late Cretaceous Maps, PALEOMAP Atlas for ArcGIS, Maps 16–22 Vol. 2, The Cretaceous, Mollweide Projection*; PALEOMAP Project, Evanston, USA, 2014; 13 pp.
273. Scotese, C.R. *Atlas of Paleogene Paleogeographic Maps, PALEOMAP Atlas for ArcGIS, Maps 8-15, Vol. 1, The Cenozoic, Mollweide Projection*; PALEOMAP Project, Evanston, USA, 2014; 12 pp.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.