
The REPLUMLESS Transcription Factor Controls the Expression of the RECEPTOR-LIKE CYTOPLASMIC KINASE VI_A2 Gene Involved in Shoot and Fruit Patterning of *Arabidopsis thaliana*

[Erzsébet Kenesi](#) , [Orsolya Beöthy-Fehér](#) , [Réka Szöllősi](#) , [Ildikó Domonkos](#) , Ildikó Valkai , [Attila Fehér](#) *

Posted Date: 9 July 2024

doi: 10.20944/preprints202407.0524.v1

Keywords: phyllotaxis; stem thickness; BELLRINGER transcription factor; replum; vascular bundles; ROP GTPase-activated kinase



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

The REPLUMLESS Transcription Factor Controls the Expression of the *RECEPTOR-LIKE CYTOPLASMIC KINASE VI_A2* Gene Involved in Shoot and Fruit Patterning of *Arabidopsis thaliana*

Erzsébet Kenesi ¹, Orsolya Beöthy-Fehér ^{1,2}, Réka Szöllösi ³, Ildikó Domonkos ¹, Ildikó Valkai ¹ and Attila Fehér ^{1,3,*}

¹ Institute of Plant Biology, HUN-REN Biological Research Centre, H-6726, Szeged, Hungary

² Doctoral School in Biology, Faculty of Science and Informatics, University of Szeged, H-6726 Szeged, Hungary

³ Department of Plant Biology, Faculty of Science and Informatics, University of Szeged, H-6726 Szeged, Hungary

Abstract: The promoter of the *RECEPTOR-LIKE CYTOPLASMIC KINASE VI_A2* gene contains nine binding sites for the *REPLUMLESS* transcription factor. In agreement, the expression of the kinase gene was strongly downregulated in the *rpl-4* mutant. Comparing phenotypes of loss-of-function mutants, it was revealed that both genes are involved in stem growth, phyllotaxis, organization of the vascular tissues, and the replum, highlighting potential functional interactions. Expression of the *RLCKVI_A2* gene from the constitutive 35S promoter could not complement the *rpl-4* phenotypes, but exhibited a dominant positive effect on stem growth and affected vascular differentiation and organization. The results also indicated that the number of vascular bundles is independently regulated from stem thickness. Although our study cannot demonstrate the direct link between the *RPL* and *RLCKVI_A2* genes, it highlights the significance of the proper developmental regulation of the *RLCKVI_A2* promoter for balanced stem development.

Keywords: phyllotaxis; replum; stem thickness; *BELLRINGER* transcription factor; vascular bundles; ROP GTPase-activated kinase

Introduction

In plants, Rho-of-plants (ROP) GTPases play a central role in morphogenesis by regulating cell shape, polarity, hormonal responses, as well as defense mechanisms [1]. The signaling activity of Rho-type GTPases depends on their GTP-bound conformation that facilitates their binding to downstream effector proteins. Recent studies have demonstrated the link of ROPs via their positive regulator guanine nucleotide exchange factors to receptor kinase signaling, but our knowledge on the downstream effectors in these signaling pathways is still limited [2,3]. As ROP effectors, a specific class of receptor-like cytoplasmic kinases (RLCKs) have been identified in several plant species [2,4–8]. In *Arabidopsis thaliana*, these kinases belong to the group A of the RLCK VI class (*RLCKVI_A1-7*) [9] and have been implicated in the control of plant growth, morphogenesis, and defence [2,5–7,10].

Intriguingly, the *Arabidopsis RLCKVI_A2* gene contains multiple *REPLUMLESS* (RPL) homeobox transcription factor (TF) binding sites in its promoter. The RPL TF, also named as *VAAMANA*, *PENNYWISE*, *LARSON* or *BELLRINGER*, has diverse roles in plant development, including the control of shoot meristem structure and integrity [11–14], the spatial arrangement of lateral organs (phyllotaxis of leaves, flowers, and siliques) [15–17], vascular differentiation [13], and flower [18–21] as well as fruit [22,23] development.

Among the RPL-regulated processes, ROP signaling has been associated with those that depend on cell wall composition, namely the secondary cell wall development during vascular differentiation [13,24] and the pectin methylesterase activity-dependent phyllotactic arrangement of lateral organs developing from the shoot meristem [15,25]. These observations prompted us to investigate the

possible role of the ROP-activated RLCKVI_A2 kinase in these processes downstream of the RPL TF. It was first verified that the RPL TF is required for *RLCKVI_A2* expression. Next, the *RLCKVI_A2* cDNA was expressed under the control of the 35S promoter in wild type, *rpl4* and *rlckvi_a2* plants. Morphological, and histological comparisons revealed that the RPL TF and the RLCKVI_A2 kinase are involved in similar processes. Interestingly, the *rpl4* and *rlckvi_a2* mutants often exhibited contrasting phenotypes. Furthermore, the strong ectopic expression of the kinase gene had the most profound impact on the investigated processes independent of the genetic background, suggesting that tight control of *RLCKVI_A2* expression by its own promoter is required for the correct organisation/development of the investigated tissues/organs.

Materials and Methods

Promoter Cloning and Analysis

All plasmids/vectors (except pBSK) were obtained from the Arabidopsis Biological Resource Center; <http://www.Arabidopsis.org/>; last visited 18.01.2024).

The promoter of the Arabidopsis gene At2G18890 coding for the RLCKVI_A2 kinase was obtained from the F12F24 Arabidopsis genomic sequence cloned in the pBleoBAC vector. Based on the sequence information, we identified and isolated the 10410 bp PstI fragment of the BAC clone, which contained the entire At2G18890 gene. It was cloned into the pBSK vector (Agilent Technologies Stratagene Products Division, La Jolla, CA, USA) to facilitate further cloning. The HindIII-XbaI fragment carrying 2928 bp upstream and 512 bp downstream of the start codon was ligated to the EcoRI-XbaI fragment of the pMDC162 vector [26] encoding the *Escherichia coli* β -glucuronidase (GUS) enzyme. The resulting chimeric gene codes for the first 147 amino acids of the RLCK VI_A2 protein in fusion with the GUS reporter under the control of the At2G18890 gene promoter. The 147 amino acid-coding sequence also contains the first intron of the At2G18890 gene. The chimeric gene was inserted into the EcoRI-XbaI sites of the binary vector pCXGFP-P ([27]; GeneBank accession FJ905225) from which the BamHI fragment had been previously deleted. For the map of the final vector molecule see Figure S1A.

The promoter sequence of At2G18890 gene was searched for transcription factor (TF) binding sites at the AtcisDB database (<https://agris-knowledgebase.org/AtcisDB>; last accessed 18.01.2024).

The cDNA of the AtRLCKVI_A2 gene was cloned as an EcoRI-XhoI fragment from pET28a:RLCKVI_A2 [4] into the appropriate sites of the pK7F2WG binary Agrobacterium vector in fusion with GFP [28] (for the map see FigS1 B).

Arabidopsis Transformation

The vector molecules were introduced into *Agrobacterium tumefaciens* (GV3101; pMP90) cells [29] by electroporation [30]. Plant transformation took place in the following way. The 4-6-week-old flowering Arabidopsis plants were immersed in an Agrobacterium solution containing 5% sucrose and 0.01% Silwet L77 (Crompton Europe Ltd. Kenneth House. 4Langley Quay. Slough. Berkshire SL3 6EH. United Kingdom) and then stored covered overnight [31]. This step was repeated after 7-10 days with the new flowers appearing in the meantime, to increase the efficiency of the transformation. After that, the plants were allowed to flower, and the mature seeds were collected for the selection of transformants. Selection was performed on ½ AR (half-strength Arabidopsis Regeneration medium) plates containing 2.2 g/L Murashige Skoog powder (DUCHEFA BIOCHEMIE B.V, Haarlem, The Netherlands), 5 g/L sucrose, and 0.8% agar, supplemented with 20 μ g/mL hygromycin for the selection of lines harboring pCXGUS-P, or with 50 μ g/mL kanamycin for the selection of pK7F2WG-transformed plants. Plant lines used in further investigations were tested for GFP expression by Western blot analysis (see further).

Western Blot Analysis of GFP:RLCKVI_A2 Protein Abundance

Western blot analysis was carried out according to standard protocols using 1:2000 dilution of anti-GFP rat monoclonal antibody (3H9; ChromoTek GmbH, Planegg/Martinsried Germany) as primary, and 1:10 000 dilution of horse radish peroxidase conjugated anti-IgG(H,L) rat antibody

(BI2411, Abliance, Compiègne, FRANCE) as secondary antibody. Approximately equal amounts of total protein extracts were loaded on 10% SDS–polyacrylamide gels. After separation, proteins were transferred onto PVDF membranes (Immobilon-P, Merck KGaA, Darmstadt, Germany). The blocking solution was TBS (150 mM NaCl, 50 mM Tris-HCl, pH 7.5) with 0.2% Tween and 5% non-fat milk. Immunoreactive bands were visualized using the Immobilon Western chemiluminescent HRP substrate (WBKLS0500, Merck KGaA) and the iBright™ CL1500 Imaging System (Thermo Fischer Scientific, Waltham, MA USA). Before performing the Western blot, membranes were stained with Ponceau S solution (100mg Ponceau S in 5% acetic acid) as protein loading control.

GUS Staining

Histochemical analyses were performed with one-week-old seedlings and four to eight-week-old flowering plants (leaves, flowers, and siliques). GUS activity was detected histochemically by submerging the plant materials in GUS histochemical buffer (100 mM sodium phosphate, 0.5 mM $K_3Fe(CN)_6$, 0.5 mM $K_4Fe(CN)_6$, 10 mM Na_2-EDTA , 0.1% Triton X-100, and 1 mg/ml 5-bromo-4-chloro-3-indoyl glucuronide (X-gluc)) and incubating them at 37°C until colour developed (12h÷O/N). Samples were cleared with 70% v/v ethanol and were mounted on microscope slides and observed under stereomicroscope (Olympus SZX12 stereozoom microscope, Evident Corporation, Tokyo, Japan).

Phyllotactic Pattern Measurements

Plants were grown in greenhouse conditions at 16h/18h light/night cycles at 22°C until siliques were fully developed. Stems were then cut right above rosette leaves; photos of all stems were taken, and divergence angles were measured between the insertion points of two successive siliques according to Peaucelle A et al. [15]. The top 5 cm of the stem was not assessed, as elongation was incomplete. The phyllotactic orientation was set to the direction giving the smallest average divergence angle. For graphical representation divergence angles between two successive siliques on the stem were allocated into twelve 30° classes and the percentage of total measurements (n) falling into each class were displayed.

Scanning Electron Microscopy (SEM)

For SEM measurements shoot apical meristems were collected when plants reached the bolting phase with 1-2 cm long stems, while replums were collected when stems were fully developed. Plant samples were vacuum infiltrated then fixed in 2.5% glutaraldehyde containing phosphate buffer (pH 7.4) during 3 hours or overnight, dehydrated in aqueous solutions of increasing ethanol concentrations, critical point dried, covered with 15 nm gold by a Quorum Q150T ES sputter (Quorum Technologies, Laughton, East Sussex, UK) and observed in a JEOL JSM-7100F/LV scanning electron microscope (JEOL Europe BV, Nieuw-Vennep, The Netherlands) by using 3-5 kV accelerating voltage.

Preparation and Microscopy of Cross Sections of Arabidopsis siliques and Inflorescence Stems

Small (generally 0.5 cm long) pieces of samples derived from Arabidopsis siliques, or the axes of the inflorescences were fixed in 4% (w/v) paraformaldehyde according to the protocol of (Barroso et al., 2006). After the fixation, silique or inflorescence stem samples were washed in distilled water and embedded in 5% agarose based on [32]. Then 100 µm thick cross sections were prepared using a vibratome (Microm HM 650 V, Thermo Scientific, Waltham, MA USA). The sections were placed on a microscopic slide with a drop of water and were stained with toluidine blue [TB; 0.02% (w/v)] to stain the cell walls [33]. To visualize lignified cells in the tissue samples, sections were incubated in phloroglucinol-HCl (Wiesner's reagent), too [34]. The sections were observed by a light microscope and an inverted fluorescent microscope (Zeiss Axiovert 200 M, Carl Zeiss, Oberkochen Germany) equipped with a digital camera (AxiocamHR, HQ CCD, Carl Zeiss), and applying filter set 9 (exc.: 450–490 nm, em.: 515–∞ nm) and filter set 49 (exc.: 365 nm, em.: 445/50 nm).

Image PROCESSING, STATISTICS and REPRODUCIBILITY

Scanning electron microscopic images as well as fluorescent and light microscopic images were analysed with the aim of ImageJ software [35]. Statistical analyses were carried out using the Excell software (Microsoft Corporation, Redmond, WA, USA), and treated statistically by Student's t-test pairwise comparing treated and control samples. Statistical significance was deemed significant at a minimum threshold of $P < 0.05$. Non-quantitative experiments have been performed at least three times with independent biological samples and were included if produced similar results.

Results

The RPL HOMEBOX TF CONTROLS RLCKVI_A2 GENE EXPRESSION

The RLCKVI_A2 gene has a diversity of TF binding sites in its promoter. The gene is likely under the control of a range of transcription factors belonging to the bZIP, GATA, MYB or MYB-related, LFY, ABI3, WRKY, E2F/DP, and homeobox families (TableS 1). The homeobox family is represented by nine binding motifs for the RPL TF.

To verify the significance of RPL in the regulation of RLCKVI_A2 expression, a fragment of the gene containing 928 bp upstream and 512 bp downstream regions from its start codon was fused to the β -glucuronidase (GUS) reporter gene. In this way, the chimeric gene contained the first exon, the first intron and a short fragment of the second exon of the RLCKVI_A2 gene in addition to the promoter. The predicted translation product thus has the first 147 amino acids of the kinase in fusion with the GUS enzyme (Figure S1A). This chimeric gene was introduced into wild type and *rpl-4* mutant Arabidopsis plants. The *rpl-4* mutant has been previously shown to lack RPL transcripts [36]. In young wild-type seedlings, the promoter expressed the fusion protein in all organs with strong signals in the shoot, especially in its meristematic region and the vasculature, but no signal in the root meristem (Figure 1 a-h). In eight-week-old flowering plants, the expression was strongest in the veins of leaves, the petals and carpels of flowers, and the siliques (Figure 1 i-l). In the *rpl-4* genetic background, the expression of the construct was barely detectable, indicating that RPL is required for proper RLCKVI_A2 expression in all organs from the seedling stage to maturity.




Figure 1. Comparison of the expression of the At2G18890 (RLCKVI_A2) promoter driving the GUS marker gene in wild-type (A,C,E,G,I,K,M) and *rpl-4* mutant (B,D,F,H,J,L,N) genetic backgrounds.

The RLCKVI_A2 and the RPL Genes are Involved in Similar Processes

The T-DNA insertion mutant *rlckvi_a2* has previously been reported to control cell and hypocotyl elongation, as well as overall plant growth [10]. Here we report a more detailed phenotypic analysis of the same *rlckvi_a2* mutant, focusing on features compromised in the *rpl-4* plants. First, the phyllotactic pattern of siliques was determined measuring the divergence angles of two successive

siliques on the stem (Figure 2). The optimal divergence angle in spiral phyllotaxis is 137.5 degrees [37]. Despite the rarity of individual angles aligning exactly at 137.5 degrees even in wild-type plants, their average tends to approximate this optimal value. In agreement, we found that in wild-type Col-0 plants, the divergence angles showed a rather uniform distribution around the average of 136.98 degrees (Figure 2A). In contrast, the *rpl-4* mutant exhibited a high variation in divergence angles ranging from 10-320 degrees (average 150.98; Figure 2B). The phyllotaxis of the *rlckvi_a2* mutant was disturbed at a lower frequency, resulting in an average degree of 140.73 (Fig2.C). Nevertheless, the frequency of angles falling into the category close to the optimal value (120-149 degrees) decreased from 50% in the wild type to 35% in the mutant, indicating the likely contribution of the kinase to proper phyllotactic patterning.




Figure 2. The phyllotactic pattern of siliques of wild type (wt), *rlckvi_a2*, and *rpl-4* mutants without and with ectopic expression (OX1 and OX2) of the *RLCKVI_A2* gene under the control of the 35S promoter, respectively, is shown. The divergence angles of two successive siliques on the stem were determined for T4 generation plants. The distribution of measured angles falling into the indicated angle size categories is shown.

As its name indicates, the RPL homeodomain protein is required for replum development (the replum is an abaxial structure of the *Arabidopsis* fruit that remains attached to the plant even after the valves of the mature fruit have fallen) [22]. The identity of the replum is maintained by RPL repressing the valve margin cell fate. Therefore, in *rpl* mutants, the valve margin is extended to the region where the replum would normally develop [22]. In agreement, in mature *rpl-4* fruits, only a narrow cell file can be observed instead of the well-developed replum of the wild-type ones (Figure 3 A and B). The mutation in the *RLCKVI_A2* gene did not result in a considerably thinner replum compared to the control (Figure 3C). Microscopical cross sections of mature fruits verified the compromised development of the smaller replum in the *rpl-4* mutant (Figure 3 A-E). In contrast, the inner replum of the *rlckvi_a2* mutant was larger in size compared to the control (Figure 3 D, F, G, H). Fluorescent images revealed that the sizes of phloroglucinol-stained cells having lignified walls are bigger, but the overall number of these cells are fewer in the replum of the *rpl-4* and *rlckvi_a2* mutants compared to the wild type (FigS. 2). These lignified cells are present in the separation zone as well as in the inner replum (FigS. 2). The cells in the kinase mutant are notably larger in size, thereby leading to the overall enlargement of its inner replum despite a decrease in cell quantity.




Figure 3. The replum of wild type (wt), *rpl-4*, and *rlckvi_a2* fruits. Scanning electron microscopy of the fruit surface (A-C) and cross sectioning (D-F) verified a thinner visible and smaller inner replum for the *rpl-4* mutant compared to the other two lines. Overlaying cross sections for three replums per line shows that the inner replum of the *rlckvi_a2* mutant is larger than that of the wild type, although the visible replum of these two lines is about the same size (G).

Investigating the cross sections of the inflorescence stem verified the roles of both the RPL TF and the RLCKVI_A2 kinase in vascular development and organization. The *rpl-4* mutant has a thinner stem with fewer vascular bundles than the wild type (Figure 4A,B), in agreement with previous studies [13]. In contrast, the *rlckvi_a2* mutant has a stem cross section area that is more than double that of the wild type, with approximately 50% more vascular bundles (Figure 4A-C). Nevertheless, the vascular bundles occupy a smaller area of the stem of the *rlckvi_a2* mutant compared to that of the other two lines (Figure S3A), in agreement with the more extended pith of the *rlckvi_a2* stem (Figure 4A).

In the *rpl-4* mutant, the vascular bundles exhibit a flattened arc of procambial cells and a smaller xylem with fewer vessels (Figure 4D-G), as described earlier [13]. In contrast, the line of procambial cells is more dome-shaped, and the vascular bundles have some extra-large xylem vessels in the *rlckvi_a2* mutant compared to the other two lines (Figure 4F,G).




Figure 4. Stem thickness and vascular organization of wild type (wt), *rlckvi_a2*, and *rpl-4* mutant plants. Cross sections of the inflorescence stems of the three investigated lines (A) were analysed for stem cross section area (B), the number of vascular bundles per stem (C), xylem size (D; see dashed lines in G), the number of xylem vessels per xylems (E), and the size of xylem vessels (F). Stem parameters were determined for ten plants per line. Xylem parameters were measured for all xylems in four plants per line. The distribution of the measured values is represented as a box plot for each line. Significant differences from the wild-type were determined by Student's t-test ($p < 0.05 = *$; $p < 0.01 = **$). The organization of vascular bundles is exemplified in G. vb – vascular bundle; xv – xylem vessel; pc – arc of procambium.

In summary, the *rpl-4* and *rlckvi_a2* mutations both disturb the normal development of the inflorescence stem and the replum. However, the phenotypic changes the two mutations cause are contrary in many cases (e.g. replum size, stem thickness, number of large xylem vessels).

The Ectopic Expression of the RLCKVI_A2 Gene Interferes with RPL4-Regulated Processes but Cannot Complement the rpl-4 Mutation

To test the original hypothesis that the lack of *RLCKVI_A2* expression contributes to *rpl-4* phenotypes, we have expressed 35S promoter-driven *RLCKVI_A2* cDNA in the *rpl-4* mutant as well as in the wild-type and *rlckvi_a2* genetic backgrounds (FigS. 4). Plant lines overexpressing the *RLCKVI_A2* cDNA under the control of the 35S promoter were designated as OX (standing for “overexpressor”). The phyllotactic divergence angles were increased by overexpression of the kinase in the wild type, while hardly affecting those in the *rlckvi_a2* mutant background (Figure 2A,B,D-G). The 35S promoter-driven expression of *RLCKVI_A2* could not complement the *rpl-4* phenotype either (Figure 2 C,H,I). A possible reason is that the 35S promoter is unable to provide expression for the *RLCKVI_A2* gene in the same developmental context as its own promoter.

Peaucelle et al. [15] observed the formation of ectopic primordia on *blr6* meristems. We could observe this defect also at a few meristems of the *rpl-4* line. Interestingly, the ectopic expression of the *RLCKVI_A2* gene resulted in the more frequent formation of ectopic organ primordia in the shoot meristem of all investigated genetic background (37±11% in comparison to 10±9% in the controls; $n < 18$) (Figure S5). The ectopic activity of the kinase, therefore, might compromise processes involved in early phyllotactic patterning in the shoot meristem.

Overexpression of *RLCKVI_A2* did not significantly alter the overall size of the replum in either genetic background (Figure S6A). In the wild type, ectopic expression of the kinase gene increased the size but decreased the number of lignified cells, while it had the opposite effect in the *rlckvi_a2* background, partly compensating for the consequences of the mutation (Figure S6B). In the *rpl-4* mutant, both the size and number of lignified cells were decreased by overexpression of the kinase

gene (Figure S6B). Therefore, ectopic expression of *RLCKVI_A2* did not complemented, but rather enhanced the *rpl-4* replum phenotype.

The ectopic expression of the *RLCKVI_A2* gene has a profound impact on stem thickness and vascular organization (Figures 5 and 6). The stem cross-sectional area was significantly increased in all three genetic backgrounds (Figures 5A,D,G, and 6). Interestingly, the *rlckvi_a2* mutation and the ectopic *RLCKVI_A2* overexpression both increased stem thickness. The effects of the mutation and the ectopic overexpression seem to be independent since they were additive in the *rlckvi_a2* mutant expressing 35S:*RLCKVI_A2* (Figure 5G). This highlights the significance of proper transcriptional control of *RLCKVI_A2* expression during stem growth. The ectopic expression of the kinase augmented the number of vascular bundles as well as the size of the xylem in the wild-type such as in the *rlckvi_a2* and *rpl-4* backgrounds (Figure 5 B,E,H and C,F,I, respectively). It was also observed that the ectopic expression of the kinase resulted in the clustering/fusion of vascular bundles (Figure 6). This could sporadically be observed in control (non-transgenic) plants as well. In contrast, all investigated *RLCKVI_A2* overexpressing plants, regardless of the genetic background, had at least one instance of closely placed or even clustered vascular bundles. The strength of this phenotype seems to depend on stem thickness. A higher number of vascular bundles, caused by the ectopic expression of the kinase, can be placed at more or less regular intervals in the thick stems of *rlckvi_a2* plants. However, this is not possible in thinner wild-type stems, and especially not in the very thin *rpl-4* stems. This indicates that the number of vascular bundles and stem thickness are independently regulated by the *RLCKVI_A2* kinase.




Figure 5. Effect of ectopic *RLCKVI_A2* expression on stem thickness and the vasculature. Stem cross section area (A, D, G) the number of vascular bundles per stem (B, E, H), and the average size of xylems (C, F, I) were determined for wild-type (wt; A-C), *rpl-4* (D-F), or *rlckvi_a2* (G-I) plants without and with overexpression (OX1 and OX2) of the 35S:*RLCKVI_A2* gene. Ten plants per line were measured. The distribution of the measured values is represented as a box plot for each line. The data of the overexpressor lines (OX1 and OX2) were compared to their respective control using Student's t-test ($p < 0.05 = *$; $p < 0.01 = **$).




Figure 6. Stem cross sections of wild-type (wt) and mutant plants (*rpl-4* or *rlckvi_a2*) without and with overexpression (OX1 and OX2) of the 35S:RLCKVI_A2 gene. Red asterisks indicate closely placed/clustering vascular bundles.

The ectopic overexpression of RLCKVI_A2 led to an increase in the number of vessel elements in the xylem of all backgrounds (FigS. 7A), although its impact on vessel size remained ambiguous (Figure S7B).

Discussion

The broad impact of the REPLUMLESS (RPL) TF (also named VAAMANA, PENNYWISE, LARSON, or BELLRINGER) on shoot meristem maintenance, vegetative as well as reproductive shoot morphogenesis, and organ patterning has been clearly evidenced [11,17,18,23,38–45]. These studies highlighted its complex regulatory interactions with several other TFs. Among others, the direct binding of RPL to the promoter of AGAMOUS, thereby repressing its expression, has been reported [19]. We noticed that the promoter of the At2G18890 Arabidopsis gene, which encodes the ROP GTPase-activated receptor-like cytoplasmic kinase, RLCKVI_A2, involved in plant morphogenesis [10], contains nine copies of the same binding elements (Table S1). Since our understanding is limited concerning the downstream cytoplasmic effectors of RPL, we decided to investigate if RPL controls the expression of *RLCKVI_A2*. Introducing the At2G18890 promoter-controlled GUS marker into the *rpl-4* mutant background revealed that RPL is required for the proper expression of the *RLCKVI_A2* gene that was mainly confined to the shoot apex, vasculature, flower organs, fruit, and seeds (Figure 1). The absence of RPL almost fully abolished the expression of the chimeric construct (Figure 1). Whether this regulation is direct or indirect cannot be known at present. Bencivenga et al. identified potential target genes and binding sites of RPL in Arabidopsis inflorescence apices [46]. The At2G18890 gene was not included in the high confidence list of RPL-regulated genes. Our results indicate under-threshold, tissue-specific, or indirect interactions of RPL and *RLCKVI_A2*.

Nevertheless, we investigated the potential involvement of the RLCKVI_A2 kinase in RPL-controlled processes such as replum development [22], phyllotaxis [15], stem morphogenesis [46], and the differentiation of the vasculature [13]. Detailed analysis of the *rlckvi_a2* T-DNA insertion mutant, previously characterized as being affected in cell/hypocotyl elongation and overall plant growth [10], revealed moderate phenotypic alterations in all of the above processes (Figures 2-4). The RLCKVI_A2 kinase belongs to a small protein family with members potentially having overlapping expressions and functions [9] that may explain the relatively weak phenotypes of the mutant. It is of note that although the kinase seems to be involved in similar processes that of RPL, their mutations sometimes resulted in opposite phenotypes such as in the cross-section area of the stem and the replum (Figures 3 and 4). In other cases, they acted into the same direction such as increasing the

number of vascular bundles per stem and decreasing the number but increasing the size of lignified cells in the replum or of xylem vessels (Figure S. 2, Figure 4). Increasing the number of vascular bundles and indirectly controlling cell wall lignification have been previously described for RPL [18,47].

Although the RLCKVI_A2 kinase is unlikely to be one of the main targets of RPL, we investigated the possibility of expressing the RLCKVI_A2 gene using the constitutive 35S promoter to partially complement some phenotypes caused by the *rpl-4* mutation. This assumption was not validated by our observations. Some phenotypes of *rpl-4*, such as the phyllotactic aberrations, were not affected (Figure 2) while others, such as the higher number of vascular bundles, were even enhanced (Figure 5E) by ectopic RLCKVI_A2 expression. Considering that 35S:RLCKVI_A2 expression could not complement even the *rlckvi_a2* mutation, but resulted in very similar phenotypes in *wild type*, *rpl-4*, or *rlckvi_a2* plants, one can draw the conclusion that the 35S promoter cannot replace the RLCKVI_A2 promoter. The strong phenotypes caused by ectopic RLCKVI_A2 expression also highlights the importance of the proper control of RLCKVI_A2 expression by its own promoter during plant development. The ectopic expression of RLCKVI_A2 had a profound impact on stem thickness including the increase in the size of the pith and the number of vascular bundles (Figure 5). Interestingly the *rlckvi_a2* mutation had similar effects on these parameters (Figure 4) and their effects were additive (Figure 5). It suggests that the phosphorylation of some its targets by RLCKVI_A2 is a requirement for balanced stem growth and development, but it may have negative impact outside of the expression domain of the RLCKVI_A2 gene. There are no known substrates for Arabidopsis RLCKVI_A2 at present. However, the mutation of the RLCK VI_A2 has been reported to alter the expression of genes the products of which are associated with cell membranes, act at the cell periphery or in the apoplast, and function in cellular transport and/or cell wall reorganisation [10]. It is of note that both the RPL and RLCKVI_A2 proteins have been reported to control cell wall organisation and cell size although in an opposite way (Peaucelle et al., 2011; Etchells et al., 2012; Valkai et al., 2020). The size of parenchymatic and epidermal cells has been observed to be smaller in the case of RPL overexpression, while the RLCKVI_A2 kinase was shown to be required for proper cell elongation in the epidermis. Instances where the *rpl-4* and *rlckvi_a2* mutations have opposite phenotypes, such as in stem thickness, may indicate a negative feedback relationship between the transcription factor and the downstream kinase. The RPL transcription factor has been shown to control threedimensional patterns of cell division in the rib meristem directly repressing organ boundary genes [46]. Even though the RLCKVI_A2 gene was not identified in a genome-wide screen as a direct target of RPL, it also seems to control stem thickness. It is of note that the RLCKVI_A2 gene is also expressed in the shoot meristem stem cell niche [48].

Conclusions

The RPL TF is required for the tissue-specific expression of the ROP GTPase-activated RLCKVI_A2 kinase gene. Although previous studies could not identify RLCKVI_A2 as a direct target of RPL, the kinase and the TF seem to influence similar processes, including stem growth, phyllotaxis, and replum size. Our results highlight that the expression of the kinase outside of its expression domain alters shoot/fruit morphogenesis and vascular differentiation. This highlights the importance of proper developmental control of its promoter, for which the RPL TF might be partly responsible.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

Acknowledgments: The authors acknowledge the contribution of the Core Facility (Cellular Imaging Laboratory) of the HUN-REN Biological Research Centre. The equipment and expertise provided by the Microbial- and Plant Genomics Research Unit at the HUN-REN Biological Research Center and Szilárd Kovács for the preparation of cross sections for microscopic analyses are highly appreciated.

Funding: This work was supported by a grant from the National Research, Development, and Innovation Fund (grant no. K 132486).

References

- Berken, A. ROPs in the Spotlight of Plant Signal Transduction. *Cell. Mol. Life Sci. CMLS* **2006**, *63*, 2446–2459, doi:10.1007/s00018-006-6197-1.
- Fehér, A.; Lajkó, D.B. Signals Fly When Kinases Meet Rho-of-Plants (ROP) Small G-Proteins. *Plant Sci.* **2015**, *237*, 93–107.
- Miyawaki, K.N.; Yang, Z. Extracellular Signals and Receptor-like Kinases Regulating ROP GTPases in Plants. *Front. Plant Sci.* **2014**, *5*.
- Dorjgotov, D.; Jurca, E.; Fodor, C.; Szűcs, A.; Ötvös, K.; Klement, É.; Bíró, J.; Fehér, A. Plant Rho-Type (Rop) GTPase-Dependent Activation of Receptor-like Cytoplasmic Kinases in Vitro. *FEBS Lett.* **2009**, *583*, 1175–1182, doi:10.1016/j.febslet.2009.02.047.
- Huesmann, C.; Reiner, T.; Hoefle, C.; Preuss, J.; Jurca, M.E.; Domoki, M.; Fehér, A.; Hüchelhoven, R. Barley ROP Binding Kinase1 Is Involved in Microtubule Organization and in Basal Penetration Resistance to the Barley Powdery Mildew Fungus. *Plant Physiol.* **2012**, *159*, 311–320.
- Reiner, T.; Hoefle, C.; Huesmann, C.; Ménesi, D.; Fehér, A.; Hüchelhoven, R. The Arabidopsis ROP-Activated Receptor-like Cytoplasmic Kinase RLCK VI_A3 Is Involved in Control of Basal Resistance to Powdery Mildew and Trichome Branching. *Plant Cell Rep.* **2015**, *34*, 457–468.
- Enders, T.A.; Frick, E.M.; Strader, L.C. An Arabidopsis Kinase Cascade Influences Auxin-Responsive Cell Expansion. *Plant J. Cell Mol. Biol.* **2017**, *92*, 68–81, doi:10.1111/tpj.13635.
- Lajkó, D.B.; Valkai, I.; Domoki, M.; Ménesi, D.; Ferenc, G.; Ayaydin, F.; Fehér, A. In Silico Identification and Experimental Validation of Amino Acid Motifs Required for the Rho-of-Plants GTPase-Mediated Activation of Receptor-like Cytoplasmic Kinases. *PLANT CELL Rep.* **2018**, *37*, 627–639, doi:10.1007/s00299-018-2256-y.
- Jurca, M.E.; Bottka, S.; Fehér, A. Characterization of a Family of Arabidopsis Receptor-like Cytoplasmic Kinases (RLCK Class VI). *Plant Cell Rep.* **2008**, *27*, 739–748.
- Valkai, I.; Kénesi, E.; Domonkos, I.; Ayaydin, F.; Tarkowská, D.; Strnad, M.; Faragó, A.; Bodai, L.; Fehér, A. The Arabidopsis RLCK VI_A2 Kinase Controls Seedling and Plant Growth in Parallel with Gibberellin. *Int. J. Mol. Sci.* **2020**, *21*, 7266.
- Bhatt, A.M.; Etchells, J.P.; Canales, C.; Lagodienko, A.; Dickinson, H. VAAMANA—a BEL1-like Homeodomain Protein, Interacts with KNOX Proteins BP and STM and Regulates Inflorescence Stem Growth in Arabidopsis. *Gene* **2004**, *328*, 103–111, doi:10.1016/j.gene.2003.12.033.
- Cole, M.; Nolte, C.; Werr, W. Nuclear Import of the Transcription Factor SHOOT MERISTEMLESS Depends on Heterodimerization with BLH Proteins Expressed in Discrete Sub-Domains of the Shoot Apical Meristem of Arabidopsis Thaliana. *Nucleic Acids Res.* **2006**, *34*, 1281–1292, doi:10.1093/nar/gkl016.
- Etchells, J.P.; Moore, L.; Jiang, W.Z.; Prescott, H.; Capper, R.; Saunders, N.J.; Bhatt, A.M.; Dickinson, H.G. A Role for BELLRINGER in Cell Wall Development Is Supported by Loss-of-Function Phenotypes. *BMC Plant Biol.* **2012**, *12*, 212, doi:10.1186/1471-2229-12-212.
- Hepworth, S.R.; Pautot, V.A. Beyond the Divide: Boundaries for Patterning and Stem Cell Regulation in Plants. *Front. Plant Sci.* **2015**, *6*, 1052, doi:10.3389/fpls.2015.01052.
- Peaucelle, A.; Louvet, R.; Johansen, J.N.; Salsac, F.; Morin, H.; Fournet, F.; Belcram, K.; Gillet, F.; Höfte, H.; Laufs, P.; et al. The Transcription Factor BELLRINGER Modulates Phyllotaxis by Regulating the Expression of a Pectin Methyltransferase in Arabidopsis. *Development* **2011**, *138*, 4733–4741, doi:10.1242/dev.072496.
- Ung, N.; Lal, S.; Smith, H.M.S. The Role of PENNYWISE and POUND-FOOLISH in the Maintenance of the Shoot Apical Meristem in Arabidopsis. *Plant Physiol.* **2011**, *156*, 605–614, doi:10.1104/pp.110.171462.
- Byrne, M.E.; Groover, A.T.; Fontana, J.R.; Martienssen, R.A. Phyllotactic Pattern and Stem Cell Fate Are Determined by the Arabidopsis Homeobox Gene BELLRINGER. *Development* **2003**, *130*, 3941–3950, doi:10.1242/dev.00620.
- Smith, H.M.S.; Hake, S. The Interaction of Two Homeobox Genes, BREVIPEDICELLUS and PENNYWISE, Regulates Internode Patterning in the Arabidopsis Inflorescence. *Plant Cell* **2003**, *15*, 1717–1727, doi:10.1105/tpc.012856.
- Bao, X.; Franks, R.G.; Levin, J.Z.; Liu, Z. Repression of AGAMOUS by BELLRINGER in Floral and Inflorescence Meristems. *Plant Cell* **2004**, *16*, 1478–1489, doi:10.1105/tpc.021147.
- Kanrar, S.; Bhattacharya, M.; Arthur, B.; Courtier, J.; Smith, H.M.S. Regulatory Networks That Function to Specify Flower Meristems Require the Function of Homeobox Genes PENNYWISE and POUND-FOOLISH in Arabidopsis. *Plant J.* **2008**, *54*, 924–937, doi:10.1111/j.1365-313X.2008.03458.x.
- Andrés, F.; Romera-Branchat, M.; Martínez-Gallegos, R.; Patel, V.; Schneeberger, K.; Jang, S.; Altmüller, J.; Nürnberg, P.; Coupland, G. Floral Induction in Arabidopsis by FLOWERING LOCUS T Requires Direct Repression of BLADE-ON-PETIOLE Genes by the Homeodomain Protein PENNYWISE. *Plant Physiol.* **2015**, *169*, 2187–2199, doi:10.1104/pp.15.00960.
- Roeder, A.H.K.; Ferrándiz, C.; Yanofsky, M.F. The Role of the REPLUMLESS Homeodomain Protein in Patterning the Arabidopsis Fruit. *Curr. Biol.* **2003**, *13*, 1630–1635, doi:10.1016/j.cub.2003.08.027.

23. González-Reig, S.; Ripoll, J.J.; Vera, A.; Yanofsky, M.F.; Martínez-Laborda, A. Antagonistic Gene Activities Determine the Formation of Pattern Elements along the Mediolateral Axis of the Arabidopsis Fruit. *PLOS Genet.* **2012**, *8*, e1003020, doi:10.1371/journal.pgen.1003020.
24. Oda, Y.; Fukuda, H. Emerging Roles of Small GTPases in Secondary Cell Wall Development. *Front. Plant Sci.* **2014**, *5*, 428, doi:10.3389/fpls.2014.00428.
25. Li, H.; Shen, J.-J.; Zheng, Z.-L.; Lin, Y.; Yang, Z. The Rop GTPase Switch Controls Multiple Developmental Processes in Arabidopsis. *Plant Physiol.* **2001**, *126*, 670–684, doi:10.1104/pp.126.2.670.
26. Curtis, M.D.; Grossniklaus, U. A Gateway Cloning Vector Set for High-Throughput Functional Analysis of Genes in Planta. *Plant Physiol.* **2003**, *133*, 462–469, doi:10.1104/pp.103.027979.
27. Chen, S.; Songkumarn, P.; Liu, J.; Wang, G.-L. A Versatile Zero Background T-Vector System for Gene Cloning and Functional Genomics. *Plant Physiol.* **2009**, *150*, 1111–1121, doi:10.1104/pp.109.137125.
28. Karimi, M.; Inzé, D.; Depicker, A. GATEWAY™ Vectors for *Agrobacterium*-Mediated Plant Transformation. *Trends Plant Sci.* **2002**, *7*, 193–195, doi:10.1016/S1360-1385(02)02251-3.
29. Koncz, C.; Martini, N.; Szabados, L.; Hroudá, M.; Bachmair, A.; Schell, J. Specialized Vectors for Gene Tagging and Expression Studies. In *Plant Molecular Biology Manual*; Gelvin, S.B., Schilperoort, R.A., Eds.; Springer Netherlands: Dordrecht, 1994; pp. 53–74 ISBN 978-94-011-0511-8.
30. Wise, A.A.; Liu, Z.; Binns, A.N. Three Methods for the Introduction of Foreign DNA into *Agrobacterium*. In *Agrobacterium Protocols*; Wang, K., Ed.; Methods in Molecular Biology; Humana Press: Totowa, NJ, 2006; pp. 43–54 ISBN 978-1-59745-130-7.
31. Clough, S.J.; Bent, A.F. Floral Dip: A Simplified Method for *Agrobacterium* -Mediated Transformation of Arabidopsis Thaliana. *Plant J.* **1998**, *16*, 735–743, doi:10.1046/j.1365-313x.1998.00343.x.
32. Zelko, I.; Lux, A.; Sterckeman, T.; Martinka, M.; Kollárová, K.; Lišková, D. An Easy Method for Cutting and Fluorescent Staining of Thin Roots. *Ann. Bot.* **2012**, *110*, 475–478, doi:10.1093/aob/mcs046.
33. Pradhan Mitra, P.; Loqué, D. Histochemical Staining of Arabidopsis Thaliana Secondary Cell Wall Elements. *J. Vis. Exp. JoVE* **2014**, 51381, doi:10.3791/51381.
34. Ghanati, F.; Morita, A.; Yokota, H. Induction of Suberin and Increase of Lignin Content by Excess Boron in Tobacco Cells. *Soil Sci. Plant Nutr.* **2002**, *48*, 357–364, doi:10.1080/00380768.2002.10409212.
35. Schneider, C.A.; Rasband, W.S.; Eliceiri, K.W. NIH Image to ImageJ: 25 Years of Image Analysis. *Nat. Methods* **2012**, *9*, 671–675, doi:10.1038/nmeth.2089.
36. Xu, M.; Wang, X.; Liu, J.; Jia, A.; Xu, C.; Deng, X.W.; He, G. Natural Variation in the Transcription Factor REPLUMLESS Contributes to Both Disease Resistance and Plant Growth in Arabidopsis. *Plant Commun.* **2022**, *3*, 100351, doi:10.1016/j.xplc.2022.100351.
37. Kuhlemeier, C. Phyllotaxis. *Trends Plant Sci.* **2007**, *12*, 143–150, doi:10.1016/j.tplants.2007.03.004.
38. Kanrar, S.; Onguka, O.; Smith, H.M.S. Arabidopsis Inflorescence Architecture Requires the Activities of KNOX-BELL Homeodomain Heterodimers. *Planta* **2006**, *224*, 1163–1173, doi:10.1007/s00425-006-0298-9.
39. Alonso-Cantabrana, H.; Ripoll, J.J.; Ochando, I.; Vera, A.; Ferrándiz, C.; Martínez-Laborda, A. Common Regulatory Networks in Leaf and Fruit Patterning Revealed by Mutations in the Arabidopsis ASYMMETRIC LEAVES1 Gene. *Development* **2007**, *134*, 2663–2671, doi:10.1242/dev.02864.
40. Ragni, L.; Truernit, E.; Pautot, V. KNOXing on the BELL: TALE Homeobox Genes and Meristem Activity. *Int. J. Plant Dev. Biol.* **2007**, *1*, 42–48.
41. Ragni, L.; Belles-Boix, E.; Günl, M.; Pautot, V. Interaction of KNAT6 and KNAT2 with BREVIPEDICELLUS and PENNYWISE in Arabidopsis Inflorescences. *Plant Cell* **2008**, *20*, 888–900, doi:10.1105/tpc.108.058230.
42. Yu, L.; Patibanda, V.; Smith, H.M.S. A Novel Role of BELL1-like Homeobox Genes, PENNYWISE and POUND-FOOLISH, in Floral Patterning. *Planta* **2009**, *229*, 693–707, doi:10.1007/s00425-008-0867-1.
43. Arnaud, N.; Pautot, V. Ring the BELL and Tie the KNOX: Roles for TALEs in Gynoecium Development. *Front. Plant Sci.* **2014**, *5*, 93, doi:10.3389/fpls.2014.00093.
44. Sehra, B.; Franks, R.G. Redundant CARG Box Cis-Motif Activity Mediates SHATTERPROOF2 Transcriptional Regulation during Arabidopsis Thaliana Gynoecium Development. *Front. Plant Sci.* **2017**, *8*, 1712, doi:10.3389/fpls.2017.01712.
45. Simonini, S.; Stephenson, P.; Østergaard, L. A Molecular Framework Controlling Style Morphology in Brassicaceae. *Development* **2018**, *145*, doi:10.1242/dev.158105.
46. Bencivenga, S.; Serrano-Mislata, A.; Bush, M.; Fox, S.; Sablowski, R. Control of Oriented Tissue Growth through Repression of Organ Boundary Genes Promotes Stem Morphogenesis. *Dev. Cell* **2016**, *39*, 198–208, doi:10.1016/j.devcel.2016.08.013.
47. Khan, M.; Xu, M.; Murmu, J.; Tabb, P.; Liu, Y.; Storey, K.; McKim, S.M.; Douglas, C.J.; Hepworth, S.R. Antagonistic Interaction of BLADE-ON-PETIOLE1 and 2 with BREVIPEDICELLUS and PENNYWISE Regulates Arabidopsis Inflorescence Architecture. *Plant Physiol.* **2012**, *158*, 946–960, doi:10.1104/pp.111.188573.
48. Yadav, R.K.; Tavakkoli, M.; Xie, M.; Girke, T.; Reddy, G.V. A High-Resolution Gene Expression Map of the Arabidopsis Shoot Meristem Stem Cell Niche. *Dev. Camb. Engl.* **2014**, *141*, 2735–2744, doi:10.1242/dev.106104.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.