

Review

Not peer-reviewed version

Bioactive Polysaccharides from *Gracilaria lemaneiformis*: Preparation, Structures, and Therapeutic Insights

[Min Wang](#) , [Zhen Zhu](#) , Xiaocheng Wu , Kit-Leong Cheong , Xiaohua Li , Wanli Yu , [Yinlin Yao](#) , [Jiang Wu](#) , [Zhanhui Cao](#) *

Posted Date: 1 August 2024

doi: 10.20944/preprints202408.0022.v1

Keywords: *Gracilaria lemaneiformis*; polysaccharides; extraction; biological activities; gut microbiota



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

Bioactive Polysaccharides from *Gracilaria lemaneiformis*: Preparation, Structures, and Therapeutic Insights

Min Wang ¹, Zhen Zhu ¹, Xiaocheng Wu ¹, Kit-Leong Cheong ², Xiaohua Li ¹, Wanli Yu ¹, Yinlin Yao ¹, Jiang Wu ¹ and Zhanhui Cao ^{2,*}

¹ College of Coastal Agriculture Sciences, Guangdong Ocean University, Zhanjiang 524088, Guangdong, China; wangmin@gdou.edu.cn (M.W.); Zhuzhen156243@163.com (Z.Z.); 952082384@qq.com (X.W.); li1563549668@126.com (X.L.); 1003978102@qq.com (W.Y.); yaoyiinlin_2021@qq.com (Y.Y.); wuj@gdou.edu.cn (W.J.)

² College of Food Science and Technology, Guangdong Ocean University, Zhanjiang 524088, Guangdong, China; klcheong@gdou.edu.cn (K.C.)

* Correspondence: 57113124@qq.com

Abstract: *Gracilaria lemaneiformis*, a red seaweed, is an abundant source of bioactive polysaccharides with significant health-promoting properties. Nevertheless, the broad application of *G. lemaneiformis* in the nutraceutical and pharmaceutical sectors remains constrained due to the absence of comprehensive data. This review provides a detailed examination of the preparation methods, structural characteristics, and biological activities of *G. lemaneiformis* polysaccharides (GLP). We explore both conventional and advanced extraction techniques, highlighting the efficiency and yield improvements achieved through methods such as microwave-, ultrasonic-, and enzyme-assisted extraction. The structural elucidation of GLP using modern analytical techniques, including high-performance liquid chromatography, gas chromatography, and nuclear magnetic resonance spectroscopy, is discussed, providing comprehensive insights into their molecular composition and configuration. Furthermore, we critically evaluate the diverse biological activities of GLP, including their antioxidant, anti-inflammatory, antitumor, and gut microbiota modulation properties. This review underscores the therapeutic potential of GLP and suggests future research directions to fully harness their health benefits.

Keywords: *Gracilaria lemaneiformis*; polysaccharides; extraction; biological activities; gut microbiota

1. Introduction

The growing interest in marine resources for bioactive compounds has led to the exploration of various seaweeds [1,2], including *Gracilaria lemaneiformis*. This red algae species is widely distributed in coastal regions and has been traditionally valued in different cultures for its nutritional and medicinal benefits [3]. In China, the cultivation of *G. lemaneiformis* is primarily driven by its economic and environmental advantages. As a fast-growing seaweed, it is often used in integrated multi-trophic aquaculture systems to mitigate the environmental impact of fish and shrimp farming by absorbing excess nutrients, thereby improving water quality and the sustainability of aquaculture operations [4,5]. Besides its environmental benefits, *G. lemaneiformis* is a valuable resource in the food and nutraceutical industries due to its abundance of bioactive components, particularly polysaccharides [6].

G. lemaneiformis polysaccharides (GLP) have garnered significant interest due to their unique structural characteristics and biological functions. The processes of extracting and purifying GLP are crucial as they affect yield, purity, and biological efficacy. Traditional thermal extraction methods are commonly employed but are often inefficient and time-consuming [7]. To overcome these limitations, advanced techniques such as ultrasound-assisted extraction (UAE), microwave-assisted extraction

(MAE), and enzyme-assisted extraction (EAE) have been developed [8]. These methods aim to enhance extraction efficiency, preserve bioactivity, and increase the overall yield of polysaccharides. Structural characterization of GLP involves analyzing their monosaccharide composition, molecular weight, and structural configuration. Techniques like high-performance liquid chromatography (HPLC), gas chromatography (GC), and nuclear magnetic resonance (NMR) are widely used for this purpose [9]. These analytical methods provide detailed insights into the molecular structure of GLP, which is crucial for understanding their biological activities. GLP are primarily sulfated galactans, known for their diverse biological activities, including antioxidant, immunomodulatory, antitumor, and intestinal effects [10–12]. Understanding these structural features is essential for utilizing the therapeutic potential of GLP.

This review aims to deliver a thorough examination of the preparation, structural features, and biological activities of GLP. We will explore a range of extraction methods, including both traditional and innovative techniques. The structural characterization of GLP using contemporary analytical methods will be outlined, providing insights into their molecular composition and configuration. Additionally, we will critically assess the biological activities of GLP, focusing on their potential therapeutic applications in health and disease management. This review will underscore the notable advancements in GLP research and suggest future pathways for their development and application.

2. Extraction of GLP

The extraction of GLP is essential for their use in various industrial and health-related applications. Different extraction techniques, each with its own advantages and limitations, are summarized in Figure 1. The objective of extraction is to break down the cell wall and solubilize the polysaccharides in aqueous extract. The choice of extraction method, along with its unique mechanism and operating conditions, significantly affects the yield, purity, and bioactivity of the extracted polysaccharides. This section provides an overview of commonly used techniques, including traditional, EAE, MAE, and UAE methods. Understanding and optimizing extraction parameters can greatly enhance the process. The most influential factors are extraction temperature and time, which crucially affect the solubility of polysaccharides [13]. Higher temperatures generally improve extraction efficiency by enhancing solvent penetration, reducing viscosity, and increasing the dissolution of polysaccharides into the solvent [14]. However, excessively high temperatures can degrade thermolabile polysaccharides, leading to a loss of bioactivity and structural integrity [15]. Therefore, it is essential to balance these factors to maximize yield while preserving the quality of the polysaccharides.




Figure 1. The pros and cons of the conventional extraction method and advanced extraction techniques, including microwave-assisted, ultrasonic-assisted, and enzymatic-assisted extraction.

2.1. Traditional Extraction Method

The traditional extraction of polysaccharides from *G. lemaneiformis* typically involves a straightforward and widely used hot water extraction process. In this method, dried and ground algae are submerged in hot water at temperatures ranging from 90°C to 100°C for an extended period. For example, *G. lemaneiformis* powder is soaked and extracted using hot water at 90°C with a solid-to-liquid ratio of 1:45 for a duration of 4 h [16]. Acid solutions, such as citric acid (pH 2.0) at a solid-to-liquid ratio of 1:50 (*w/v*) at 100°C for 3 h, are also used in extraction [17]. Additionally, alkali solutions are employed; semi-dried pieces of *G. lemaneiformis* are soaked in alkali solution (0.3 mol/L NaOH) at 25°C for 2 h before being extracted with cold water three times, each extraction lasting for 2 h [12]. The heat in the extraction process facilitates the breakdown of cell walls and the extracting polysaccharides into the water. This method is advantageous due to its simplicity, cost-effectiveness, and minimal need for specialized equipment, making it particularly suitable for large-scale operations [18]. However, hot water extraction has notable limitations. It is time-consuming and may not efficiently extract all polysaccharides present in the biomass [19]. Additionally, the lack of specificity means other soluble components may be co-extracted, necessitating further purification steps. Despite these limitations, hot water extraction remains a foundational technique in the initial stages of polysaccharide research and production.

2.2. Microwave-Assisted Extraction Method

MAE is an advanced technique that utilizes microwave radiation to improve the extraction of polysaccharides from *G. lemaneiformis*. The principle behind MAE involves using microwaves to heat the solvent and biomass simultaneously. Microwaves cause rapid oscillation of polar molecules, generating heat through friction and dielectric heating [20]. This localized heating disrupts cell walls more efficiently than conventional methods, facilitating the release of polysaccharides into the solvent [21]. Process parameters, such as microwave power, exposure time, and solvent type, are carefully controlled to optimize extraction efficiency and yield. MAE provides several benefits compared to traditional thermal extraction methods. It considerably shortens extraction time and reduces solvent usage, making the process eco-friendlier and more cost-effective. For example, comparing MAE with hot water extraction under the same liquid-to-solid ratio of 1:20 and extraction temperature of 70°C, the MAE using a 500 W microwave yielded a higher polysaccharide content (9.618%) than hot water extraction (9.143%). Additionally, MAE required only 20 min, compared to 60 min for hot water extraction [22]. The rapid and uniform heating provided by microwaves minimizes the thermal degradation of sensitive polysaccharides, preserving their bioactivity and structural integrity, and ensuring better reproducibility [23]. However, the initial setup costs for microwave extraction can be high due to the specialized equipment required. Additionally, scaling up MAE from the laboratory to an industrial scale can be challenging due to differences in microwave penetration and heating uniformity in larger volumes [24,25].

2.3. Ultrasonic-Assisted Extraction Method

UAE is a cutting-edge technique that employs ultrasonic waves to improve the extraction of polysaccharides from *G. lemaneiformis*. This method involves applying high-frequency sound waves to a mixture of solvent and biomass, inducing cavitation bubbles within the liquid [26]. These bubbles collapse rapidly, releasing intense localized energy that disrupts the algae's cell walls [27]. This mechanical disruption facilitates the release of polysaccharides into the solvent. Parameters such as ultrasound frequency, power, and duration are carefully optimized to maximize extraction efficiency. Tang et al. utilized UAE to extract *G. lemaneiformis* polysaccharides using an ultrasonic device at 65°C for 30 min [28]. Furthermore, a combined technique employing ultrasound and microwave-assisted extraction was applied, with extraction conditions set at 87°C, ultrasonic power of 50 W, microwave power of 800 W, extraction duration of 31.7 min, and a solid-to-water ratio of 1.0:60.7, resulting in a polysaccharide yield of 34.8%, compared to 29.7% for hot water extraction [29]. UAE offers significant benefits over conventional methods. It shortens extraction time and solvent usage, enhancing cost-

effectiveness and environmental sustainability. The cavitation effect ensures efficient disruption of cell walls and improved solvent penetration, leading to higher yields. Moreover, the lower temperatures used in UAE help preserve the bioactive and structural stability of thermally sensitive polysaccharides. However, the initial investment in ultrasonic equipment may be prohibitive for smaller operations considering adopting this technology.

2.4. Enzymatic-Assisted Extraction Method

EAE leverages specific enzymes to dismantle the cell walls of *G. lemaneiformis*, facilitating the release of polysaccharides in a mild and efficient manner. This process involves cell wall-degrading enzymes, such as cellulases, amylase, hemicellulases, and pectinases, which hydrolyze the complex carbohydrates and proteins that make up the algal cell walls [30,31]. The enzymatic action disrupts the structural integrity of the cells, allowing polysaccharides to be solubilized and extracted into the surrounding solvent [32]. Key extraction parameters include the type and concentration of enzymes, temperature, and pH to maximize enzymatic activity. For example, enzymatic extraction using a thermostable α -amylase from *Thermococcus* sp. HJ21 at a high temperature of 95°C and pH 5 resulted in a yield of *G. lemaneiformis* polysaccharides reaching 49.15% on a dry weight [33]. Additionally, some reports indicate that extraction efficiency can be increased by adding enzymes to the initial hot water extract mixture, supplementing with 1% papain and 0.5% cellulase (*w/v*) and incubating at 60°C for 2 h to obtain more polysaccharides [16]. Chen et al. reported that H₂O₂-assisted cellulase extraction of agar from *G. lemaneiformis* preserves the sulfate content, achieving about 3.56%, which is higher than the 1.80% sulfate content obtained with traditional alkali-extracted agar [34]. The benefits of enzymatic extraction include its high specificity, targeting only cell wall components and minimizing the extraction of unwanted substances, thus improving the purity of the polysaccharides [35]. Additionally, enzymatic extraction operates under mild conditions, which helps preserve the bioactivity and structural integrity of the polysaccharides, making them suitable for sensitive applications such as pharmaceuticals and nutraceuticals. However, the cost of enzymes and the need for careful control of process parameters can be seen as limitations [36].

2.5. Other Extraction Techniques

Liquid-phase pulsed discharge is a non-thermal technique developed based on electrical breakdown in water. This method generates UV light emission, radical species, high-amplitude shock waves, and cavitation, all of which promote the release of intracellular polysaccharides [37,38]. Ju and Xi reported the liquid-phase pulsed discharge extraction of *G. lemaneiformis* polysaccharides, using an electric field strength of 80 kV/cm and a flow velocity of 14 mL/min. Although the extraction yields were not significantly different—140.19 mg/g for hot water extraction and 147.94 mg/g for liquid-phase pulsed discharge extraction—the latter method offered notable advantages. Specifically, it required a shorter extraction time (79 min versus 300 min) and operated at a lower temperature (25°C versus 60°C) [39].

3. Purification of GLP

Purifying GLP is vital for producing high-quality, consistent polysaccharide preparations for research and industrial use. Effective purification methods are essential to remove impurities that might interfere with the structural analysis and biological evaluation of GLP. Achieving high-purity polysaccharides is crucial for accurately determining structure-function relationships, which are key to understanding their biological activities and ensuring they meet the rigorous quality standards required for pharmaceuticals, nutraceuticals, and functional foods. A summary of the advantages and disadvantages of different purification techniques has been shown in Figure 2. Ethanol precipitation is a commonly employed technique for purifying GLP due to its simplicity, cost-effectiveness, and efficiency in removing contaminants. This process involves adding ethanol to an aqueous solution of crude polysaccharide extract, resulting in the precipitation of polysaccharides while smaller molecules and impurities remain dissolved in the ethanol-water mixture [40,41].

Researchers typically adjust the ethanol concentration to between 30% and 80% to isolate GLP fractions [12,42]. However, ethanol precipitation has limitations, including the potential co-precipitation of low molecular weight impurities, which reduces the overall purity of the final product and necessitates additional purification steps [43,44]. Additionally, the requirement for large volumes of ethanol can be both costly and hazardous.




Figure 2. Summary of the advantages and disadvantages of different purification techniques for GLP, including ethanol precipitation, column chromatography, and membrane technology.

Membrane separation techniques use semi-permeable membranes to separate molecules by size, making them effective for purifying polysaccharides [45,46], including those from *G. lamaneiformis*. This method offers several benefits, such as low energy consumption, scalability, and the ability to operate at ambient temperatures. Membranes with a molecular weight cut-off of 1 kDa are typically used to dialyze and remove low molecular weight impurities from GLP [47]. Ultrafiltration, a specific membrane purification method, utilizes membranes with various pore sizes to purify polysaccharides of different molecular weights [48]. By choosing membranes with particular pore sizes, researchers can effectively separate polysaccharides according to their size. For instance, membranes with larger pores (e.g., 100 kDa) can retain high molecular weight polysaccharides while allowing smaller molecules and impurities to pass through. Conversely, membranes with smaller pores (e.g., 10 kDa) can isolate low molecular weight polysaccharides by excluding even smaller contaminants [49,50]. Membrane separation techniques are continuous processes, making them suitable for large-scale industrial applications [51]. However, membrane fouling, where impurities or the polysaccharides themselves block the membrane pores, can be a significant drawback, reducing efficiency and increasing maintenance requirements [52].

Chromatographic techniques like ion-exchange chromatography and size-exclusion chromatography are highly selective and efficient for purifying GLP, resulting in highly purified fractions. Ion-exchange chromatography separates GLP based on their charge properties [53,54]. By gradually adjusting the ionic strength or pH of the elution buffer, polysaccharides with different charge densities can be selectively eluted [55]. Neutral and acidic GLP are purified using DEAE Sephadex A-50 resin, eluted with distilled water and 0.3-0.6 mol/L NaCl solution [29]. DEAE-52 and DEAE-Sepharose fast-flow column resins can also be used to obtain sulfated polysaccharides, eluted with 0.4 mol/L NaCl solution [56,57]. Size-exclusion chromatography or gel filtration chromatography, purifies GLP based on their molecular size [58,59]. Using Sephadex G-100 size exclusion chromatography, eluted with distilled water, results in high-purity homogeneous peaks for GLP [60,61]. Despite their effectiveness in achieving high-purity GLP, these chromatographic techniques present challenges such as high costs for purification materials and the time-consuming nature of the processes [62].

4. Structural Characterization of GLP

Structural characterization is vital in studying GLP as it provides insights into their functional properties, biological activities, and potential applications. This analysis includes determining the monosaccharide composition, glycosidic linkages, molecular weight, and chain conformation of the polysaccharides [63,64]. Chromatography methods like HPLC and GC, along with spectroscopy techniques such as Fourier transform infrared spectrometer and NMR, are essential for obtaining detailed structural characterization.

4.1. Monosaccharide Composition

Monosaccharide composition is a vital structural feature of GLP, offering key insights into the types and proportions of monosaccharides present. To determine this composition, polysaccharides are initially hydrolyzed into their constituent monosaccharides. These monosaccharides are then separated and quantified using chromatographic techniques, such as HPLC or GC [65,66]. For example, polysaccharides from *G. lamaneiformis* are hydrolyzed with trifluoroacetic acid at 100°C for 2 h. The resulting monosaccharide solution is subjected to PMP derivatization and analyzed by HPLC. This analysis revealed that GLP is predominantly composed of galactose (82.76%), rhamnose (9.01%), mannose (6.25%), and glucose (1.97%) [67]. GLP was hydrolyzed using 4 mol/L TFA at 105°C for 2 h. The resulting hydrolyzed product was derivatized with hydroxylamine hydrochloride and pyridine, followed by acetylation with acetic anhydride. Subsequent analysis by GC revealed the monosaccharide composition of GLP, showing molar ratios of 45.84% galactose, 35.17% glucose, 18.12% xylose, and 0.33% mannose [57].

The anhydrogalactose residues are acid-labile and convert to galactose residues or 5-hydroxymethyl-furfural under strong acid hydrolysis procedures [68]. Therefore, a reductive hydrolysis of GLP using a methylmorpholine–borane complex under acidic conditions is employed, followed by acetylation of the alditols and subsequently determined by GC [69]. The results showed that different red seaweed polysaccharides had varying monosaccharide compositions. *Porphyra haitanensis*, *Gracilaria blodgettii*, and *Gracilaria chouae* consisted of anhydrogalactose and galactose with molar ratios of 1:1.4, 1:1.5, and 1:1.6, respectively. In contrast, *G. lamaneiformis* and *Eucheuma galetinae* had relatively high amounts of galactose, with molar ratios of 1:3.0 and 1:3.1, respectively [70].

4.2. Molecular Weight

Determining the molecular weight of GLP is typically done using HPLC coupled with a size-exclusion column. In this method, polysaccharide molecules are separated based on their size as they move through a column packed with porous beads. The molecular weight is calculated using a series of dextran standards with known molecular weights. A study determined that a low molecular weight GLP had an average molecular weight of 2.7×10^3 Da using the HPLC dextran calibration method [71]. Additionally, HPLC has been used to investigate the impact of various extraction and degradation methods on the molecular weight of GLP. Liao et al. found that GLP with a molecular weight of 1.2189×10^5 Da could be chemically degraded to yield two polysaccharides with molecular weights of 57.02×10^3 Da and 14.29×10^3 Da [72]. Regarding extraction methods, the enzymatically extracted agar had a larger molecular weight of 1.936×10^6 Da, while the traditional alkali-extracted agar had a molecular weight of 7.91×10^5 Da, and H_2O_2 -assisted enzymatic extraction yielded agar with a molecular weight of 8.52×10^5 Da [34].

High-performance size-exclusion chromatography coupled with multi-angle laser light scattering (HPSEC-MALLS) is a sophisticated method for determining the molecular weight of GLP without the need for dextran calibration. This technique not only provides precise and accurate molecular weight measurements but also determines the radius of gyration and intrinsic viscosity, giving a detailed understanding of the polysaccharide's chain conformation. Using HPSEC-MALLS combined with a viscometer, Veeraperumal et al. found that the molecular weight and intrinsic viscosity of GLP were 1.570×10^5 Da and 133.94 mL/g, respectively. Furthermore, the exponent α of

the Mark-Houwink equation $[\eta] = kMw^\alpha$ for GLP in sodium nitrate solution was calculated to be 0.737, indicating that GLP has a flexible chain in investigated aqueous solution [56].

4.3. Glycosidic Linkages

The glycosidic linkages of GLP involve identifying the specific connections between monosaccharide units in the polysaccharide chain. Methylation analysis, followed by gas chromatography-mass spectrometry (GC-MS), is commonly used to determine these linkage positions. For instance, Shi et al. used this method to confirm that GLP consists of linear linkages of 1,3-linked galactopyranose and 1,6-linked galactopyranose [29]. Similarly, Bajwa et al. found that *Gracilariopsis* sp. polysaccharides contained a significant amount of 3-linked galactopyranose (40%) and 4-linked anhydro-galactopyranose (27%) through methylation analysis [73].

NMR spectroscopy is a powerful tool for investigating detailed structural information about polysaccharides, including the sequence of monosaccharides and the types of glycosidic linkages present. Studies have shown that the chemical structure of GLP is primarily composed of galactose. For example, Li et al. described GLP as having a linear backbone of β -(1 \rightarrow 3)- and α -(1 \rightarrow 4)-linked galactopyranose residues and anhydro-galactose units [42]. Similarly, Wang et al. described a GLP backbone consisting of alternating α -(1 \rightarrow 3)-galactopyranose and β -(1 \rightarrow 4)-galactopyranose units [71]. Another study investigated the chemical structure of GLP using various NMR techniques, including ^1H NMR, ^{13}C NMR, COSY, HSQC, and HMBC, revealing that the structure consists mainly of glucose and galactose, specifically \rightarrow 4)- α -D-Glcp-(1 \rightarrow 4)- α -D-Glcp-(1 \rightarrow 4)- β -D-Galp-(1 \rightarrow 6)- α -D-Glcp-(1 \rightarrow 6)- α -D-Glcp-(1 \rightarrow [74].

5. Biological Activities of GLP

5.1. Antioxidant Activities

GLP possess a multitude of bioactive properties, with their antioxidant activity being one of the most reported. GLP exhibit robust antioxidant activities through several well-documented mechanisms, primarily by scavenging free radicals—unstable molecules that cause significant cellular damage through oxidative stress [75,76]. The biological activities, related mechanisms of action, and structural information of GLP are summarized in Table 1. Reactive oxygen species (ROS), including superoxide anions, hydroxyl radicals, and hydrogen peroxide, as well as reactive nitrogen species such as nitric oxide and peroxynitrite, pivotal in the onset and progression of various diseases, including cancer, cardiovascular disorders, and neurodegenerative conditions [77,78]. GLP can donate hydrogen atoms or electrons to these reactive species, neutralizing them and mitigating their harmful effects. Wang et al. demonstrated the radical scavenging activities of GLP in vitro using assays such as the superoxide radical assay, hydroxyl radical assay, DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging assay, and reducing power assay. [79]. In another study, the scavenging ability of GLP showed an IC_{50} of 9.62 mg/mL for ABTS (2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid)) radicals, 23.85 mg/mL for DPPH, 4.97 mg/mL for ferric ion, and 3.56 mg/mL for superoxide radicals [80]. Additionally, in cell culture studies on human keratinocyte cells (HaCaT) exposed to H_2O_2 , a common inducer of oxidative stress, GLP significantly reduced ROS levels and increased the expression of Nrf-2/Keap-1, a key transcription factor regulating antioxidant responses [80]. This radical scavenging capability of GLP is crucial for maintaining cellular homeostasis and preventing oxidative damage to tissue and organ proteins, lipids, and nucleic acids.

In addition to directly neutralizing free radicals, GLP enhance the body's antioxidant defense systems by upregulating the activity of endogenous antioxidant enzymes, including superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx). SOD catalyzes the dismutation of superoxide radicals into oxygen and hydrogen peroxide, which CAT then breaks down into water and oxygen. GPx further reduces hydrogen peroxide to water, using glutathione as a substrate. By increasing the levels and activity of these enzymes, GLP bolster the cellular antioxidant defense mechanism, enhancing the overall capacity to detoxify harmful ROS. Fang et al. investigated the

antioxidant activity of GLP in in vitro cell culture assays and reported that GLP alleviated H₂O₂-induced oxidative injury in human fetal lung fibroblast 1 cells by protecting them from oxidative damage and increasing the activities of SOD, CAT, and GPx [81].

Aging and diabetes are strongly connected to oxidative stress, which occurs when there is a disparity between the production of ROS and the body's ability to counteract them with antioxidant defenses [82,83]. Antioxidant enzyme activity typically declines with age and chronic hyperglycemia, exacerbating oxidative damage [84]. In vivo studies indicate that GLP can mitigate oxidative stress related to both aging and diabetes. Zhang et al. found that GLP markedly restored the levels of antioxidant enzymes—SOD, CAT, and GSH-Px—in the serum and brain of D-galactose-induced aging mice, in a dose-dependent manner [85]. Similarly, in streptozotocin-induced diabetic mice, GLP treatment significantly increased the levels of SOD, GSH-Px, and CAT [86]. In addition, ultraviolet (UV) radiation in the environment can induce oxidative stress in living organisms. UV radiation, particularly UVA (320-400 nm) and UVB (280-320 nm) wavelengths, penetrates the skin and interacts with cellular components, generating ROS. In an in vitro assay, UVB-induced damage decreased intracellular levels of GSH and SOD in HaCaT cells, while GLP treatment increased these levels, demonstrating that GLP protected the cells from UVB-induced damage [87].

5.2. Immuno-Modulatory Activity

The immune system is essential for overall health and well-being, serving as the body's primary defense against pathogens and disease [88]. It consists of a complex network of cells, tissues, and organs that work together to detect and neutralize harmful invaders such as bacteria, viruses, fungi, and parasites [89]. GLP have demonstrated significant immunomodulatory activities by activating macrophages. These cells are vital to the innate immune system, as they detect, engulf, and destroy pathogens and apoptotic cells, thereby enhancing the body's immune responses [90]. Ren et al. reported that treatment with GLP has been shown to increase the phagocytic capability of RAW264.7 cells and increase the production of nitric oxide and ROS. Additionally, GLP stimulate the secretion of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF- α) and interleukin-6 (IL-6), further amplifying the immune response [57]. Proinflammatory cytokines such as tumor necrosis factor-alpha (TNF- α), interleukin-1 beta (IL-1 β), and interleukin-6 (IL-6) are produced by immune cells, including macrophages, dendritic cells, and T-lymphocytes, in response to pathogens or cellular damage [91,92]. These cytokines are crucial mediators of the immune response, promoting inflammation to defend against infections and injuries. However, the overproduction or prolonged presence of proinflammatory cytokines can lead to chronic inflammation, contributing to the progression of various inflammatory diseases such as rheumatoid arthritis, inflammatory bowel disease, and cardiovascular diseases [93,94]. Thus, maintaining a balanced cytokine response is critical for effective immune defense while preventing the detrimental effects of chronic inflammation. In a lipopolysaccharide-induced IEC-6 cell model, GLP co-culture with IEC-6 cells decreased the release and inhibited the gene expression of TNF- α , IL-6, and IL-1 β , demonstrating significant anti-inflammatory effects [42,95].

Another mechanism of GLP's immunomodulatory effect involves modulating lymphocyte proliferation and differentiation. Lymphocytes, including T cells and B cells, play pivotal roles in adaptive immunity [96]. In individuals with food allergies, the immune system mistakenly identifies harmless food proteins as threats [97], triggering the activation of T-helper 2 (Th2) cells. These cells release cytokines such as IL-4 and IL-13. Liu et al. investigated GLP's potential to alleviate food allergies using tropomyosin-sensitized mice. They found that GLP significantly reduced Th2-dependent tropomyosin-specific IgE and IgG1 serum levels and decreased the production of IL-4 and IL-13. The immunosuppressive mechanism may be related to a reduction in p38 MAPK activity, contributing to the mitigation of food allergy symptoms [11,98,99]. The oligosaccharide derived from GLP has been demonstrated to modulate type 1 immunity by suppressing T cell activation, as evidenced by studies conducted both in vivo and in vitro. In vivo, this oligosaccharide reduced the production of interferon-gamma (IFN- γ) in mice immunized with ovalbumin. In vitro, using OT-II

CD4⁺ T cells, the oligosaccharide inhibited mTOR activity, glycolysis, cell cycle progression, and DNA replication [100].

5.3. Anti-Tumor Activity

The significance of polysaccharides derived from natural resources in cancer research is notably accentuated by their low or non-toxicity, distinguishing them from many conventional chemotherapeutic agents [101,102]. GLP exhibit anti-tumor activity primarily through significant immune modulation. They enhance immune cell activity by stimulating the proliferation and activation of immune cells such as T lymphocytes, natural killer cells, macrophages, and [103]. Ji et al. investigated the antitumor effects of GLP on Kunming mice transplanted with S180 tumor cells in the armpit of the right hind limbs. They found that GLP suppressed the aggressive growth of solid S180 tumors by enhancing the proliferation of splenocytes, increasing the cytotoxic activity of natural killer cells, and elevating the serum cytokine levels of IL-2, IFN- γ , and TNF- α [104]. This cytokine modulation not only boosts overall immune surveillance and attack on tumor cells but also helps reduce the immunosuppressive environment often created by tumors [105].

Moreover, GLP exhibit antitumor activity through the direct inhibition of tumor growth. Kang et al. reported that GLP showed inhibitory effects on the growth of three cancer cell lines: human gastric cancer cell line MKN45, cervical carcinoma cell line HeLa, and non-small cell lung cancer cell line A549, and. Their transcriptome analysis demonstrated that GLP regulated apoptosis, the cell cycle, nuclear division, and cell death-related genes [106]. Shi et al. also reported similar findings, demonstrating that GLP inhibit the proliferation of various cancer cell lines *in vitro*. These include human breast cancer cells MCF-7, HeLa, and human hepatocellular carcinoma cells HepG2 [29]. Another study demonstrated that GLP inhibit the proliferation of various cancer cell lines, including gastric cancer cell line MKN28, human lung cancer cell line A549, and mouse melanoma cell line B16. This suppression is associated with increased expression of the Fas/FasL signaling pathway [61]. Fas (CD95) is a death receptor located on the cell surface, and its ligand, FasL, binds to it, triggering a cascade of events that lead to programmed cell death [107]. GLP enhance the expression of both Fas and FasL or increase their interaction. This activation of the Fas/FasL pathway promotes the formation of the death-inducing signaling complex, which further facilitates apoptosis. Cai et al. demonstrated that GLP effectively inhibits tumor growth in a mouse model bearing colon-26 carcinoma. This anti-tumor effect is primarily mediated through the induction of ferroptosis and the regulation of ferroptosis-related metabolic pathways [108]. Ferroptosis is a type of programmed cell death characterized by the accumulation of lipid peroxides and the depletion of intracellular glutathione, is crucial for the antitumor activity of investigated samples [109]. The study found that GLP promotes ferroptosis by inhibiting glutathione synthesis and disrupting the function of glutathione peroxidase 4, an enzyme essential for protecting cells from lipid peroxidation. Additionally, GLP increases the levels of 4-hydroxy-2-nonenal, a marker of lipid peroxidation, further driving ferroptosis. This mechanism highlights the potential of GLP as a therapeutic agent that leverages ferroptosis to enhance its efficacy in targeting cancer cells, offering a promising approach for cancer treatment [108].

5.4. Intestinal Health and Gut Microbiota

The gut microbiota is a diverse population of microorganisms residing in the gastrointestinal tract, and it is essential for maintaining intestinal health and overall well-being [110]. A healthy gut ensures efficient digestion and nutrient absorption, supports a robust immune system, and maintains the integrity of the gut barrier, preventing harmful pathogens and toxins from entering the bloodstream [111,112]. Disruptions in gut health can lead to various diseases. Conditions like inflammatory bowel disease (IBD), which encompasses ulcerative colitis and Crohn's disease, involve chronic inflammation of the gut lining, causing severe abdominal pain, diarrhea, and malnutrition [113,114]. Furthermore, gut dysbiosis has been linked to metabolic diseases such as obesity, type 2 diabetes, heart disorders, and non-alcoholic fatty liver disease [115–117]. Beyond intestinal and metabolic health, the gut microbiota also significantly influences mental health through the gut-brain

axis, a bidirectional communication network connecting the central nervous system and the gastrointestinal tract [118]. Therefore, maintaining a balanced gut microbiota is essential for promoting intestinal health and preventing disease [119].

Prebiotics are non-digestible food components that positively impact the host by promoting the growth or activity of specific beneficial microorganisms in the colon [120,121]. GLP can serve as prebiotics because they remain undigested in the upper gastrointestinal tract. [122]. Studies have shown that GLP resists digestion by simulated gastric conditions, including exposure to gastric juice containing pepsin, gastric lipase, and HCl at pH 3.0. Additionally, GLP is not digested by amylase or by components of intestinal juice such as pancreatin, trypsin, and bile salts [17]. These findings indicate that GLP remains intact in the upper gastrointestinal tract and may reach the large intestine, where it can be fermented and utilized by gut microbiota [123].

The intestinal flora consists of a diverse community of microorganisms, predominantly from four phyla: Firmicutes, Bacteroidetes, Actinobacteria, and Proteobacteria [124]. GLP can be fermented by Bacteroidetes. An in vitro simulated human fecal fermentation assay showed that GLP increased Bacteroidetes' relative abundance while decreasing that of Firmicutes [10]. Bacteroidetes utilize various carbohydrate-active enzymes (CAZymes) to break down complex polysaccharides. These include glycoside hydrolases, which cleave glycosidic bonds in polysaccharide chains [125]. CAZymes are categorized by their specific functions and substrate preferences, allowing Bacteroidetes to degrade a wide range of polysaccharides such as xylans, glucan, mannan, arabinans, and agarans [63,126]. It has been established that *Bacteroides plebeius* possesses an endo-type β -agarase, BpGH16A, which is part of the glycoside hydrolase (GH) family 16. This enzyme breaks β -1,4-glycosidic bonds in agarose, generating neoagarooligosaccharides. [127]. Co-culturing red seaweed polysaccharides with *Bacteroides thetaiotaomicron* revealed the presence of a GH16 family enzyme that catalyzes the hydrolysis of β -D-galactopyranose-(1 \rightarrow 4)- α -L-galactopyranose- β -6-sulfate linkages [128]. Additionally, the bacterium *Aquimarina* sp. AD1, which belongs to the Bacteroidetes phylum, contains GH96 family enzymes with specific activity targeting the α -1,3 linkage in LA α 1 \rightarrow 3G6S of funoran [129]. Bacteroidetes engage in a synergistic approach to break down complex polysaccharides. For instance, *Bifidobacteria* and *Lactobacillus* further process oligosaccharides that Bacteroidetes initially degrade. This interspecies cooperation involves exchanging intermediate metabolites, enhancing the breakdown of dietary fibers and improving the efficiency of carbohydrate fermentation. This is why GLP can increase the relative abundance of both *Bacteroidetes* spp. and *Bifidobacteria* spp., while reducing the relative abundance of *Escherichia* [10].

The gut microbiota breaks down GLP into smaller carbohydrates and monosaccharides, leading to the production of short-chain fatty acids (SCFAs) [130,131]. The main SCFAs are acetate, propionate, and butyrate, each contributing uniquely to gut health and overall physiological functions [132]. Acetate, the most prevalent SCFAs, is used by peripheral tissues for energy and helps regulate cholesterol levels [133]. Propionate, though less abundant, aids in glucose regulation and is linked to a lower risk of metabolic disorders such as obesity and type 2 diabetes [134]. Butyrate is vital for colon health, serving as the primary energy source for colonocytes, enhancing gut barrier integrity through mucus production and tight junction formation, and providing anti-inflammatory benefits [135,136]. SCFAs exert their physiological effects through specific receptors on the surface of various cell types in the gut and other tissues. The primary SCFAs receptors are G-protein coupled receptors (GPCRs), namely GPR41 (also known as free fatty acid receptor 3, FFAR3), GPR43 (FFAR2), and GPR109A [137,138]. Research by Han et al. demonstrated that GLP alleviated DSS-induced colitis in mice. The study found that increased expression of SCFAs receptors, such as GPR43, GPR109A, and the olfactory receptor Olfr78, along with elevated SCFAs levels in feces, indicated substantial SCFAs absorption in the colon. This upregulation significantly enhanced the integrity of intestinal tight junctions, including zonula occludens-1, claudin-1, and mucin, thereby strengthening the intestinal barrier. Furthermore, the beneficial effects of SCFAs on the intestinal barrier may be linked to the presence of bacteria such as *Enterorhabdus*, *Desulfovibrio*, *Alistipes*, and *Bacteroides acidifaciens* [139]. SCFAs are also vital in regulating the immune response, maintaining immune tolerance, and preventing excessive inflammation [140,141]. GLP can alleviate DSS-induced colitis by increasing

SCFAs levels. Additionally, GLP treatment resulted in decreased levels of CC chemokine ligand-25 (CCL-25) and CC-chemokine receptor-9 (CCR-9), while CD40 and TGF- β 1 levels were elevated [142]. CCR-9, the receptor for CCL-25, is primarily expressed on T cells that migrate to the gut. By inhibiting CCL-25 and CCR-9, GLP reduces the infiltration of inflammatory T cells into the gut, thereby mitigating intestinal inflammation [143,144]. CD40, a co-stimulatory protein on antigen-presenting cells, interacts with CD40 ligand on T cells to activate and proliferate both T cells and B cells, thus enhancing adaptive immunity [145].

Table 1. Biological activities of GLP, related mechanisms, and structural information.

Bioactivities	Mechanism	Structural information of GLP	Ref
Antioxidant	Radicals (hydroxyl, DPPH, and ABTS) scavenging capacity, regulation of antioxidant enzyme levels (MDA, SOD) in HK-2 cells, and decreased ROS levels.	Mw: 106, 49.6, 10.5, 6.14, 5.06, 3.71 and 2.42 kDa	[146]
Antioxidant	Radicals (ABTS, hydroxyl, and nitrite) scavenging capacity.	Glc: Gal: Xyl: Fuc= 4.48: 18.76: 1.811: 5.968; Mw: 591 kDa	[22]
Antioxidant	Decreased senescence-associated β -galactosidase activity and suppression of p21 and p53 gene expression.	Mw: 2.7 kDa alternating α -(1 \rightarrow 3)- and β -(1 \rightarrow 4)-Gal.	[71]
Antitumor	Inhibition of tumor cell proliferation in vitro through the apoptosis-related Fas/FasL signaling pathway.	Mw: 123.06, 14.29, 64.78, and 57.02 kDa Gal: Glc= 11.68: 1	[61]
Antitumor	Inhibition of tumor cell proliferation in vitro, enhancement of NK cell activity, and increased levels of serum cytokines <i>in vivo</i> .	Mw: 20958 Da Ara: Xyl: Man: Glc: Gal= 0.22: 0.23: 0.11: 0.17: 1.00	[104]
Hypoglycemic	Regulation of blood sugar levels. Increases in SOD, GSH-Px, and total antioxidant capacity.	Mw: 121.89, 57.02, and 14.29 kDa	[72]
Hypoglycemic	Inhibition of α -glucosidase activity.	Mw: 1478 kDa 34.35% Glc and 57.37% Gal	[147]
Intestinal health	Modulation of gut microbiota and an increase in short-chain fatty acids (SCFAs).	alternating 4-linked 3,6-anhydro- α -L-Gal and 3-linked β -D-Gal units with sulfate residues	[142]
Intestinal health	Modulate gut microbiota, increase short-chain fatty acids (SCFAs), and enhance the expression of tight junction proteins and MUC-2.	Mw: 22.38 kDa Ara: Fuc: Xyl: Man: Glc: Gal= 1.14: 1.45: 7.92: 4.23: 13.48: 71.78	[17,139]
Anti-influenza virus	Inhibit viral replication and decrease viral adsorption ability.	Mw: 4.9, 52, and 67 kDa AnGal and Gal	[148]
Wound healing	Promotes cell proliferation and migration through activation of the PI3K/aPKC signaling pathway. Enhances epithelial layer thickness and collagen deposition <i>in vivo</i> .	Mw: 157 kDa alternating 4-linked 3,6-anhydro- α -L-Gal and 3-linked β -D-Gal units with sulfate residues	[56]

* Mw: Molecular weight; Glc: glucose; Gal: galactose; Ara: arabinose; Fuc: fucose; Xyl: xylose; Man: mannose; AnGal: Anhydrogalactose.

6. Conclusion

GLP is a promising focus of research due to their diverse and potent biological activities. This review underscores the significant progress made in extracting and characterizing GLP. Traditional thermal extraction methods have been pivotal for isolating these polysaccharides, but advanced techniques such as UAE, MAE, and EAE have greatly enhanced yield and extraction efficiency. The structural analysis of GLP, using techniques like HPLC, GC, and NMR spectroscopy, has provided detailed insights into their monosaccharide composition, molecular weight, and structural configurations.

The biological activities of GLP are well-documented, with strong evidence supporting their antioxidant, immunomodulatory, antitumor, and intestinal health properties. These bioactivities underscore the potential applications of GLP in health and disease management. Their antioxidant properties can help prevent oxidative stress-related diseases, while their anti-inflammatory effects can benefit conditions such as inflammatory bowel disease and ulcerative colitis. The antitumor activities of GLP, including their ability to induce apoptosis, inhibit tumor cell proliferation, and modulate immune responses, highlight their potential in cancer therapy. Additionally, the prebiotic effects of GLP on gut microbiota suggest their potential in improving gut health and managing metabolic disorders.

Future research should focus on several key areas to fully harness the therapeutic potential of GLP. Clinical trials are essential to validate the health benefits observed in preclinical studies and establish effective dosages for human consumption. Mechanistic studies are needed to elucidate the pathways through which GLP exert their biological effects, aiding in the development of targeted therapies. Moreover, exploring the synergistic effects of GLP with other bioactive compounds could lead to the development of more effective functional foods and nutraceuticals. Investigating the sustainability and scalability of GLP production will also be crucial for their widespread application. Overall, continued research and development in this field hold great promise for the integration of GLP into health and wellness products, contributing to the prevention and treatment of various diseases.

Author Contributions: Conceptualization, M.W., Z.C; resources, Z.Z., X.W., K.C., X.L., W.Y., Y.Y., J.W.; writing—original draft preparation, M.W.; writing—review and editing, Z.Z., X.W., K.C., X.L., W.Y., Y.Y., J.W.; supervision, Z.C.; funding acquisition, Z.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by the National College Student Innovation and Entrepreneurship Training Program (CXXL2024055).

Data Availability Statement: No new data were created or analyzed in this study

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Hamed, I.; Özogul, F.; Özogul, Y.; Regenstein, J.M., Marine Bioactive Compounds and Their Health Benefits: A Review. *Comprehensive Reviews in Food Science and Food Safety* **2015**, *14*, 446-465.
2. Yao, W.; Zhang, Y.; Zhang, G., Marine peptides as potential anti-aging agents: Preparation, characterization, mechanisms of action, and future perspectives. *Food Chemistry* **2024**, *460*, 140413.
3. Wang, X.; He, L.; Ma, Y.; Huan, L.; Wang, Y.; Xia, B.; Wang, G., Economically important red algae resources along the Chinese coast: History, status, and prospects for their utilization. *Algal Research* **2020**, *46*, 101817.
4. Wei, Z.; You, J.; Wu, H.; Yang, F.; Long, L.; Liu, Q.; Huo, Y.; He, P., Bioremediation using *Gracilaria lemaneiformis* to manage the nitrogen and phosphorous balance in an integrated multi-trophic aquaculture system in Yantian Bay, China. *Marine Pollution Bulletin* **2017**, *121*, 313-319.
5. Ashkenazi, D.Y.; Israel, A.; Abelson, A., A novel two-stage seaweed integrated multi-trophic aquaculture. *Reviews in Aquaculture* **2019**, *11*, 246-262.

6. Torres, P.; Santos, J.P.; Chow, F.; dos Santos, D.Y.A.C., A comprehensive review of traditional uses, bioactivity potential, and chemical diversity of the genus *Gracilaria* (Gracilariales, Rhodophyta). *Algal Research* **2019**, *37*, 288-306.
7. Pang, Y.; Peng, Z.; Ding, K., An in-depth review: Unraveling the extraction, structure, bio-functionalities, target molecules, and applications of pectic polysaccharides. *Carbohydrate Polymers* **2024**, *343*, 122457.
8. Jönsson, M.; Allahgholi, L.; Sardari, R.R.R.; Hreggviðsson, G.O.; Nordberg Karlsson, E., Extraction and Modification of Macroalgal Polysaccharides for Current and Next-Generation Applications. *Molecules* **2020**, *25*, 930.
9. Ji, X.; Hou, C.; Shi, M.; Yan, Y.; Liu, Y., An Insight into the Research Concerning *Panax ginseng* C. A. Meyer Polysaccharides: A Review. *Food Reviews International* **2022**, *38*, 1149-1165.
10. Zhang, X.; Aweya, J.J.; Huang, Z.-X.; Kang, Z.-Y.; Bai, Z.-H.; Li, K.-H.; He, X.-T.; Liu, Y.; Chen, X.-Q.; Cheong, K.-L., In vitro fermentation of *Gracilaria lemaneiformis* sulfated polysaccharides and its agaro-oligosaccharides by human fecal inocula and its impact on microbiota. *Carbohydrate Polymers* **2020**, *234*, 115894.
11. Liu, Q.-M.; Yang, Y.; Maleki, S.J.; Alcocer, M.; Xu, S.-S.; Shi, C.-L.; Cao, M.-J.; Liu, G.-M., Anti-Food Allergic Activity of Sulfated Polysaccharide from *Gracilaria lemaneiformis* is Dependent on Immunosuppression and Inhibition of p38 MAPK. *Journal of Agricultural and Food Chemistry* **2016**, *64*, 4536-4544.
12. Fan, Y.; Wang, W.; Song, W.; Chen, H.; Teng, A.; Liu, A., Partial characterization and anti-tumor activity of an acidic polysaccharide from *Gracilaria lemaneiformis*. *Carbohydrate Polymers* **2012**, *88*, 1313-1318.
13. Sun, Y.; He, H.; Wang, Q.; Yang, X.; Jiang, S.; Wang, D., A Review of Development and Utilization for Edible Fungal Polysaccharides: Extraction, Chemical Characteristics, and Bioactivities. *Polymers* **2022**, *14*, 4454.
14. Leong, Y.K.; Yang, F.-C.; Chang, J.-S., Extraction of polysaccharides from edible mushrooms: Emerging technologies and recent advances. *Carbohydrate Polymers* **2021**, *251*, 117006.
15. Steiner, J.; Franke, K.; Kießling, M.; Fischer, S.; Töpfl, S.; Heinz, V.; Becker, T., Influence of hydrothermal treatment on the structural modification of spent grain specific carbohydrates and the formation of degradation products using model compounds. *Carbohydrate Polymers* **2018**, *184*, 315-322.
16. Long, X.; Hu, X.; Xiang, H.; Chen, S.; Li, L.; Qi, B.; Li, C.; Liu, S.; Yang, X., Structural characterization and hypolipidemic activity of *Gracilaria lemaneiformis* polysaccharide and its degradation products. *Food Chemistry: X* **2022**, *14*, 100314.
17. Han, R.; Pang, D.; Wen, L.; You, L.; Huang, R.; Kulikouskaya, V., In vitro digestibility and prebiotic activities of a sulfated polysaccharide from *Gracilaria Lemaneiformis*. *Journal of Functional Foods* **2020**, *64*, 103652.
18. He, L.; Yan, X.; Liang, J.; Li, S.; He, H.; Xiong, Q.; Lai, X.; Hou, S.; Huang, S., Comparison of different extraction methods for polysaccharides from *Dendrobium officinale* stem. *Carbohydrate Polymers* **2018**, *198*, 101-108.
19. Tang, W.; Liu, D.; Yin, J.-Y.; Nie, S.-P., Consecutive and progressive purification of food-derived natural polysaccharide: Based on material, extraction process and crude polysaccharide. *Trends in Food Science & Technology* **2020**, *99*, 76-87.
20. Hu, Q.; He, Y.; Wang, F.; Wu, J.; Ci, Z.; Chen, L.; Xu, R.; Yang, M.; Lin, J.; Han, L.; Zhang, D., Microwave technology: a novel approach to the transformation of natural metabolites. *Chinese Medicine* **2021**, *16*, 87.
21. Gharibzadeh, S.M.T.; Marti-Quijal, F.J.; Barba, F.J.; Altintas, Z., Current emerging trends in antitumor activities of polysaccharides extracted by microwave- and ultrasound-assisted methods. *International Journal of Biological Macromolecules* **2022**, *202*, 494-507.
22. He, J.; Xu, Y.; Chen, H.; Sun, P., Extraction, Structural Characterization, and Potential Antioxidant Activity of the Polysaccharides from Four Seaweeds. *International Journal of Molecular Sciences* **2016**, *17*, 1988.
23. Hashemifesharaki, R.; Xanthakis, E.; Altintas, Z.; Guo, Y.; Gharibzadeh, S.M.T., Microwave-assisted extraction of polysaccharides from the marshmallow roots: Optimization, purification, structure, and bioactivity. *Carbohydrate Polymers* **2020**, *240*, 116301.
24. Guzik, P.; Kulawik, P.; Zajac, M.; Migdał, W., Microwave applications in the food industry: an overview of recent developments. *Critical Reviews in Food Science and Nutrition* **2022**, *62*, 7989-8008.
25. Arrutia, F.; Adam, M.; Calvo-Carrascal, M.Á.; Mao, Y.; Binner, E., Development of a continuous-flow system for microwave-assisted extraction of pectin-derived oligosaccharides from food waste. *Chemical Engineering Journal* **2020**, *395*, 125056.
26. Wu, J.; Chen, R.; Tan, L.; Bai, H.; Tian, L.; Lu, J.; Gao, M.; Bai, C.; Sun, H.; Chi, Y., Ultrasonic disruption effects on the extraction efficiency, characterization, and bioactivities of polysaccharides from *Panax notoginseng* flower. *Carbohydrate Polymers* **2022**, *291*, 119535.
27. Carreira-Casais, A.; Otero, P.; Garcia-Perez, P.; Garcia-Oliveira, P.; Pereira, A.G.; Carpena, M.; Soria-Lopez, A.; Simal-Gandara, J.; Prieto, M.A., Benefits and Drawbacks of Ultrasound-Assisted Extraction for the Recovery of Bioactive Compounds from Marine Algae. *International Journal of Environmental Research and Public Health* **2021**, *18*.

28. Tang, L.; Luo, X.; Wang, M.; Wang, Z.; Guo, J.; Kong, F.; Bi, Y., Synthesis, characterization, in vitro antioxidant and hypoglycemic activities of selenium nanoparticles decorated with polysaccharides of *Gracilaria lemaneiformis*. *International Journal of Biological Macromolecules* **2021**, 193, 923-932.
29. Shi, F.; Yan, X.; Cheong, K.-L.; Liu, Y., Extraction, purification, and characterization of polysaccharides from marine algae *Gracilaria lemaneiformis* with anti-tumor activity. *Process Biochemistry* **2018**, 73, 197-203.
30. Giovannoni, M.; Gramegna, G.; Benedetti, M.; Mattei, B., Industrial Use of Cell Wall Degrading Enzymes: The Fine Line Between Production Strategy and Economic Feasibility. *Frontiers in Bioengineering and Biotechnology* **2020**, 8, 356.
31. Fournière, M.; Latire, T.; Lang, M.; Terme, N.; Bourgougnon, N.; Bedoux, G., Production of Active Poly- and Oligosaccharidic Fractions from *Ulva* sp. by Combining Enzyme-Assisted Extraction (EAE) and Depolymerization. *Metabolites* **2019**, 9, 182.
32. Łubek-Nguyen, A.; Ziemichód, W.; Olech, M., Application of Enzyme-Assisted Extraction for the Recovery of Natural Bioactive Compounds for Nutraceutical and Pharmaceutical Applications. *Applied Sciences* **2022**, 12, 3232.
33. Wu, S.; Lu, M.; Wang, S., Amylase-assisted extraction and antioxidant activity of polysaccharides from *Gracilaria lemaneiformis*. *3 Biotech* **2017**, 7, 38.
34. Chen, H.; Xiao, Q.; Weng, H.; Zhang, Y.; Yang, Q.; Xiao, A., Extraction of sulfated agar from *Gracilaria lemaneiformis* using hydrogen peroxide-assisted enzymatic method. *Carbohydrate Polymers* **2020**, 232, 115790.
35. Dobrinčić, A.; Balbino, S.; Zorić, Z.; Pedisić, S.; Bursać Kovačević, D.; Elez Garofulić, I.; Dragović-Uzelac, V., Advanced Technologies for the Extraction of Marine Brown Algal Polysaccharides. *Marine Drugs* **2020**, 18, 168.
36. Das, S.; Nadar, S.S.; Rathod, V.K., Integrated strategies for enzyme assisted extraction of bioactive molecules: A review. *International Journal of Biological Macromolecules* **2021**, 191, 899-917.
37. Boussetta, N.; Turk, M.; De Taeye, C.; Larondelle, Y.; Lanoisellé, J.L.; Vorobiev, E., Effect of high voltage electrical discharges, heating and ethanol concentration on the extraction of total polyphenols and lignans from flaxseed cake. *Industrial Crops and Products* **2013**, 49, 690-696.
38. Ju, T.; Deng, Y.; Xi, J., Optimization of Circulating Extraction of Polysaccharides from *Gracilaria Lemaneiformis* Using Pulsed Electrical Discharge. *ACS Sustainable Chemistry & Engineering* **2019**, 7, 3593-3601.
39. Ju, T.; Xi, J., Continuous extraction optimization, molecular structures and antioxidant activities of polysaccharide from *Gracilariopsis lemaneiformis* using liquid-phase pulsed discharge. *Separation and Purification Technology* **2020**, 236, 116241.
40. Niu, G.; You, G.; Zhou, X.; Fan, H.; Liu, X., Physicochemical properties and in vitro hypoglycemic activities of hsian-tsao polysaccharide fractions by gradient ethanol precipitation method. *International Journal of Biological Macromolecules* **2023**, 231, 123274.
41. Zou, P.; Yang, X.; Huang, W.-W.; Zhao, H.-T.; Wang, J.; Xu, R.-B.; Hu, X.-L.; Shen, S.-Y.; Qin, D., Characterization and Bioactivity of Polysaccharides Obtained from Pine Cones of *Pinus koraiensis* by Graded Ethanol Precipitation. *Molecules* **2013**, 18, 9933-9948.
42. Li, X.; Gong, Y.; Yao, W.; Chen, X.; Xian, J.; You, L.; Fardim, P., Structural characterization and protective effects of polysaccharide from *Gracilaria lemaneiformis* on LPS-induced injury in IEC-6 cells. *Food Chemistry: X* **2021**, 12, 100157.
43. Shi, L., Bioactivities, isolation and purification methods of polysaccharides from natural products: A review. *International Journal of Biological Macromolecules* **2016**, 92, 37-48.
44. Ren, Y.; Bai, Y.; Zhang, Z.; Cai, W.; Del Rio Flores, A., The Preparation and Structure Analysis Methods of Natural Polysaccharides of Plants and Fungi: A Review of Recent Development. *Molecules* **2019**, 24, 3122.
45. Liu, H.-B.; Li, B.; Guo, L.-W.; Pan, L.-M.; Zhu, H.-X.; Tang, Z.-S.; Xing, W.-H.; Cai, Y.-Y.; Duan, J.-A.; Wang, M.; Xu, S.-N.; Tao, X.-B., Current and Future Use of Membrane Technology in the Traditional Chinese Medicine Industry. *Separation & Purification Reviews* **2022**, 51, 484-502.
46. Shao, W.; Zhang, H.; Duan, R.; Xie, Q.; Hong, Z.; Xiao, Z., A rapid and scalable integrated membrane separation process for purification of polysaccharides from *Enteromorpha prolifera*. *Natural Product Research* **2019**, 33, 3109-3119.
47. Li, X.; Huang, S.; Chen, X.; Xu, Q.; Ma, Y.; You, L.; Kulikouskaya, V.; Xiao, J.; Piao, J., Structural characteristic of a sulfated polysaccharide from *Gracilaria Lemaneiformis* and its lipid metabolism regulation effect. *Food & Function* **2020**, 11, 10876-10885.
48. Eder, S.; Zueblin, P.; Diener, M.; Peydayesh, M.; Boulos, S.; Mezzenga, R.; Nyström, L., Effect of Polysaccharide Conformation on Ultrafiltration Separation Performance. *Carbohydrate Polymers* **2021**, 260, 117830.
49. Zhang, Y.; Teng, J.; Zou, H.; Zhang, W.; Cheng, S.; Zhang, M.; Lin, H., The molecular weight-fouling matrix: A novel dissection of polysaccharide interactions in ultrafiltration processes. *Separation and Purification Technology* **2024**, 345, 127340.

50. Sun, H.; Qi, D.; Xu, J.; Juan, S.; Zhe, C., Fractionation of polysaccharides from rapeseed by ultrafiltration: Effect of molecular pore size and operation conditions on the membrane performance. *Separation and Purification Technology* **2011**, 80, 670-676.
51. Xie, J.-H.; Shen, M.-Y.; Nie, S.-P.; Zhao, Q.; Li, C.; Xie, M.-Y., Separation of water-soluble polysaccharides from *Cyclocarya paliurus* by ultrafiltration process. *Carbohydrate Polymers* **2014**, 101, 479-483.
52. Pinelo, M.; Jonsson, G.; Meyer, A.S., Membrane technology for purification of enzymatically produced oligosaccharides: Molecular and operational features affecting performance. *Separation and Purification Technology* **2009**, 70, 1-11.
53. Kravtchenko, T.P.; Voragen, A.G.J.; Pilnik, W., Studies on the intermolecular distribution of industrial pectins by means of preparative ion-exchange chromatography. *Carbohydrate Polymers* **1992**, 19, 115-124.
54. Chen, Y.; Yao, F.; Ming, K.; Wang, D.; Hu, Y.; Liu, J., Polysaccharides from Traditional Chinese Medicines: Extraction, Purification, Modification, and Biological Activity. *Molecules* **2016**, 21, 1705.
55. Huang, X.; Ai, C.; Yao, H.; Zhao, C.; Xiang, C.; Hong, T.; Xiao, J., Guideline for the extraction, isolation, purification, and structural characterization of polysaccharides from natural resources. *eFood* **2022**, 3, e37.
56. Veeraperumal, S.; Qiu, H.-M.; Zeng, S.-S.; Yao, W.-Z.; Wang, B.-P.; Liu, Y.; Cheong, K.-L., Polysaccharides from *Gracilaria lemaneiformis* promote the HaCaT keratinocytes wound healing by polarised and directional cell migration. *Carbohydrate Polymers* **2020**, 241, 116310.
57. Ren, Y.; Zheng, G.; You, L.; Wen, L.; Li, C.; Fu, X.; Zhou, L., Structural characterization and macrophage immunomodulatory activity of a polysaccharide isolated from *Gracilaria lemaneiformis*. *Journal of Functional Foods* **2017**, 33, 286-296.
58. Lecacheux, D.; Brigand, G., Preparative fractionation of natural polysaccharides by size exclusion chromatography. *Carbohydrate Polymers* **1988**, 8, 119-130.
59. Wang, X.; Zhang, Y.; Liu, Z.; Zhao, M.; Liu, P., Purification, Characterization, and Antioxidant Activity of Polysaccharides Isolated from Cortex Periplocae. *Molecules* **2017**, 22, 1866.
60. Yang, X.Q.; Liu, M.Q.; Qi, B.; Li, L.H.; Deng, J.C.; Hu, X.; Wu, Y.Y.; Hao, S.X., Extraction, purification and partial characterizations of polysaccharides from *Gracilaria lemaneiformis*. *Advanced Materials Research* **2014**, 881, 776-780.
61. Kang, Y.; Wang, Z.-J.; Xie, D.; Sun, X.; Yang, W.; Zhao, X.; Xu, N., Characterization and Potential Antitumor Activity of Polysaccharide from *Gracilariopsis lemaneiformis*. *Marine Drugs* **2017**, 15, 100.
62. Garcia-Vaquero, M.; Rajauria, G.; O'Doherty, J.V.; Sweeney, T., Polysaccharides from macroalgae: Recent advances, innovative technologies and challenges in extraction and purification. *Food Research International* **2017**, 99, 1011-1020.
63. Cheong, K.-L.; Liu, K.; Chen, W.; Zhong, S.; Tan, K., Recent progress in *Porphyra haitanensis* polysaccharides: Extraction, purification, structural insights, and their impact on gastrointestinal health and oxidative stress management. *Food Chemistry: X* **2024**, 22, 101414.
64. Yao, W.; Gong, Y.; Li, L.; Hu, X.; You, L., The effects of dietary fibers from rice bran and wheat bran on gut microbiota: An overview. *Food Chemistry: X* **2022**, 13, 100252.
65. Yan, J.; Shi, S.; Wang, H.; Liu, R.; Li, N.; Chen, Y.; Wang, S., Neutral monosaccharide composition analysis of plant-derived oligo- and polysaccharides by high performance liquid chromatography. *Carbohydrate Polymers* **2016**, 136, 1273-1280.
66. Villanueva-Suárez, M.J.; Redondo-Cuenca, A.; Rodríguez-Sevilla, M.D.; de las Heras Martínez, M., Characterization of Nonstarch Polysaccharides Content from Different Edible Organs of Some Vegetables, Determined by GC and HPLC: Comparative Study. *Journal of Agricultural and Food Chemistry* **2003**, 51, 5950-5955.
67. Bakky, M.A.H.; Tran, N.T.; Zhang, M.; Zhang, Y.; Liang, H.; Wang, Y.; Zhang, Y.; Ma, H.; Zheng, H.; Li, S., In vitro fermentation of *Gracilaria lemaneiformis* and its sulfated polysaccharides by rabbitfish gut microbes. *International Journal of Biological Macromolecules* **2023**, 246, 125561.
68. Jol, C.N.; Neiss, T.G.; Penninkhof, B.; Rudolph, B.; De Ruiter, G.A., A Novel High-Performance Anion-Exchange Chromatographic Method for the Analysis of Carrageenans and Agars Containing 3,6-Anhydrogalactose. *Analytical Biochemistry* **1999**, 268, 213-222.
69. Navarro, D.A.; Stortz, C.A., Determination of the configuration of 3,6-anhydrogalactose and cyclizable α -galactose 6-sulfate units in red seaweed galactans. *Carbohydrate Research* **2003**, 338, 2111-2118.
70. Xie, X.-T.; Zhang, X.; Liu, Y.; Chen, X.-Q.; Cheong, K.-L., Quantification of 3,6-anhydro-galactose in red seaweed polysaccharides and their potential skin-whitening activity. *3 Biotech* **2020**, 10, 189.
71. Wang, X.; Xu, X.; Mao, G.; Guo, Y.; Wang, G.; Sun, X.; Xu, N.; Zhang, Z., Structural Characterization of *Gracilariopsis lemaneiformis* Polysaccharide and Its Property in Delaying Cellular Senescence. *Frontiers in Nutrition* **2022**, 9, 876992.
72. Liao, X.; Yang, L.; Chen, M.; Yu, J.; Zhang, S.; Ju, Y., The hypoglycemic effect of a polysaccharide (GLP) from *Gracilaria lemaneiformis* and its degradation products in diabetic mice. *Food & Function* **2015**, 6, 2542-2549.

73. Bajwa, B.; Xing, X.; Terry, S.A.; Gruninger, R.J.; Abbott, D.W., Methylation-GC-MS/FID-Based Glycosidic Linkage Analysis of Unfractionated Polysaccharides in Red Seaweeds. *Marine Drugs* **2024**, *22*, 192.
74. Feng, Z.; Deng, L.; Chen, X.; Ouyang, P.; Li, Y.; Tang, X.; Fan, L.; Tan, H.; Mei, S.; Ye, H.; Wu, K.; Luo, H., Structural characterization and anti-pigmentation of a novel heteropolysaccharide from *Gracilaria lemaneiformis* via α -MSH/MC1R pathway. *Journal of Functional Foods* **2023**, *107*, 105650.
75. Aslan, M.; Dogan, S.; Kucuksayan, E., Oxidative stress and potential applications of free radical scavengers in glaucoma. *Redox Report* **2013**, *18*, 76-87.
76. Poprac, P.; Jomova, K.; Simunkova, M.; Kollar, V.; Rhodes, C.J.; Valko, M., Targeting Free Radicals in Oxidative Stress-Related Human Diseases. *Trends in Pharmacological Sciences* **2017**, *38*, 592-607.
77. Simpson, D.S.A.; Oliver, P.L., ROS Generation in Microglia: Understanding Oxidative Stress and Inflammation in Neurodegenerative Disease. *Antioxidants* **2020**, *9*, 743.
78. Islam, M.T., Oxidative stress and mitochondrial dysfunction-linked neurodegenerative disorders. *Neurological Research* **2017**, *39*, 73-82.
79. Wang, X.; Zhang, Z.; Wu, Y.; Sun, X.; Xu, N., Synthesized sulfated and acetylated derivatives of polysaccharide extracted from *Gracilariopsis lemaneiformis* and their potential antioxidant and immunological activity. *International Journal of Biological Macromolecules* **2019**, *124*, 568-572.
80. Long, X.; Hu, X.; Pan, C.; Xiang, H.; Chen, S.; Qi, B.; Liu, S.; Yang, X., Antioxidant Activity of *Gracilaria lemaneiformis* Polysaccharide Degradation Based on Nrf-2/Keap-1 Signaling Pathway in HepG2 Cells with Oxidative Stress Induced by H₂O₂. *Marine Drugs* **2022**, *20*, 545.
81. Fang, T.; Zhang, X.; Hu, S.; Yu, Y.; Sun, X.; Xu, N., Enzymatic Degradation of *Gracilariopsis lemaneiformis* Polysaccharide and the Antioxidant Activity of Its Degradation Products. *Marine Drugs* **2021**, *19*, 270.
82. Hajam, Y.A.; Rani, R.; Ganie, S.Y.; Sheikh, T.A.; Javaid, D.; Qadri, S.S.; Pramodh, S.; Alsulimani, A.; Alkhanani, M.F.; Harakeh, S.; Hussain, A.; Haque, S.; Reshi, M.S., Oxidative Stress in Human Pathology and Aging: Molecular Mechanisms and Perspectives. *Cells* **2022**, *11*, 552.
83. Volpe, C.M.O.; Villar-Delfino, P.H.; dos Anjos, P.M.F.; Nogueira-Machado, J.A., Cellular death, reactive oxygen species (ROS) and diabetic complications. *Cell Death & Disease* **2018**, *9*, 119.
84. Okoduwa, S.I.R.; Umar, I.A.; Ibrahim, S.; Bello, F.; Habila, N., Age-dependent alteration of antioxidant defense system in hypertensive and type-2 diabetes patients. *Journal of Diabetes & Metabolic Disorders* **2015**, *14*, 32.
85. Zhang, Z.; Li, X.; Xu, X.; Mao, G.; Sun, X.; Xu, N.; Wang, X., *Gracilariopsis lemaneiformis* Polysaccharide Attenuates D-Galactose-Induced Aging of Mice by Regulating Oxidative Stress and Gut Microbiota. *Frontiers in Marine Sciences* **2022**, *9*.
86. Wen, L.; Zhang, Y.; Sun-Waterhouse, D.; You, L.; Fu, X., Advantages of the polysaccharides from *Gracilaria lemaneiformis* over metformin in antidiabetic effects on streptozotocin-induced diabetic mice. *RSC Advances* **2017**, *7*, 9141-9151.
87. Lu, Y.; Mei, S.; Wang, P.; Ouyang, P.; Liao, X.; Ye, H.; Wu, K.; Ma, X., Protective effects of *Gracilaria lemaneiformis* extract against ultraviolet B-induced damage in HaCaT cells. *Pharmacognosy Magazine* **2020**, *16*.
88. Miller, A.H.; Haroon, E.; Felger, J.C., The Immunology of Behavior—Exploring the Role of the Immune System in Brain Health and Illness. *Neuropsychopharmacology* **2017**, *42*, 1-4.
89. Antushevich, H., Interplays between inflammasomes and viruses, bacteria (pathogenic and probiotic), yeasts and parasites. *Immunology Letters* **2020**, *228*, 1-14.
90. Lemke, G., How macrophages deal with death. *Nature Reviews Immunology* **2019**, *19*, 539-549.
91. Mosser, D.M.; Hamidzadeh, K.; Goncalves, R., Macrophages and the maintenance of homeostasis. *Cellular & Molecular Immunology* **2021**, *18*, 579-587.
92. Becher, B.; Spath, S.; Goverman, J., Cytokine networks in neuroinflammation. *Nature Reviews Immunology* **2017**, *17*, 49-59.
93. Plaza-Díaz, J.; Ruiz-Ojeda, F.J.; Vilchez-Padial, L.M.; Gil, A., Evidence of the Anti-Inflammatory Effects of Probiotics and Synbiotics in Intestinal Chronic Diseases. *Nutrients* **2017**, *9*, 555.
94. Alfaddagh, A.; Martin, S.S.; Leucker, T.M.; Michos, E.D.; Blaha, M.J.; Lowenstein, C.J.; Jones, S.R.; Toth, P.P., Inflammation and cardiovascular disease: From mechanisms to therapeutics. *American Journal of Preventive Cardiology* **2020**, *4*, 100130.
95. Qiu, S.-M.; Veeraperumal, S.; Tan, K.; Zhong, S.; Cheong, K.-L., The in vitro anti-inflammatory mechanism of *Porphyra haitanensis* oligosaccharides on lipopolysaccharide-induced injury in IEC-6 cells. *Journal of Functional Foods* **2024**, *112*, 106005.
96. Cronkite, D.A.; Strutt, T.M., The Regulation of Inflammation by Innate and Adaptive Lymphocytes. *Journal of Immunology Research* **2018**, *2018*, 1467538.
97. Zhang, Y.; Che, H.; Li, C.; Jin, T., Food Allergens of Plant Origin. *Foods* **2023**, *12*, 2232.
98. Chen, H.-Y.; Zhou, Y.-C.; Liu, Y.; Huang, J.-Y.; Liu, H.; Liu, C.-F.; Liu, W.-H.; Liu, G.-M.; Liu, Q.-M., Fermented *Gracilaria lemaneiformis* polysaccharides alleviate food allergy by regulating Treg cells and gut microbiota. *International Journal of Biological Macromolecules* **2024**, *269*, 132215.

99. Liu, Q.; Zhang, Y.; Shu, Z.; Liu, M.; Zeng, R.; Wang, Y.; Liu, H.; Cao, M.; Su, W.; Liu, G., Sulfated oligosaccharide of *Gracilaria lemaneiformis* protect against food allergic response in mice by up-regulating immunosuppression. *Carbohydrate Polymers* **2020**, 230, 115567.
100. Liu, Q.; Zhou, Y.; Ma, L.; Gu, F.; Liao, K.; Liu, Y.; Zhang, Y.; Liu, H.; Hong, Y.; Cao, M.; Liu, W.-H.; Liu, C.; Liu, G., Sulfate oligosaccharide of *Gracilaria lemaneiformis* modulates type 1 immunity by restraining T cell activation. *Carbohydrate Polymers* **2022**, 288, 119377.
101. Hou, C.; Yin, M.; Lan, P.; Wang, H.; Nie, H.; Ji, X., Recent progress in the research of *Angelica sinensis* (Oliv.) Diels polysaccharides: extraction, purification, structure and bioactivities. *Chemical and Biological Technologies in Agriculture* **2021**, 8, 13.
102. Yao, W.; Qiu, H.-M.; Cheong, K.-L.; Zhong, S., Advances in anti-cancer effects and underlying mechanisms of marine algae polysaccharides. *International Journal of Biological Macromolecules* **2022**, 221, 472-485.
103. Wang, Y.; Zhang, Q.; Chen, Y.; Liang, C.-L.; Liu, H.; Qiu, F.; Dai, Z., Antitumor effects of immunity-enhancing traditional Chinese medicine. *Biomedicine & Pharmacotherapy* **2020**, 121, 109570.
104. Ji, H.-y.; Yu, J.; Dong, X.-d.; Liu, A.-j., Preparation of soluble dietary fibers from *Gracilaria lemaneiformis* and its antitumor activity *in vivo*. *Journal of Food Measurement and Characterization* **2019**, 13, 1574-1582.
105. Kondoh, N.; Mizuno-Kamiya, M., The Role of Immune Modulatory Cytokines in the Tumor Microenvironments of Head and Neck Squamous Cell Carcinomas. *Cancers* **2022**, 14, 2884.
106. Kang, Y.; Li, H.; Wu, J.; Xu, X.; Sun, X.; Zhao, X.; Xu, N., Transcriptome Profiling Reveals the Antitumor Mechanism of Polysaccharide from Marine Algae *Gracilariopsis lemaneiformis*. *PLoS One* **2016**, 11, e0158279.
107. Seyrek, K.; Ivanisenko, N.V.; Wohlfromm, F.; Espe, J.; Lavrik, I.N., Impact of human CD95 mutations on cell death and autoimmunity: a model. *Trends in Immunology* **2022**, 43, 22-40.
108. Cai, B.; Luo, L.; Chen, X.; Zhao, X.; He, J.; Chen, H.; Wan, P.; Chen, D.; Pan, J., Structural characterization of a sulfated polysaccharide from *Gracilariopsis lemaneiformis* and its potentiation of cisplatin antitumor activity in Colon-26 carcinoma tumor-bearing mice by inducing ferroptosis. *Food & Function* **2023**, 14, 3712-3721.
109. Li, J.; Cao, F.; Yin, H.-l.; Huang, Z.-j.; Lin, Z.-t.; Mao, N.; Sun, B.; Wang, G., Ferroptosis: past, present and future. *Cell Death & Disease* **2020**, 11, 88.
110. Fan, Y.; Pedersen, O., Gut microbiota in human metabolic health and disease. *Nature Reviews Microbiology* **2021**, 19, 55-71.
111. Zhang, P., Influence of Foods and Nutrition on the Gut Microbiome and Implications for Intestinal Health. *International Journal of Molecular Sciences* **2022**, 23, 9588.
112. Colella, M.; Charitos, I.A.; Ballini, A.; Cafiero, C.; Topi, S.; Palmirotta, R.; Santacroce, L., Microbiota revolution: How gut microbes regulate our lives. *World journal of gastroenterology* **2023**, 29, 4368-4383.
113. Kudelka, M.R.; Stowell, S.R.; Cummings, R.D.; Neish, A.S., Intestinal epithelial glycosylation in homeostasis and gut microbiota interactions in IBD. *Nature Reviews Gastroenterology & Hepatology* **2020**, 17, 597-617.
114. Haneishi, Y.; Furuya, Y.; Hasegawa, M.; Picarelli, A.; Rossi, M.; Miyamoto, J., Inflammatory Bowel Diseases and Gut Microbiota. *International Journal of Molecular Sciences* **2023**, 24, 3817.
115. Lu, S.-Y.; Tan, K.; Zhong, S.; Cheong, K.-L., Marine algal polysaccharides as future potential constituents against non-alcoholic steatohepatitis. *International Journal of Biological Macromolecules* **2023**, 250, 126247.
116. Yang, G.; Wei, J.; Liu, P.; Zhang, Q.; Tian, Y.; Hou, G.; Meng, L.; Xin, Y.; Jiang, X., Role of the gut microbiota in type 2 diabetes and related diseases. *Metabolism* **2021**, 117, 154712.
117. Chakaroun, R.M.; Massier, L.; Kovacs, P., Gut Microbiome, Intestinal Permeability, and Tissue Bacteria in Metabolic Disease: Perpetrators or Bystanders? *Nutrients* **2020**, 12, 1082.
118. Margolis, K.G.; Cryan, J.F.; Mayer, E.A., The Microbiota-Gut-Brain Axis: From Motility to Mood. *Gastroenterology* **2021**, 160, 1486-1501.
119. Wang, M.; Veeraperumal, S.; Zhong, S.; Cheong, K.-L., Fucoindan-Derived Functional Oligosaccharides: Recent Developments, Preparation, and Potential Applications. *Foods* **2023**, 12, 878.
120. Divyashri, G.; Karthik, P.; Murthy, T.P.K.; Priyadarshini, D.; Reddy, K.R.; Raghu, A.V.; Vaidyanathan, V.K., Non-digestible oligosaccharides-based prebiotics to ameliorate obesity: Overview of experimental evidence and future perspectives. *Food Science and Biotechnology* **2023**, 32, 1993-2011.
121. Li, Y.-F.; Wu, B.; Chen, J.-p.; Veeraperumal, S.; Wei, J.-C.; Tan, K.-s.; Zhong, S.; Cheong, K.-L., Prebiotic characteristics of added-value polysaccharides from jackfruit peel waste during *in vitro* digestion and fecal fermentation. *LWT* **2023**, 187, 115330.
122. Xie, X.-T.; Cheong, K.-L., Recent advances in marine algae oligosaccharides: structure, analysis, and potential prebiotic activities. *Critical Reviews in Food Science and Nutrition* **2022**, 62, 7703-7717.
123. Xu, S.-Y.; Chen, X.-Q.; Liu, Y.; Cheong, K.-L., Ultrasonic/microwave-assisted extraction, simulated digestion, and fermentation *in vitro* by human intestinal flora of polysaccharides from *Porphyra haitanensis*. *International Journal of Biological Macromolecules* **2020**, 152, 748-756.
124. Koliada, A.; Moseiko, V.; Romanenko, M.; Lushchak, O.; Kryzhanovska, N.; Guryanov, V.; Vaiserman, A., Sex differences in the phylum-level human gut microbiota composition. *BMC Microbiology* **2021**, 21, 131.

125. Wardman, J.F.; Bains, R.K.; Rahfeld, P.; Withers, S.G., Carbohydrate-active enzymes (CAZymes) in the gut microbiome. *Nature Reviews Microbiology* **2022**, 20, 542-556.
126. McKee, L.S.; La Rosa, S.L.; Westereng, B.; Eijssink, V.G.; Pope, P.B.; Larsbrink, J., Polysaccharide degradation by the Bacteroidetes: mechanisms and nomenclature. *Environmental Microbiology Reports* **2021**, 13, 559-581.
127. Hehemann, J.-H.; Kelly, A.G.; Pudlo, N.A.; Martens, E.C.; Boraston, A.B., Bacteria of the human gut microbiome catabolize red seaweed glycans with carbohydrate-active enzyme updates from extrinsic microbes. *Proceedings of the National Academy of Sciences* **2012**, 109, 19786-19791.
128. Yu, B.; Lu, Z.; Zhong, S.; Cheong, K.-L., Exploring potential polysaccharide utilization loci involved in the degradation of typical marine seaweed polysaccharides by *Bacteroides thetaiotaomicron*. *Frontiers in Microbiology* **2024**, 15, 1332105.
129. Zhang, Y.; Zheng, L.; Liu, G.; Shen, J.; Chen, G.; Mei, X.; Chang, Y.; Xue, C., The α -linkage in funoran and agarose could be hydrolyzed by a GH96 family enzyme: Discovery of the α -funoranase. *Carbohydrate Polymers* **2024**, 338, 122201.
130. Ho Do, M.; Seo, Y.S.; Park, H.-Y., Polysaccharides: bowel health and gut microbiota. *Critical Reviews in Food Science and Nutrition* **2021**, 61, 1212-1224.
131. Sun, X.; Duan, M.; Liu, Y.; Luo, T.; Ma, N.; Song, S.; Ai, C., The beneficial effects of *Gracilaria lemaneiformis* polysaccharides on obesity and the gut microbiota in high fat diet-fed mice. *Journal of Functional Foods* **2018**, 46, 48-56.
132. Kim, C.H., Microbiota or short-chain fatty acids: which regulates diabetes? *Cellular & Molecular Immunology* **2018**, 15, 88-91.
133. van der Hee, B.; Wells, J.M., Microbial Regulation of Host Physiology by Short-chain Fatty Acids. *Trends in Microbiology* **2021**, 29, 700-712.
134. Mandaliya, D.K.; Patel, S.; Seshadri, S., The Combinatorial Effect of Acetate and Propionate on High-Fat Diet Induced Diabetic Inflammation or Metaflammation and T Cell Polarization. *Inflammation* **2021**, 44, 68-79.
135. Fu, X.; Liu, Z.; Zhu, C.; Mou, H.; Kong, Q., Nondigestible carbohydrates, butyrate, and butyrate-producing bacteria. *Critical Reviews in Food Science and Nutrition* **2019**, 59, S130-S152.
136. Cheong, K.-L.; Xie, X.-T.; Zhou, T.; Malairaj, S.; Veeraperumal, S.; Zhong, S.; Tan, K., Exploring the therapeutic potential of porphyran extracted from *Porphyra haitanensis* in the attenuation of DSS-induced intestinal inflammation. *International Journal of Biological Macromolecules* **2024**, 271, 132578.
137. Feng, Z.; Sun, R.; Cong, Y.; Liu, Z., Critical roles of G protein-coupled receptors in regulating intestinal homeostasis and inflammatory bowel disease. *Mucosal Immunology* **2022**, 15, 819-828.
138. Felizardo, R.J.F.; de Almeida, D.C.; Pereira, R.L.; Watanabe, I.K.M.; Doimo, N.T.S.; Ribeiro, W.R.; Cenedeze, M.A.; Hiyane, M.I.; Amano, M.T.; Braga, T.T.; Ferreira, C.M.; Parmigiani, R.B.; Andrade-Oliveira, V.; Volpini, R.A.; Vinolo, M.A.R.; Mariño, E.; Robert, R.; Mackay, C.R.; Camara, N.O.S., Gut microbial metabolite butyrate protects against proteinuric kidney disease through epigenetic- and GPR109a-mediated mechanisms. *The FASEB Journal* **2019**, 33, 11894-11908.
139. Han, R.; Ma, Y.; Xiao, J.; You, L.; Pedisić, S.; Liao, L., The possible mechanism of the protective effect of a sulfated polysaccharide from *Gracilaria Lemaneiformis* against colitis induced by dextran sulfate sodium in mice. *Food and Chemical Toxicology* **2021**, 149, 112001.
140. Kim, C.H., Complex regulatory effects of gut microbial short-chain fatty acids on immune tolerance and autoimmunity. *Cellular & Molecular Immunology* **2023**, 20, 341-350.
141. Gonçalves, P.; Araújo, J.R.; Di Santo, J.P., A Cross-Talk Between Microbiota-Derived Short-Chain Fatty Acids and the Host Mucosal Immune System Regulates Intestinal Homeostasis and Inflammatory Bowel Disease. *Inflammatory Bowel Diseases* **2018**, 24, 558-572.
142. Lu, S.-Y.; Liu, Y.; Tang, S.; Zhang, W.; Yu, Q.; Shi, C.; Cheong, K.-L., *Gracilaria lemaneiformis* polysaccharides alleviate colitis by modulating the gut microbiota and intestinal barrier in mice. *Food Chemistry: X* **2022**, 13, 100197.
143. Xie, X.-T.; Zheng, L.-X.; Duan, H.-M.; Liu, Y.; Chen, X.-Q.; Cheong, K.-L., Structural characteristics of *Gracilaria lemaneiformis* oligosaccharides and their alleviation of dextran sulphate sodium-induced colitis by modulating the gut microbiota and intestinal metabolites in mice. *Food & Function* **2021**, 12, 8635-8646.
144. Zhang, S.; Zhang, M.; Li, W.; Ma, L.; Liu, X.; Ding, Q.; Yu, W.; Yu, T.; Ding, C.; Liu, W., Research progress of natural plant polysaccharides inhibiting inflammatory signaling pathways and regulating intestinal flora and metabolism to protect inflammatory bowel disease. *International Journal of Biological Macromolecules* **2023**, 253, 126799.
145. Chand Dakal, T.; Dhabhai, B.; Agarwal, D.; Gupta, R.; Nagda, G.; Meena, A.R.; Dhakar, R.; Menon, A.; Mathur, R.; Mona; Yadav, V.; Sharma, A., Mechanistic basis of co-stimulatory CD40-CD40L ligation mediated regulation of immune responses in cancer and autoimmune disorders. *Immunobiology* **2020**, 225, 151899.

146. Liu, J.-H.; Zheng, Y.-Y.; Ouyang, J.-M., Antioxidant Activities and Cytotoxicity of the Regulated Calcium Oxalate Crystals on HK-2 Cells of Polysaccharides from *Gracilaria lemaneiformis* with Different Molecular Weights. *Foods* **2023**, 12, 1031.
147. Long, X.; Hu, X.; Zhou, S.; Xiang, H.; Chen, S.; Li, L.; Liu, S.; Yang, X., Optimized Degradation and Inhibition of α -glucosidase Activity by *Gracilaria lemaneiformis* Polysaccharide and Its Production *In Vitro*. *Marine Drugs* **2022**, 20, 13.
148. Chen, M.-z.; Xie, H.-g.; Yang, L.-w.; Liao, Z.-h.; Yu, J., In vitro anti-influenza virus activities of sulfated polysaccharide fractions from *Gracilaria lemaneiformis*. *Virologica Sinica* **2010**, 25, 341-351.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.