

Article

Not peer-reviewed version

Molecular Characterization of Sterol C4-Methyl Oxidase in *Leishmania major*

Yu Ning , [Somrita Basu](#) , [Fong-fu Hsu](#) , [Mei Feng](#) , [Michael Zhuo Wang](#) , [Kai Zhang](#) *

Posted Date: 2 August 2024

doi: 10.20944/preprints202408.0195.v1

Keywords: CYP5122A1; ERG25; *Leishmania*; Sterol; Demethylase



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Molecular Characterization of Sterol C4-Methyl Oxidase in *Leishmania major*

Yu Ning¹, Somrita Basu¹, Fong-fu Hsu², Mei Feng³, Michael Zhuo Wang³, and Kai Zhang^{1,*}

¹ Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409, USA.

² Mass Spectrometry Resource, Division of Endocrinology, Diabetes, Metabolism, and Lipid research, Department of Internal Medicine, Washington University School of Medicine, St. Louis, MO, 63110, USA.

³ Department of Pharmaceutical Chemistry, School of Pharmacy, The University of Kansas, Lawrence, KS 66047, USA.

* Correspondence: kai.zhang@ttu.edu; Tel.: 1-806-834-0550

Abstract: Sterol biosynthesis requires the oxidative removal of two methyl groups from the C-4 position by sterol C-4-demethylase and one methyl group from the C-14 position by sterol C-14-demethylase. In *Leishmania donovani*, a CYP5122A1 protein was recently identified as the bona fide sterol C-4 methyl oxidase catalyzing the initial steps of C-4-demethylation. Besides CYP5122A1, *Leishmania* parasites possess orthologs to ERG25, the canonical sterol C-4 methyl oxidase in *Saccharomyces cerevisiae*. To determine the contribution of CYP5122A1 and ERG25 in sterol biosynthesis, we characterized these genes in *Leishmania major* which cause cutaneous leishmaniasis. Like in *L. donovani*, CYP5122A1 in *L. major* could only be deleted in the presence of a complementing episome. Even with strong negative selection, *L. major* chromosomal CYP5122A1-null mutants retained the complementing episome in both promastigote and amastigote stages, demonstrating its essentiality. In contrast, the *L. major* ERG25-null mutants were fully viable and replicative in culture and virulent in mice. Deletion and overexpression of ERG25 did not affect the sterol composition, indicating that ERG25 is not required for C-4-demethylation. These findings suggest that CYP5122A1 is the dominant and possibly only sterol C-4 methyl oxidase in *Leishmania*, and inhibitors of CYP5122A1 may have strong therapeutic potential against multiple *Leishmania* species.

Keywords: CYP5122A1; ERG25; *Leishmania*; sterol; demethylase

1. Introduction

Leishmania parasites are trypanosomatid protozoans transmitted through the bite of sandflies. During their life cycle, these protozoans alternate between extracellular flagellated promastigotes in sandfly gut and intracellular non-flagellated amastigotes in mammalian macrophages. *Leishmania* parasites cause a group of diseases in humans with symptoms ranging from self-limiting skin lesions (cutaneous leishmaniasis) to life-threatening infections of the bone marrow, liver, and spleen (visceral leishmaniasis) [1]. Current treatments are plagued with high toxicity, low efficacy, and increasing resistance [2]. Thus, new drugs are needed to control leishmaniasis.

The sterol biosynthesis pathway presents multiple targets for antileishmanial drugs. Among them is sterol C14-demethylase (C14DM, also known as CYP51) which is the primary target of azoles [3,4]. C14DM is a heme-containing enzyme catalyzing the oxidative removal of the C14-methyl group from sterol intermediates (Figure 1)[5]. Genetic or chemical inhibition of C14DM in *Leishmania* leads to increased plasma membrane fluidity, hypersensitivity to heat, and compromised mitochondrial functions [6,7]. In addition to C14-demethylation, two more methyl groups need to be removed from the C4 position during sterol synthesis (Figure 1). In *Saccharomyces cerevisiae*, three enzymes catalyze sterol C4-demethylation: an O₂-dependent C4 sterol methyl oxidase (ERG25) that sequentially generates hydroxy, aldehyde and carboxylate intermediates; a C4 decarboxylase (ERG26) removing CO₂ from carboxylate intermediates to yield 3-ketosterol intermediates; and a 3-ketosterol reductase (ERG27) which regenerates the C3-hydroxy group [8–10]. While *S. cerevisiae* and vertebrates use the

same C4 sterol methyl oxidase (ERG25) on both C4 methyl groups, plants have two distinct ERG25 homologs that oxidize 4,4-dimethylsterols and 4 α -methylsterols, respectively [11,12] (Figure 1).




Figure 1. Demethylation steps catalyzed by C4DM and C14DM in *Leishmania* sterol synthesis pathway. I: lanosterol; II: 4,14-dimethylzymosterol; III: 14-methylzymosterol; IV: 4,4-dimethylzymosterol; V: 4-methylzymosterol; VI: zymosterol.

A recent study has identified *Leishmania donovani* CYP5122A1 (XP_003861867.1) as a non-canonical sterol C4-methyl oxidase [13]. CYP5122A1 is well conserved among multiple trypanosomatid species. Recombinant LdCYP5122A1 protein binds and converts C4-methylated sterols into oxidized metabolites in the presence of NADPH and oxygen. Importantly, CYP5122A1 is essential for both promastigotes and intracellular amastigotes in *L. donovani*. Heterozygous CYP5122A1 mutants (half knockout) in which one of the two CYP5122A1 alleles is deleted showed reduced infectivity in mice [13,14]. In addition, inhibitors of both CYP5122A1 and C14DM display superior antileishmanial activity against *L. donovani* in comparison to C14DM-selective inhibitors [13,15]. These findings suggest that C4DM is a promising drug target.

In addition to CYP5122A1, *Leishmania* parasites possess a putative ERG25 in their genomes. It is not known whether the leishmanial ERG25 protein is involved in the C4-demethylation reaction or whether its function overlaps with CYP5122A1. In this study, we investigated the roles of ERG25 in growth, virulence and sterol synthesis by characterizing genetic mutants of ERG25 in *Leishmania major*. We also explored the essentiality of CYP5122A1 in *L. major*. Our findings argue that CYP5122A1 is the sole sterol C4-methyl oxidase in *Leishmania*.

2. Results

2.1. Identification and Cellular Localization of ERG25 and CYP5122A1 (22A1) in *L. major*

We recently characterized the *Leishmania donovani* CYP5122A1 protein (TriTrypDB: LdBPK_270090.1; GenBank: XP_003861867.1) as a sterol C4-methyl oxidase catalyzing the sequential oxidation of lanosterol to form C4-oxidation metabolites [13] (Figure 1). *L. major* CYP5122A1 or 22A1 (TriTrypDB: LmjF.27.0090; GenBank: XP_003721815.1) is 94.4% identical to *L. donovani* CYP5122A1 with 606 amino acids and one transmembrane domain (AA51-74) (supplemental Figure S1). In addition to 22A1, *L. major* genome contains an ortholog of the *Saccharomyces cerevisiae* ERG25 gene (TriTrypDB: LmjF.36.2540) which encodes the sterol C4-methyl oxidase in yeast. *L. major* ERG25 has 402 amino acids with two transmembrane domains (AA67-90 and 136-159) showing a modest 26% identity to ScERG25 from AA194-384 (supplemental Figure S2). Based on information from TriTrypDB, CYP122A1 orthologs are present in multiple *Leishmania* and *Trypanosoma* species, but ERG25 orthologs are not found among *Trypanosoma* spp.

To determine the cellular localization of 22A1, a 22A1-GFP fusion protein gene was introduced into *L. major* wild type (WT) parasites. The resulting WT +22A1-GFP promastigotes were labeled with an anti-*T. brucei* BiP antibody to mark the endoplasmic reticulum (ER) [16]. Imaging analysis showed a ~72% overlap between 22A1-GFP and BiP (Figure 2A and 2C). Similar observations were obtained when a GFP-ERG25 fusion protein was expressed in *L. major* WT parasites (Figure 2B and 2C; ~80% overlap with BiP). Thus, 22A1 and ERG25 are primarily located in the ER.




Figure 2. Cellular localization of Cyp5122A1 and Erg25 in *L. major*. (A-B) Log phase promastigotes of WT +pXG-22A1-GFP (A) or WT + pXG-GFP-ERG25 (B) were labeled with an anti-*T. brucei* BiP antibody (ER marker) and processed for immunofluorescence microscopy. Merge images were the overlay of GFP and BiP. Overlap coefficients were determined from 30 cells each (average \pm standard deviations). Scale bars: 10 μ m. (C) Log phase promastigotes of WT, WT +pXG-22A1-GFP, WT + pXG-C14DM-GFP or WT + pXG-GFP-ERG25 were analyzed by western blot using antibodies against GFP (top) or α -tubulin.

2.2. Successful Generation of *ERG25*-Null but not *22A1*-Null Mutant in *L. major*

Using the targeted gene deletion approach based on homologous recombination, we replaced the two endogenous alleles of *ERG25* with drug resistant genes for puromycin (*PAC*) and hygromycin (*HYG*) [17]. The resulting *erg25*⁻ clones were verified by Southern blot (Figure 3A and supplemental Figure S3). Expression of *ERG25* was restored in the add-back strain *erg25*^{-/+}*ERG25* which overexpresses *ERG25* from pXG-*ERG25*.




Figure 3. Southern blots to confirm the deletions of chromosomal *ERG25* and *22A1* alleles. Genomic DNA samples from WT, *ERG25*^{+/-}, *erg25*⁻ (three clones), *22A1*^{+/-}, and *22A1*⁻+pXNG4-*22A1* (two clones) parasites were digested with restriction enzymes and hybridized with radiolabeled probes for the open reading frames (top) or upstream flanking regions (bottom) of *22A1* (A) or *Erg25* (B). Bands corresponding to *22A1*, pXNG4-*22A1*, *Erg25*, and antibiotic-resistant genes (*BSD*/*HYG*/*PAC*) are indicated. Details of the Southern blots were included in supplemental Figure. S3-S4.

With the same homologous recombination approach, we generated the heterozygous *22A1*^{+/-} parasites (half knockout). However, repeated attempts to generate *22A1*-null mutant were unsuccessful which was reminiscent of *CYP5122A1* in *L. donovani* [13]. To delete the chromosomal *22A1* alleles, pXNG4-*22A1* was introduced into the heterozygous *22A1*^{+/-} parasites. In the presence of episomal pXNG4-*22A1*, we were able to remove the second chromosomal *22A1* to generate *22A1*⁻+pXNG4-*22A1* (Figure 3B and supplemental Figure S4).

2.3. *ERG25* is not Required for the Growth or Virulence of *L. major* Promastigotes.

As shown in Figure 4A, the *erg25*⁻ and *erg25*^{-/+}*ERG25* promastigotes replicated at the same rate as WT parasites during the log phase. After entering the stationary stage, *erg25*⁻ had similar percentages of round cells and dead cells as WT and add-back parasites (Figure 4B-C). Thus, deletion or episomal overexpression of *ERG25* does not affect the survival or proliferation of *L. major* promastigotes in culture.




Figure 4. ERG25 is not required for proliferation or virulence in *L. major*. (A-C) WT, *erg25*⁻ and *erg25*⁻/*+ERG25* promastigotes were inoculated in complete M199 media at 1.0×10^5 cells/ml and culture densities were determined daily (A). After cultures entered the stationary stage (when densities reached 2.0×10^7 cells/ml), percentages of round cells (B) and dead cells (C) were determined daily. (D-E) Day 3 stationary phase promastigotes were injected into the footpads of female BALB/c mice. The development of footpad lesions was monitored weekly (D) and parasite numbers in infected footpads were determined at 4- and 7-weeks (one mouse each) post infection via limiting dilution assay (E). Error bars represent standard deviations.

To investigate the virulence of *erg25*⁻ in a mouse model, late stationary phase promastigotes were injected into the footpads of female BALB/c mice. As shown in Figure 4D, mice infected by *erg25*⁻ or *erg25*⁻/*+ERG25* developed lesions at similar rates as WT-infected mice. *ERG25* mutants appeared to have a slower growth rate in mice than WT based on parasite number analyses at 4- and 7-weeks post infection (Figure 4E). These findings suggest that while *ERG25* is not essential for virulence in *L. major*, endogenous *ERG25* expression is needed for the optimal growth in mice. In addition, the abundance and cellular localization of LPG and GP63, two major virulence factors in *Leishmania*, appeared to be unaltered in *erg25*⁻ by immunofluorescence microscopy analysis (supplemental Figure S5).

2.4. *ERG25* Deletion or Overexpression Does Not Affect Sterol Composition in *L. major*

To determine if *ERG25* is involved in sterol biosynthesis, we analyzed the sterol contents in log phase promastigotes of WT and *erg25*⁻ by electron ionization (EI) gas chromatography-mass spectrometry (GC-MS) [6,18]. We also examined the sterols in *erg25*⁻/*ERG25* and *erg25*⁻/*GFP-ERG25* parasites which overexpress *ERG25* or *GFP-ERG25* proteins from high copy number plasmids [19]. As previously reported, the four major sterol species in *L. major* WT promastigotes consisted of 5-dehydroepisterol, ergosterol, cholesta-5,7,24-trienol and cholesterol, identified based on molecular weights, retention times and EI spectra (Figure 5A) [6]. Similar sterol compositions (27-31% ergosterol, 42-51% 5-dehydroepisterol, 5-7% cholesterol and 15-24% cholesta-5,7,24-trienol) were detected in *erg25*⁻, *erg25*⁻/*ERG25* and *erg25*⁻/*GFP-ERG25* parasites (Figure 5B-E). Furthermore, the cellular level of major sterols in *erg25*⁻ were similar to WT and add-back parasites based on comparison of endogenous sterol peaks to that of the internal standard cholesta-3,5-diene (retention time: 8.78) (Figure 5A-D). Overall, these results indicate that *ERG25* is not required for sterol synthesis in *L. major*.




Figure 5. *ERG25* deletion does not affect sterol composition in *L. major*. (A-D) Partial GC chromatograms of free sterols from WT (A), *erg25*⁻ (B), *erg25*⁻/*ERG25* (C) and *erg25*⁻/*GFP-ERG25* (D) promastigotes. The peaks represented ergosterol (1), 5-dehydroepisterol (2), cholesterol (3), cholesta-5,7,24-trienol (4) and cholesta-3,5-diene standard (std). Sterol compositions were summarized in E.

2.522. *A1* is Indispensable during the Promastigote Stage of *L. major*

To assess the essentiality of 22A1 in culture promastigotes, we tested whether the pXNG4-22A1 episome can be removed from the 22A1⁻ +pXNG4-22A1 parasites. The presence of a thymidine kinase gene on pXNG4-22A1 made these parasites susceptible to negative selection by ganciclovir (GCV). With GCV treatment, parasites tend to lose pXNG4-22A1 if the plasmid is not essential [13,20]. When

22A1^{+/-} +pXNG4-22A1 and 22A1⁻ +pXNG4-22A1 parasites were cultivated in the presence of nourseothricin (SAT), 80-85% of them had high GFP expression indicating a high pXNG4-22A1 level (Figure 6A and supplemental Figure S6). Without SAT, 22A1^{+/-} +pXNG4-22A1 promastigotes (which had one chromosomal 22A1 allele) gradually lost the plasmid as the GFP-high cells decreased from ~80% to ~10% in six passages (Figure 6A). With GCV treatment, the plasmid loss was accelerated and near complete in 5-6 passages due to the negative selection (Figure 6A). Therefore, in the presence of endogenous 22A1, the pXNG4-22A1 episome is dispensable.




Figure 6. 22A1 is essential for *L. major* promastigotes in culture. (A) Promastigotes were continuously cultivated in the presence or absence of GCV or nourseothricin (SAT) and percentages of GFP-high cells were determined by flow cytometry for each passage. Error bars represent standard deviations from three replicates. (B–D) After 6 passages, single clones were isolated from 22A1⁻+pXNG4-22A1 cells grown in the presence of GCV by serial dilution and allowed to proliferate. Flow cytometry analyses were performed on the original 22A1⁻+pXNG4-22A1 GCV population (B) and two representative clones (C and D). In B–D, percentages of GFP-high cells were indicated as R3. (E) Promastigotes were inoculated at 1.0×10^5 cells/ml in the presence or absence of GCV or SAT and culture densities were determined daily (***: $p < 0.001$).

Notably, the GFP-high cells in 22A1⁻ +pXNG4-22A1 (which are chromosomal null for 22A1) sustained at 79-83% in the absence of GCV and 59-64% in the presence of GCV after six passages

(Figure 6A), suggesting that these promastigotes could not lose the episome even with negative selection. When single cells (clones) were isolated from the *22A1*⁻+pXNG4-*22A1* GCV population and allowed to proliferate, they showed a mixture of GFP-high (51-65%) cells and GFP-low (35-49%) cells similar to original population, suggesting that the pXNG4-*22A1* plasmid cannot be eliminated (Figure 6B-D). Together, these findings indicate that *22A1* is required for the survival and replication of *L. major* promastigotes.

Additionally, we compared the growth rates of *22A1*^{+/-}+pXNG4-*22A1* and *22A1*⁻+pXNG4-*22A1* with WT promastigotes (Figure 6E). When cultivated in the absence of GCV, these parasites proliferated at very similar rates. However, in the presence of GCV, *22A1*⁻+pXNG4-*22A1* showed a significant growth delay in comparison to WT and *22A1*^{+/-}+pXNG4-*22A1* cells at day 1-2 post inoculation (Figure 6E, two clones of *22A1*⁻+pXNG4-*22A1* were tested). It is possible that the GCV-induced cytotoxicity affected *22A1*⁻+pXNG4-*22A1* more due their inability to lose the plasmid.

2.622. *A1* is Essential for *L. major* during the Amastigote Stage

To determine if *22A1* is required during the amastigote stage, stationary phase promastigotes were injected into the footpad of BALB/c mice. Half of the mice were treated with GCV daily for 14 days, while the other half received an equal volume of sterile PBS as controls. As indicated in Figure 7A-B, mice infected by WT and *22A1*^{+/-} showed similar lesion progression and had equivalent parasite numbers. Thus, losing one chromosomal *22A1* allele does not compromise *L. major* virulence or growth in mice. Also, GCV treatment did not affect footpad lesion development or parasite loads in WT- and *22A1*^{+/-}-infected mice




Figure 7. 22A1 is indispensable for *L. major* amastigotes. BALB/c mice were infected in the footpads with stationary phase promastigotes and treated with either GCV or PBS daily for 14 days. (A) Footpad lesions were recorded weekly. (B) Average parasite numbers per infected footpad were determined at the indicated times post infection by qPCR. (C) Average pXNG4-22A1 copy numbers in amastigotes were determined at the indicated times post infection by qPCR. ND: not determined. Error bars represent standard deviations from 4-5 mice (A) or 2-3 mice (B and C). **: $p < 0.01$.

Mice infected by 22A1^{+/-} +pXNG4-22A1 or 22A1⁻ +pXNG4-22A1 and received PBS treatment also showed similar lesion progression and parasite numbers as WT-infected mice (Figure 7A-B). In contrast, GCV treatment significantly delayed lesion development and amastigote replication in these mice. This observation is in line with the effect of GCV on those parasites in culture (Figure 6E). The reduced growth of 22A1^{+/-} +pXNG4-22A1 + GCV (four weeks delay in comparison to WT) may be due to the cytotoxic effect of GCV during the initial phase post infection when parasites contained a high level of pXNG4-22A1 (supplemental Figure S6). The delay was much more pronounced with the 22A1⁻ +pXNG4-22A1 + GCV group (nine weeks delay), which could arise if these parasites were unable to lose the plasmid.

To examine the retention of pXNG4-22A1, we performed qPCR analyses on the genomic DNA extracted from lesion derived amastigotes. As shown in Figure 7C, the plasmid copy number was 0.45-0.75 per cell for 22A1^{+/-} +pXNG4-22A1 amastigotes with PBS treatment at 5-7 weeks post infection. With GCV treatment, less than 0.1 copy per cell was retained for 22A1^{+/-} +pXNG4-22A1 at

7-11 weeks post infection. Meanwhile, the 22A1⁻+pXNG4-22A1 amastigotes contained 1.3-2.2 copies per cell with PBS treatment and 0.8-1.8 copies per cell with GCV treatment even after 16 weeks post infection. Based on these findings, we conclude that 22A1 is essential for the survival of *L. major* in the amastigotes stage.

2.722. A1 Overexpression Confers Resistance to Posaconazole and DB766 but Does Not Affect Sterol Composition or Stress Response in *L. major*

In *L. donovani*, CYP5122A1 overexpression (from pXNG4-LdCYP5122A1) led to increased tolerance to stress conditions such as high temperature, acidic pH, and starvation [13]. To test if the 22A1 had a similar impact in *L. major*, stationary phase promastigotes of 22A1^{+/-}+pXNG4-22A1 and 22A1⁻+pXNG4-22A1 were exposed to 37 °C or pH5.0 conditions and their tolerance was measured as previously described [6,13]. As summarized in supplemental Figure S7, 22A1 overexpressors showed similar level of survival as WT and 22A1^{+/-} parasites. We also examined whether 22A1 half knockout and overexpression alter the sterol composition in *L. major* by liquid chromatograph tandem mass spectrometry (LC-MS/MS). While the major sterol species were very similar between WT and 22A1 mutants (supplemental Figure S8), we detected a slight accumulation of squalene, a sterol biosynthesis intermediate, in 22A1^{+/-}+pXNG4-22A1 parasites (supplemental Figure S9). The reason for this minor change is not clear. Compared to CPY5122A1 mutants in *L. donovani* which showed altered levels of 4,14-methylated sterol intermediates [13], 22A1 overexpression and half knockout in *L. major* had little influence on sterol composition in *L. major*.

Sterol synthesis inhibitors such as ketoconazole, posaconazole, and DB766 target C14DM and/or C4DM [6,13,21]. To test if 22A1 half knockout and overexpressors have altered sensitivity to these inhibitors, we cultivated promastigotes in various concentrations of ketoconazole (0-15 μM), posaconazole (0-6 μM) or DB766 (0-32 nM) and measured culture densities after 48 hours. The effective concentrations to inhibit 25% and 50% of growth (referred to as EC25 and EC50, respectively) were calculated using cells grown in the absence of drugs as controls (Table 1). As illustrated in Figure 8A, these parasites showed similar levels of susceptibility to ketoconazole except that at higher concentrations (10-15 μM), WT and 22A1^{+/-} parasites were slightly more sensitive. For posaconazole and DB766, 22A1^{+/-}+pXNG4-22A1 and 22A1⁻+pXNG4-22A1 were more resistant than WT and 22A1^{+/-} parasites as their EC50s were 1.5-3.0-fold higher (Figure 8B-C and Table 1). These findings suggest that 22A1 overexpression leads to modest resistance to C4DM inhibitors.




Figure 8. Effects of 22A1 overexpression on promastigotes' sensitivity to sterol synthesis inhibitors.

Log phase promastigotes were inoculated in various concentrations of ketoconazole (A), posaconazole (B) or DB766 (C). Cell densities were determined after 48 hours, and percentages of growth were calculated using cells grown in the absence of inhibitors as controls. Error bars represent standard deviations from three experiments.

Table 1. EC25 and EC50 values for posaconazole and DB766.

Inhibitor	Effective concentration (EC)	WT	22A1+/-	22A1+/- +pXNG4-22A1	22A1 ⁻ +pXNG4-22A1
Posaconazole	EC25 ($\mu\text{M} \pm \text{SD}$)	0.29 \pm 0.089	0.21 \pm 0.041	0.48 \pm 0.15	0.47 \pm 0.14
	EC50 ($\mu\text{M} \pm \text{SD}$)	1.0 \pm 0.096	0.74 \pm 0.18	3.0 \pm 1.0*	1.7 \pm 0.38*
DB766	EC25 (nM \pm SD)	0.90 \pm 0.12	0.96 \pm 0.16	1.6 \pm 0.33	1.9 \pm 0.094
	EC50 (nM \pm SD)	1.9 \pm 0.31	2.04 \pm 0.41	3.8 \pm 0.68**	3.5 \pm 0.19**

Student's *t* test was performed to analyze the difference between WT and 22A1 overexpressing cells (*: $p < 0.05$; **: $p < 0.01$).

2.8. Genetic Manipulation of 22A1 or ERG25 Do Not Affect the Expression Level of C14DM

To determine if 22A1 half knockout or overexpression affects the expression of C14DM (CYP51), western blots were conducted using antibodies against *L. donovani* CYP51^{22A1} and CYP51 (Figure 9). Clearly, genetic manipulation of 22A1 had little impact on C14DM (CYP51) expression at the protein level. Conversely, C14DM-null and add-back parasites contained similar levels of 22A1 as WT. As expected, 22A1 half knockout and overexpressors showed reduced and elevated levels of 22A1, respectively. Furthermore, deletion and add-back of ERG25 did not affect the levels of 22A1 or C14DM (CYP51) (Figure 9). These findings indicate that expressions of 22A1, C14DM and ERG 25 are not linked.




Figure 9. Expression levels of 22A1 and C14DM in *L. major* promastigotes. (A) Whole cell lysates from log phase promastigotes were denatured at 65 °C and analyzed by western blot using antibodies against C14DM (top), 22A1 (middle), or α -tubulin (bottom). (B) The relative abundances of 22A1 and C14DM were determined after three experiments using α -tubulin as the loading control. Error bars represent standard deviations. *** $p < 0.001$.

3. Discussion

Sterol biosynthesis requires the formation of lanosterol followed by the removal of three methyl groups: two from the C4 position and one from the C14 position (Figure 1). Compared to the C14DM/CYP51 which is well characterized biochemically and genetically as the main target for azole drugs, the C4DM is less studied. In *L. donovani* (the etiological agent for visceral leishmaniasis), we recently identified CYP5122A1 as a bona fide sterol C4-methyl oxidase capable of converting C4-methylated sterols into mono-hydroxy, aldehyde and carboxylate intermediates in vitro [13]. Chromosomal CYP5122A1-null mutants cannot survive without a complementing CYP5122A1-episome, indicating that it is required for *L. donovani* promastigotes and amastigotes. In this report, we investigated whether the essentiality of CYP5122A1 can be extended to *L. major* which causes cutaneous leishmaniasis. Previous studies from us and others have revealed differences in the requirement of sterol and sphingolipid metabolism among *Leishmania* species [17,31–33]. For example, while C14DM/CYP51 can be removed from *L. major*, it is indispensable in *L. donovani* [17,32]. In addition, the SL-degrading enzyme ISCL is required for the virulence of *L. major* but not *L. amazonensis* [31,33]. Thus, it is important to determine if different *Leishmania* species have distinct requirements for C4DM.

Using the same plasmid segregation method as described for *L. donovani* CYP5122A1 [13,20], we showed that *L. major* 22A1⁺+pXNG4-22A1 promastigotes retained a high level of episome even when

facing strong negative selection from GCV treatment, whereas 22A1+/- +pXNG4-22A1 lost the episome after 6 passages. When single clones isolated from 22A1- +pXNG4-22A1 GCV treatment group were allowed to proliferate, they showed a mix of GFP-low and GFP-high cells, suggesting that GFP-low cells (lacking pXNG4-22A1) could not perpetuate by themselves. Thus, like in *L. donovani*, 22A1 is essential for *L. major* promastigotes. Given the fact that *L. major* C14DM/CYP51-null mutants are viable [6], these findings suggest that removal of the two C4 methyl groups is more crucial than the removal of C14-methyl group. It is possible that the accumulation of C4-methyl intermediates is more harmful to *Leishmania* membrane than the accumulation of C14-methyl intermediates [22].

In the intracellular mammalian stage, *L. major* amastigotes are less dependent on *de novo* lipid synthesis and can salvage and remodel host lipids to support their activity [6,23–25]. To determine whether 22A1 is dispensable for amastigotes, BALB/c mice infected with 22A1- +pXNG4-22A1 were treated with GCV or PBS for 14 days. As illustrated in Figure 7, GCV treatment dramatically reduced the virulence and growth of 22A1- +pXNG4-22A1 in mice, yet these parasites retained the pXNG4-22A1 plasmid at 16 weeks post infection. These results are consistent with the notion that 22A1 is required during the amastigote stage. For mice infected by 22A1+/- +pXNG4-22A1, GCV treatment caused an initial delay in footpad lesion formation which might arise from the elevated level of pXNG4-22A1 in the early stage of infection. After 7 weeks, these parasites lost the episome and displayed normal levels of virulence and growth in mice, indicating that the plasmid is not essential for these parasites which contain one of chromosomal 22A1. These findings are largely consistent with the data from *L. donovani* CYP5122A1. However, one distinction is that *L. major* 22A1+/- (half knockout) shows WT-level of infectivity while *L. donovani* half knockout has attenuated virulence [13,14]. Also, *L. major* 22A1-overexpression and half knockout did not affect stress response or bulk sterol composition. These results suggest that *L. major* parasites can tolerate the impact of 22A1 over- or under-expression better than *L. donovani*. We did detect a slight accumulation of squalene in 22A1-overexpressing cells, suggesting that its overproduction can affect the early steps of sterol synthesis.

Overall, our results demonstrate that 22A1 is essential for *L. major* amastigotes just like it is for *L. donovani* amastigotes. This is notable because most sterols in intracellular amastigotes consist of host-derived cholesterol and only a small portion was synthesized *de novo* [6,26]. These findings mark the potential of 22A1 as a pan-*Leishmania* drug target. In culture promastigotes, the overexpression of 22A1 led to 2-3-fold resistance to several 22A1 and CYP51 inhibitors including DB766 and posaconazole (Figure 8). Future studies will aim to identify more potent and selective inhibitors of 22A1.

Despite these findings, it is not clear whether CYP5122A1 is the sole sterol C4 methyl oxidase in *Leishmania*. In *S. cerevisiae* and animals, the two C-4 methyl groups are removed consecutively, and a single sterol C4 methyl oxidase catalyzes the initial oxygenation reactions [8–10,27]. The *S. cerevisiae* ERG25-null mutants are ergosterol auxotroph and accumulate C4,4-dimethylzymosterol [8]. Unlike *S. cerevisiae*, *Aspergillus fumigatus* has the two functional sterol C4 methyl oxidases (ERG25A and ERG25B) with ERG25A being the dominant enzyme. ERG25A mutants are viable but show susceptibility to hypoxia and ER stress [28]. *Candida albicans* also has two sterol C4 methyl oxidase genes and both appear to be essential suggesting that they work in tandem for efficient activity [29]. Similarly, plant cells possess two sterol C4 methyl oxidases each removing one C4-methyl group in a subsequent, but biochemically distinct mechanism [11,30,31]. In *Arabidopsis thaliana*, removal of each of the two C4-methyl groups requires a different sterol C4 methyl oxidase and mutants of the individual enzymes primarily accumulate their respective sterol intermediates [28]. Finally, the methanotrophic gram-negative bacterium *Methylococcus capsulatus* contains two C-4 demethylases that are distinct from the eukaryotic enzymes [32]. Overall, these studies demonstrate diverse mechanisms involved in the C-4 demethylation step in different organisms.

In addition to 22A1, there are other genes in *Leishmania* parasites encoding proteins homologous to *S. cerevisiae* sterol C4 methyl oxidase: the previously characterized lathosterol oxidase (LSO, LmjF.23.1300) and a putative ERG25 (LmjF.36.2540). These orthologs show 29% and 26% identity to *S. cerevisiae* sterol C4 methyl oxidase respectively. *L. major* LSO-null mutants synthesize sterols

without the C-5-C-6 double bond but show no defects in C4-demethylation [33]. In this study, we characterized the putative ERG25 in *L. major*. Like other sterol biosynthetic enzymes, ERG25 is largely localized in the ER. *L. major* ERG25-null mutants were fully viable with growth rate and virulence similar to WT parasites (Figure 4). These mutants synthesize ergosterol and 5-dehydroepisterol as their main sterols and their overall sterol composition is very similar to WT and add-back parasites (Figure 5). These observations suggest that ERG25 is not involved in the oxidative removal of C4-methyl groups. Unlike other sterol biosynthetic enzymes, the deletion or overexpression of ERG25 had no impact on the cellular levels or localization of LPG and GP63 [6,18,33].

While the precise function of ERG25 in *Leishmania* awaits further investigation, it is important to note that ERG25 is not well conserved among trypanosomatids. According to Tritypdb, ERG25 orthologs are present in *Leishmania* spp., *Angomonas deanei*, *Bodo saltans*, *Crithidia fasciculata*, *Endotrypanum monterogeii*, *Leptomonas* spp., *Paratrypanosoma confusum*, and *Porcisia hertigi*, but absent in *Trypanosoma* spp. In contrast, CYP5122A1 orthologs are identified in all of the above organisms plus *Trypanosoma* spp. including *T. brucei* and *T. cruzi* [34]. Given the dispensable nature of ERG25 in *L. major* sterol synthesis, it is possible that ERG25 has evolved to diverge from the sterol C4 methyl oxidase function and was lost in some trypanosomatid species. Meanwhile, CYP5122A1 was conserved by most trypanosomatids as the only or dominant sterol C4 methyl oxidase.

4. Materials and Methods

4.1. Materials

M199 media and cholesta-3,5-diene were purchased from Sigma-Aldrich (St. Louis, MO). Ketoconazole and posaconazole were purchased from TCI (Portland, OR) and Carbosynth (San Diego, CA), respectively. DB766 was synthesized as previously described [21]. Rat anti-LdCYP5122A1 and anti-LdC14DM (CYP51) polyclonal antisera were generated at the University of Kansas Medical Center by Dr. Jianming Qiu using purified recombinant proteins as antigens. All other chemicals were purchased from Thermo Fisher Scientific unless otherwise specified.

4.2. Molecular Constructs

The predicted open reading frames (ORFs) of *L. major* ERG25 (LmjF.36.2540) and CYP5122A1 (LmjF.27.0090) were amplified from *L. major* LV39 (Rho/SU/59/P) genomic DNA by PCR and digested with restriction enzymes. The ERG25 ORF was cloned into a *Leishmania* expression vector pXG to generate pXG-ERG25 or a GFP-tagged expression vector pXG-GFP2+' to generate pXG-GFP-ERG25 for localization studies. The CYP5122A1 ORF (referred to as 22A1) was cloned into the pXNG4 vector or pXG-'GFP+' vector to generate pXNG4-22A1 or pXG-22A1-GFP, respectively. All molecular constructs were verified by restriction enzyme digestion and Sanger sequencing.

To generate knockout constructs, ~600 bp of the 5'- and 3'-flanking sequences of ERG25 ORF were PCR-amplified and cloned into the pUC18 vector, followed by the insertion of drug resistance genes for puromycin (PAC) or hygromycin B (HYG) to generate pUC18-KO-ERG25::PAC or pUC18-KO-ERG25::HYG, respectively. Similar knockout constructs for 22A1 were generated which include ~1000 bp of the 5'- and 3'-flanking sequences and drug resistance genes for blasticidin (BSD) or HYG. Oligonucleotides used in this study are summarized in supplemental Table S1.

4.3. *Leishmania* Promastigote Culture and Growth Rate Measurement

L. major strain LV39 clone 5 promastigotes were cultivated in M199 media (pH7.4) with 10% fetal bovine serum and other supplements at 27 °C as previously described [35]. To determine growth rates, log phase parasites ($1.0-10.0 \times 10^6$ cells/ml) were inoculated at $1.0-2.0 \times 10^5$ cells/ml and cell densities were determined daily using a hemocytometer. During the stationary phase ($2.0-4.0 \times 10^7$ cells/ml), percentage of round cells whose long axis was less than twice of the short axis was measured by microscopy and percentage of dead cells was assessed by flow cytometry after propidium iodide staining as previously described [36].

To measure the antileishmanial activity of sterol synthesis inhibitors (prepared as 10 mM stocks in DMSO), log phase promastigotes were inoculated in complete M199 media at 2.0×10^5 cells/ml in 24-well plates (1 ml/well). Inhibitors were added to various concentrations and control wells contained DMSO only (0.1–0.5%). Culture densities were determined after 48 h using a Beckman Z2 Cell Counter.

4.4. *Leishmania* Genetic Manipulation and Southern Blot

To delete the endogenous *ERG25* alleles, pUC18-KO-*ERG25::PAC* and pUC18-KO-*ERG25::HYG* were digested with EcoRI and HindIII to generate linear DNA fragments which were sequentially introduced into *L. major* wild type (WT) promastigotes by electroporation. The resulting *ERG25*-null mutant *erg25⁻* (Δ *ERG25::PAC*/ Δ *ERG25::HYG*) was selected based on resistance to puromycin and hygromycin. The replacement of *ERG25* alleles with *HYG* and *PAC* in *erg25⁻* was confirmed by Southern blot using ³²P-labeled probes for the ORF and an upstream flanking region of *ERG25* [37]. The add-back line *erg25⁻/+Erg25* was generated by introducing pXG-*ERG25* into *erg25⁻*. A similar approach was used to produce the heterozygous *22A1* mutant *22A1+/-* (Δ *22A1::BSD*/*22A1*). To acquire the chromosomal *22A1*-null mutant, *22A1+/-* parasites were transfected with pXNG4-*22A1* and then the second chromosomal *22A1* allele was replaced with *HYG*. The resulting chromosomal *22A1*-null mutant was confirmed by Southern blot and referred to as *22A1⁻/+pXNG4-22A1*.

4.5. Sterol Analysis by Gas Chromatography and Mass Spectrometry (GC-MS) and Liquid Chromatograph Tandem Mass Spectrometry (LC-MS/MS)

For GC-MS analysis, total lipids were extracted from log phase promastigotes using the Folch method [38]. An internal standard, cholesta-3,5-diene (FW = 368.84), was added to cell lysate at 2.0×10^7 molecules/cell during lipid extraction. Lipid samples were dissolved in methanol at 1.0×10^9 cells/ml. Electron ionization GC-MS analyses of sterols and related lipids were performed as previously described [6]. For LC-MS/MS analysis, free sterols were isolated using a modified hexane/isopropanol extraction method and analyzed by high performance liquid chromatography tandem mass spectrometry (HPLC-MS/MS) using a triple quadrupole mass spectrometer as described previously [39].

4.6. Immunofluorescence Microscopy and Western Blot

As previously described [18], log phase promastigotes were loaded onto poly-lysine coated coverslips by centrifugation and fixed with 3.7% paraformaldehyde. Cells were then permeabilized with ice-cold ethanol followed by rehydration with phosphate buffered saline (PBS). To stain the endoplasmic reticulum (ER), cells were labeled with a rabbit anti-*Trypanosoma brucei* BiP antiserum (1:2000) [16] followed by goat anti-rabbit IgG-Texas Red antibody (1:2000). To label LPG or GP63, cells were stained with mouse anti-LPG antibody WIC79.3 (1:2000) [40] or mouse anti-GP63 antibody 96/26 (1:2000), followed by goat anti-mouse IgG-FITC (1:1000). Hoechst 33342 (2.5 μ g/ml) was used to stain DNA. Fluorescence microscopy was performed using a BX51 fluorescence microscope or FV3000 laser scanning confocal microscope.

For western blots, whole cell lysates were resolved by SDS-PAGE and transferred to polyvinylidene difluoride membranes. Blots were probed with rabbit anti-GFP (Abcam ab6556, 1:1000), rat anti-LdCYP5122A1 (1:500), rat anti-LdCYP51 (1:500), or mouse anti-tubulin antisera (1:1000), followed by appropriate HRP-conjugated secondary antibodies (1:2000). Signals were detected using a Fuji phosphorimager.

4.7. GCV Treatment of Promastigotes

22A1+/- +pXNG4-22A1 and *22A1⁻/+pXNG4-22A1* promastigotes were cultivated in the presence or absence of ganciclovir (GCV, 50 μ g/ml) or nourseothricin (SAT, 150 μ g/ml) and re-inoculated into fresh media containing the same negative or positive selection agents at 1.0×10^5 cells/ml every 3 days [13]. The retention of pXNG4-*22A1* plasmid was assessed by measuring the level of GFP fluorescence

on the third day of each passage using an Attune NxT Acoustic Flow Cytometer. After six passages, individual clones of 22A1-/+ pXNG4-22A1 were isolated by serial dilution in 96-well plates and expanded in the presence of GCV. The GFP levels of selected clones were determined by flow cytometry.

4.8. Mouse Footpad Infections

To assess parasite virulence, 1.0×10^6 stationary phase promastigotes (day 3) were resuspended in 50 μ l of DMEM and injected into the footpad of 8 weeks old female BALB/c mouse (5 mice per group). The development of footpad lesions was measured weekly using a Vernier Caliper by comparing the infected footpad with the uninfected one. Parasite numbers in infected footpad were estimated by the limiting-dilution assay [41]. During the footpad injection and measurement, an isoflurane chamber was used to anesthetize mice. Mice euthanasia was achieved through a controlled flow of CO₂ asphyxiation.

To determine whether 22A1 is required during the amastigote stage, one-half of the infected mice received GCV at 7.5 mg/kg/day for 14 consecutive days (0.5 ml each, intraperitoneal injection) while the other half received an equivalent volume of sterile PBS [42]. Mice were euthanized at the indicated time points and genomic DNA was extracted from footpads for quantitative PCR (qPCR) analyses described as follows.

4.9. QPCR Analyses

To determine parasite numbers in infected footpads, qPCR reactions were run in triplicates using primers targeting the 28S rRNA gene of *L. major* [42]. Cycle threshold (Ct) values were determined from melt curve analysis. A standard curve of Ct values was generated using serially diluted genomic DNA samples from *L. major* promastigotes (from 0.1 cell/reaction to 10⁵ cells/reaction) and Ct values >30 were considered negative. Parasite numbers in infected footpads were calculated using the standard curve. Control reactions included sterile water and DNA extracted from uninfected mouse footpads.

To determine pXNG4-22A1 plasmid copy numbers in promastigotes and amastigotes, a similar standard curve was generated using serially diluted pXNG4-22A1 plasmid DNA (from 0.1 copy/reaction to 10⁵ copies/reaction) and primers targeting the *GFP* gene. QPCR was performed using the same set of primers on DNA samples from promastigotes or infected footpads and the average plasmid copy number per cell was determined by dividing the total plasmid copy number by the total parasite number based on Ct values [42].

4.10. Statistical Analyses

All experiments were repeated at least 3 times and error bars represent standard deviations. Graphs were made using Sigmaplot 11.0 (Systat Software Inc, San Jose, CA). Differences between experimental groups were evaluated using the Student's *t* test (for two groups) or one-way ANOVA test (for three or more groups). *P* values indicating statistical significance were grouped into values of <0.05, <0.01 and <0.001.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Figure S1. Pair-wise alignment of *L. major* CYP5122A (LmjF.27.0090) and *L. donovani* CYP5122A1 (LdBPK_270090.1) using EMBOSS Needle. Identity: 94.4%. Similarity: 95.9%. Figure S2. The alignment of ERG25 proteins (Lm: *L. major*; At: *Arabidopsis thaliana*; Sc: *Saccharomyces cerevisiae*; Mm: *Mus musculus*) using Clustal Omega. Asterisks (*): fully conserved residues; colons (:): highly similar residues; periods (.): weakly similar residues. *L. major* ERG25 shows 23-25% identity to other ERG25s. Figure S3. Schematic depiction of the Southern blot for *L. major* ERG 25. (A) The genomic locus of ERG25 in WT *L. major* along with its open reading frame (ORF), 5' and 3' flanking regions (FR), recognition sites of Age I, BseE II and Sca I, and the positions of ORF probe and upstream probe. (B-C) Replacement of ERG25 ORF with hygromycin (HYG) and puromycin (PAC) resistance genes. The expected sizes of DNA fragments recognized by the ORF probe or upstream probe were indicated. Figure S4. Schematic depiction of the Southern blot for *L. major* CYP5122A1 (22A1). (A) The genomic locus of CYP5122A1 in WT *L. major* along with its open reading frame

(ORF), 5' and 3' flanking regions (FR), recognition sites of Sac I and Spe I, and the positions of ORF probe and upstream probe. (B-C) Replacement of 22A1 ORF with blasticidin (BSD) and puromycin (PAC) resistance genes. (D) Episomal expression of 22A1 from pXNG4-5122A1. TK: thymidine kinase, GFP: green fluorescent protein, SAT: nourseothricin resistance gene. The expected sizes of DNA fragments recognized by the ORF probe or upstream probe were indicated. Figure S5. The expression of LPG and GP63 is not affected in *erg25*⁻ promastigotes. Log phase promastigotes were labeled with mouse anti-LPG (A) or anti-GP63 (B) monoclonal antibody followed by goat anti-mouse IgG-FITC. Scale bars: 10 μ m. Figure S6. Plasmid levels in promastigotes. Stationary phase promastigotes of 22A1^{+/-} +pXNG4-22A1 and 22A1⁻ +pXNG4-22A1 were cultivated in the presence of SAT and their DNA was extracted to determine the average pXNG4-Lm22A1 plasmid copy numbers per cell by qPCR. Figure S7. Stress response of 22A1 mutants. Day 1 stationary phase promastigotes were incubated under various conditions and percentages of dead cells were determined by flow cytometry at the indicated times. A: in complete M199, pH7.4, 37 °C. B: in complete M199, 27 °C, pH5.0. Error bars represent standard deviations from three experiments. Figure S8. 22A1 half knockout and overexpression do not affect bulk sterol composition in *L. major*. Partial LC-MS chromatograms of free sterols from WT, 22A1^{+/-} #4, 22A1^{+/-} #4 +pXNG4-22A1, 22A1^{+/-} #5, and 22A1^{+/-} #5 +pXNG4-22A1 promastigotes. Figure S9. Squalene was detected in 22A1 overexpressing cells. (A) Partial GC chromatograms of total lipids from WT, 22A1^{+/-} #4, 22A1^{+/-} #4 +pXNG4-22A1, 22A1^{+/-} #5, and 22A1^{+/-} #5 +pXNG4-22A1 promastigotes showed the presence of squalene in the boxed region. (B) The relative abundance of squalene was determined. Table S1. List of oligonucleotides used in this study

Author Contributions: Conceptualization, Michael Wang and Kai Zhang; Data curation, Yu Ning, Somrita Basu and Kai Zhang; Formal analysis, Fong-fu Hsu and Mei Feng; Funding acquisition, Fong-fu Hsu, Michael Wang and Kai Zhang; Methodology, Yu Ning, Somrita Basu, Fong-fu Hsu and Mei Feng; Project administration, Michael Wang; Writing – original draft, Yu Ning and Kai Zhang; Writing – review & editing, Yu Ning, Somrita Basu, Fong-fu Hsu, Michael Wang and Kai Zhang. All authors have read and agreed to the published version of the manuscript.

Funding: Please add: This research was funded by the National Institutes of Health grants R01AI139198 (MZW), P20GM113117 (MZW), P30GM110761 (MZW), R15AI156746 (KZ), P41-GM103422 (FH), P60-DK20579 (FH) and P30-DK56341 (FH). Opinions, interpretations, conclusions, and recommendations are those of the authors and do not necessarily represent the official views of the National Institutes of Health.

Institutional Review Board Statement: Not applicable

Animal use: The animal study protocol was approved by Texas Tech University Institutional Animal Care and Use Committee (#22008-03, approval date 10/27/2023).

Informed Consent Statement: Not applicable

Data Availability Statement: We encourage all authors of articles published in MDPI journals to share their research data. In this section, please provide details regarding where data supporting reported results can be found, including links to publicly archived datasets analyzed or generated during the study. Where no new data were created, or where data is unavailable due to privacy or ethical restrictions, a statement is still required. Suggested Data Availability Statements are available in section “MDPI Research Data Policies” at <https://www.mdpi.com/ethics>.

Acknowledgments: The original contributions presented in this study are included in this article and Supplementary Material. Further inquiries can be directed to the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Pace, D. Leishmaniasis. *The Journal of infection* 2014, 69 Suppl 1, S10-18, doi:10.1016/j.jinf.2014.07.016.
2. Croft, S.L.; Olliaro, P. Leishmaniasis chemotherapy--challenges and opportunities. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* 2011, 17, 1478-1483, doi:10.1111/j.1469-0691.2011.03630.x.
3. Beach, D.H.; Goad, L.J.; Holz, G.G., Jr. Effects of antimycotic azoles on growth and sterol biosynthesis of Leishmania promastigotes. *Molecular and biochemical parasitology* 1988, 31, 149-162.
4. Lepesheva, G.I.; Waterman, M.R. Sterol 14 α -demethylase (CYP51) as a therapeutic target for human trypanosomiasis and leishmaniasis. *Curr Top Med Chem* 2011, 11, 2060-2071, doi:10.2174/156802611796575902.
5. Lepesheva, G.I.; Waterman, M.R. Structural basis for conservation in the CYP51 family. *Biochimica et biophysica acta* 2011, 1814, 88-93, doi:10.1016/j.bbapap.2010.06.006.

6. Xu, W.; Hsu, F.F.; Baykal, E.; Huang, J.; Zhang, K. Sterol Biosynthesis Is Required for Heat Resistance but Not Extracellular Survival in *Leishmania*. *PLoS pathogens* 2014, 10, e1004427, doi:10.1371/journal.ppat.1004427.
7. Mukherjee, S.; Moitra, S.; Xu, W.; Hernandez, V.; Zhang, K. Sterol 14-alpha-demethylase is vital for mitochondrial functions and stress tolerance in *Leishmania major*. *PLoS pathogens* 2020, 16, e1008810, doi:10.1371/journal.ppat.1008810.
8. Bard, M.; Bruner, D.A.; Pierson, C.A.; Lees, N.D.; Biermann, B.; Frye, L.; Koegel, C.; Barbuch, R. Cloning and characterization of ERG25, the *Saccharomyces cerevisiae* gene encoding C-4 sterol methyl oxidase. *Proceedings of the National Academy of Sciences of the United States of America* 1996, 93, 186-190, doi:10.1073/pnas.93.1.186.
9. Gachotte, D.; Barbuch, R.; Gaylor, J.; Nickel, E.; Bard, M. Characterization of the *Saccharomyces cerevisiae* ERG26 gene encoding the C-3 sterol dehydrogenase (C-4 decarboxylase) involved in sterol biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* 1998, 95, 13794-13799, doi:10.1073/pnas.95.23.13794.
10. Gachotte, D.; Sen, S.E.; Eckstein, J.; Barbuch, R.; Krieger, M.; Ray, B.D.; Bard, M. Characterization of the *Saccharomyces cerevisiae* ERG27 gene encoding the 3-keto reductase involved in C-4 sterol demethylation. *Proceedings of the National Academy of Sciences of the United States of America* 1999, 96, 12655-12660, doi:10.1073/pnas.96.22.12655.
11. Rahier, A. Dissecting the sterol C-4 demethylation process in higher plants. From structures and genes to catalytic mechanism. *Steroids* 2011, 76, 340-352, doi:10.1016/j.steroids.2010.11.011.
12. Pascal, S.; Taton, M.; Rahier, A. Plant sterol biosynthesis. Identification and characterization of two distinct microsomal oxidative enzymatic systems involved in sterol C4-demethylation. *The Journal of biological chemistry* 1993, 268, 11639-11654.
13. Jin, Y.; Basu, S.; Feng, M.; Ning, Y.; Munasinghe, I.; Joachim, A.M.; Li, J.; Madden, R.; Burks, H.; Gao, P.; et al. CYP5122A1 encodes an essential sterol C4-methyl oxidase in *Leishmania donovani* and determines the antileishmanial activity of antifungal azoles. *Res Sq* 2023, doi:10.21203/rs.3.rs-3185204/v1.
14. Verma, S.; Mehta, A.; Shaha, C. CYP5122A1, a novel cytochrome P450 is essential for survival of *Leishmania donovani*. *PloS one* 2011, 6, e25273, doi:10.1371/journal.pone.0025273.
15. La Rosa, C., et al. N-substituted-4-(pyridin-4-ylalkyl)piperazine-1-carboxamides and Related Compounds as *Leishmania* CYP51 and CYP5122A1 Inhibitors *Bioorganic & medicinal chemistry* 2024.
16. Bangs, J.D.; Uyetake, L.; Brickman, M.J.; Balber, A.E.; Boothroyd, J.C. Molecular cloning and cellular localization of a BiP homologue in *Trypanosoma brucei*. Divergent ER retention signals in a lower eukaryote. *Journal of cell science* 1993, 105 (Pt 4), 1101-1113.
17. Cruz, A.; Beverley, S.M. Gene replacement in parasitic protozoa. *Nature* 1990, 348, 171-173.
18. Mukherjee, S.; Xu, W.; Hsu, F.F.; Patel, J.; Huang, J.; Zhang, K. Sterol methyltransferase is required for optimal mitochondrial function and virulence in *Leishmania major*. *Molecular microbiology* 2019, 111, 65-81, doi:10.1111/mmi.14139.
19. Ha, D.S.; Schwarz, J.K.; Turco, S.J.; Beverley, S.M. Use of the green fluorescent protein as a marker in transfected *Leishmania*. *Molecular and biochemical parasitology* 1996, 77, 57-64.
20. Murta, S.M.; Vickers, T.J.; Scott, D.A.; Beverley, S.M. Methylene tetrahydrofolate dehydrogenase/cyclohydrolase and the synthesis of 10-CHO-THF are essential in *Leishmania major*. *Molecular microbiology* 2009, 71, 1386-1401.
21. Wang, M.Z.; Zhu, X.; Srivastava, A.; Liu, Q.; Sweat, J.M.; Pandharkar, T.; Stephens, C.E.; Riccio, E.; Parman, T.; Munde, M.; et al. Novel arylimidamides for treatment of visceral leishmaniasis. *Antimicrobial agents and chemotherapy* 2010, 54, 2507-2516, doi:10.1128/AAC.00250-10.
22. Ohvo-Rekila, H.; Ramstedt, B.; Leppimaki, P.; Slotte, J.P. Cholesterol interactions with phospholipids in membranes. *Prog Lipid Res* 2002, 41, 66-97.
23. Zhang, K.; Hsu, F.F.; Scott, D.A.; Docampo, R.; Turk, J.; Beverley, S.M. *Leishmania* salvage and remodelling of host sphingolipids in amastigote survival and acidocalcisome biogenesis. *Molecular microbiology* 2005, 55, 1566-1578.
24. Boitz, J.M.; Gilroy, C.A.; Olenyik, T.D.; Paradis, D.; Perdeh, J.; Dearman, K.; Davis, M.J.; Yates, P.A.; Li, Y.; Riscoe, M.K.; et al. Arginase Is Essential for Survival of *Leishmania donovani* Promastigotes but Not Intracellular Amastigotes. *Infection and immunity* 2017, 85, doi:10.1128/IAI.00554-16.
25. Ilg, T. Proteophosphoglycans of *Leishmania*. *Parasitology today (Personal ed)* 2000, 16, 489-497.
26. Zhang, K. Balancing de novo synthesis and salvage of lipids by *Leishmania* amastigotes. *Current opinion in microbiology* 2021, 63, 98-103, doi:10.1016/j.mib.2021.07.004.
27. He, M.; Kratz, L.E.; Michel, J.J.; Vallejo, A.N.; Ferris, L.; Kelley, R.I.; Hoover, J.J.; Jukic, D.; Gibson, K.M.; Wolfe, L.A.; et al. Mutations in the human SC4MOL gene encoding a methyl sterol oxidase cause psoriasisform dermatitis, microcephaly, and developmental delay. *The Journal of clinical investigation* 2011, 121, 976-984, doi:10.1172/JCI42650.

28. Blosser, S.J.; Merriman, B.; Grahl, N.; Chung, D.; Cramer, R.A. Two C4-sterol methyl oxidases (Erg25) catalyse ergosterol intermediate demethylation and impact environmental stress adaptation in *Aspergillus fumigatus*. *Microbiology (Reading)* 2014, *160*, 2492-2506, doi:10.1099/mic.0.080440-0.
29. Kennedy, M.A.; Johnson, T.A.; Lees, N.D.; Barbuch, R.; Eckstein, J.A.; Bard, M. Cloning and sequencing of the *Candida albicans* C-4 sterol methyl oxidase gene (ERG25) and expression of an ERG25 conditional lethal mutation in *Saccharomyces cerevisiae*. *Lipids* 2000, *35*, 257-262, doi:10.1007/s11745-000-0521-2.
30. Benveniste, P. Biosynthesis and accumulation of sterols. *Annu Rev Plant Biol* 2004, *55*, 429-457, doi:10.1146/annurev.arplant.55.031903.141616.
31. Darnet, S.; Bard, M.; Rahier, A. Functional identification of sterol-4alpha-methyl oxidase cDNAs from *Arabidopsis thaliana* by complementation of a yeast erg25 mutant lacking sterol-4alpha-methyl oxidation. *FEBS Lett* 2001, *508*, 39-43, doi:10.1016/s0014-5793(01)03002-2.
32. Lee, A.K.; Banta, A.B.; Wei, J.H.; Kiemle, D.J.; Feng, J.; Giner, J.-L.; Welander, P.V. C-4 sterol demethylation enzymes distinguish bacterial and eukaryotic sterol synthesis. *Proceedings of the National Academy of Sciences* 2018, *115*, 5884-5889.
33. Ning, Y.; Frankfater, C.; Hsu, F.F.; Soares, R.P.; Cardoso, C.A.; Nogueira, P.M.; Lander, N.M.; Docampo, R.; Zhang, K. Lathosterol Oxidase (Sterol C-5 Desaturase) Deletion Confers Resistance to Amphotericin B and Sensitivity to Acidic Stress in *Leishmania major*. *mSphere* 2020, *5*, doi:10.1128/mSphere.00380-20.
34. Cosentino, R.O.; Agüero, F. Genetic profiling of the isoprenoid and sterol biosynthesis pathway genes of *Trypanosoma cruzi*. *PLoS one* 2014, *9*, e96762.
35. Kapler, G.M.; Coburn, C.M.; Beverley, S.M. Stable transfection of the human parasite *Leishmania major* delineates a 30-kilobase region sufficient for extrachromosomal replication and expression. *Mol Cell Biol* 1990, *10*, 1084-1094.
36. Zhang, O.; Wilson, M.C.; Xu, W.; Hsu, F.F.; Turk, J.; Kuhlmann, F.M.; Wang, Y.; Soong, L.; Key, P.; Beverley, S.M.; et al. Degradation of host sphingomyelin is essential for *Leishmania* virulence. *PLoS pathogens* 2009, *5*, e1000692, doi:10.1371/journal.ppat.1000692.
37. Zhang, K., Barron, T., Turco, S. J., and Beverley, S. M. The *LPG1* gene family of *Leishmania major*. *Mol. Biochem. Parasitol.* 2004, *136*, 11-23. PMC Journal – In Process. PMID: 15138063.
38. Folch, J.; Lees, M.; Sloane Stanley, G.H. A simple method for the isolation and purification of total lipides from animal tissues. *The Journal of biological chemistry* 1957, *226*, 497-509.
39. Feng, M.; Jin, Y.; Yang, S.; Joachim, A.M.; Ning, Y.; Mori-Quiroz, L.M.; Fromm, J.; Perera, C.; Zhang, K.; Werbovetz, K.A.; et al. Sterol profiling of *Leishmania* parasites using a new HPLC-tandem mass spectrometry-based method and antifungal azoles as chemical probes reveals a key intermediate sterol that supports a branched ergosterol biosynthetic pathway. *Int J Parasitol Drugs Drug Resist* 2022, *20*, 27-42, doi:10.1016/j.ijpddr.2022.07.003.
40. de Ibarra, A.A.; Howard, J.G.; Snary, D. Monoclonal antibodies to *Leishmania tropica major*: specificities and antigen location. *Parasitology* 1982, *85 (Pt 3)*, 523-531.
41. Titus, R.G.; Marchand, M.; Boon, T.; Louis, J.A. A limiting dilution assay for quantifying *Leishmania major* in tissues of infected mice. *Parasite Immunol* 1985, *7*, 545-555.
42. Basu, S.; Pawlowic, M.C.; Hsu, F.F.; Thomas, G.; Zhang, K. Ethanolaminephosphate cytidyltransferase is essential for survival, lipid homeostasis and stress tolerance in *Leishmania major*. *PLoS pathogens* 2023, *19*, e1011112, doi:10.1371/journal.ppat.1011112.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.