
Selenomethionine Mitigates Effects of *Nocardia cyriacigeorgica*-Induced Inflammation, Oxidative Stress and Apoptosis in Bovine Mammary Epithelial Cells

[Talgat Assabayev](#) , Jinge Han , Halihaxi Bahetijiang , [Venera Abdrassilova](#) , [Muhammad Asfandyar Khan](#) , [Herman W. Barkema](#) , [Gang Liu](#) , [John P. Kastelic](#) , [Xueying Zhou](#) ^{*} , [Bo Han](#) ^{*}

Posted Date: 9 September 2024

doi: 10.20944/preprints202409.0635.v1

Keywords: bovine mastitis; *Nocardia cyriacigeorgica*; selenomethionine; oxidative stress; apoptosis



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Selenomethionine Mitigates Effects of *Nocardia cyriacigeorgica*-Induced Inflammation, Oxidative Stress and Apoptosis in Bovine Mammary Epithelial Cells

Talgat Assabayev ¹, Jinge Han ², Halihaxi Bahetijiang ¹, Venera Abdrasilova ³, Muhammad Asfandyar Khan ¹, Herman W. Barkema ⁴, Gang Liu ¹, John P. Kastelic ⁴, Xueying Zhou ^{1,*} and Bo Han ^{1,*}

¹ College of Veterinary Medicine, China Agricultural University, Beijing 100193, P.R. China; talgat.asabaev@mail.ru; 1563828570@qq.com; asfand@cau.edu.cn; gangliu@cau.edu.cn; xueying@cau.edu.cn; hanbo@cau.edu.cn

² College of Animal Science and Veterinary Medicine, Tianjin Agricultural University, Tianjin, 300384, P. R. China; 2203020138@stu.tjau.edu.cn

³ Department of normal physiology with biophysics course, Asfendiyarov Kazakh National Medical University, Almaty 050012, Kazakhstan; Cholpan_69@mail.ru

⁴ Faculty of Veterinary Medicine, University of Calgary, Calgary, AB, Canada, T2N 4N1; barkema@ucalgary.ca; jpkastel@ucalgary.ca

* Correspondence: xueying@cau.edu.cn (X.Z.); hanbo@cau.edu.cn (B.H.), Tel.: 0086-1062733801

Abstract: *Nocardia cyriacigeorgica* causes bovine mastitis, reduces milk quantity and quality, and is often resistant to antimicrobials. Selenomethionine (SeMet) is a form of selenium, an essential trace element. Although SeMet reduced ROS-mediated apoptosis and intramammary infections. However, protective effects of SeMet on *N. cyriacigeorgica*-infected bovine mammary epithelial cells (bMECs) are unclear. The objective was to evaluate whether SeMet mitigated *N. cyriacigeorgica*-induced inflammatory injury, oxidative damage and apoptosis in bMECs. Cells were cultured with or without being pretreated 40 μ M of SeMet for 12 h, then challenged *N. cyriacigeorgica* (MOI=5:1) for 6 h. Although *N. cyriacigeorgica* was resistant to lincomycin, erythromycin, enrofloxacin, penicillin, amoxicillin, cephalonium, cephalexin, and ceftriaxone, 40 μ M SeMet increased cell viability and inhibited LDH release in infected bMECs. Furthermore, *N. cyriacigeorgica* significantly induced mRNA production and protein expression of TNF- α , IL-1 β , IL-6, and IL-8 at 6 h. Cell membrane rupture, cristae degeneration and mitochondria swelling were evident with transmission electron microscopy. SOD and GSH-px activities were down regulated after 3, 6 or 12 h, whereas MDA and ROS contents were significantly upregulated, with cell damage and apoptosis rapidly evident (the latter increased significantly in a time-dependent manner). In contrast, bMECs pretreated with 40 μ M SeMet before infection, SOD and GSH-px activities were upregulated ($P < 0.05$), MDA and ROS concentrations were downregulated ($P < 0.05$), and apoptosis was reduced ($P < 0.05$). In conclusion, 40 μ M SeMet alleviated inflammation, oxidative stress and apoptosis induced by *N. cyriacigeorgica* in bMECs cultured *in vitro*.

Keywords: bovine mastitis; *Nocardia cyriacigeorgica*; selenomethionine; oxidative stress; apoptosis

1. Introduction

Bovine mastitis affects dairy cows worldwide, with *Nocardia* spp. causing mastitis in dairy cows and enormous economic losses [1]. Furthermore, *Nocardia* can cause various local and systemic infections in humans and animals and is transmitted in a variety of ways, including inhalation, ingestion or traumatic implantation, with subsequent dissemination via lymphatics and blood [2]. *Nocardia cyriacigeorgica* causes chronic inflammation that does not respond to therapy [3] and is the

species of *Nocardia* most commonly linked to bovine mastitis [4–6]. *Nocardia* mastitis in cattle causes granulomatous inflammation, initially acute and eventually chronic [3,7].

As *Nocardia* spp. are resistant to many antimicrobials commonly used in veterinary medicine, mastitis caused by these bacteria result in chronic intramammary infection, mammary dysfunction, and cow culling [8]. Longstanding antibiotic use has increased antimicrobial resistance in *N. cyriacigeorgica*, creating global concerns for human and animal health [9]. However, antimicrobial resistance data on isolates associated with mastitis are limited.

Bacterial mastitis causes oxidative stress by promoting production of reactive oxygen species (ROS). Malondialdehyde (MDA) is frequently used to assess lipid peroxidation [10], which is counteracted by specific enzymes [11]. Oxidative stress can damage udder tissues and decrease milk production [12]. With cells cultured in vitro, damaged membranes release intracellular lactate dehydrogenase (LDH), an indicator of cytopathic effects [13]. In addition, ROS generated during oxidative stress could result in cytotoxic effects through protein and amino acid oxidation, DNA damage, altered intracellular signaling, inhibition of antioxidant defense systems, and tissue damage.

Apoptosis occurs in response to various stimuli, including infection and injury [14]. Apoptosis has characteristic ultrastructural changes, including cell shrinkage, chromatin condensation, nuclear fragmentation, and apoptotic bodies [15]. The role of apoptosis in the context of infection or injury, particularly in bovine mastitis caused by *N. cyriacigeorgica*, is an area of ongoing research [3,15]. Although apoptosis has a role in host defense mechanisms against microbial invasion, specific interactions between *N. cyriacigeorgica* and bMECs regarding apoptosis are not well studied. Investigations on bovine mastitis caused by *N. cyriacigeorgica* have primarily focused on prevalence, molecular characterization, immunogenic proteins, and virulence determinants [3,14,15]. Further research on the apoptotic response of bMECs to *N. cyriacigeorgica* infection may provide insights into host-pathogen interactions and pathogenesis.

Selenomethionine (SeMet) is an important micronutrient that can prevent mastitis in dairy cows through various biological pathways, including suppressing apoptosis [16]. A deficiency of selenium increases susceptibility to various diseases and decreases productivity globally [17]. Nevertheless, effects of SeMet on apoptosis in *N. cyriacigeorgica*-infected bMECs are unknown. The objective was to evaluate whether SeMet mitigated *N. cyriacigeorgica*-induced inflammatory injury, oxidative damage, and apoptosis in bMECs.

2. Results

2.1. Antimicrobial Susceptibility

The *N. cyriacigeorgica* was resistant to most antibiotics tested, but it was susceptible to gentamycin (66.7%) and ceftiofur (93.3%) (Table 1).

Table 1. Antimicrobials used for MIC assay.

Antimicrobial	<i>Nocardia cyriacigeorgica</i> – MIC (µg/mL)											Resistance rate (%)	MIC90 (µg/mL)
	0.5	1	2	4	8	16	32	64	128	256	512		
Gentamycin				5	2			1	2		5	10/15 (66.7)	512
Enrofloxacin				1		1					13	15/15 (100)	256
Penicillin											15	15/15 (100)	512
Amoxicillin								2			13	15/15 (100)	256
cephalonium							1	2			12	15/15 (100)	256
Cefalexin											15	15/15 (100)	512
Ceftriaxone				1					2	12		15/15 (100)	256
Ceftiofur			1						2	12		14/15 (93.3)	256
Lincomycin											15	15/15 (100)	-
Erythromycin											14	15/15 (100)	-

2.2. Cytotoxicity Effects of SeMet on bMECs

Based on the CCK-8 assay, bMECs treated with 40 μM of SeMet had higher viability ($P < 0.01$; Figure 1A) compared to the control group. However, 60 or 80 μM of SeMet significantly damaged bMECs at 12 h. In bMECs infected with *N. cyriacigeorgica*, 40 μM SeMet increased cell viability ($P < 0.05$; Figure 1B). Although 40 μM SeMet inhibited LDH release, > 60 μM SeMet increased LDH release ($P < 0.01$; Figure 1C). Furthermore, 40 μM SeMet reduced LDH release ($P < 0.01$) in bMECs infected with *N. cyriacigeorgica* (Figure 1D) without cytotoxicity.




Figure 1. Effects of SeMet with or without *N. cyriacigeorgica* on viability of bMECs and cytotoxicity of *N. cyriacigeorgica*. Cell viability of bMECs exposed to various concentrations of SeMet for 12 h, followed by *N. cyriacigeorgica* for 6 h. (A) Cell viability was determined with a CCK-8 assay in SeMet-treated bMECs. (B) Cell viability was determined with a CCK-8 assay in SeMet-treated bMECs subsequently infected with *N. cyriacigeorgica*. (C) LDH release detected by an LDH assay kit in SeMet-treated bMECs. (D) LDH activities determined by an LDH assay kit in SeMet-treated bMECs that were subsequently infected with *N. cyriacigeorgica*. Data are means \pm SD of 3 independent experiments. * $P < 0.05$; ** $P < 0.01$.

2.3. Transmission Electron Microscopy (TEM) of bMECs

In bMECs, at 6 h post-infection with *N. cyriacigeorgica*, ultrastructure changes included mitochondria and endoplasmic reticulum swelling, organelle loss, cell membrane rupture, perinuclear space, and cellular dysfunction (Figure 2). Moreover, Nocardia were in the nucleus and

cytoplasm, with indications that this bacterium caused apoptosis. In contrast, the SeMet-treated group had fewer intracellular bacteria and cell morphology was similar to the untreated group.




Figure 2. Ultrastructural changes of bMECs infected with *N. cyriacigeorgica* and examined with transmission electron microscopy. (A) Control group bMECs had abundant mitochondria (green arrow) in cytoplasm, nucleus (blue arrow) and normal endoplasmic reticulum (black arrow); (B) In the SeMet-treated group, all bMECs components appeared unaltered; (C) *N. cyriacigeorgica* infection group, the red arrow indicates *N. cyriacigeorgica* (NC); note that the endoplasmic reticulum (black arrow) were swollen, indicating cells were under stress. The green arrow indicates mitochondria, which appeared enlarged, with disruption of mitochondrial cristae (compared to the control group), whereas the blue arrow indicates the nucleus; (D) Pre-treatment with SeMet before *N. cyriacigeorgica* infection. Red arrow indicates *N. cyriacigeorgica* (NC), the green arrow indicates enlarged mitochondria and black arrows indicate swollen endoplasmic reticulum which are under stress. Note the protective effects of SeMet on bMECs, including mitochondria (green arrow), endoplasmic reticulum (black arrow), and nucleus (blue arrow), with less changes compared to the infection group.

2.4. Gene Expression and Production of Pro-Inflammatory Cytokines

Compared to the control group, both mRNA and protein expression levels of TNF- α , IL-1 β , IL-6, and IL-8 in bMECs were significantly downregulated and decreased in cells pretreated with 40 μ M of SeMet and then infected with *N. cyriacigeorgica* (Figure 3, $P < 0.01$; Figure 4, $P < 0.05$).




Figure 3. Effects of SeMet on the gene expression of proinflammatory cytokines. Expression of: (A) TNF- α ; (B) IL-1 β ; (C) IL-6; and (D) IL-8. The bMECs were pretreated with 40 μ M SeMet or nothing for 6 h before stimulation with *N. cyriacigeorgica* (MOI = 5:1) for 4 h. The mRNA expression of TNF- α , IL-1 β , IL-6 and IL-8 were downregulated compared to the control group. Data are means \pm SD of 3 independent experiments. * P < 0.05; ** P < 0.01.




Figure 4. SeMet alleviated *N. cyriaciorgica*-induced inflammatory responses in bMECs. The bMECs were treated with 40 μM of SeMet for 6 h and then infected with *N. cyriaciorgica* for 4 h. SeMet pretreatment alleviated *N. cyriaciorgica* -induced release of inflammatory cytokines in bMECs. (A) TNF-α; (B) IL-1β; (C) IL-6; and (D) IL-8. Data are means ± SD of 3 independent experiments. * $P < 0.05$; ** $P < 0.01$.

2.5. SOD and GSH-px Activities and MDA Concentration

Activities of antioxidant enzymes SOD and GSH-px were decreased ($P < 0.01$), whereas MDA concentration was increased ($P < 0.01$) in infected cells versus Controls. However, in cells pretreated with 40 μM SeMet and later challenged with *N. cyriaciorgica*, activities of both SOD and GSH-px increased ($P < 0.05$) whereas MDA content decreased ($P < 0.01$) compared to the infection group (Figure 5).




Figure 5. Effects of SeMet on *N. cyriacigeorgica* induced oxidative stress indicators in bMECs. (A) Superoxide dismutase (SOD) activities of bMECs infected *N. cyriacigeorgica* for 3, 6 or 12 h were decreased gradually ($P < 0.01$) but increased ($P < 0.05$) with 40 μ M SeMet pretreatment; **(B)** glutathione peroxidase GSH-px activity for 1, 2, 3, 4 and 5 min were decreased ($P < 0.01$) but increased ($P < 0.05$) with 40 μ M SeMet pretreatment; **(C)** malondialdehyde (MDA) content in bMECs infected with *N. cyriacigeorgica* for 3, 6, 9, or 12h were increased ($P < 0.01$) but were decreased ($P < 0.05$) with 40 μ M SeMet pretreatment. Means \pm SD of 3 independent experiments. Data are means \pm SD of 3 independent experiments. * $P < 0.05$, ** $P < 0.01$.

2.6. *N. Cyriacigeorgica* Stimulated Reactive Oxygen Species (ROS)

At 6 h after *N. cyriacigeorgica* infection, ROS concentrations in bMECs increased ($P < 0.01$) compared to the control group. However, in cells pretreated with 40 μ M SeMet, ROS concentrations were decreased ($P < 0.01$) compared to the infection group. Furthermore, ROS concentration was not different between the group pretreated with SeMet and uninfected bMECs ($P > 0.05$, Figure 6).




Figure 6. Effects of SeMet on *N. cyriacigeorgica*-induced increased in ROS concentrations in bMECs. In bMECs infected *N. cyriacigeorgica*, ROS was increased ($P < 0.01$). However, in bMECs pretreated with 40 μ M SeMet for 6 h before infection with *N. cyriacigeorgica* (MOI = 5:1), the ROS concentration was decreased ($P < 0.01$). In addition, ROS concentration of bMECs only pretreated with 40 μ M SeMet for 6 h was same as control. Data are means \pm SD of 3 independent experiments. ** $P < 0.01$.

2.7. *N. cyriacigeorgica* Induced Apoptosis in bMECs

Percentage of apoptosis in bMECs infected with *N. cyriacigeorgica* was increased in a time-dependent manner. At 3 h, apoptosis of the infection group was higher than the control group ($P < 0.05$, Figure 7). In addition, apoptosis in the infection group was higher ($P < 0.01$) than in the SeMet-treated group.




Figure 7. Apoptosis of bMECs treated with 40 uM SeMet for 12 h and subsequently infected with *N. cyriacigeorgica*. The bMECs were treated with 40 uM SeMet 12 h and then infected with *N. cyriacigeorgica* at MOI of 0 and 5 for 3, 6, 9, or 12 h. Percentage of apoptosis in bMECs infected with *N. cyriacigeorgica* was increased in a time-dependent manner. Apoptosis of the infection group was higher ($P < 0.01$) at various time points compared to the group given 40 uM SeMet. Therefore, SeMet mitigated apoptosis induced by *N. cyriacigeorgica*. Data are means \pm SD of 3 independent experiments. * $P < 0.05$; ** $P < 0.01$.

3. Discussion

Outbreaks of bovine mastitis caused by *Nocardia* spp. are common worldwide, including in China [3]. Key factors contributing to *Nocardia* bovine mastitis include poor management and hygiene, such as inadequate disinfection of teats before and after milking, lack of proper sanitation, and use of contaminated drugs or containers during milking [18]. Additionally, factors such as decreased immunity due to high milk production, calving and other diseases can increase susceptibility of cows to *Nocardia* infection.

As *N. cyriacigeorgica* adheres to and penetrate bMECs, causing cell damage and apoptosis [19], it is necessary to seek effective treatments. Although *Nocardia* spp. have diverse sensitivity to antibiotics, which can be used for phenotypic identification [20], those affecting animals are usually resistant to common veterinary antimicrobials. Antimicrobial susceptibility (MIC method) of *N. cyriacigeorgica* to 10 antimicrobials was determined. The *N. cyriacigeorgica* were susceptible only to gentamycin and ceftiofur, but resistant to 8 other microbials. *N. puris*, *N. veterana*, *N. cyriacigeorgica*, *N. arthritidis* and *N. africana* were resistant to cloxacillin (75%), ampicillin (55%) and cefoperazone (47%), but less resistant to amikacin, cefuroxime or gentamicin [7]. However, there are limited reports characterizing antimicrobial sensitivity/resistance of *Nocardia* spp. from bovine mastitis [21]. Moreover, based on the intracellular location of the pathogen and virulence characteristics that cause a pyogranulomatous reaction, treatment of *Nocardia* mastitis is usually unsuccessful. Therefore, early identification of clinical and subclinical cases of bovine mastitis by microbiological identification and drug sensitivity tests *in vitro* is crucial to improve response to treatment or promote timely culling [22]. Although ceftriaxone and imipenem were used for treatment of moderate and severe nocardiosis in Australia, only 58.8% of isolates were sensitive to ceftriaxone and 41.0% to imipenem

[2]. A similar susceptibility pattern was reported in other Australian studies for ceftriaxone (44-60%) and imipenem (33.7-48%) based on a bone mineral density test [8,23].

In this study, effects of various concentrations of SeMet on cell viability, LDH assay, antioxidant status and oxidative stress in bMECs were determined. Although 40 μ M SeMet treatment had protective effects on bMECs, higher concentrations (i.e., 60 or 80 μ M) were toxic. It was reported that SeMet alleviated ESBL *E. coli*-induced inflammation by activating the selenoprotein S-mediated TLR4/NF- κ B signaling pathway in bMECs [24], consistent with the present results. Also, SeMet enhanced expression of glutathione peroxidase 1 and 3 *in vivo* and *in vitro* and improved growth and viability of bMECs [25], whereas selenium concentrations from 50 to 100 nmol/L improved antioxidant function but did not affect milk protein synthesis in healthy bMECs. In addition, selenium mitigated lipopolysaccharide-induced damage by increasing antioxidant markers and stimulating milk protein synthesis and expression of milk protein-related genes in bMECs [26].

Oxidative stress impacts physiological functions of cells, including the immune response and the reproductive system [27,28]. Selenomethionine is regarded as an important element of the antioxidant system, as selenium enhances bMECs antioxidant function and protects cells from diethylenetriamine/NO-induced oxidative damage, mainly by increasing TrxR activity and reducing NO concentration via modulation of Nrf2 and MAPK signalling pathways [29]. Dietary intake of selenomethionine is directly correlated with the activity of various antioxidant enzymes (GSH-px, SOD, MDA and ROS). In this study, SOD and GSH-px activities were downregulated by *N. cyriacigeorgica* infection. However, pretreatment with 40 μ M SeMet upregulated SOD and GSH-px activities. Furthermore, in bMECs, MDA is considered a major product of lipid peroxidation and a marker of oxidative stress and antioxidant status [28]. In this study, whereas MDA and ROS contents were upregulated by *N. cyriacigeorgica* infection, SeMet pretreatment downregulated MDA and ROS, implying SeMet had a protective effect. Similarly, dairy cows given selenium yeast supplements (SeMet is the predominant form of selenium) had increased serum GSH-px activity compared to cows given sodium selenite supplements [30]. Furthermore, 0.3 mg/kg DM of hydroxy-selenomethionine enhanced GSH-px activity in dairy cow serum compared to the same dose of sodium selenite [31]. In studies using primary rainbow trout hepatocytes to determine effects of sodium selenite and SeMet on arsenite cytotoxicity, SeMet attenuated oxidative stress by increasing SOD [31,32]. Furthermore, SeMet increased capacity of bMECs to withstand oxidative stress, as indicated by decreased ROS and MDA [31], consistent with current results. Also, selenium suppressed *S. aureus*-induced inflammation by inhibiting the NLRP3 inflammasome and ROS activation in bMECs [33].

The current study was to investigate whether SeMet mitigated inflammatory damage and oxidative damage in bMECs caused by *N. cyriacigeorgica* strains isolated from mastitic milk. Expression of TNF- α , IL-1 β , IL-6, and IL-8 were significantly upregulated in bMECs infected by *N. cyriacigeorgica*. Nevertheless, pretreatment with 40 μ M SeMet significantly downregulated gene expression of TNF- α , IL-1 β , IL-6, and IL-8. Therefore, SeMet reduced production of inflammatory cytokines and decreased inflammatory damage to bMECs induced by *N. cyriacigeorgica*. Similarly, SeMet increased the autophagic flux and inhibited the NF- κ B-mediated inflammatory response of bMECs caused by *Klebsiella pneumoniae* [34]. Furthermore, gene expression of TNF- α , IL-1 β , IL-6 and IL-8 were also significantly upregulated 1 to 5 h after infection by *K. pneumoniae* [35], although SeMet reduced inflammation and protein expression of TLR4, Ikappa-B, NF- κ B, and TNF- α in bMECs and macrophages infected with ESBL *E. coli* [16]. Increased selenium concentrations decreased expression of IL-1 β and IL-6 [36]. In addition, selenium reduced gene expression of inflammatory mediators TNF- α , IL-1 β and IL-6 through the TLR2, NF- κ B and MAPK signaling pathway in *S. aureus*-infected bMECs, implying this was the basis of the anti-inflammatory effect of selenium [37]. Pre-treatment with various concentrations of SeMet significantly downregulated protein expression in ESBL-producing *E. coli*-infected bMECs [16], consistent with current findings. IL-1 β triggers neutrophil migration and its expression level is related to severity of inflammation. Moreover, IL-6 promotes cell proliferation and differentiation and induces acute gene expression in hepatocytes, increasing inflammation [38].

In the current study, SeMet clearly had a protective effect against apoptosis induced by *N. cyriacigeorgica* infection of bMECs. Although apoptosis in the *N. cyriacigeorgica* infection group was significantly upregulated in a time-dependent manner, that prevented by pre-treatment with 40 μ M SeMet, indicating that SeMet alleviated *N. cyriacigeorgica*-induced cell apoptosis, as this bacterium induces both apoptotic and necrotic changes in bMECs after 6 h [19]. Based on transcription levels of apoptosis-associated genes, overexpression of heme-binding protein could induce cell apoptosis [39]. MTSP3141, a secretory protein from *N. seriolae*, could target mitochondria and induce apoptosis in host cells, and function as a virulence factor [40]. However, selenium-related treatments protected against apoptosis, whereas selenium-enriched plant supplements significantly downregulated apoptosis of ovarian cells [41]. SeMet supplementation protected bMECs from H₂O₂-induced apoptosis and enhanced cell proliferation and viability under oxidative stress [42]. Furthermore, 500 nM of Na₂ SeO₃ or 30 μ M of L-selenium-methylselenocysteine significantly suppressed apoptosis and reduced mitochondrial membrane potential in murine liver cells, whereas 0.45 mg/kg dietary selenium also remarkably ameliorated liver damage in mice fed high-fat diets [43]. Extracellular vesicles derived from selenium-deficient bMECs induced oxidative stress, provoking apoptosis and inflammation via endoplasmic reticulum stress and PI3K/Akt/mTOR signaling pathway, contributing to explain susceptibility to mastitis due to selenium deficiency [44]. SeMet decreased apoptosis by increasing overall antioxidant capacity and reducing cellular ROS [45], consistent with current results.

4. Materials and Methods

4.1. Cell Culture

A bovine mammary epithelial cell line (MAC-T) was cultured in Dulbecco's modified Eagle medium (DMEM) containing 10% fetal bovine serum and 1% antibiotic mix (100 μ g/mL streptomycin and 100 U/mL penicillin), in an incubator with 5% CO₂ at 37°C. At 70-80% confluence, cells were detached and seeded into 6- or 96-well plates for assays (Figure 8).




Figure 8. Schematic diagram of the experimental protocol.

4.2. Bacterial Culture and Basic Tests

N. cyriacigeorgica previously isolated in our laboratory from milk samples with bovine mastitis, was activated from frozen stocks by culturing on tryptose soya agar (Difco™, Becton Dickson, Sparks, MD USA) supplemented with 5% defibrinated sheep blood. It was maintained in aerobic conditions at 37°C for 72 h, then sub-cultured in Brain Heart Infusion (BHI) broth for the following experiments.

4.3. Minimum Inhibitory Concentration (MIC) Determination

Antimicrobial susceptibility of *N. cyriacigeorgica* was assessed with a broth microdilution assay, recommended by the Clinical and Laboratory Standards Institute (CLSI, 2017) [46]. *Staphylococcus aureus* ATCC 29213 was used for quality control [47]. The MIC was the lowest concentration of antimicrobial that totally inhibited bacterial growth. For MIC analyses, 96-well microplates were used, with 200 μ L of BHI and antimicrobial added. After incubation of *N. cyriacigeorgica* in BHI broth for 48 h, MIC values were the lowest concentrations of the antimicrobial without visible bacterial growth. We tested the following antimicrobial agents: gentamycin, lincomycin, erythromycin, enrofloxacin, penicillin, amoxicillin, cephalonium, cephalixin, ceftriaxone, and ceftiofur. With MIC values based on visual observation. Concentrations needed to inhibit visible growth of 50 and 90% of isolates were defined as MIC₅₀ and MIC₉₀, respectively.

4.4. Assessment of Cell Viability by CCK-8

Cell Counting Kit-8 (CCK-8 kit; Beyotime Institute of Biotechnology, Beijing, China) is a rapid and highly sensitive method for measuring cell proliferation and cytotoxicity. MAC-T cells (~5,000 cells/well) were plated in 96-well plates with 0, 5, 10, 20, 40, 60, or 80 μ mol/L SeMet for 12 h. After treatment, cells were infected with *N. cyriacigeorgica* (MOI=5:1) and incubated at 37°C for 6 h and a CCK-8 kit used to assess cell viability, in accordance with label protocols. After 10 μ L of CCK-8 solution was added in each well, cells were incubated at 37°C for 2 h. Absorbance was recorded at 450 nm in a DNM 9602 microplate reader. Cell viability was evaluated as a percentage (%) relative to the control.

4.5. Assessment of Lactate Dehydrogenase (LDH) Release

Cytotoxic effects of *N. cyriacigeorgica* on bMECs were evaluated using an LDH assay kit (Beyotime Institute of Biotechnology). The bMECs cells were cultured at 37°C with 5% CO₂ in 96-well plates and incubated with 0, 5, 10, 20, 40, 60, or 80 μ mol/L SeMet for 12 h. Afterwards, treated cells were infected with *N. cyriacigeorgica* at a MOI of 5:1 for 4 or 6 h. After incubation, the cell culture plate was centrifuged at 400 \times g for 5 min, and then 120 μ L of supernatant was transferred to a new 96-well plate, with 60 μ L of reaction mixture added to each well, followed by incubation at room temperature in the dark for 30 min on a horizontal shaker (150 rpm). Optical density (OD) was measured by a spectrophotometer at 490 nm.

4.6. Transmission Electron Microscopy (TEM) of BMECs

For TEM, bMECs were seeded on 6-well plates and infected with *N. cyriacigeorgica* (MOI=5:1) for 6 h. After infection, cells were collected with a cell scraper and centrifuged at 1000 \times g for 5 min. They were washed twice with PBS, fixed in 2.5% glutaraldehyde for 2 h and 1% osmium tetroxide for 2 h at 4°C, dehydrated in a graded ethanol series, embedded in epoxy resin acetone mixtures (2:1) for 2 h, then immersed in pure resin for 12 h at 37°C. After polymerization, ultrathin sections (50 ~ 70 nm) were cut, stained with 1% uranyl acetate and lead citrate, and examined with transmission electron microscopy (JEM-1400, Tokyo, Japan) at 200 KeV.

4.7. Determination of Pro-Inflammatory Cytokines

BMECs were infected with *N. cyriacigeorgica* (MOI=5:1) for 4 h and remained in culture medium with antibiotics and lysozyme. The supernatant was collected by centrifugation at 12000 \times g for 10 min at 4°C. An ELISA kit (Jiangsu Meimian Industrial Co., Ltd., Jiangsu, China) with a detection range of 0.1 to 20.0 ng/mL was used (according to manufacturer's instructions) to assess concentrations of pro-inflammatory cytokines TNF- α , IL-1 β , IL-6, and IL-8 in cell supernatant. Absorbance at 450 nm was measured on a DNM 9602 microplate reader. All measurements were done 3 times, and the mean of 3 independent readings used for statistical analyses.

4.8. RNA Extraction and Reverse Transcriptase Quantitative Polymerase Chain Reaction (RT-qPCR)

Briefly, bMECs were washed once with PBS and harvested with 1 mL of TransZol Up lysis solution to extract total RNA, according to label instructions (TransGen Biotech Co. Ltd., Beijing, China). The cDNA was synthesized by TransScript® II All-in-One First-Strand cDNA Synthesis SuperMix for qPCR (One-Step gDNA Removal; TransGen Biotech). Concentrations of total RNA and cDNA were determined by absorbance at 260 and 280 nm on a NanoDrop-2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). A 20- μ L RT-qPCR mixture was used, comprised of 10 μ L of 2 \times RealStar Fast SYBR qPCR mix (GenStar, Beijing, China), 10 μ M (0.5 μ L) of each primer, 1 μ L cDNA, and 8 μ L of nuclease-free water. Expression levels of mRNA were tested by real-time PCR as follows: pre-denaturation at 94°C for 2 min; 40 cycles of denaturation at 95°C for 5 s; and annealing at 60°C for 30 s using an Applied Biosystems StepOnePlus Real Time PCR system (Thermo Fisher Scientific). Primers for a housekeeping gene (GAPDH) and genes encoding TNF- α , IL-1 β , IL-6, and IL-8 were used (Table 2). Cycle threshold (Ct) values and 2 $^{-\Delta\Delta$ Ct values were calculated to determine expression.

Table 2. Primer sequences of bovine genes used for quantitative PCR.

Gene		Primer sequence	Reference
TNF- α	Upstream	5'-ACGGGCTTTACCTCATCTACTC	Lin et al., 2020 [48]
	Downstream	3'-GCTCTTGATGGCAGACAGG	
IL-1 β	Upstream	5'-AGGTGGTGTTCGGTCATCGT	Lin et al., 2020[48]
	Downstream	3'-GCTCTCTGTCCTGGAGTTTGC	
IL-6	Upstream	5'-ATCAGAACACTGATCCAGATCC	Lin et al., 2020[48]
	Downstream	3'-CAAGGTTTCTCAGGATGAGG	
IL-8	Upstream	5'-ACACATTCCACACCTTTCCA	Lin et al., 2020[48]
	Downstream	3'-GGTTTAGGCAGACCTCGTTT	
GAPDH	Upstream	5'-CATTGACCTTCACTACATGGT	Lin et al., 2020[48]
	Downstream	3'-ACCCTTCAAGTGAGCCCCAG	

4.9. Determination of MDA Concentration, SOD and GSH-px Activities

The bMECs were cultured in 6-well plates and incubated with *N. cyriacigeorgica* (MOI 0 and 5) for 6 h. After infection, bMECs were washed 3 times with ice-cold PBS, lysed in SOD sample preparation solution, then centrifuged at 12,000 \times g at 4 °C for 3-5 min and supernatant collected for a subsequent assay. Then, MDA concentrations and activities of GSH-px and SOD were detected using Cell MDA, GSH-px and SOD assay kits (Cat. no. S0131, S0056 and S0101, respectively; Beyotime Biotechnology), according to manufacturer's instructions. Readings were recorded with a DNM 9602 microplate reader at 450 and 340 nm. Each group was assayed 30 times (10 replicates per group and each sample assayed in triplicate).

4.10. Measurement of Reactive Oxygen Species (ROS) Concentrations

Intracellular ROS were measured by the reactive oxygen species assay kit (Beyotime Institute of Biotechnology, Shanghai, China) with an oxidation-sensitive fluorescent probe (DCFH-DA). Cells were seeded into 6-well plates with glass coverslips, treated, then cells were washed once using PBS and stained in darkness with 100 μ M of diluted DCFH-DA for 20 min at 37°C and then washed 3 times with DMEM. The DCFH-DA was diluted 1000 times with serum-free DMEM/F12 culture medium, according to manufacturer's instructions. Finally, cells were stained with DAPI for 10 min. After washing 3 times in ice-cold PBS, samples were viewed with a laser confocal microscope (NIKON A1 HD25, Japan).

4.11. Determination of Apoptosis

Apoptosis induced by *N. cyriacigeorgica* in bMECs was determined with an Annexin V-FITC/propidium iodide (PI) analysis kit (Yeasen Biotechnology (Shanghai) Co., Ltd.) according to

manufacturer's instructions. Cells were plated into 6-well plates and infected with *N. cyriaciageorgica* at MOI of 0 or 5 for 3, 6, 9, or 12 h. After infection, cell culture medium was discarded and cells were harvested with 0.25% trypsin without EDTA and collected by pipetting into 1.5 ml PCR tubes. Cells were centrifuged at $1000 \times g$ $4^{\circ}C$ for 5 min, washed twice with ice-cold PBS, re-suspended in 100 μ L of $1 \times$ binding buffer, and stained with 5 μ L of Annexin V (FITC) and 10 μ L of PI. After 15 min in the dark at room temperature, cells were assessed (FAC Scan Flow Cytometer; Becton, Dickinson and Co., East Rutherford, NJ, USA), as follows: Q1 = represents necrotic cells; Q2 = late apoptotic cells; Q3 = early apoptotic cells; and Q4 = late live cells or normal cells.

4.12. Statistical Analyses

Results were expressed as the mean \pm standard deviation (SD) from at least 3 independent experiments. GraphPad Prism 8.0 (GraphPad Software, Inc., San Diego, CA, USA) was used and significance was assessed using SPSS 26.0 (IBM Corp., Armonk, NY, USA). Statistical differences were assessed with one-way analysis of variance (ANOVA), followed by the Tukey's multiple comparison *post hoc* test. Significance was accepted at $*P < 0.05$; $**P < 0.01$.

5. Conclusion

In conclusion, *Nocardia cyriaciageorgica* infection of bMECs induced structural disruption and mitochondrial degeneration. However, pretreatment with 40 μ M SeMet had the following protective actions: upregulated antioxidant enzyme activities, improved mRNA expressions and protein synthesis rates, reduced oxidative stress and apoptosis, and protected against inflammatory damage. An optimal dose of selenomethionine had cytoprotective roles in bMECs, although higher doses were cytotoxic. These findings highlighted the potential therapeutic benefits of selenomethionine supplementation in mitigating oxidative stress, inflammation, apoptosis and structural damage induced by bacterial infection in bMECs.

Author Contributions: Conceptualization, B.H.; methodology, A.T. and V.A.; software, A.T. and V.A.; validation, A.T., J.H., M.A.K., H. B. and G.L.; formal analysis, A.T., J.H., V.A. and M.A.K.; investigation, A.T.; resources, G.L.; data curation, A.T., J.H. and M.A.K.; writing—original draft preparation, A.T., J.H., V.A., M.A.K., G.L. and X.Z.; writing—review and editing, J.K. X.Z. and H.B.; visualization, A.T.; supervision, X.Z., B.H.; project administration, B.H.; funding acquisition, B.H. All authors have read and agreed to the submitted version of the manuscript.

Institutional Review Board Statement: The research was conducted in accordance with the Ethical Committee at the College of Veterinary Medicine, China Agricultural University (CAU), Beijing (Protocol AW01703202-2-1), and adhered to CAU's established ethical standards.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The raw data supporting the conclusions of this manuscript will be made available by the authors, without undue reservation, to any qualified researcher.

Acknowledgments : This study was supported financially by: National Natural Science Foundation of China (No. 32172928) and the High-End Foreign Experts Recruitment Program (No. G2022108009L).

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

bMECs: bovine mammary epithelial cells; MOI: multiplicity of infection; PBS: phosphate-buffered saline; *N. cyriaciageorgica*: *Nocardia cyriaciageorgica*; mTOR: Mammalian target of rapamycin; ESBL: extended-spectrum β -lactamase; TLR4/NF- κ B: Toll-like receptor 4/nuclear factor kappa B; TLR2: Toll-like receptor 2; Nrf2: nuclear factor-E2-related factor 2; MAPK: mitogen-activated protein kinase; NLRP3: NOD-like receptor protein 3; TNF- α : tumor necrosis factor- α ; IL-1 β : interleukin-1 β ; IL-6: interleukin-6; IL-8: interleukin-8; PI3K/Akt/mTOR: phosphatidylinositol 3-kinase/protein kinase B/mammalian target of rapamycin; MDA: malondialdehyde; GSH-px: glutathione peroxidase; SOD: superoxide dismutase; ROS: reactive oxygen species; LDH: Lactate

Dehydrogenase; SeMet: Selenomethionine; TEM: transmission electron microscope; MIC: minimum inhibitory concentration.

References

1. Mushtaq, S., Shah, A.M., Shah, A., Lone, S.A., Hussain, A., Hassan, Q.P., Ali, M.N., Bovine mastitis: An appraisal of its alternative herbal cure. *Microb. Pathog.* **2018**, *114*, 357-361.
2. Yeoh, K., Globan, M., Naimo, P., Williamson, D.A., Lea, K., Bond, K., Identification and antimicrobial susceptibility of referred *Nocardia* isolates in Victoria, Australia 2009-2019. *J. Med. Microbiol.* **2022**, *71(8)*, 1-7.
3. Chen, W., Liu, Y.X., Barkema, H.W., Gao, J., De Buck, J., Kastelic, J.P., Liu, G., Ali, T., Shahid, M., Han, B., Short communication: Molecular characteristics, antimicrobial susceptibility, and pathogenicity of clinical *Nocardia cyriacigeorgica* isolates from an outbreak of bovine mastitis. *J. Dairy Sci.* **2017a**, *100*, 8414-842.
4. Lu, S.H., Qian, Z.W., Mou, P.P., Xie, L., Clinical *Nocardia* species: Identification, clinical characteristics, and antimicrobial susceptibility in Shandong, China. *Bosn. J. Basic Med. Sci.* **2020**, *20(4)*, 531-538.
5. Conville, P.S., Brown-Elliott, B.A., Smith, T., Zelazny, A.M., The complexities of *Nocardia* taxonomy and identification. *J. Clin. Microbiol.* **2017**, *6(1)*, e01419-17.
6. Stevanović, O., Milanov, D., Prošić, I., Gajdov, V., Nedić, D., Sladojević, Ž., Radalj, A., Multilocus sequence analysis (MLSA) of a *Nocardia cyriacigeorgica* strain causing severe bovine mastitis in Bosnia and Herzegovina. *Acta Vet. Hung.* **2023**, *71(2)*, 65-70.
7. Condas, L.A.Z., de Farias, M.R., Siqueira, A.K., Salerno, T., Chi, K.D., Werner, J., de Vargas, A.C., Bond, G.B., Gono, T., Matsuzawa, T., Ribeiro, M.G., Molecular identification and antimicrobial resistance pattern of *Nocardia* isolated from 14 diseased dogs and cats. *Braz. J. Microbiol.* **2023**, *54(2)*, 1287-1294.
8. Tan, Y.E., Chen, S.C.A., Halliday, C.L., Antimicrobial susceptibility profiles and species distribution of medically relevant *Nocardia* species: results from a large tertiary laboratory in Australia. *J. Glob. Antimicrob. Resist.* **2020**, *20*, 110-117.
9. Lebeaux, D., Bergeron, E., Berthet, J., Djadi-Prat, J., Mounié, D., Boiron, P., Lortholary, O., Rodriguez-Nava, V., Antibiotic susceptibility testing and species identification of *Nocardia* isolates: a retrospective analysis of data from a French expert laboratory, 2010-2015. *Clin. Microbiol. Infect.* **2019**, *25*, 489-495.
10. Li, H., Song, F., Duan, L.R., Sheng, J.J., Xie, Y.H., Yang, Q., Chen, Y., Dong, Q.Q., Zhang, B.L., Wang, S.W., Paeonol and danshensu combination attenuates apoptosis in myocardial infarcted rats by inhibiting oxidative stress: Roles of Nrf2/HO-1 and PI3K/Akt pathway. *Sci. Rep.* **2016**, *6(1)*, 23693.
11. Rojo de la Vega, M., Dodson, M., Gross, C., Mansour, H.M., Lantz, R.C., Chapman, E., Wang, T., Black, S.M., Garcia, J.G., Zhang, D.D., Role of Nrf2 and autophagy in acute lung injury. *Curr. Pharmacol. Rep.* **2016**, *2(2)*, 91-101.
12. Ayemele, A.G., Tilahun, M., Lingling, S., Elsaadawy, S.A., Guo, Z., Zhao, G., Xu, J., Bu, D., Oxidative stress in dairy cows: Insights into the mechanistic mode of actions and mitigating strategies. *Antioxidants (Basel)* **2021**, *10(12)*, 1918.
13. Kaja, S., Payne, A.J., Naumchuk, Y., Koulen, P., Quantification of lactate dehydrogenase for cell viability testing using cell lines and primary cultured astrocytes. *Curr. Protoc. Toxicol.* **2017**, *72*, 1-15.
14. Damien, B., Eike, L., Bernardo, S.F., Necroptosis, pyroptosis and apoptosis: an intricate game of cell death. *Cell. Mol. Immunol.* **2021**, *18(5)*, 1106-1121.
15. Xue, B.X., Yue, Y.L., Zi, C.H., Apoptosis and apoptotic body: disease message and therapeutic target potentials. *Bioscience Rep.* **2019**, *39(1)*, BSR20180992.
16. Zhuang, C., Gao, J., Liu, G., Zhou, M., Yang, J., Wang, D., Kastelic, J., Han, B., Selenomethionine activates selenoprotein S, suppresses Fas/FasL and the mitochondrial pathway, and reduces *Escherichia coli*-induced apoptosis of bovine mammary epithelial cells. *J. Dairy Sci.* **2021**, *104*, 10171-10182.
17. Zou, Y., Shao, J., Li, Y., Zhao, F., Liu, J., Liu, H., Protective effects of inorganic and organic selenium on heat stress in bovine mammary epithelial cells. *Oxid. Med. Cell. Long.* **2019**, *2019*, 1503478.
18. Sharun, K., Dhama, K., Tiwari, R., Gugjoo, M.B., Iqbal Yattoo, M., Patel, S.K., Pathak, M., Karthik, K., Khurana, S.K., Singh, R., Puvvala, B., Amarpal, Singh, R., Singh, K.P., Chaicumpa, W., Advances in therapeutic and management approaches of bovine mastitis: a comprehensive review. *Vet. Q.* **2021**, *41(1)*, 107-136.

19. Chen, W., Liu, Y., Zhang, L., Gu, X., Liu, G., Shahid, M., Gao, J., Ali, T., Han, B., *Nocardia cyriacigeogica* from bovine mastitis induced *in vitro* apoptosis of bovine mammary epithelial cells via activation of mitochondrial-caspase pathway. *Front. Cell. Infect. Microbiol.* **2017b**, *7*, 194.
20. Toyokawa, M., Ohana, N., Ueda, A., Imai, M., Tanno, D., Honda, M., Takano, Y., Ohashi, K., Saito, K., Shimura, H., Identification and antimicrobial susceptibility profiles of *Nocardia* species clinically isolated in Japan. *Sci. Rep.* **2021**, *11(1)*, 16742.
21. Huang, L., Chen, X., Xu, H., Sun, L., Li, C., Guo, W., Xiang, L., Luo, G., Cui, Y., Lu, B., Clinical features, identification, antimicrobial resistance patterns of *Nocardia* species in China: 2009–2017. *Diagn. Microbiol. Infect. Dis.* **2019**, *94(2)*, 165-172.
22. Amano, T., Nishikawa, T., Oka, K., Ota, K., Shimizu, T., How an antimicrobial stewardship team treated a *Nocardia farcinica*-associated brain abscess: A case report. *Cureus* **2024**, *16(2)*, e54605.
23. Paige, E.K., Spelman, D., Nocardiosis: 7-year experience at an Australian tertiary hospital. *Intern. Med. J.* **2019**, *49(3)*, 373–379.
24. Zhuang, C., Liu, G., Barkema, H.W., Zhou, M., Xu, S., ur Rahman, S., Liu, Y., Kastelic, J.P., Gao, J., Han, B., Selenomethionine suppressed TLR4/NF- κ B pathway by activating selenoprotein S to alleviate ESBL *Escherichia coli*-induced inflammation in bovine mammary epithelial cells and macrophages. *Front. Microbiol.* **2020**, *11*, 1461.
25. Miranda, S.G., Wang, Y.J., Purdie, N.G., Osborne, V.R., Coomber, B.L., Cant, J.P., Selenomethionine stimulates expression of glutathione peroxidase 1 and 3 and growth of bovine mammary epithelial cells in primary culture. *J. Dairy Sci.* **2009**, *92(6)*, 2670-2683.
26. Zhang, B., Guo, Y., Yan, S., Guo, X., Zhao, Y., Shi, B., The protective effect of selenium on the lipopolysaccharide-induced oxidative stress and depressed gene expression related to milk protein synthesis in bovine mammary epithelial cells, *Biol. Trace Elem. Res.* **2020**, *197(1)*, 141-148.
27. Mavangira, V., Sordillo, L.M., Role of lipid mediators in the regulation of oxidative stress and inflammatory responses in dairy cattle, *Res. Vet. Sci.* **2018**, *116*, 4–14.
28. Sordillo, L.M., Nutritional strategies to optimize dairy cattle immunity, *J. Dairy Sci.* **2016**, *99(6)*, 4967–4982.
29. Guo, Y., Guo, X., Yan, S., Zhang, B., Shi, B., Mechanism underlying the protective effect of selenium on NO-induced oxidative damage in bovine mammary epithelial cells, *Biol. Trace Elem. Res.* **2019**, *191(1)*, 104-114.
30. Gong, J., Ni, L., Wang, D., Shi, B., Yan, S., Effect of dietary organic selenium on milk selenium concentration and antioxidant and immune status in midlactation dairy cows. *Livest. Sci.* **2014**, *70*, 84–90.
31. Sun, P., Wang, J., Liu, W., Bu, D.P., Liu, S.J., Zhang, K.Z., Hydroxy-selenomethionine: A novel organic selenium source that improves antioxidant status and selenium concentrations in milk and plasma of mid-lactation dairy cows. *J. Dairy Sci.* **2017**, *100(12)*, 9602–9610.
32. Jamwal, A., Niyogi, S., Dose and chemical species-specific effects of selenium against arsenite toxicity in cultured hepatocytes of rainbow trout (*Oncorhynchus mykiss*). *Metallomics* **2017**, *9(6)*, 744–756.
33. Yang, Y., Lv, S., Wang, Z., Liu, J., Selenium ameliorates *S. aureus*-induced inflammation in bovine mammary epithelial cells by regulating ROS-induced NLRP3 inflammasome, *Biol. Trace Elem. Res.* **2022**, *200(7)*, 3171-3175.
34. Xu, S., Miao, Y., Dong, J., Cui, L., Liu, K., Li, J., Meng, X., Zhu, G., Wang, H., Selenomethionine inhibits NF- κ B-mediated inflammatory responses of bovine mammary epithelial cells caused by *Klebsiella pneumoniae* by increasing autophagic flux, *Biol. Trace Elem. Res.* **2024**, *202(4)*, 1568-1581.
35. Ma, X.M., Xu, S.Y., Li, J.J., Cui, L.Y., Dong, J.S., Meng, X., Zhu, G.Q., Wang, H., Selenomethionine protected bMECs from inflammatory injury and oxidative damage induced by *Klebsiella pneumoniae* by inhibiting the NF- κ B and activating the Nrf2 signaling pathway. *Inter. Immunopharmacol.* **2022**, *110*, 109027.
36. Mahmoodpoor, A., Hamishehkar, H., Shadvar, K., Ostadi, Z., Sanaie, S., Saghaleini, S.H., Nader, N.D., The effect of intravenous selenium on oxidative stress in critically ill patients with acute respiratory distress syndrome, *Immunol. Invest.* **2019**, *48(2)*, 147-159.
37. Wang, H., Bi, C., Wang, Y., Sun, J., Meng, X., Li, J., Selenium ameliorates *Staphylococcus aureus*-induced inflammation in bovine mammary epithelial cells by inhibiting activation of TLR2, NF- κ B and MAPK signaling pathways, *BMC Vet. Res.* **2018**, *14(1)*, 197.
38. Ulasov, A.V., Rosenkranz, A.A., Georgiev, G.P., Sobolev, A.S., Nrf2/Keap1/are signaling: Towards specific regulation, *Life Sci.* **2022**, *291*, 120111.
39. Wen, Y., Kang, X., Li, Z., Xia, L., Lu, Y., Identification of a secretory heme-binding protein from *Nocardia seriolae* involved in cell apoptosis. *J. Fish Dis.* **2022**, *45(8)*, 1189-1199.

40. Chen, J., Xia, L., Wang, W., Wang, Z., Hou, S., Xie, C., Cai, J., Lu, Y., Identification of a mitochondrial-targeting secretory protein from *Nocardia seriolae* which induces apoptosis in fathead minnow cells. *J. Fish Dis.* **2019**, *42(11)*, 1493-1507.
41. Wang, Y., Liu, B.B., Wu, P.X., Chu, Y., Gui, S.S., Zheng, Y.Z., Chen, X.D., Dietary selenium alleviated mouse liver oxidative stress and NAFLD induced by obesity by regulating the KEAP1/NRF2 pathway. *Antioxidants (Basel)* **2022**, *11(2)*, 349.
42. Miranda, S.G., Purdie, N.G., Osborne, V.R., Coomber, B.L., Cant, J.P., Selenomethionine increases proliferation and reduces apoptosis in bovine mammary epithelial cells under oxidative stress. *J. Dairy Sci.* **2011**, *94(1)*, 165-173.
43. Wang, H., Cong, X., Qin, K., Yan, M.K., Xu, X.F., Liu, M.K., Xu, X., Zhang, Y., Gao, Q.Y., Cheng, S.Y., Zhao, J.C., Zhu, H., Liu, Y., Se-enriched cardamine violifolia improves laying performance and regulates ovarian antioxidative function in aging laying hens. *Antioxidants (Basel)* **2023**, *12(2)*, 450.
44. Chen, Y., Zhang, X., Yang, J., Feng, W., Deng, G., Xu, S., Guo, M., Extracellular vesicles derived from selenium-deficient MAC-T cells aggravated inflammation and apoptosis by triggering the endoplasmic reticulum (ER) stress/PI3K-AKT-mTOR pathway in bovine mammary epithelial cells. *Antioxidants (Basel)*. **2023**, *12(12)*, 2077.
45. Xie, L., Xu, Y., Ding, X., Li, K., Liang, S., Li, D., Wang, Y., Fu, A., Yu, W., Zhan, X., Selenomethionine attenuated H₂O₂-induced oxidative stress and apoptosis by Nrf2 in chicken liver cells. *Antioxidants (Basel)* **2023**, *12(9)*, 1685.
46. CLSI M07, Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. 12th edition, **2024**.
47. Ross, J.E., Scangarella-Oman, N.E., Flamm, R.K., Jones, R.N., Determination of disk diffusion and MIC quality control guidelines for GSK2140944, a novel bacterial type II topoisomerase inhibitor antimicrobial agent. *J. Clin. Microbiol.* **2014**, *52(7)*, 2629-2632.
48. Lin, J., Li, G., Xu, C., Lu, H., Zhang, C., Pang, Z., Liu, Z., Monocyte chemotactic protein 1-induced protein 1 is highly expressed in inflammatory bowel disease and negatively regulates neutrophil activities. *Mediators Inflamm.* **2020**, *2020*, 8812020.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.