

---

# Abundance, Size Structure, and Growth of the Invasive Blue Crab *Callinectes sapidus* in the Lesina Lagoon, Southern Adriatic Sea

---

[Giorgio Mancinelli](#) , [Nicola Lago](#) , Tommaso Scirocco , Oscar Antonio Lillo , Raffaele De Giorgi , [Lorenzo Doria](#) , [Emanuele Mancini](#) , [Francesco Mancini](#) , [Luigi Potenza](#) , and [Lucrezia Cilenti](#) \*

Posted Date: 1 October 2024

doi: 10.20944/preprints202410.0018.v1

Keywords: invasive species; fishery biology; CPUE; sex ratio; size structure; morphological maturity; growth dynamics; Mediterranean Sea



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

# Abundance, size structure, and growth of the invasive blue crab *Callinectes sapidus* in the Lesina Lagoon, southern Adriatic Sea

Giorgio Mancinelli <sup>1,2,3</sup>, Nicola Lago <sup>4</sup>, Tommaso Scirocco <sup>5</sup>, Oscar Antonio Lillo <sup>6,7</sup>, Raffaele De Giorgi <sup>1,2</sup>, Lorenzo Doria <sup>1</sup>, Emanuele Mancini <sup>1,2</sup>, Francesco Mancini <sup>8</sup>, Luigi Potenza <sup>1</sup> and Lucrezia Cilenti <sup>9,\*</sup>

<sup>1</sup> Department of Biological and Environmental Sciences and Technologies (DiSTeBA), University of Salento, 73100 Lecce, Italy; giorgio.mancinelli@unisalento.it

<sup>2</sup> NBFC, National Biodiversity Future Center, 90133 Palermo, Italy

<sup>3</sup> CoNISMa, National Inter-University Consortium for Marine Sciences, 00196 Roma, Italy

<sup>5</sup> National Research Council (CNR), Institute for Biological Resources and Marine Biotechnologies (CNR-IRBIM), 71010, Lesina (FG), Italy; tommaso.scirocco@cnr.it

<sup>4</sup> Istituto Zooprofilattico Sperimentale delle Venezie (IZSVe), 35020 Legnaro (PD), Italy; NLago@izsvenezie.it

<sup>6</sup> Metaponto Research Centre, Regional Agency for the Environmental Prevention and Protection of Basilicata (ARPAB), 75012 Metaponto, Italy

<sup>7</sup> Environmental Research Foundation of Basilicata Region (FARBAS), 85100 Potenza, Italy; antoniooscar.lillo@farbas.it

<sup>8</sup> International Centre for Advanced Mediterranean Agronomic Studies (CIHEAM-Bari), 70010 Valenzano (BA), Italy; mancini@iamb.it

<sup>9</sup> National Research Council (CNR), Institute of Sciences of Food Production (ISPA), 71121 Foggia, Italy

\* Correspondence: lucrezia.cilenti@cnr.it

**Simple Summary:** The exploitation of the Atlantic blue crab *Callinectes sapidus* has recently gained credit for effectively mitigating the impacts of the species in invaded Mediterranean waters. A several of studies have focused on the fishery biology of the species in the Aegean Sea and in the Eastern Mediterranean, while for other sectors of the basin information are lacking. Here fyke net catches of *C. sapidus* in the Lesina Lagoon (Adriatic Sea, SE Italy) were analysed on a monthly frequency between February 2021 and January 2022. A comprehensive assessment of the crab's life history parameters in terms of abundance, sex ratio, size-frequency distribution, morphological maturation, growth, and mortality was provided adopting a suite of analytical techniques. The results of the present study may provide a valuable knowledge basis for future comparative investigations on the fishery biology of *C. sapidus* in Mediterranean waters.

**Abstract:** The fishery biology of the invasive Atlantic blue crab *Callinectes sapidus* in the Mediterranean Sea outside the eastern sectors of the basin has been only recently investigated. Here we studied the population of *C. sapidus* in the Lesina Lagoon (Adriatic Sea, SE Italy). In total, 838 crabs were captured monthly between February 2021 and January 2022 using fyke nets. Abundances varied seasonally with catches per unit effort ranging between 0 and 1.76 crabs fyke nets<sup>-1</sup> d<sup>-1</sup> in winter and summer. Spatial abundances estimated in summer by a Carle-Strub procedure ranged between 0.06 and 0.64 crabs m<sup>-2</sup>. Sex ratio ( $\sigma/\varphi$ ) was close to 1:1; males prevailed only in August and September; ovigerous females occurred from April to August. Males' size at morphological maturity was smaller than females (110.6–112.3 mm vs. 122.1–123.1 mm). Seasonal von Bertalanffy growth parameters indicated that, compared with males, females showed a shorter maximum lifespan (5 vs. 8 years), a higher growth coefficient K (0.6 vs. 0.4 y<sup>-1</sup>) and growth performance index  $\Phi'$  (4.6 vs. 4.3), while maximum sizes CW $\infty$  (237.8 vs. 232.6 mm) and seasonality indices C (0.62 vs. 0.57) were similar. Furthermore, females showed higher natural and fishing mortalities and exploitation rate. We discussed the results of the present study in the context of the available literature to provide a valuable basis for the implementation of standardised Mediterranean-scale management plans, matching exploitation of *C. sapidus* with sustainable conservation of coastal ecosystems.

**Keywords:** invasive species; fishery biology; CPUE; sex ratio; size structure; morphological maturity; growth dynamics; Mediterranean Sea

---

## 1. Introduction

The Atlantic blue crab *Callinectes sapidus* Rathbun, 1896 (“blue crab” hereafter) is a portunid brachyuran characterized among other members of the genus by a trans-hemispheric distribution along the western coasts of the Atlantic Ocean, ranging from the Gulf of Maine through the Gulf of Mexico to Venezuela, Brazil, and Argentina [1–3]. In its native range, ecological and economic importance, acting as a keystone species regulating trophic cascades and the transfer of energy between the benthic and pelagic compartments [4,5], as well as representing the object of a major commercial and recreational fishery [6–9]. Hence, a huge body of information exists on the biology and ecology of the species primarily from the USA - among the thousands of publications available to date, the book “the Blue Crab: *Callinectes sapidus*” [10] represents an outstanding example and provides a comprehensive research reference - and from central and south American countries [9,11–15].

In the past century, unsuccessful attempts have been made to deliberately introduce the blue crab in California, Hawaii, and Japan [16]. In Europe the species was recorded for the first time in 1901 along the Atlantic coasts of France, probably introduced accidentally by ballast waters, while in the Mediterranean Sea it appeared in 1947 in the Aegean Sea [3,16]. In the following years the blue crab progressively extended its invasion range, with a remarkable expansion observed in the last decade in the Mediterranean and Black Seas as well as along the European and African Atlantic coasts [3,17,18].

A number of life-history and ecological traits likely contributed to the invasion success of the *C.sapidus* [16,19–21], yet a crucial role might be attributed to its high fecundity - females produce 2 to 8 million eggs per spawning event and brood repeatedly during the year depending on water temperature [22–24] and high dispersal capability of larval stages in coastal environments [25,26]. In general, the high environmental connectivity of marine habitats represents one of the main factors hampering the actions generally put in place to control bioinvaders, such as those aiming at their complete eradication [27]. Functional eradication, i.e., targeting a population to decrease its density to levels determining acceptable impacts on the invaded system, is presently acknowledged as a more appropriate mitigation strategy [28,29], in particular when the bioinvaders are species of economic interest [8,30]. The opportunity to develop an economically valuable fishery while controlling the ecological impact of the blue crab has long been emphasized [8]. Recently, the General Fisheries Commission for the Mediterranean (GFCM) identified adaptation strategies to cope with the potential effects of invasive species on fisheries as target outputs of the mid-term strategy (2017–2020) [31]. Noticeably, among the strategies developed, the Recommendation GFCM/42/2018/7 established a regional program to fill scientific and research gaps on *C. sapidus* and the Lessepsian *Portunus segnis* [32]; see also <https://www.fao.org/gfcm/researchprogramme-bluecrabs>. The underlying assumption of the research program is that the integration of fishery management actions within a context of functional eradication and impact mitigation requires an advanced, context-specific understanding of the ecology and biology of the species across invaded Mediterranean countries.

A diverse set of impact-related ecological information is starting to become available at a whole Mediterranean scale on the blue crab as related to e.g., its distribution and invasion dynamics [3,18,20], multi-specific effects on recipient communities [33], trophic habits [21,34–37], contamination by pollutants [38–43], and pathogens [44–46]. In contrast, studies focused on the fishery biology of the species have been performed prevalently in the eastern sectors of the Mediterranean basin [47,48]. Recent investigations from Greece [48–50], Southern Italy [51], and Montenegro [52] are valuable exceptions, yet additional research efforts are urgently needed on the biological characteristics of blue crab populations to allow for an effective management of their Mediterranean stocks.

Accordingly, our primary aim was to contribute to fill this knowledge gap and perform a comprehensive investigation of the abundance, size structure, sex ratio, reproductive biology, and growth dynamics of the species in the Lesina Lagoon, an Italian brackish system located on the


southwestern coasts of the Adriatic Sea. The blue crab was first recorded in the lagoon in 2007 and its occurrence has been repeatedly confirmed in the following years [24,53] and the species currently represents a predominant component of the by-catch of local artisanal fisheries [54]. Blue crabs were captured using fyke nets on a monthly frequency over one year; a suite of analytical approaches and tools based on size-frequency data were used to provide an in-depth scrutiny of crucial parameters characterizing the fishery biology of the blue crab population in the Lesina Lagoon, including natural and fishing mortality rates as well exploitation rates.

## 2. Materials and Methods

### 2.1. Study Area

The Lesina Lagoon (41.88°N, 15.45°E) is a shallow (0.7 m mean depth, 1.6 m maximum depth), micro-tidal lagoon located northward of the Gargano promontory along the southwestern Adriatic coasts of Italy (Figure 1a). The basin has a surface area of 51.4 km<sup>2</sup> and is separated from the Adriatic Sea by an 18 km-long sandy bar. Seawater inputs are assured by the Acquarotta and Schiapparo channels, located respectively at the western and eastern ends of the basin, respectively, while the Lauro and Zannella streams, together with several intermittent creeks and ditches, contribute freshwater in particular along the south-eastern coast of the lagoon (Figure 1a) [24,55]. These hydrological conditions determine permanent, east-west temperature and salinity gradients [56,57]; however, a remarkable seasonal variability is generally observed in both temperature and salinity values, with minima close to 10°C and 10 PSU observed in winter between December and February and maxima of 25-29°C and 28 PSU occurring in summer between June and August [56,57].

Four experimental sites were identified in January 2021 in the eastern sector of the lagoon (Figure 1a) where higher blue crab abundances are generally observed as compared with the rest of the basin [58]. The sector is included in a natural reserve covering an area of 930 ha (DMAF 27/04/1981) and is a Natura 2000 site (IT9110031). The common reed *Phragmites australis* (Cav.) Trin. ex Steud dominates the riparian vegetation while the interplay between freshwater and seawater inputs affects the local salinity conditions and ultimately influences the occurrence and abundance of aquatic angiosperms [*Zostera noltei* Hornemann, 1832 and *Ruppia cirrhosa* (Petagna) Grande, 1918] and macroalgae [*Zannichellia palustris* L., *Chaetomorpha linum* (O.F.Müller) Kützing, 1845, *Myriophyllum spicatum* L., 1753, *Chara globularis* Thuiller, 1799] [59].



**Figure 1.** (a) The Lesina Lagoon. Red circles and consecutive numbers indicate the location of the four fixed installations used to capture *Callinectes sapidus* specimens during the study period. (b) Illustration of a typical fyke similar to those used in the study. See the text for additional details on dimensions and sampling procedures. Image credit: <https://www.fao.org/fishery/en/geartype/226/en> [60]. (c) Schematic representation of the fixed installation located at Site 1. Please note that the figure is not in scale. Identical installations were located at each of the remaining sampling sites identified in Figure 1a.

## 2.2. Sampling procedures

Fishers operate in the Lesina Lagoon during fall and winter (September to February) using fixed gillnets, locally known as “paranze”, placed perpendicularly to the shores of the basin and conveying mobile prey into fyke-nets (see Figure 1b for an example) located at regular intervals along them [61]. In this study, an experimental fixed installation based on the general operative principles of a “paranza” was assembled at each study site on February 2021. In brief, the device was made of a 6 mm mesh size, 25 m long gill net perpendicular to the shore (leader) and 3 5 m long, 1 m wide cylindrical fyke nets with a 8 mm mesh size of the final net chamber (Figure 1c). After 24 h, captured prey from each site/fyke net combination was collected and transferred to the laboratory in refrigerated containers. At each site, sampling operations were reiterated for six consecutive days; the soaking time of the fyke nets was set at 24 h, yet it varied between 24 and 72 h during the investigation due to bad weather conditions preventing field operations. In each month, on the first sampling day a multi-parametric probe (Hydrolab DS5) was used to measure close to each installation water temperature (°C) and salinity (PSU) in triplicate. Identical sampling procedures were repeated monthly until January 2022.

## 2.3. Laboratory procedures

Collected brachyurans were identified to the species level; blue crabs were enumerated and sexed according to the shape of the abdominal tergites (the “apron”) [22]. The morphological

maturity status of individuals was determined by inspecting the apron [62,63]: males were recognized as mature when they had an inverted T-shaped apron, while mature females had round aprons, carried eggs or a mat of remnant brood setae. Individuals of both sexes showing triangular-shaped aprons adhering to thoracic sternites were identified as juveniles. In addition, for ovigerous females the developmental stage of the eggs based on a three-level colour scale (i.e., stage 1 = orange, stage 2 = brown, and stage 3 = black) was determined using the Virginia Marine Resources Commission colour coding system (<http://www.mrc.state.va.us/regulations/sponge.shtm>).

Consequently, for each blue crab the cephalothorax width (i.e., the distance between the two outermost anterolateral spines of the cephalothorax; CW hereafter) and length (i.e., the distance between the centre of the anterior interorbital margin and the centre of the posterior margin; hereafter), were measured to the nearest mm using a vernier caliper.

#### 2.4. Climatic parameters

Air temperature and precipitation data measured at 4 m above the ground level were downloaded from the website of the Agenzia Regionale per la Prevenzione e la Protezione dell'Ambiente of the Puglia Region (ARPA-Puglia; <http://www.webgis.arpa.puglia.it/lizmap/index.php/view/map?repository=1&project=meteo>; accessed on August 11<sup>th</sup>, 2024). Data were obtained from the climatic station in the city of San Severo (FG), located approximately 20 km southward of the Lesina Lagoon (41.695006N, 15.379159E). Hourly air temperatures (in °C) and precipitation (in mm) were obtained for the period February 1<sup>st</sup>, 2021 - January 31<sup>st</sup>, 2022. For the sake of clarity, temperatures were averaged per month; precipitation data were cumulated over the same temporal resolution.

#### 2.5. Data analysis

When not stated otherwise, values in the text are expressed as mean  $\pm$  1 SE. All data treatments and statistical procedures were implemented in the R package [64]. For univariate parametric analyses, data normality and homoscedasticity were preliminarily checked using Shapiro-Wilks and Levene tests, respectively; when necessary, data were log- or square root-transformed to meet the assumptions.

Parameters repeatedly determined during the investigation (e.g., water parameters or CPUEs) were treated as non-independent, time-correlated variables; accordingly, differences between sampling months were tested using paired t-tests. Differences in ratios and deviations from 1:1 were tested for significance using  $\chi^2$  tests. Statistical significance was evaluated at  $\alpha = 0.05$ ; when tests were reiterated,  $\alpha$  values were adjusted by performing sequential Benjamini-Hochberg corrections for multiple tests to reduce the risk of a type-I error [65].

##### 2.5.1. Abundance

The overall abundance of blue crabs recorded during the investigation was estimated by cumulating the number of captured individuals for each site over the different sampling times on a monthly basis. Data were ultimately normalized by the total number of fyke nets per site (3) and total soak time (in h), and expressed in Catch Per Unit Effort (CPUE hereafter) as N. of individuals fyke net<sup>-1</sup> d<sup>-1</sup>.

In June and July, when CPUE values reached maximum values (see Results), the spatial abundance of blue crabs was estimated for each of the four sampling sites using the Carle-Strub depletion method [66]. The method overcomes the limitations characterizing other approaches such as the Zippin removal method [67] when captures from passes two or three are higher than the first pass. The Carle-Strub method was implemented using the FSA R package [68] on four consecutive catch data with soak time = 24 h, and normalized by the expected area of action of each fixed installation (i.e., 20 m  $\times$  9.5 m = 190 m<sup>2</sup>; Figure 1c).

##### 2.5.2. Size cohorts

The occurrence of distinct cohorts in the size distribution of the blue crabs collected over the whole investigation was verified using the Bhattacharya's method [69]. The procedure was

implemented in the *TropFishR* R package [70,71] on CW data by fitting Gaussian components to mode means, and decompose the size-frequency distribution into a series of normal curves representing the cohorts. Modes were accepted as distinct cohorts if their separation index was above a critical value of 2.0 [72].

### 2.5.3. Growth dynamics and mortality

In temperate regions, seasonality significantly affects the growth of decapod crustaceans, including the blue crab [73,74]. Here we used Somers' seasonalised version of the von Bertalanffy growth function (sVBGF hereafter) to calculate the growth parameters of the crab in the study area [75,76]:

$$CW_t = CW_\infty \left[ 1 - e^{\left[ -K(t-t_0) + \left( \frac{C}{2\pi} \right) \sin 2\pi(t-t_s) - \left( \frac{C}{2\pi} \right) \sin 2\pi(t_0-t_s) \right]} \right]$$

where  $CW_t$  is the carapace width at age  $t$  (in mm),  $CW_\infty$  is the asymptotic carapace width (in mm),  $K$  is the annual growth coefficient ( $\text{yr}^{-1}$ ),  $t_0$  is the theoretical time of the year when  $CW = 0$  and spawning occurs [77],  $t_s$  is the time of the year when the growth rate is highest,  $C$  is the magnitude of the seasonal variation in growth, varying between 0 (no seasonal variation) and 1 (maximum variation).

Growth parameters were estimated using the *TropFishR* R package [70,71]. Specifically, CW frequency data were aggregated at a monthly level, binned into 5 mm size class intervals, and restructured after a square root-transformation to identify unobservable frequency peaks [78] adopting a moving average span  $M$  of 9, 7, and 5 for the total number of individuals, and for females and males separately.  $M$  was determined considering the number of bins spanning the width of the smallest-sized cohort observed during the investigation (see Results)[79]. The Electronic Length Frequency ANalysis (ELEFAN) procedure was implemented using an advanced optimization approach based on a genetic algorithm and 1,000 resamples [79]; the procedure has been already proven reliable for other brachyuran species [80,81]. The fit of the models was assessed by calculating the fitting score index  $R_n$  as:

$$R_n = 10^{ESP/ASP/10}$$

where  $ESP$  is the estimated sum of peaks, computed by summing the scored bins intersected by the sVBGF curve, and  $ASP$  is the available sum of peaks, computed as the maximum sum of positive scored bins a single curve could get [78]. The procedure allowed the estimation of the parameters of the sVBGF function (i.e.,  $CW_\infty$ ,  $K$ ,  $t_0$ ,  $t_s$ , and  $C$ ), the maximum age, and the growth performance index ( $\Phi'$ ) computed as:

$$\Phi = \text{Log}(K) + 2 \times \text{Log}(CW_\infty)$$

We estimated the total instantaneous mortality rate  $Z$  adopting the length-converted catch curve analysis and the graphical approach proposed by Pauly [82] using the CW frequency data. To calculate an approximation of the natural mortality  $M$ , due to e.g., predation or senescence, we used the sVBGF growth parameters  $CW_\infty$  and  $K$  [83], while the instantaneous fishing mortality rate,  $F$ , was estimated as  $F = Z - M$ . the exploitation rate  $E$  was calculated as  $E = F/Z$  [84].

### 2.5.4. Size at maturity


To determine the carapace width at 50% morphometric maturity ( $L_{50}$ ) in male and female crabs we compared two different procedures. First we directly fitted a binomial logistic regression model to CW measurements and maturity stage as classified by the inspection of the apron using the R package *AquaticLifeHistory* [85]. The second procedure was indirect and assumed the occurrence of statistically significant breaks in allometric relationships related with morphometric maturity. In the blue crabs, the morphological maturity is known to be associated with variations in the shape of the carapace in terms of length/width ratio [86,87]. Here we used the R package *sizeMat* [88] to classify the individuals in two groups (immature and mature) using a Principal Components Analysis with two allometric variables ( $CW$  = independent variable,  $CL$  = dependent variable) in log base. Individuals were assigned to each group using Ward's hierarchical clustering procedure applied to Euclidean distances [89]. Subsequently, a linear discriminant analysis was performed to classify individuals based on their CW and CL values. The  $L_{50}$  was ultimately estimated using a frequentist

logit procedure. Segmented regression was applied to identify breakpoints and confidence intervals at which biometric data indicated morphometric maturity in relation to the allometric relationships [88].

### 3. Results

#### 3.1. Water temperature and salinity

Water temperatures in the study area varied three-fold during the investigation, ranging between  $10.4 \pm 0.1$  °C (January 2022) and  $30 \pm 0.4$  °C (July 2021), and generally mirroring the atmospheric seasonality (Figures 2a and 2b; Pearson  $R^2 = 0.96$ ,  $P < 0.0001$ , 10 d.f.). In particular, during winter months (December, January, and February) similar temperatures lower than  $11.1 \pm 0.1$  °C were recorded (max  $t = 1.86$ ,  $P = 0.16$ , 3 d.f. for the comparison December 2021 vs. January 2022). In summer, temperatures measured in June ( $26.7 \pm 0.3$  °C) were significantly lower than in July and August 2021 ( $30 \pm 0.4$  °C and  $29.6 \pm 0.5$  °C respectively; min  $t = 9.24$ ,  $P = 0.003$ , 3 d.f. for the comparison June vs. July) the latter two months showing no significant differences ( $t = 1.27$ ,  $P = 0.29$ , 3 d.f.).



**Figure 2.** (a) Variation in mean ( $\pm 1SE$ ,  $n = 4$ ) water temperature (°C) and salinity (PSU) in the study area during the investigation. (b) Air temperatures (°C) and precipitations (mm) measured at the ARPA-Puglia climatic station located in the city of San Severo (FG, Italy) during the study period. See the text for additional details. Air temperatures are expressed as monthly means of hourly measurements  $\pm 1SE$  (boxes); whiskers are 95% confidence intervals. Precipitation data are cumulated at a monthly scale.

In contrast, considerable temperature shifts occurred in spring when values doubled from  $11.1 \pm 0.2$  to  $24.3 \pm 0.3$  °C between March and May (min  $t = 9.01$ ,  $P = 0.003$ , 3 d.f. for the comparison March vs. April). Similarly, in autumn temperatures dropped significantly from  $21 \pm 0.1$  °C to  $13.8 \pm 0.1$  °C between September and November (min  $t = 4.65$ ,  $P = 0.02$ , 3 d.f. for the comparison October vs. November).

Water salinities ranged between  $16.1 \pm 1.2$  PSU (April) and  $29.6 \pm 1.5$  PSU (August) (Figure 2a). A nearly significant co-variation was observed with temperature (Pearson  $r = 0.56$ ,  $P = 0.06$ , 10 d.f.); however, a less-pronounced seasonality was observed. In August, salinities were significantly higher than those observed in all the remaining months (min  $t = 4.19$ ,  $P = 0.02$ , 3 d.f. for the comparison with November) with the exception of July ( $24.6 \pm 2.1$  PSU;  $t = 2.67$ ,  $P = 0.07$ , 3 d.f.). In April the lowest salinities were measured, close to those determined in May ( $17.1 \pm 1.4$  PSU;  $t = 1.41$ ,  $P = 0.25$ , 3 d.f.). June, July, and November showed similar values (max  $t = 2.08$ ,  $P = 0.13$ , 3 d.f., for June vs. November) ranging between  $23.5 \pm 1.7$  PSU (June) and  $25 \pm 1.4$  PSU (November), significantly higher than the values determined in September, October, and December, when negligible differences were observed (max  $t = 3.02$ ,  $P = 0.06$ , 3 d.f. September vs. October). February, March, and January showed similar values (max  $t = 1.38$ ,  $P = 0.26$ , 3 d.f.), ranging between  $18.9 \pm 0.7$  PSU (March) and  $19.7 \pm 0.1$  PSU

(January). In winter, significant differences were observed only between December 2021 and January 2022 or February 2021 (min  $t = 5.96$ ,  $P = 0.03$ , 3 d.f. for the comparison October vs. November).


### 3.2. Abundance patterns

Overall, 838 blue crabs were collected during the investigation (Table 1). Mean total CPUE estimations ranged between 0 in December 2021 and January 2022 when no specimens were captured, and  $1.76 \pm 0.37$  ( $\pm 1SE$ ) crabs fyke nets<sup>-1</sup> d<sup>-1</sup> in July.

**Table 1.** Number of blue crabs captured during the investigation. Mean carapace width and lengths are included; standard errors are in parentheses while min-max ranges are in square brackets.

	<b>Total</b>	<b>Females</b>	<b>Males</b>
N. individuals	838	427	411
Carapace width (mm)	146.4 (1.4) [14 - 245]	150.6 (1.6) [20 - 230]	142.1 (2.3) [14 - 245]
Carapace length (mm)	66.4 (0.6) [5 - 120]	66.3 (0.6) [11 - 90]	66.4 (0.9) [5 - 120]

Overall, total catches followed a seasonal pattern (Figure 3), significantly correlated with water temperature (Pearson  $R^2 = 0.57$ ,  $P = 0.01$ , 10 d.f.), but not with salinity ( $R^2 = 0.34$ ,  $P = 0.28$ ). CPUEs progressively increased from February ( $0.15 \pm 0.02$  crabs fyke nets<sup>-1</sup> d<sup>-1</sup>) to maximum, similar values in June ( $1.51 \pm 0.69$  crabs fyke nets<sup>-1</sup> d<sup>-1</sup>) and July ( $1.76 \pm 0.37$  crabs fyke nets<sup>-1</sup> d<sup>-1</sup>;  $t = 0.31$ ,  $P = 0.78$ , 3 d.f.). May showed low, apparently off-pattern CPUEs compared with those determined in April and June ( $0.3 \pm 0.2$  vs.  $1.3 \pm 0.4$  and  $1.5 \pm 0.7$  crabs fyke nets<sup>-1</sup> d<sup>-1</sup>, respectively; Figure 3); however, no statistically significant differences were observed (max  $t = 2.71$ ,  $P = 0.07$ , 3 d.f., for the comparison May vs. June). Subsequently, a progressive decrease occurred from August ( $1.18 \pm 0.1$  crabs fyke nets<sup>-1</sup> d<sup>-1</sup>) to November ( $0.1 \pm 0.01$  crabs fyke nets<sup>-1</sup> d<sup>-1</sup>), when CPUE values showed values similar to February ( $t = 0.57$ ,  $P = 0.61$ , 3 d.f.). As observed for total catches, CPUEs of females and males showed a remarkable seasonal pattern of variation (Figure 3) and a significant co-variation with water temperatures was observed (Pearson  $R^2 = 0.66$ ,  $P = 0.002$  and  $R^2 = 0.34$ ,  $P = 0.04$  for males and females respectively; 10 d.f.). Noticeably, while females resulted completely independent from salinity ( $R^2 = 0.003$ ,  $P = 0.85$ ), males showed a nearly significant relationship ( $R^2 = 0.31$ ,  $P = 0.06$ ).



**Figure 3.** Variation in mean (+1SE,  $n = 4$ ) monthly catches per unit effort (CPUE, measured as N. individuals fyke nets<sup>-1</sup> d<sup>-1</sup>) of blue crabs determined during the investigation. CPUE values are reported for sexes combined and for the two sexes separately.

An overall spatial abundance of 0.21 blue crabs per square meter ( $\pm 0.07$  SE,  $n = 8$ ) was estimated between June and July using the Carle-Strub depletion method on consecutive catch data (Appendix A, Table A1).


**Table 2.** Mean spatial abundance of blue crabs (crabs m<sup>2</sup>; standard error in parenthesis) captured at the four sampling sites in June and July estimated using the Carle-Strub depletion method (see text for details).

Site	June	July
1	0.64 (0.11)	0.19 (0.04)
2	0.06 (0.01)	0.08 (0.01)
3	0.14 (0.01)	0.12 (0.04)
4	0.09 (0.02)	0.25 (0.04)

Estimations varied ten-fold across sampling sites and months between 0.06 and 0.64 crabs m<sup>-2</sup> (site 1 and 2 in June, respectively; Table 2) yet no significant effect of the factor “month” was observed ( $t = 0.56$ ,  $P = 0.61$ , 3 d.f.).

### 3.3. Sex ratio and ovigerous females

Of the 838 blue crabs captured during the investigation, 411 were males and 427 females (Table 1), with a total sex ratio of 0.96:1 ( $\sigma:\text{♀}$ ). The sex ratio remained close to 1 in February, March, June, July, October, and November (max  $\chi^2 = 2.98$ ,  $P = 0.08$ ; Figure 4a), while males dominated only in August ( $\sigma:\text{♀} = 3.05$ ;  $\chi^2 = 13.63$ ,  $P = 0.0002$ ) and September ( $\sigma:\text{♀} = 2.55$ ;  $\chi^2 = 4.51$ ,  $P = 0.03$ ). Conversely, females outnumbered males in April ( $\sigma:\text{♀} = 0.61$ ;  $\chi^2 = 6.52$ ,  $P = 0.01$ ) and May ( $\sigma:\text{♀} = 0.41$ ;  $\chi^2 = 4.86$ ,  $P = 0.03$ ).




**Figure 4.** (a) Sex ratio of blue crabs captured during the investigation. The blue dashed line identifies the 1:1 ratio. Red bars indicate significant departures from a 1:1 ratio ( $\chi^2$  test,  $P < 0.05$  after correction for multiple tests) indicated by the blue dashed line. Statistical results for February and November 2021 should be considered with caution given the low number of total specimens collected (<10). (b) Number of ovigerous females at three colour-determined egg stages (stage 1 = yellow; stage 2 = brown; stage 3 = black) captured during the investigation.

Ovigerous females ( $N = 45$ ) were captured for the first time in April ( $N = 13$ ), when a prevalence of individuals at egg stage 1 ( $N = 12$ ) and only one female at egg stage 3 were collected (Figure 4b). The highest number of ovigerous females were captured in May ( $N = 16$ ) when females at egg stage 2 and 3 dominated over those at egg stage 1 ( $N = 5$  and  $9$  vs.  $N = 2$ , respectively). The number of ovigerous females progressively decreased in the following summer months until August, with an increase in the dominance of individuals at stage 3; no ovigerous females were collected in the autumn and winter months (Figure 4b).


### 3.4. Size structure

Overall, blue crabs captured during the investigation showed a mean CW ( $\pm 1SE$ ) of  $146.43 \pm 1.39$  mm, with values ranging between 14 mm and 245 mm (Table 1). The mean CW of females ( $150.93 \pm 30.6$  mm; Table 1), was significantly larger than that of males ( $142.86 \pm 45.11$  mm; t-test with separate variance estimates:  $t = 3.07$ ,  $P = 0.002$ , 747.77 d.f.). Conversely, males showed a higher variability in size, with values varying between 14 and 245 mm ( $\Delta = 231$  mm) while females ranged between 20 and 230 mm ( $\Delta = 210$  mm; Table 1). The analysis of the CW-frequency distribution (Figure 5) of the total number of captured blue crabs highlighted the occurrence of three main modes at CW values corresponding to  $25 \pm 0.67$  mm (Age 0),  $95.62 \pm 1.45$  mm (Age 1), and  $171.46 \pm 1.61$  mm (Age 2+) (mean  $\pm 1SD$ ). Small-sized individuals represented by the first mode showed a non-seasonal patterns, as they occurred mainly in February and in November but were also captured in May and June (Figure 6). Medium-sized individuals included in the second mode occurred mainly in summer between June and July, while large-sized crabs belonging to the third mode were captured in spring with a peak in April and throughout the summer and autumn months (Figure 6).

The first mode was not recognized in females, given the exiguous number of small-sized individuals with CW  $< 50$  mm ( $N = 5$ ) captured only in February and June (Figure 6); thus, only two modes at CW =  $111.37 \pm 1.95$  mm and  $169.06 \pm 1.11$  mm were identified. Conversely, 14 small-sized males with CW values  $< 50$  mm were collected mainly in February and episodically in May, June, September and November (Figure 6); accordingly, their frequency distribution was characterized by three modes at  $2.01 \pm 0.49$  mm,  $104.94 \pm 2.81$  mm, and  $179.97 \pm 1.75$  mm. No remarkable sex-related differences in the monthly frequency distributions were evident for medium- and large sized crabs belonging to the second and third modal classes (Figure 6).



**Figure 5.** Frequency distributions of the carapace widths of blue crabs captured during the investigation for sexes combined (a), females only (b) and males only (c). Frequency distribution histograms include the modes identified by the modal analysis performed using the Bhattacharya method (bottom plots). See text for additional details.



**Figure 6.** Monthly frequency distributions of the carapace widths of blue crabs captured during the investigation expressed in terms of sexes combined (a), females (b), and males (c).

### 3.5. Morphological maturity

On the basis of the visual classification performed on the shape of the apron, 92 females and 122 males were identified as immature, representing 21.54% and 29.68% of the total number of captured individuals as per respective sex. Immature and mature females were characterized by a mean CW ( $\pm 1$ SE) of  $99.03 \pm 2.66$  mm and  $164.79 \pm 17.86$  mm respectively, while the mean CW of immature and mature males were  $81.89 \pm 2.06$  mm and  $167.48 \pm 1.45$  mm. Immatures represented in February almost the totality of the catches independently from the sex; subsequently, the number of immature individuals peaked in June and July in both sexes (Table 3); in June, in particular, they

represented 63.16% of the total number of captured females and 87.38% of males. Contributions > 50% were also observed in August for females and in November for males (Table 3).

**Table 3.** Number of immature female and male blue crabs as identified by the shape of the apron collected during the investigation. The contribution (in %) to the total captures as per the respective sexes are reported in parentheses.


Month	Females	Males
February 2021	4 (80)	6 (100)
March	1 (3.13)	4 (14.29)
April	1 (0.81)	1 (1.33)
May	2 (5.41)	4 (28.57)
June	48 (63.16)	90 (87.38)
July	24 (28.24)	10 (14.71)
August	11 (52.38)	---
September	---	2 (6.06)
October	---	---
November	1 (25)	5 (55.56)
December	---	---
January 2022	---	---

The mean carapace width at morphological maturity ( $L50_{Apron}$ ) of female and male blue crabs was estimated using a bootstrapped binomial model at 122.1 mm and 110.6 mm, respectively (Appendix B, Figure B1; see also Table 4 for SE and 95% bootstrapped CI).


Similar mean CL values were observed for females and males (Table 1; t-test with separate variance estimates,  $t = 0.09$ ,  $P = 0.92$ , 703.07 d.f.). The classification procedure performed on CW-CL data successfully identified in both female and male blue crabs significant discontinuities that allowed an effective discrimination of two groups of individuals (Figures 7 and 8).  $L50$  values calculated on the basis of the results of the analysis ( $L50_{CW-CL}$ ) were fully consistent with  $L50_{Apron}$  estimations, resulting in 123.1 mm for females and in 112.3 mm for males (Figures 8a and 8b), fully overlapping in terms of 95% bootstrapped confidence intervals (Table 4).

**Table 4.** Mean carapace widths at morphological maturity of female and male blue crabs (SE in square brackets; 95% bootstrapped confidence intervals are included) estimated through the visual classification performed on the shape of their apron ( $L50_{Apron}$ ) and through the analysis of carapace width-length relationships ( $L50_{CW-CL}$ ). The total variance explained by the logistic models with a binomial error structure fitted to the data ( $R^2$ ) is reported.

	Sex	$R^2$	Mean (mm)	95% CI (mm)
$L50_{Apron}$	Females	0.82	122.1 [2.2]	115.7 - 125.8
	Males	0.93	110.6 [1.6]	107.2 - 113.9
$L50_{CW-CL}$	Females	0.95	123.1 [1.8]	120.2 - 126.5
	Males	0.96	112.3 [1.7]	110 - 114.8



**Figure 7.** (a) Relationship between carapace widths and lengths (in mm) in female blue crabs; linear regression models are fitted to individual data subjected to a classification procedure (see text for details). The regression equation and the total explained variance ( $R^2$ ) are provided for each linear model; the results of an ANCOVA testing for statistically significant differences in the slopes of the two models are included. (b) The mean size at maturity (L50) of female blue crabs estimated using a bootstrapped binomial model applied to individuals classified as mature or immature through the analysis of carapace width-length relationships ( $L50_{CW-CL}$ ). Shaded areas represent 95% bootstrapped confidence intervals. L50 values estimated using a Bayesian procedure are reported in square brackets.



**Figure 8.** (a) Relationship between carapace widths and lengths (in mm) in male blue crabs; linear regression models are fitted to individual data subjected to a classification procedure (see text for details). The regression equation and the total explained variance ( $R^2$ ) are provided for each linear model; the results of an ANCOVA testing for statistically significant differences in the slopes of the two models are included. (b) The mean size at maturity (L50) of male blue crabs estimated using a bootstrapped binomial model applied to individuals classified as mature or immature through the analysis of carapace width-length relationships ( $L50_{CW-CL}$ ). Shaded areas represent 95% bootstrapped confidence intervals. L50 values estimated using a Bayesian procedure are reported in square brackets.

### 3.5. Growth and mortality

Table 5 summarizes the seasonal von Bertalanffy growth parameters estimated from carapace width-frequency distribution data of female and male blue crabs (Appendix C, Figure C1).

**Table 5.** Seasonalised Von Bertalanffy growth parameters determined for female and male blue crabs in the Lesina Lagoon. Estimations for sexes combined are included.

Sex	$t_s$	$CW_{\infty}$ (mm)	Age max (y)	K ( $y^{-1}$ )	$t_0$	C	$\Phi'$	Rn
Females	0.51	237.8	5	0.63	0.53	0.62	4.55	0.77
Males	0.52	232.6	8	0.36	0.75	0.57	4.29	0.58
Total	0.84	236.7	4.8	0.66	0.62	0.42	4.57	0.87

Females showed a larger  $CW_{\infty}$  (237.78 mm CW) than males (232.56 mm CW); similarly, the growth coefficient K was higher in females (0.63) than males (0.36). A seasonal growth oscillation was more evident in females than in males ( $C = 0.62$  vs.  $0.57$ ). The start of the slowest growth period was estimated in January for males ( $WP = 1$ ). The growth performance index  $\Phi'$  was higher for females than for males ( $\Phi' = 2.55$  vs.  $2.31$ ). The overall potential lifespan of the population was assessed in 4.8 years (Table 2), yet a remarkable sex-related difference was observed, as for females a maximum age of 5 years was estimated, as compared to 8 years for males.

A total instantaneous mortality rate  $Z$  of  $2.77 y^{-1}$  was estimated for the whole population, yet an almost three-fold difference was observed between sexes, with females showing the highest  $Z$  as compared with males (Table 6). Similarly, natural mortality rates  $M$  confirmed a sex-related differentiation; however, the largest dissimilarity occurred for the fishing mortality  $F$ , six times higher in females than in males (Table 6). Exploitation rates were estimated in 0.57 for the whole the population, with females twice as higher than in males ( $0.7$  vs.  $0.32$ ; Table 6).

**Table 6.** Estimated total, natural, and fishing instantaneous mortalities ( $Z$ ,  $M$ , and  $F$  respectively; in  $y^{-1}$ ) for blue crabs in the Lesina Lagoon, as referred to sexes combined, and for females and males alone. Estimations of exploitation rates ( $E$ ) are included.

Parameter	Total	Females	Males
Z	$2.77 \pm 0.33$	$3.88 \pm 0.43$	$1.28 \pm 0.14$
M	1.19	1.16	0.87
F	1.58	2.72	0.41
E	0.57	0.7	0.32

## 4. Discussion

### 4.1. Abundance, sex ratio, and size structure

The biology of *Callinectes sapidus* has long been investigated in its native range and a seasonally-dependent pattern has been generally identified, even though considerable latitudinal variations may occur in the timing of key events [90,91]. Temperature is a critical factor influencing the phenology of the crab throughout its life cycle [90,92,93]. In addition, to complete its life cycle the species requires both marine and brackish environments, with habitat use significantly affected by sex and ontogeny [22,94]. The results of our analyses on blue crab catches in the eastern sector of the Lesina Lagoon are coherent with this framework and reflect the interplay of seasonal variations in water temperature with the crab's behaviour and demography [94]. Indeed, the significant relationship observed between CPUE values and water temperature, the latter in turn influenced by local climate, confirms the results of a number of investigations carried out in native and invaded temperate coastal systems, ultimately identifying in seasonally-related variations in climatic conditions a driver of blue crab abundance fluctuations [6,48,95]. More specifically, in late spring (i.e., April and May) we observed a prevalence of large-sized adult females with a CW comprised between 150 and 200 mm, likely belonging to the third modal class presumably including individuals of age 2+ (Figure 5b and 7b). The female population was predominant, and the majority of individuals were in an ovigerous condition (Figure 4a and 4b). In the following months until September peak abundances of females were observed, with ovigerous females progressively shifting to different egg stages, decreasing in abundance to the point that the sex ratio  $\sigma:\text{♀}$  reached values between 2.5 and 3. During this period together with females of age 2+, also specimens from the second modal class of age 1+ (CW ranging between 80 and 130 mm) constituted a significant aliquot of the captures. In temperate environments,

while male blue crabs generally reside in estuaries and other brackish waters independently from their ontogenetic stage, in late spring adult females migrate after mating to neighbouring marine waters to spawn [23,94]. Once released, planktonic larvae (zoea stage) are dispersed offshore; subsequently, at the megalopa stage they return to brackish systems where they settle as juveniles and complete their life cycle [23,94]. In addition, females store sperm from a single mating event, and may have a lifetime reproductive potential of at least 8 broods distributed over more than 2 years [96–98]. Our observations conducted during the late spring and summer months corroborate the migration of two age classes of females inseminated in preceding years and prepared to leave the lagoon, as evidenced by their presence at the Schiapparo mouth. In June and July, when the maximum number of blue crabs was captured, spatial abundances were determined using the Carle-Strub depletion method on consecutive catch data. A number of efforts have been made to provide quantitative estimations of the spatial abundance of blue crab populations, and Taylor and Fehon [91] summarized available data for shallow-water, unvegetated, soft-bottom habitats from the Middle Atlantic, South Atlantic, and Gulf of Mexico. A considerable heterogeneity of results was observed, due to differences in local conditions, in the ontogenetic stage targeted in the investigations, and in sampling methodologies. However, summer densities ranged between 0.1 and 0.5 crabs  $m^{-2}$  on average, with maxima between 0.12 and 2 crabs  $m^{-2}$ . Our estimations were consistent with these data, as an overall spatial abundance of 0.21 crabs  $m^{-2}$  was determined, even though estimations varied ten-fold across sampling sites and months between 0.06 and 0.64 crabs  $m^{-2}$ .

In the following autumn months, the steep decrease in water temperature was mirrored by a decrease in catches, motivated by the departure of mated females but also by the movement of juveniles and mature males shifting to deeper waters to overwinter [6,94]. Indeed, the average water depth of the Lesina Lagoon is 0.7 m yet the eastern sector is generally shallower, with maximum water depths of 1.6 m occurring in the central and western sectors [99]. Thus, it is likely that a partial redistribution of individuals towards other sectors of the lagoon might have taken place from September to early winter, when minimum CPUEs were observed in November. Catches in subsequent months reduced to zero when water temperatures fell under 11°C (Figure 2). Winter temperatures exert significant control over blue crab abundance and overall population dynamics of blue crabs [92,100]. For example, moulting, and thus growth ceases when the temperature falls below a minimum temperature threshold of 9–11°C [101,102]. Accordingly, at temperate mid-latitudes where winter temperatures regularly fall below this threshold, blue crabs overwinter burrowing in the sediment in a state of dormancy, when an increase in mortality rates can be observed [100,103–105]. The immediate consequence is that no crabs are captured using fishing gears whose effectiveness relies on active movement of the prey, as this is the case here. Noticeably, while dormancy in winter months appears to be the norm in temperate coastal systems in the United States such as the Chesapeake Bay, in invaded Mediterranean systems blue crabs have been shown to remain active, even though at low abundances, when water temperatures fall below 9–11°C [48]. Accordingly, here we observed in February 2021 at water temperatures close to 11°C (Figure 2) low abundances of juveniles belonging to the age 0 class and large-sized individuals from the age 2+ class (Figure 3 and 7). Indeed, here we compared abundance patterns against monthly temperature means, yet a more advanced analysis performed at a daily scale and based on degree-days might have provided a more accurate picture of the events occurring at the onset of spring. Moreover, a differential thermal sensitivity across age classes, ultimately reflecting in an asynchronous duration of the dormancy period might be hypothesised here, yet this specific aspect of the biology of the blue crab remains to date unexplored, deserving further investigations in both the native and invaded ranges.

The results of our analyses provided a clear and consistent picture of the seasonal dynamics of blue crabs in the Lesina Lagoon, yet some concluding methodological remarks are necessary. Here we used fyke nets adopting a fixed-site design, generally acknowledged to provide reliable results for population-scale studies on brachyurans [91]. In addition, compared with other fishing gears (e.g., baited traps), fyke nets are considered non-selective as they capture any specimen moving along the shoreline and is intercepted by the leader regardless of its size and sex [106,107]. It is evident that the differing movement behaviours between migrating females and less active males may have

determine an increase in catchability of the former, at least during spring and summer months. Further lagoon-wide investigations based on non-selective dredge or trawl surveys and adopting a stratified random design are necessary to corroborate our estimations of blue crab abundances, size structure, and sex ratio. Specifically:

1) here we had no information regarding the catchability of blue crabs by fyke nets, thus our estimations of abundances, both in terms of CPUE and spatial density by the Carle-Strub method, might be underestimated. The catchability of a target species by a given type of fishing gear depends critically on the absolute spatial density of the species itself [108]. Here the natural density of blue crabs was unknown, and we cautiously assumed it proportional to the catches;

2) overall, the CW frequency distribution of crabs had a tri-modal shape with three age classes assigned to age 0 (young-of-year crabs hatched during the preceding summer), age 1 (crabs one year old) and age 2+ (crabs two or more year old; Figure 5a). Blue crab populations consist of two-three year classes and possibly six to eight [62,109–111] as confirmed by the results of our seasonal von Bertalanffy analysis (Appendix C, Figure C1a). However, contrary to males here for females the age 0 class was not identified due to the paucity of sampled individuals. Since the sex ratio is presumed to be balanced for blue crabs in the juvenile stage [112,113], sex-related differences in movement behaviour might have influenced the catchability of fyke nets, ultimately biasing our size structure assessments;

3) we observed a sex ratio close to 1:1, with females dominating in April, May, and July (Figure 4a). Information from system-wide dredge surveys in the US performed in winter generally indicated balanced sex ratios [6,112]. Yet, a dominance of males has been reported in a variety of habitats along the Atlantic coast [84,85] and in the Mediterranean Sea [48,50,106,114–116], with an exception represented by studies performed in Turkey, where females prevailed [47,117]. As for the proceeding point, we argue that, beside local factors that might differently affect the survival of males and females (see further in this section), migration behaviour in females coupled with variations in catchability of different fishing gears may bias the estimation of sex ratios [118], representing the ultimate determinant of the disparity of results presented in the literature.

#### 4.2. Size at morphological maturity

The assessment of the size at which organisms reach sexual maturity represents a key step in fisheries management, as it sets the minimum size for individuals to breed at least once before being harvested, ultimately protecting the reproductive potential of exploited populations [119,120]. In addition, in invasive species management it may provide useful information for the implementation of effective actions for functional eradication and the identification of harvest-driven trait changes in the bioinvader of interest [121,122].

Here, the results of the analyses based on the visual inspection of the apron shape indicated for females a larger  $L_{50\text{Apron}}$  than males (122.1 mm vs. 110.6 mm). The relatively scant body of information available for the blue crab confirms the occurrence of dissimilarities between sexes consistent with our findings, even though with varying degree of differentiation [123–125]. A number of factors might be at the origin of the observed differences, including sex-specific strategies of energy allocation to somatic and reproductive tissues during growth [126], as well as a differential sensitivity to variations in local biotic and abiotic factors such as e.g., temperature, salinity or quality and availability of trophic resources [97,127]. Local environmental conditions may also be the determinants of the huge variation in the size at morphological maturity observed across locations in both sexes [47,48,123]; ultimately, fishing and overexploitation may further affect the size at maturity of females, as evidenced in native US systems [128].

Noticeably, we showed that  $L_{50\text{Apron}}$  values matched those calculated using the classification procedure based on the detection of discontinuities in CW-CL data ( $L_{50\text{CW-CL}}$ : 123.1 mm and 112.3 mm for females and males, respectively). In crustaceans, growth is essentially continuous, while the increase in external dimensions proceeds by a discontinuous series of moults mirrored by sudden changes in the relative growth rates of body parts [126]. The strongest variations usually coincide with morphometric sexual maturity when a metabolic and physiological trade-off between investment in somatic or reproductive tissues related with gonadal maturation occurs [129]. To our knowledge, this represents the first attempt to compare the two approaches for *Callinectes sapidus* (but see [130] for other congeners). Additionally, while several studies on *C. sapidus* as well as the

congeners *C. danae* and *C. ornatus* successfully adopted a similar allometric approach comparing carapace width measurements against those of chelipeds, or the abdomen [123,130–133], no attempt has been made to date to directly use CW-CL data. This is despite the fact that Newcombe and Gray as early as 1938 [134] reported on changes in the shape of the carapace in blue crabs with the onset of maturity, and that an abrupt variation in length/width ratio characterizes both females and males after the pubertal moult [63,86,87,134]. Here we did not performed a geometric analysis of the carapace shape in captured blue crabs as those presented in other morphometric studies on *C. sapidus* and congeners [135,136]. Yet, given that the blue crab is characterized by prominent lateral spines whose length relative to the total carapace width increases during growth [22,137], it is likely that the change in length/width ratio characterizing sexual maturity may coincide with a transition from a short- to a long-spined shape form [63].

The generality of our results must be necessarily confirmed by additional studies to verify the occurrence of morphological variability in carapace shape across different populations as that observed e.g., for *Carcinus maenas* [138]. However, we provided a clear and statistically robust indication that allometric analysis on CW-CL data has the potential to assess morphological maturity in blue crabs with an effectiveness similar to that provided by other conventional, time-consuming procedures based on the examination of the apron or measurement of body parts. 4.3. Growth and mortality

Sumer et al. [47] summarized von Bertalanffy growth parameters (i.e.,  $CW_{\infty}$ ,  $K$ ,  $t_0$ , and  $\Phi'$ ) for the blue crab obtained from a number of sources. These data should be taken with caution, as they were obtained on different sexes or sexes combined, and, most importantly, using mainly non-seasonal von Bertalanffy models. In this study we analysed the growth dynamics of the blue crab using a seasonal von Bertalanffy model, explicitly acknowledging the dependence of the phenology and demography of the species upon seasonal variations in local abiotic and biotic factors such as temperature and food availability [139]. Accordingly, here we discuss our results only considering studies adopting a similar methodological procedure. Of the three investigations to date available, one is a pond study performed in the USA [140] and two are on wild blue crab populations from invaded Turkish [141] and Egyptian waters [142] (Table 6). In comparison to the aforementioned references, our study showed for both sexes combined higher  $CW_{\infty}$  and potential longevity (Age max in Table 6), the lowest  $K$ , and intermediate  $\Phi'$  values (Table 4 and Table 6). The higher  $CW_{\infty}$  and potential longevity, and the lower  $K$  values were confirmed for females and males separately. The comparison evidently does not allow for the identification of any general pattern, yet some issues can be addressed in terms of differences between sexes within the Lesina Lagoon.

**Table 6.** Summary of literature information on seasonalised Von Bertalanffy growth parameters determined for the blue crab. Data on sexes combined and for female and male only are reported.

Sex	$CW_{\infty}$ (mm)	Age max (y)	$K$ ( $y^{-1}$ )	$t_0$	$C$	$\Phi'$	Ref.
Total	201.6	3.7	0.81	0.22	0	4.52*	[19]
	207.5	---	1.19/1.71	0.15-0.31	0.94/1.0	4.71-4.86	[140]
	226.4***	2.11	1.38	0.15	---	4.85*	[142]
Females	206.6	4	0.74	0.24	0	4.50*	[19]
	181.9	2.72**	1.1	0.85	0.93	4.52	[140]
	227.6***	2.07	1.51	0.15	---	4.89*	[142]
Males	209.5	6	0.5	0.36	0.2	4.34*	[19]
	230.1	3.49**	0.86	0.16	0.29	4.72	[140]
	215.5***	1.96	1.59	0.14	---	4.87*	[142]

\*  $CW_{\infty}$  originally expressed in cm; values were re-calculated using the equation  $F' = \text{Log}(K) + 2 \times \text{Log}(CW_{\infty})$ ; \*\* calculated as  $3/K$  [143]; \*\*\* calculated from  $CL_{\infty}$  data and CW-CL relationships provided in the publication

Summer represented the period with peak growth rates for both sexes ( $t_s = 0.51-0.52$ ), with an evident seasonal variation of growth for females ( $C = 0.62$ ) and males ( $C = 0.57$ ) comprised within July and August. Noticeably, in agreement with previous studies in US waters reporting longevities varying between 3 and 8 years in relation with local environmental conditions and exploitation pressures [110,111,144,145], our estimation of the potential longevity of blue crabs was 4.8 years for

both sexes combined, with females showing a lower longevity than males (5 vs. 8 years). It can be hypothesized that higher predation rates during breeding migrations, higher sensitivity to low winter temperatures, and male-induced injury during copulation [62,103] may all contribute to an increase in the natural mortality rate of females (1.16 vs. 0.87; Table 5), and ultimately a reduction in their longevity. Females also showed a fishing mortality more than 6 times higher than males, that can be ascribed, as previously discussed, to the interplay of female-specific behaviour and fyke net catchability in the sector of the Lesina Lagoon where the present study was carried out. Accordingly, for females an exploitation rate twice as higher than that of males was determined (0.7 vs. 0.32). This depicts a condition definitely opposite to what is done in native areas to protect blue crab populations [146]. Thus, while the whole blue crab population in the Lesina Lagoon does not appear to be overexploited ( $E = 0.57$ ), our results indicate that the fishing activities focused in the areas close to connections with the sea may actually determine a remarkable impact on females. From a conservation perspective, this represents an undeniable positive condition, as the capture of mated females might effectively reduce the reproductive potential of the population and ultimately its abundance. In the perspective of exploiting the species as a shellfish product, however, it implies an evident risk of overfishing. A thorough assessment of the advantages and disadvantages of alternative strategies of exploitation of the blue crab in the lagoon is necessary, and we strongly recommend the implementation of an integrated strategy identifying an Optimum Blue Crab Yield (OBCY) taking as a reference what has been proposed for the invasive lionfish *Pterois* spp. in the Gulf of Mexico [147].

OBCY may set a target for catches of the entire population and for females only, exceeding the conventional maximum sustainable yield (MSY) but still providing a relatively high sustainable yield. This approach contributes to population suppression and the mitigation of its impacts on the recipient system, exceeding the achievable results of targets set at or below MSY.

## 5. Conclusions

For invasive marine species of economic interest, an in-depth understanding of the ecological and biological factors regulating their abundance and dynamics in recipient systems represents a mandatory precondition to integrate fishery management with functional eradication and impact mitigation. In the present study, we performed a comprehensive scrutiny of the biological characteristics and life history traits of the population of *Callinectes sapidus* in the Lesina Lagoon. At the same time, by using a suite of analytical approaches based on size-frequency data, we provided important clues on the fishery biology of the crab while verifying the effectiveness of alternative methodologies for estimating crucial parameters in fisheries management, such as size at morphological maturity. Lastly, we casted the results of the investigation in an integrated perspective acknowledging the need of controlling the abundance of the species to mitigate its ecological impact. We are confident that the present study may provide a useful basis for the implementation of standardised, cost-effective, Mediterranean-scale management plans focused on *C. sapidus* as well as other invasive portunid brachyurans such as *Portunus segnis* [148], matching exploitation with sustainable conservation and protection of coastal ecosystems.

**Author Contributions:** Conceptualization, G.M. and L.C.; methodology, G.M., N.L., and L.C.; formal analysis, G.M., R.D.G., L.D., E.M., L.P., and F.M.; investigation, N.L., T.S., O.A.L., and L.C.; data curation, L.C., T.S., N.L., A.O.L., and G.M.; writing-original draft preparation, G.M. and L.C.; writing - review and editing, all authors; funding acquisition, L.C. and G.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Apulia Region through Po-FEAMP Misura 1.44 (Art. 44, par. 1 lett. c del Reg. UE n. 508/2014) "CatchUpFish—Sviluppo di metodologie innovative per lo sfruttamento sostenibile delle risorse biologiche nella Laguna di Lesina". Additional financial support was provided to R.D.G. and E.M. by the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – Next Generation EU. Project code CN\_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP D33C22000960007, Project title "National Biodiversity Future Center - NBFC".

**Institutional Review Board Statement:** No ethical issues related with the use of animals in the performed analyses were involved.

**Informed Consent Statement:** Not applicable

**Data Availability Statement:** Data are available upon request from the corresponding author; they are not publicly available due to ongoing comparative analyses.

**Acknowledgments:** The authors wish to thank the associations of fishers operating in the Lesina Lagoon for their help during field operations.


**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## Appendix A

**Table A1.** Number of blue crabs captured in June and July on consecutive dates across the four sampling sites.


Month	Date	Site 1	Site 2	Site 3	Site 4
June	21/06/2021	34	5	11	5
	22/06/2021	21	3	8	6
	23/06/2021	15	1	4	4
	24/06/2021	17	2	2	1
July	14/07/2021	11	7	5	17
	15/07/2021	8	4	6	9
	16/07/2021	5	2	4	6
	17/07/2021	5	2	3	7

## Appendix B



**Figure B1.** The mean size at maturity (L50) of female (a) and male (b) blue crabs estimated using a bootstrapped binomial model applied to individuals classified as mature or immature by the shape of their apron. Shaded areas represent 95% bootstrapped confidence intervals.

## Appendix C



**Figure C1.** Restructured CW-frequency distributions with seasonal von Bertalanffy growth curves of blue crabs captured during the investigation expressed in terms of both sexes (a), females (b), and males (c).

## References

1. Williams, A.B. The swimming crabs of the genus *Callinectes* (Decapoda: Portunidae). *Fisheries Bulletin* **1971**, *72*, 685-798.
2. Johnson, D.S. The savory swimmer swims north: a northern range extension of the blue crab *Callinectes sapidus*? *J. Crust. Biol.* **2015**, *35*, 105-110.
3. Mancinelli, G.; Bardelli, R.; Zenetos, A. A global occurrence database of the Atlantic blue crab *Callinectes sapidus*. *Sci. Data* **2021**, *8*, 1-10.
4. Silliman, B.R.; Bertness, M.D. A trophic cascade regulates salt marsh primary production. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 10500-10505.
5. Boudreau, S.A.; Worm, B. Ecological role of large benthic decapods in marine ecosystems: a review. *Mar. Ecol. Prog. Ser.* **2012**, *469*, 195-213.
6. Sharov, A.F.; Vølstad, J.H.; Davis, G.R.; Davis, B.K.; Lipcius, R.N.; Montane, M.M. Abundance and exploitation rate of the blue crab (*Callinectes sapidus*) in Chesapeake Bay. *Bull. Mar. Sci.* **2003**, *72*, 543-565.
7. Mendonça, J.T.; Verani, J.R.; Nordi, N. Evaluation and management of blue crab *Callinectes sapidus* (Rathbun, 1896) (Decapoda-Portunidae) fishery in the Estuary of Cananéia, Iguape and Ilha Comprida, São Paulo, Brazil. *Braz. J. Biol.* **2010**, *70*, 37-45.
8. Mancinelli, G.; Chainho, P.; Cilenti, L.; Falco, S.; Kapiris, K.; Katselis, G.; Ribeiro, F. On the Atlantic blue crab (*Callinectes sapidus* Rathbun 1896) in southern European coastal waters: Time to turn a threat into a resource? *Fish. Res.* **2017**, *194*, 1-8.
9. Morales-Azpeitia, R.; Balmori-Ramírez, A.; Seefoo-Ramos, A.A.; García-Caudillo, J.M. Evaluation and estimation of reference points for the blue crab, *Callinectes sapidus* (Decapoda: Portunidae) of the Gulf of Mexico. *Hidrobiologica* **2021**, *31*, 231-243.
10. Kennedy, V.S.; Cronin, L.E. *The Blue Crab: Callinectes sapidus*; Maryland Sea Grant College: College Park, Maryland, **2007**; p. 774.
11. Villegas-Hernández, H.; Poot-López, G.R.; López-Rocha, J.A.; González-Salas, C.; Guillen-Hernández, S. Abundance and catchability estimates of the Atlantic blue crab *Callinectes sapidus* based on mark-recapture data from the northern Yucatan Peninsula. *J. Mar. Biol. Assoc. U.K.* **2017**, *98*, 1455-1463.
12. de Pasquier, G.A.; Pinto, L.G.; Buonocore, R.; Méndez, Y. Relaciones biométricas y proporción de sexos del cangrejo azul, *Callinectes sapidus* (Rathbun, 1896), en el Lago de Maracaibo, Venezuela. *Ciencia* **2012**, *20*, 151-158.
13. Lacerda, A.L.F.; Kersanach, R.; Cortinhas, M.C.S.; Prata, P.F.S.; Dumont, L.F.C.; Proietti, M.C.; Maggioni, R.; D'Incao, F. High connectivity among blue crab (*Callinectes sapidus*) populations in the Western South Atlantic. *PLoS ONE* **2016**, *11*, e0153124.

14. de Carvalho-Souza, G.F.; Medeiros, D.V.; Silva, R.d.A.; González-Ortegón, E. Width/length–weight relationships and condition factor of seven decapod crustaceans in a Brazilian tropical estuary. *Reg. Stud. Mar. Sci.* **2023**, *60*, 102880.
15. Cesar, I.I.; Armendáriz, L.C.; Olalla, N.; Tablado, A. The blue crab, *Callinectes sapidus* Rathbun, 1896 (Decapoda, Portunidae) in the Río de la Plata, Argentina. *Crustaceana* **2003**, *76*, 377-384.
16. Nehring, S. Invasion History and Success of the American Blue Crab *Callinectes sapidus* in European and Adjacent Waters. In *In the Wrong Place - Alien Marine Crustaceans: Distribution, Biology and Impacts*, Galil, B.S., Clark, P.F., Carlton, J.T., Eds.; *Invading Nature - Springer Series in Invasion Ecology*; Springer Netherlands: **2011**; Volume 6, pp. 607-624.
17. Cerri, J.; Chiesa, S.; Bolognini, L.; Mancinelli, G.; Grati, F.; Dragičević, B.; Dulčić, J.; Azzurro, E. Using online questionnaires to assess marine bio-invasions: a demonstration with recreational fishers and the Atlantic blue crab *Callinectes sapidus* (Rathbun, 1986) along three Mediterranean countries. *Mar. Pollut. Bull.* **2020**, *156*, 111209.
18. Castriota, L.; Falautano, M.; Perzia, P. When nature requires a resource to be used - The case of *Callinectes sapidus*: distribution, aggregation patterns, and spatial structure in northwest Europe, the Mediterranean Sea, and adjacent waters. *Biology* **2024**, *13*, 279.
19. Türeli, C.; Miller, T.; Gündoğdu, S.; Yesilyurt, I.N. Growth and mortality of blue crab (*Callinectes sapidus*) in the north-eastern Mediterranean Sea. *J FisheriesSciences.com* **2016**, *10*, 55-62.
20. Mancinelli, G.; Chainho, P.; Cilenti, L.; Falco, S.; Kapisir, K.; Katselis, G.; Ribeiro, F. The Atlantic blue crab *Callinectes sapidus* in southern European coastal waters: distribution, impact and prospective invasion management strategies. *Mar. Pollut. Bull.* **2017**, *119*, 5-11.
21. Mancinelli, G.; Guerra, M.T.; Alujević, K.; Raho, D.; Zotti, M.; Vizzini, S. Trophic flexibility of the Atlantic blue crab *Callinectes sapidus* in invaded coastal systems of the Apulia region (SE Italy): a stable isotope analysis. *Estuar. Coast. Shelf Sci.* **2017**, *198*, 421-431.
22. Millikin, M.R.; Williams, A.B. *Synopsis of biological data on blue crab, Callinectes sapidus Rathbun*; FAO Fisheries Synopsis 38: **1984**; p. 39.
23. Jivoff, P.; Hines, A.H.; Quackenbush, L.S. Reproduction Biology and Embryonic Development. In *The Blue Crab: Callinectes sapidus*, Kennedy, V.S., Cronin, L.E., Eds.; Maryland Sea Grant College: College Park, Maryland, **2007**; pp. 255-298.
24. Cilenti, L.; Paziienza, G.; Scirocco, T.; Fabbrocini, A.; D'Adamo, R. First record of ovigerous *Callinectes sapidus* (Rathbun, 1896) in the Gargano Lagoons (south-west Adriatic Sea). *Bioinvasions Rec.* **2015**, *4*, 281-287.
25. Johnson, D.R.; Perry, H.M. Blue crab larval dispersion and retention in the Mississippi Bight. *Bull. Mar. Sci.* **1999**, *65*, 129-149.
26. Png-Gonzalez, L.; Papiol, V.; Balbín, R.; Cartes, J.E.; Carbonell, A. Larvae of the blue crab *Callinectes sapidus* Rathbun, 1896 (Decapoda: Brachyura: Portunidae) in the Balearic Archipelago (NW Mediterranean Sea). *Mar. Biodivers. Rec.* **2021**, *14*, 21.
27. Havel, J.E.; Kovalenko, K.E.; Thomaz, S.M.; Amalfitano, S.; Kats, L.B. Aquatic invasive species: challenges for the future. *Hydrobiologia* **2015**, *750*, 147-170.
28. Grosholz, E.; Ashton, G.; Bradley, M.; Brown, C.; Ceballos-Osuna, L.; Chang, A.; de Rivera, C.; Gonzalez, J.; Heineke, M.; Marraffini, M.; et al. Stage-specific overcompensation, the hydra effect, and the failure to eradicate an invasive predator. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2003955118.
29. Green, S.J.; Grosholz, E.D. Functional eradication as a framework for invasive species control. *Front. Ecol. Environ.* **2021**, *19*, 98-107.
30. Cerveira, I.; Baptista, V.; Teodósio, M.A.; Morais, P. What's for dinner? Assessing the value of an edible invasive species and outreach actions to promote its consumption. *Biol. Invasions* **2022**, *24*, 815-829.
31. FAO. *Mid-term Strategy (2017-2020) towards the Sustainability of Mediterranean and Black Sea Fisheries*. Food and Agriculture Organization of the United Nations; Rome, **2017**; p. 24.
32. FAO. *Recommendation GFCM/42/2018/7 on a regional research programme on blue crab in the Mediterranean Sea*; Rome, **2018**; p. 5.
33. Clavero, M.; Franch, N.; Bernardo-Madrid, R.; López, V.; Abelló, P.; Queral, J.M.; Mancinelli, G. Severe, rapid and widespread impacts of an Atlantic blue crab invasion. *Mar. Pollut. Bull.* **2022**, *176*, 113479.
34. Rady, A.; Sallam, W.S.; Abdou, N.E.I.; El-Sayed, A.A.M. Food and feeding habits of the blue crab, *Callinectes sapidus* (Crustacea: Decapoda: Portunidae) with special reference to the gastric mill structure. *Egypt. J. Aquat. Biol. Fish.* **2018**, *22*, 417-431.
35. Mancinelli, G.; Glamuzina, B.; Petrić, M.; Carrozzo, L.; Glamuzina, L.; Zotti, M.; Raho, D.; Vizzini, S. The trophic position of the Atlantic blue crab *Callinectes sapidus* Rathbun 1896 in the food web of Parila Lagoon (South Eastern Adriatic, Croatia): a first assessment using stable isotopes. *Mediterr. Mar. Sci.* **2016**, *17*, 634-643.

36. Kampouris, T.E.; Porter, J.S.; Sanderson, W.G. *Callinectes sapidus* Rathbun, 1896 (Brachyura: Portunidae): An assessment on its diet and foraging behaviour, Thermaikos Gulf, NW Aegean Sea, Greece: evidence for ecological and economic impacts. *Crustac. Res.* **2019**, *48*, 23-37.
37. Prado, P.; Peñas, A.; Ibáñez, C.; Cabanes, P.; Jornet, L.; Álvarez, N.; Caiola, N. Prey size and species preferences in the invasive blue crab, *Callinectes sapidus*: Potential effects in marine and freshwater ecosystems. *Estuar. Coast. Shelf Sci.* **2020**, *245*, 106997.
38. Zotti, M.; Del Coco, L.; De Pascali, S.A.; Migoni, D.; Vizzini, S.; Mancinelli, G.; Fanizzi, F.P. Comparative analysis of the proximate and elemental composition of the blue crab *Callinectes sapidus*, the warty crab *Eriphia verrucosa*, and the edible crab *Cancer pagurus*. *Heliyon* **2016**, *2*, 1-15.
39. Tekin, S.; Pazi, I. POP levels in blue crab (*Callinectes sapidus*) and edible fish from the eastern Mediterranean coast. *Environ. Sci. Pollut. Res.* **2017**, *24*, 509-518.
40. Renzi, M.; Cilenti, L.; Scirocco, T.; Grazioli, E.; Anselmi, S.; Broccoli, A.; Pauna, V.; Provenza, F.; Specchiulli, A. Litter in alien species of possible commercial interest: The blue crab (*Callinectes sapidus* Rathbun, 1896) as case study. *Mar. Pollut. Bull.* **2020**, *157*, 111300.
41. Compa, M.; Perelló, E.; Box, A.; Colomar, V.; Pinya, S.; Sureda, A. Ingestion of microplastics and microfibers by the invasive blue crab *Callinectes sapidus* (Rathbun 1896) in the Balearic Islands, Spain. *Environ. Sci. Pollut. Res.* **2023**, *30*, 119329-119342.
42. El Qoraychy, I.; Rharrhour, H.; Hammani, O.; Wariaghli, F.; Touhami, F.; Jaziri, H. Heavy metal accumulation in blue crabs (*Callinectes Sapidus*) in Gharb Region, Morocco. *Egypt. J. Aquat. Biol. Fish.* **2023**, *27*, 695-705.
43. De Giorgi, R.; Bardelli, R.; Cilenti, L.; Falco, S.; Fanizzi, F.P.; Guerra, M.T.; Katselis, G.; Kevrekidis, K.; Mancini, F.; Doria, L.; et al. Opportunistic omnivory impairs the use of the Atlantic blue crab *Callinectes sapidus* as a trace metal biomonitor in invaded Mediterranean coastal waters. *Mar. Pollut. Bull.* **2024**, *206*, 116715.
44. Pagliara, P.; Mancinelli, G. Parasites affect hemocyte functionality in the hemolymph of the invasive Atlantic blue crab *Callinectes sapidus* from a coastal habitat of the Salento Peninsula (SE Italy). *Mediterr. Mar. Sci.* **2018**, *19*, 193-200.
45. Marangi, M.; Lago, N.; Mancinelli, G.; Lillo Antonio, O.; Scirocco, T.; Sinigaglia, M.; Specchiulli, A.; Cilenti, L. Occurrence of the protozoan parasites *Toxoplasma gondii* and *Cyclospora cayetanensis* in the invasive Atlantic blue crab *Callinectes sapidus* from the Lesina Lagoon (SE Italy). *Mar. Pollut. Bull.* **2022**, *176*, 113428.
46. Lattos, A.; Papadopoulos, D.K.; Giantsis, I.A.; Stamelos, A.; Karagiannis, D. Histopathology and phylogeny of the dinoflagellate *Hematodinium perezii* and the epibiotic peritrich ciliate *Epistylis* sp. infecting the blue crab *Callinectes sapidus* in the eastern Mediterranean. *Microorganisms* **2024**, *12*.
47. Sumer, C.; Teksam, I.; Karatas, H.; Beyhan, T.; Aydin, C.M. Growth and Reproduction Biology of the Blue Crab, *Callinectes sapidus* Rathbun, 1896, in the Beymelek Lagoon (Southwestern Coast of Turkey). *Turk J Fish Aquat Sci* **2013**, *13*, 675-684.
48. Kevrekidis, K.; Kevrekidis, T.; Mogias, A.; Boubonari, T.; Kantaridou, F.; Kaisari, N.; Malea, P.; Dounas, C.; Thessalou-Legaki, M. Fisheries biology and basic life-cycle characteristics of the invasive blue crab *Callinectes sapidus* Rathbun in the estuarine area of the Evros River (Northeast Aegean Sea, Eastern Mediterranean). *J. Mar. Sci. Eng.* **2023**, *11*, 462.
49. Katselis, G.N.; Koutsikopoulos, C. The Establishment of Blue Crab *Callinectes sapidus* Rathbun, 1896 in the Lagoon Pogonitsa (Amvrakikos Gulf, Western Greece). In *Trends in Fisheries and Aquatic Animal Health*, Berillis, P., Ed.; Bentham Science Publishers: Sharjah, United Arab Emirates, **2017**; pp. 299-306.
50. Kevrekidis, K.; Antoniadou, C. Abundance and population structure of the blue crab (Decapoda, Portunidae) in Thermaikos Gulf (Methoni Bay), northern Aegean Sea. *Crustaceana* **2018**, *91*, 641-657.
51. Marchessaux, G.; Mangano, M.C.; Bizzarri, S.; M'Rabet, C.; Principato, E.; Lago, N.; Veysiere, D.; Garrido, M.; Scyphers, S.B.; Sarà, G. Invasive blue crabs and small-scale fisheries in the Mediterranean sea: Local ecological knowledge, impacts and future management. *Mar. Policy* **2023**, *148*, 105461.
52. Glamuzina, L.; Pešić, A.; Marković, O.; Tomanić, J.; Pećarević, M.; Dobrosravić, T.; Brailo Šćepanović, M.; Conides, A.; Grđan, S. Population structure of the invasive Atlantic blue crab, *Callinectes sapidus* on the Eastern Adriatic coast (Croatia, Montenegro). *Nase More* **2023**, *70*, 153-159.
53. Florio, M.; Breber, P.; Scirocco, T.; Specchiulli, A.; Cilenti, L.; Lumare, L. Exotic species in Lesina and Varano lakes: Gargano National Park (Italy). *Transit Water Bull* **2008**, *2*, 69-79.
54. Cilenti, L.; Lago, N.; Lillo, A.O.; Li Veli, D.; Scirocco, T.; Mancinelli, G. Soft-shell production of the invasive Atlantic blue crab *Callinectes sapidus* in the Lesina Lagoon (SE Italy): a first assessment. *J. Mar. Sci. Eng.* **2024**, *12*, 310.
55. Mancinelli, G.; Rossi, L. Indirect, size-dependent effects of crustacean mesograzers on the Rhodophyta *Gracilaria verrucosa* (Hudson) Papenfuss: evidence from a short-term study in the Lesina Lagoon (Italy). *Mar. Biol.* **2001**, *138*, 1163-1173.

56. Roselli, L.; Fabbrocini, A.; Manzo, C.; D'Adamo, R. Hydrological heterogeneity, nutrient dynamics and water quality of a non-tidal lentic ecosystem (Lesina Lagoon, Italy). *Estuar. Coast. Shelf Sci.* **2009**, *84*, 539-552.
57. Caroppo, C.; Roselli, L.; Di Leo, A. Hydrological conditions and phytoplankton community in the Lesina lagoon (southern Adriatic Sea, Mediterranean). *Environmental Science Pollution Research* **2018**, *25*, 1784-1799.
58. Azzurro, E.; Cerri, J. Participatory mapping of invasive species: a demonstration in a coastal lagoon. *Mar. Policy* **2021**, *126*, 104412.
59. Specchiulli, A.; Scirocco, T.; D'Adamo, R.; Cilenti, L.; Fabbrocini, A.; Cassin, D.; Penna, P.; Renzi, M.; Bastianoni, S. Benthic vegetation, chlorophylla and physical-chemical variables in a protected zone of a Mediterranean coastal lagoon (Lesina, Italy). *J. Coast. Conservation* **2016**, *20*, 363-374.
60. FAO. Fishing Gear types. Fyke nets. Technology Fact Sheets. In: Fisheries and Aquaculture. Rome. <https://www.fao.org/fishery/en/geartype/226/en>. **2024**.
61. Scordella, G.; Lumare, F.; Conides, A.; Papaconstantinou, C. First occurrence of the tilapia *Oreochromis niloticus niloticus* (Linnaeus, 1758) in Lesina lagoon (eastern Italian coast). *Mediterr. Mar. Sci.* **2003**, *4*, 41-48.
62. Van Engel, W.A. The blue crab and its fishery in Chesapeake Bay. Part 1. Reproduction, early development, growth and migration. *Commercial fisheries review* **1958**, *20*.
63. Olmi III, E.J.; Bishop, J.M. Variations in total width-weight relationships of blue crabs, *Callinectes sapidus*, in relation to sex, maturity, molt stage, and carapace form. *J. Crust. Biol.* **1983**, *3*, 575-581.
64. R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>. **2024**.
65. Benjamini, Y.; Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Stat. Soc. Ser. B. (Stat. Method.)* **1995**, *57*, 289-300.
66. Carle, F.L.; Strub, M.R. A new method for estimating population size from removal data. *Biometrics* **1978**, *621-630*.
67. Zippin, C. An evaluation of the removal method of estimating animal populations. *Biometrics* **1956**, *12*, 163-189.
68. Ogle, D.H.; Doll, J.C.; Powell Wheeler, A.; Dinno, A. FSA: Fisheries Stock Analysis. R package version 0.9.5. <https://CRAN.R-project.org/package=FSA>. **2023**.
69. Bhattacharya, C.G. A simple method of resolution of a distribution into Gaussian components. *Biometrics* **1967**, *115-135*.
70. Mildenerger, T.K.; Taylor, M.H.; Wolff, M. TropFishR: an R package for fisheries analysis with length-frequency data. **2017**, *8*, 1520-1527.
71. Mildenerger, T.K.; Taylor, M.H.; Wolff, M. TropFishR: Tropical Fisheries Analysis. R package version 1.6.4. <http://cran.r-project.org/web/packages/TropFishR>. **2024**.
72. Gayanilo, F.C.J.; Sparre, P.; Pauly, D. FAO-ICLARM Stock Assessment Tools II (FiSAT II). Revised version. User's guide. FAO Computerized Information Series (Fisheries). No. 8, Revised version. **2005**, *8*.
73. Chang, Y.-J.; Sun, C.-L.; Chen, Y.; Yeh, S.-Z. Modelling the growth of crustacean species. *Rev. Fish Biol. Fish.* **2012**, *22*, 157-187.
74. Puckett, B.J.; Secor, D.H.; Ju, S.-J. Validation and application of lipofuscin-based age determination for Chesapeake Bay blue crabs *Callinectes sapidus*. *Trans Am Fish Soc* **2008**, *137*, 1637-1649.
75. Von Bertalanffy, L. A quantitative theory of organic growth (inquiries on growth laws. II). *Hum Biol* **1938**, *10*, 181-213.
76. Somers, I.F. On a seasonally oscillating growth function. *Fishbyte* **1988**, *6*, 8-11.
77. Türeli, C.; Miller, T.J.; Gündođdu, S.; Yesilyurt, I.N. Growth and mortality of blue crab (*Callinectes sapidus*) in the north-eastern Mediterranean Sea. *J FisheriesSciences.com* **2016**, *10*, 55-62.
78. Pauly, D.; David, N. ELEFAN I, a BASIC program for the objective extraction of growth parameters from length-frequency data. *Meeresforschung* **1981**, *28*, 205-211.
79. Taylor, M.H.; Mildenerger, T.K. Extending electronic length frequency analysis in R. **2017**, *24*, 330-338.
80. Çiğşar, B.; Deniz, Ü.; Türeli, C. Artificial neural networks and genetic algorithm approach to determine length-weight, length frequency relationships of Lessepsian crab, *Charybdis (Goniohellenus) longicollis*, Leene, 1938 in the Iskenderun Bay, Turkey. *Acta Biologica Turcica* **2021**, *34*, 197-204.
81. Guerra-Marrero, A.; Bonino-Pérez, A.; Espino-Ruano, A.; Couce-Montero, L.; Jiménez-Alvarado, D.; Castro, J.J. Life history parameters and fishing aspects of the alien nimble spray crab *Percnon gibbesi* in a native area of the Central-East Atlantic. *Animals* **2023**, *13*, 1427.
82. Pauly, D. *Some Simple Methods for the Assessment of Tropical Fish Stocks*; Food and Agriculture Organization of the United Nations: Rome, 1983.
83. Then, A.Y.; Hoenig, J.M.; Hall, N.G.; Hewitt, D.A. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES J. Mar. Sci.* **2014**, *72*, 82-92.
84. Gulland, J.A. *Fish Stock Assessment: a Manual of Basic Methods*; Wiley: Chichester, UK, 1983; Volume 1.

85. Smart, J. AquaticLifeHistory: Life History Analysis Tools. R package version 1.0.5. <http://cran.r-project.org/web/packages/AquaticLifeHistory>. **2023**.
86. Newcombe, C.L.; Campbell, F.; Eckstine, A.M. A study of the form and growth of the blue crab *Callinectes sapidus* Rathbun. *Growth* **1949**, *13*, 71-96.
87. Tagatz, M.E. Growth of juvenile blue crabs, *Callinectes sapidus* Rathbun, in the St. Johns River, Florida. *Fish. Bull.* **1968**, *67*, 281-288.
88. Torrejon-Magallanes, J. sizeMat: Estimate Size at Sexual Maturity. R package version 1.1.2. <http://cran.r-project.org/web/packages/sizeMat>. **2020**.
89. Corgos, A.; Freire, J. Morphometric and gonad maturity in the spider crab *Maja brachydactyla*: a comparison of methods for estimating size at maturity in species with determinate growth. *ICES J. Mar. Sci.* **2006**, *63*, 851-859.
90. Rogers, T.L.; Gouhier, T.C.; Kimbro, D.L. Distinct temperature stressors acting on multiple ontogenetic stages influence the biogeography of Atlantic blue crabs. *Mar. Ecol. Prog. Ser.* **2022**, *690*, 97-111.
91. Taylor, D.L.; Fehon, M.M. Blue crab (*Callinectes sapidus*) population structure in Southern New England tidal rivers: patterns of shallow-water, unvegetated habitat use and quality. *Estuar. Coasts* **2021**, *44*, 1320-1343.
92. Bauer, L.J.; Miller, T.J. Spatial and interannual variability in winter mortality of the blue crab (*Callinectes sapidus*) in the Chesapeake Bay. *Estuar. Coasts* **2010**, *33*, 678-687.
93. Schneider, A.K.; Fabrizio, M.C.; Lipcius, R.N. Reproductive phenology of the Chesapeake Bay blue crab population in a changing climate. *Front. Ecol. Evol.* **2024**, *11*, 1304021.
94. Hines, A.H. Ecology of Juvenile and Adult Blue Crabs. In *The Blue Crab: Callinectes sapidus*, Kennedy, V.S., Cronin, L.E., Eds.; Maryland Sea Grant College: College Park, Maryland, **2007**; pp. 565-654.
95. Weiss, H.M.; Downs, J.T. Living near the edge: variability in abundance and life cycle of the blue crab *Callinectes sapidus* (Rathbun, 1896) in eastern Long Island Sound. *J. Shellfish Res.* **2020**, *39*, 461-470.
96. Dickinson, G.H.; Rittschof, D.; Latanich, C. Spawning biology of the blue crab, *Callinectes sapidus*, in North Carolina. *Bull. Mar. Sci.* **2006**, *79*, 273-285.
97. Darnell, M.Z.; Rittschof, D.; Darnell, K.M.; McDowell, R.E. Lifetime reproductive potential of female blue crabs *Callinectes sapidus* in North Carolina, USA. *Mar. Ecol. Prog. Ser.* **2009**, *394*, 153-163.
98. Ogburn, M.B.; Richie, K.D.; Jones, M.A.; Hines, A.H. Sperm acquisition and storage dynamics facilitate sperm limitation in the selectively harvested blue crab *Callinectes sapidus*. *Mar. Ecol. Prog. Ser.* **2019**, *629*, 87-101.
99. Ferrarin, C.; Zaggia, L.; Paschini, E.; Scirocco, T.; Lorenzetti, G.; Bajo, M.; Penna, P.; Francavilla, M.; D'Adamo, R.; Guerzoni, S. Hydrological regime and renewal capacity of the micro-tidal Lesina Lagoon, Italy. *Estuar. Coasts* **2014**, *37*, 79-93.
100. Bauer, L.J.; Miller, T.J. Temperature-, salinity-, and size-dependent winter mortality of juvenile blue crabs (*Callinectes sapidus*). *Estuar. Coasts* **2010**, *33*, 668-677.
101. Brylawski, B.J.; Miller, T.J. Temperature-dependent growth of the blue crab (*Callinectes sapidus*): a molt process approach. *Can. J. Fish. Aquat. Sci.* **2006**, *63*, 1298-1308.
102. Smith, S.G.; Chang, E.C. Molting and Growth. In *The Blue Crab: Callinectes sapidus*, Kennedy, V.S., Cronin, L.E., Eds.; Maryland Sea Grant College: College Park, Maryland, **2007**; pp. 197-255.
103. Rome, M.S.; Young-Williams, A.C.; Davis, G.R.; Hines, A.H. Linking temperature and salinity tolerance to winter mortality of Chesapeake Bay blue crabs (*Callinectes sapidus*). *J. Exp. Mar. Biol. Ecol.* **2005**, *319*, 129-145.
104. Glandon, H.L.; Kilbourne, K.H.; Miller, T.J. Winter is (not) coming: warming temperatures will affect the overwinter behavior and survival of blue crab. *PLoS ONE* **2019**, *14*, e0219555.
105. Molina, A.I.; Cerrato, R.M.; Nye, J.A. Population level differences in overwintering survivorship of blue crabs (*Callinectes sapidus*): a caution on extrapolating climate sensitivities along latitudinal gradients. *PLoS ONE* **2021**, *16*, e0257569.
106. Glamuzina, L.; Conides, A.; Mancinelli, G.; Glamuzina, B. A comparison of traditional and locally novel fishing gear for the exploitation of the invasive Atlantic blue crab in the eastern Adriatic Sea. *J. Mar. Sci. Eng.* **2021**, *9*, 1019.
107. Audet, D.; Miron, G.; Moriyasu, M. Biological characteristics of a newly established green crab (*Carcinus maenas*) population in the southern Gulf of St. Lawrence, Canada. *J. Shellfish Res.* **2008**, *27*, 427-441, 415.
108. Miller, R.J. Effectiveness of crab and lobster traps. *Can. J. Fish. Aquat. Sci.* **1990**, *47*, 1228-1251.
109. Fischler, K.J. The use of catch - effort, catch - sampling, and tagging data to estimate a population of blue crabs. *Trans Am Fish Soc* **1965**, *94*, 287-310.
110. Rugolo, L.J.; Knotts, K.S.; Lange, A.M.; Crecco, V.A. Stock assessment of Chesapeake Bay blue crab (*Callinectes sapidus* Rathbun). *J. Shellfish Res.* **1998**, *17*, 493-518.
111. Rugolo, L.J.; Knotts, K.; Lange, A.; Crecco, V.; Terceiro, M.; Bonzek, C.; Stagg, C.; O'Reilly, R.; Vaughn, D. *Stock assessment of Chesapeake Bay blue crab (Callinectes sapidus)*; Chesapeake Bay Stock Assessment Committee, NOAA: Annapolis, MA, **1997**.

112. Lycett, K.A.; Shields, J.D.; Chung, J.S.; Pitula, J.S. Population structure of the blue crab *Callinectes sapidus* in the Maryland Coastal Bays. *J. Shellfish Res.* **2020**, *39*, 699-713.
113. Fitz, H.C.; Wiegert, R.G. Local population dynamics of estuarine blue crabs: abundance, recruitment and loss. *Mar. Ecol. Prog. Ser.* **1992**, 23-40.
114. Carrozzo, L.; Potenza, L.; Carlino, P.; Costantini, M.L.; Rossi, L.; Mancinelli, G. Seasonal abundance and trophic position of the Atlantic blue crab *Callinectes sapidus* Rathbun 1896 in a Mediterranean coastal habitat. *Rend. Lincei Sci. Fis. Nat.* **2014**, *25*, 201-208.
115. Mancinelli, G.; Carrozzo, L.; Marini, G.; Costantini, M.L.; Rossi, L.; Pinna, M. Occurrence of the Atlantic blue crab *Callinectes sapidus* (Decapoda, Brachyura, Portunidae) in two Mediterranean coastal habitats: temporary visitor or permanent resident? *Estuar. Coast. Shelf Sci.* **2013**, *135*, 46-56.
116. Abdel Razeq, F.A.; Ismaiel, M.; Ameran, M.A.A. Occurrence of the blue crab *Callinectes sapidus*, Rathbun, 1896, and its fisheries biology in Bardawil Lagoon, Sinai Peninsula, Egypt. *Egypt. J. Aquat. Res.* **2016**, *42*, 223-229.
117. Türeli, C.; Yeşilyurt, İ.N.; Nevşat, İ.E. Female reproductive pattern of *Callinectes sapidus* Rathbun, 1896 (Brachyura: Portunidae) in Iskenderun Bay, Eastern Mediterranean. *Zoology in the Middle East* **2018**, *64*, 55-63.
118. Hines, A.H.; Lipcius, R.N.; Haddon, A.M. Population dynamics and habitat partitioning by size, sex, and molt stage of blue crabs *Callinectes sapidus* in a subestuary of central Chesapeake Bay. *Mar. Ecol. Prog. Ser.* **1987**, *36*, 55-64.
119. Herrera, D.R.; da Costa, R.C. Reproductive traits, relative growth and maturity of blue crabs *Callinectes danae* and *Callinectes ornatus* in South Atlantic waters. *Aquat. Ecol.* **2024**, *58*, 963-982.
120. Mesquita, C.; Dobby, H.; Sweeting, S.; Jones, C.S.; Pierce, G.J. Size-at-maturity of Brown Crab (*Cancer pagurus*) in Scottish waters based on gonadal and morphometric traits. *Fish. Res.* **2020**, *229*, 105610.
121. Evangelista, C.; Britton, R.J.; Cucherousset, J. Impacts of invasive fish removal through angling on population characteristics and juvenile growth rate. *Ecol. Evol.* **2015**, *5*, 2193-2202.
122. Rytwinski, T.; Taylor, J.J.; Donaldson, L.A.; Britton, J.R.; Browne, D.R.; Gresswell, R.E.; Lintermans, M.; Prior, K.A.; Pellatt, M.G.; Vis, C.; et al. The effectiveness of non-native fish removal techniques in freshwater ecosystems: a systematic review. *Environ. Rev.* **2019**, *27*, 71-94.
123. Marchessaux, G.; Gjoni, V.; Sarà, G. Environmental drivers of size-based population structure, sexual maturity and fecundity: a study of the invasive blue crab *Callinectes sapidus* (Rathbun, 1896) in the Mediterranean Sea. *PLoS ONE* **2023**, *18*, e0289611.
124. Pereira, M.J.; Branco, J.O.; Christoffersen, M.L.; Freitas, F.; Fracasso, H.A.A.; Pinheiro, T.C. Population biology of *Callinectes danae* and *Callinectes sapidus* (Crustacea: Brachyura: Portunidae) in the south-western Atlantic. *J. Mar. Biol. Assoc. U.K.* **2009**, *89*, 1341-1351.
125. Severino-Rodrigues, E.; Musiello-Fernandes, J.; Mour, Á.A.S.; Branco, G.M.P.; Canéo, V.O.C. Fecundity, reproductive seasonality and maturation size of *Callinectes sapidus* females (Decapoda: Portunidae) in the Southeast coast of Brazil. *Rev. Biol. Trop.* **2013**, *61*, 595-602.
126. Hartnoll, R.G. Growth. In *The Biology of Crustacea*, Abele, L.G., Ed.; Academic Press: New York, **1982**; Volume Vol. 2. Embryology, morphology, and genetics, pp. 111-196.
127. Fisher, M.R. Effect of temperature and salinity on size at maturity of female blue crabs. *Trans Am Fish Soc* **1999**, *128*, 499-506.
128. Lipcius, R.N.; Stockhausen, W.T. Concurrent decline of the spawning stock, recruitment, larval abundance, and size of the blue crab *Callinectes sapidus* in Chesapeake Bay. *Mar. Ecol. Prog. Ser.* **2002**, *226*, 45-61.
129. Longo, E.; Mancinelli, G. Size at the onset of maturity (SOM) revealed in length-weight relationships of brackish amphipods and isopods: an information theory approach. *Estuar. Coast. Shelf Sci.* **2014**, *136*, 119-128.
130. dos Santos Cardim, R.W.; dos Reis Souza, E.; Júnior, M.S.; Carvalho, F.L.; Rocha, S.S.J.B.d.I.d.P. Sexual maturity of *Callinectes danae* and *C. ornatus* from Paraguaçu river estuary, Bahia state, Brazil. **2022**, *48*, 1-14.
131. Marochi, M.Z.; Moreto, T.F.; Lacerda, M.B.; Trevisan, A.; Masunari, S. Sexual maturity and reproductive period of the swimming blue crab *Callinectes danae* Smith, 1869 (Brachyura: Portunidae) from Guaratuba Bay, Paraná State, southern Brazil. *Nauplius* **2013**, *21*, 43-52.
132. Araújo, M.S.L.C.; Negromonte, A.O.; Barreto, A.V.; Castiglioni, D.S. Sexual maturity of the swimming crab *Callinectes danae* (Crustacea: Portunidae) at the Santa Cruz Channel, a tropical coastal environment. *J. Mar. Biol. Assoc. U.K.* **2012**, *92*, 287-293.
133. Haefner Jr, P.A. Morphometry and size at maturity of *Callinectes ornatus* (Brachyura, Portunidae) in Bermuda. *Bull. Mar. Sci.* **1990**, *46*, 274-286.
134. Gray, E.H.; Newcombe, C.L. Studies of moulting in *Callinectes sapidus* Rathbun. *Growth* **1938**, *2*, 285-289.
135. Shinozaki-Mendes, R.A.; Lessa, R. Ontogenetic trajectories in *Callinectes danae* (Crustacea: Brachyura): sex and age polymorphism. *J. Mar. Biol. Assoc. U.K.* **2017**, *99*, 111-118.

136. Lane-Medeiros, L.; Moraes, S.A.S.N.; Alencar, C.E.R.D.; Rocha, M.A.L.; Freire, F.A.M. Body shape variations help to diminish taxonomy uncertainty in juvenile swimming crab *Callinectes* Stimpson, 1860. *Zoologischer Anzeiger* **2021**, *295*, 89-98.
137. Ogburn, M.B.; Stuck, K.C.; Heard, R.W.; Wang, S.Y.; Forward Jr, R.B. Seasonal variability in morphology of blue crab, *Callinectes sapidus*, megalopae and early juvenile stage crabs, and distinguishing characteristics among co-occurring Portunidae. *J. Crust. Biol.* **2011**, *31*, 106-113.
138. Silva, I.C.; Alves, M.J.; Paula, J.; Hawkins, S.J. Population differentiation of the shore crab *Carcinus maenas* (Brachyura: Portunidae) on the southwest English coast based on genetic and morphometric analyses. *Sci. Mar.* **2010**, *74*, 435-444.
139. García-Berthou, E.; Carmona-Catot, G.; Merciai, R.; Ogle, D.H. A technical note on seasonal growth models. *Rev. Fish Biol. Fish.* **2012**, *22*, 635-640.
140. Ju, S.J.; Secor, D.H.; Harvey, H.R. Growth rate variability and lipofuscin accumulation rates in the blue crab *Callinectes sapidus*. *Mar. Ecol. Prog. Ser.* **2001**, *224*, 197-205.
141. Türeli, C.; Yesilyurt, I. Growth of blue crab, *Callinectes sapidus*, in the Yumurtalik Cove, Turkey: a molt process approach. *Open Life Sciences* **2014**, *9*, 49-57.
142. Mehanna, S.F.; Desouky, M.G.; Farouk, A.E. Population dynamics and fisheries characteristics of the blue crab *Callinectes sapidus* (Rathbun, 1896) as an invasive species in Bardawil Lagoon, Egypt. *Egypt. J. Aquat. Biol. Fish.* **2019**, *23*, 599-611.
143. Pauly, D. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES J. Mar. Sci.* **1980**, *39*, 175-192.
144. Fogarty, M.J.; Lipcius, R.N. Population Dynamics and Fisheries. In *The Blue Crab: Callinectes sapidus*, Kennedy, V.S., Cronin, L.E., Eds.; Maryland Sea Grant College: College Park, Maryland, 2007; pp. 711-755.
145. Kahn, D.M.; Helser, T.E. Abundance, dynamics and mortality rates of the Delaware Bay stock of blue crabs, *Callinectes sapidus*. *J. Shellfish Res.* **2005**, *24*, 269-284.
146. Zohar, Y.; Hines, A.H.; Zmora, O.; Johnson, E.G.; Lipcius, R.N.; Seitz, R.D.; Eggleston, D.B.; Place, A.R.; Schott, E.J.; Stubblefield, J.D.; et al. The Chesapeake Bay blue crab (*Callinectes sapidus*): a multidisciplinary approach to responsible stock replenishment. *Rev. Fish. Sci.* **2008**, *16*, 24-34.
147. Bogdanoff, A.K.; Shertzer, K.W.; Layman, C.A.; Chapman, J.K.; Fruitema, M.L.; Solomon, J.; Sabattis, J.; Green, S.; Morris, J.A. Optimum lionfish yield: a non-traditional management concept for invasive lionfish (*Pterois* spp.) fisheries. *Biol. Invasions* **2021**, *23*, 795-810.
148. Mancinelli, G.; Dailianis, T.; Dounas, C.; Kasapidis, P.; Koulouri, P.; Skouradakis, G.; Bardelli, R.; Di Muri, C.; Guerra, M.T.; Vizzini, S.J.S. Isotopic niche and trophic position of the invasive portunid *Portunus segnis* Forskål (1775) in Elounda Bay (Crete Island, Eastern Mediterranean). *Sustainability* **2022**, *14*, 15202.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.